

DATTORRO

CONVEX
OPTIMIZATION
&
EUCLIDEAN
DISTANCE
GEOMETRY

Μεβ00

**Convex Optimization
&
Euclidean Distance Geometry**

Jon Dattorro

Μεβoo Publishing

Meboo Publishing USA
PO Box 12
Palo Alto, California 94302

Jon Dattorro, *Convex Optimization & Euclidean Distance Geometry*,
Meboo, 2005.


ISBN 0976401304 (English) ISBN 9780615193687 (International II)

This is Version 2009.04.11: available in print, as conceived, in color.

cybersearch:

- I. convex optimization
- II. semidefinite program
- III. rank constraint
- IV. convex geometry
- V. distance matrix
- VI. convex cones

PROGRAMS BY MATLAB



typesetting by *WinEdt*

with donations from **SIAM** and **AMS**.

This searchable electronic color pdfBook is click-navigable within the text by page, section, subsection, chapter, theorem, example, definition, cross reference, citation, equation, figure, table, and hyperlink. A pdfBook has no electronic copy protection and can be read and printed by most computers. The publisher hereby grants the right to reproduce this work in any format but limited to personal use.

© 2001-2009 Meboo
All rights reserved.

for *Jennie Columba*



◊ *Antonio*



& *Sze Wan*

$$\text{EDM} = \mathcal{S}_h \cap (\mathcal{S}_c^\perp - \mathcal{S}_+)$$

Prelude

The constant demands of my department and university and the ever increasing work needed to obtain funding have stolen much of my precious thinking time, and I sometimes yearn for the halcyon days of Bell Labs.

—Steven Chu, Nobel laureate [72]

Convex Analysis is the calculus of inequalities while Convex Optimization is its application. Analysis is inherently the domain of a mathematician while Optimization belongs to the engineer. Practitioners in the art of Convex Optimization engage themselves with discovery of which hard problems, perhaps previously believed nonconvex, can be transformed into convex equivalents; because once convex form of a problem is found, then a globally optimal solution is close at hand – the hard work is finished: Finding convex expression of a problem is itself, in a very real sense, the solution.

Yes there is a great race under way to determine which important problems can be posed in a convex setting. Yet, that skill acquired by understanding the geometry and application of Convex Optimization will remain more an art for some time to come; the reason being, there is generally no unique transformation of a given problem to its convex equivalent. This means, two researchers pondering the same problem are likely to formulate the convex equivalent differently; hence, one solution is likely different from the other for the same problem. Any presumption of only one right or correct solution becomes nebulous. Study of equivalence, sameness, and uniqueness therefore pervade study of Optimization.

Tremendous benefit accrues when an optimization problem can be transformed to its convex equivalent, primarily because any locally optimal solution is then guaranteed globally optimal. Solving a nonlinear system, for example, by instead solving an equivalent convex optimization problem is therefore highly preferable.^{0.1} Yet it can be difficult for the engineer to apply theory without an understanding of Analysis.

These pages comprise my journal over an eight year period bridging gaps between engineer and mathematician; they constitute a translation, unification, and cohering of about three hundred papers, books, and reports from several different fields of mathematics and engineering. Beacons of historical accomplishment are cited throughout. Much of what is written here will not be found elsewhere. Care to detail, clarity, accuracy, consistency, and typography accompanies removal of ambiguity and verbosity out of respect for the reader. Consequently there is much cross-referencing and background material provided in the text, footnotes, and appendices so as to be self-contained and to provide understanding of fundamental concepts.

– Jon Dattorro
Stanford, California
2009

^{0.1}That is what motivates a convex optimization known as *geometric programming* [52] [70] which has driven great advances in the electronic circuit design industry. [31, §4.7] [219] [342] [345] [85] [159] [168] [169] [170] [171] [172] [234] [235] [273]

Convex Optimization & Euclidean Distance Geometry

1	Overview	19
2	Convex geometry	33
2.1	Convex set	34
2.2	Vectorized-matrix inner product	46
2.3	Hulls	56
2.4	Halfspace, Hyperplane	68
2.5	Subspace representations	80
2.6	Extreme, Exposed	85
2.7	Cones	90
2.8	Cone boundary	99
2.9	Positive semidefinite (PSD) cone	106
2.10	Conic independence (c.i.)	129
2.11	When extreme means exposed	135
2.12	Convex polyhedra	135
2.13	Dual cone & generalized inequality	144
3	Geometry of convex functions	201
3.1	Convex function	202
3.2	Matrix-valued convex function	239
3.3	Quasiconvex	245
3.4	Salient properties	247

4	Semidefinite programming	249
4.1	Conic problem	250
4.2	Framework	258
4.3	Rank reduction	272
4.4	Rank-constrained semidefinite program	281
4.5	Constraining cardinality	298
4.6	Cardinality and rank constraint examples	312
4.7	Convex Iteration rank-1	345
5	Euclidean Distance Matrix	349
5.1	EDM	350
5.2	First metric properties	351
5.3	\exists fifth Euclidean metric property	351
5.4	EDM definition	356
5.5	Invariance	386
5.6	Injectivity of \mathbf{D} & unique reconstruction	391
5.7	Embedding in affine hull	397
5.8	Euclidean metric <i>versus</i> matrix criteria	402
5.9	Bridge: Convex polyhedra to EDMs	410
5.10	EDM-entry composition	417
5.11	EDM indefiniteness	420
5.12	List reconstruction	428
5.13	Reconstruction examples	432
5.14	Fifth property of Euclidean metric	439
6	Cone of distance matrices	449
6.1	Defining EDM cone	451
6.2	Polyhedral bounds	453
6.3	$\sqrt{\text{EDM}}$ cone is not convex	455
6.4	A geometry of completion	456
6.5	EDM definition in $\mathbf{11}^T$	462
6.6	Correspondence to PSD cone \mathbb{S}_+^{N-1}	470
6.7	Vectorization & projection interpretation	476
6.8	Dual EDM cone	481
6.9	Theorem of the alternative	495
6.10	Postscript	497

7 Proximity problems	499
7.1 First prevalent problem:	507
7.2 Second prevalent problem:	518
7.3 Third prevalent problem:	529
7.4 Conclusion	539
A Linear algebra	541
A.1 Main-diagonal δ operator, λ , trace, vec	541
A.2 Semidefiniteness: domain of test	546
A.3 Proper statements	549
A.4 Schur complement	561
A.5 Eigen decomposition	566
A.6 Singular value decomposition, SVD	569
A.7 Zeros	574
B Simple matrices	581
B.1 Rank-one matrix (dyad)	582
B.2 Doublet	587
B.3 Elementary matrix	588
B.4 Auxiliary V -matrices	590
B.5 Orthogonal matrix	595
C Some analytical optimal results	599
C.1 Properties of infima	599
C.2 Trace, singular and eigen values	600
C.3 Orthogonal Procrustes problem	606
C.4 Two-sided orthogonal Procrustes	608
D Matrix calculus	613
D.1 Directional derivative, Taylor series	613
D.2 Tables of gradients and derivatives	634
E Projection	643
E.1 Idempotent matrices	647
E.2 $I - P$, Projection on algebraic complement	652
E.3 Symmetric idempotent matrices	653
E.4 Algebra of projection on affine subsets	658
E.5 Projection examples	659

12 CONVEX OPTIMIZATION & EUCLIDEAN DISTANCE GEOMETRY

E.6	Vectorization interpretation,	665
E.7	Projection on matrix subspaces	673
E.8	Range/Rowspace interpretation	677
E.9	Projection on convex set	677
E.10	Alternating projection	691
F	Notation and a few definitions	711
	Bibliography	727
	Index	755

List of Figures

1	Overview	19
1	<i>Orion nebula</i>	20
2	Application of trilateration is localization	21
3	Molecular conformation	22
4	Face recognition	23
5	<i>Swiss roll</i>	24
6	USA map reconstruction	26
7	Honeycomb	27
8	Robotic vehicles	28
9	Reconstruction of David	32
10	David by distance geometry	32
2	Convex geometry	33
11	Slab	36
12	Open, closed, convex sets	38
13	Intersection of line with boundary	39
14	Tangentials	42
15	Inverse image	45
16	Linear injective mapping of Euclidean body	49
17	Linear noninjective mapping of Euclidean body	51
18	Convex hull of a random list of points	57
19	Hulls	59
20	Two Fantopes	62
21	Convex hull of rank-1 matrices	64
22	A simplicial cone	67
23	Hyperplane illustrated $\partial\mathcal{H}$ is a partially bounding line	69

24	Hyperplanes in \mathbb{R}^2	71
25	Affine independence	74
26	$\{z \in \mathcal{C} \mid a^T z = \kappa_i\}$	76
27	Hyperplane supporting closed set	77
28	Minimizing hyperplane over affine set in nonnegative orthant	84
29	Closed convex set illustrating exposed and extreme points	87
30	Two-dimensional nonconvex cone	91
31	Nonconvex cone made from lines	91
32	Nonconvex cone is convex cone boundary	92
33	Cone exterior is convex cone	92
34	Truncated nonconvex cone \mathcal{X}	93
35	Not a cone	94
36	Minimum element. Minimal element.	97
37	\mathcal{K} is a pointed polyhedral cone having empty interior	100
38	Exposed and extreme directions	105
39	Positive semidefinite cone	108
40	Convex Schur-form set	110
41	Projection of the PSD cone	113
42	Circular cone showing axis of revolution	118
43	Circular section	120
44	Polyhedral inscription	121
45	Conically (in)dependent vectors	130
46	Pointed six-faceted polyhedral cone and its dual	131
47	Minimal set of generators for halfspace about origin	134
48	Venn diagram for cones and polyhedra	137
49	Unit simplex	140
50	Two views of a simplicial cone and its dual	142
51	Two equivalent constructions of dual cone	146
52	Dual polyhedral cone construction by right angle	147
53	\mathcal{K} is a halfspace about the origin	148
54	Iconic primal and dual objective functions.	149
55	Orthogonal cones	153
56	<i>Blades</i> \mathcal{K} and \mathcal{K}^*	155
57	Shrouded polyhedral cone	167
58	Simplicial cone \mathcal{K} in \mathbb{R}^2 and its dual	172
59	Monotone nonnegative cone $\mathcal{K}_{\mathcal{M}+}$ and its dual	182
60	Monotone cone $\mathcal{K}_{\mathcal{M}}$ and its dual	184
61	Two views of monotone cone $\mathcal{K}_{\mathcal{M}}$ and its dual	185

62	First-order optimality condition	189
3	Geometry of convex functions	201
63	Convex functions having unique minimizer	203
64	1-norm ball \mathcal{B}_1 from compressed sensing/compressive sampling	207
65	Cardinality minimization, signed <i>versus</i> unsigned variable . .	209
66	Affine function	216
67	Supremum of affine functions	218
68	Epigraph	219
69	Gradient in \mathbb{R}^2 evaluated on grid	227
70	Quadratic function convexity in terms of its gradient	234
71	Contour plot of real convex function at selected levels	236
72	Iconic quasiconvex function	245
4	Semidefinite programming	249
73	Visualizing positive semidefinite cone in high dimension	254
74	2-lattice of sensors and anchors for localization example	287
75	3-lattice	288
76	4-lattice	289
77	5-lattice	290
78	Uncertainty ellipsoids orientation and eccentricity	291
79	2-lattice localization solution	296
80	3-lattice localization solution	296
81	4-lattice localization solution	297
82	5-lattice localization solution	297
83	Measurement lower-bound for cardinality problem	301
84	Signal dropout	305
85	Signal dropout reconstruction	306
86	Simplex with intersecting line problem in compressed sensing .	308
87	Permutation matrix column-norm and column-sum constraint	316
88	MAX CUT problem	323
89	Shepp-Logan phantom	329
90	MRI radial sampling pattern in Fourier domain	334
91	Aliased phantom	336
92	Neighboring-pixel candidates for estimating gradient	338
93	MIT logo	341
94	One-pixel camera	342

5	Euclidean Distance Matrix	349
95	Convex hull of three points	350
96	Complete <i>dimensionless EDM graph</i>	353
97	Fifth Euclidean metric property	355
98	Arbitrary hexagon in \mathbb{R}^3	364
99	Kissing number	366
100	<i>Trilateration</i>	370
101	This EDM graph provides unique isometric reconstruction	374
102	Two sensors \bullet and three anchors \circ	374
103	Two discrete linear trajectories of sensors	375
104	Coverage in cellular telephone network	378
105	Contours of equal signal power	378
106	Depiction of molecular conformation	380
107	Square diamond	389
108	Orthogonal complements in \mathbb{S}^N abstractly oriented	392
109	Elliptope \mathcal{E}^3	411
110	Elliptope \mathcal{E}^2 interior to \mathbb{S}_+^2	412
111	Smallest eigenvalue of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$	418
112	Some entrywise EDM compositions	418
113	Map of United States of America	431
114	Largest ten eigenvalues of $-V_{\mathcal{N}}^T O V_{\mathcal{N}}$	434
115	<i>Relative-angle inequality tetrahedron</i>	441
116	Nonsimplicial pyramid in \mathbb{R}^3	445
6	Cone of distance matrices	449
117	Relative boundary of cone of Euclidean distance matrices	452
118	Intersection of EDM cone with hyperplane	454
119	Neighborhood graph	457
120	<i>Trefoil knot</i> untied	459
121	<i>Trefoil ribbon</i>	461
122	Example of $V_{\mathcal{X}}$ selection to make an EDM	464
123	Vector $V_{\mathcal{X}}$ spirals	467
124	Three views of translated negated elliptope	474
125	Halfline \mathcal{T} on PSD cone boundary	478
126	Vectorization and projection interpretation example	479
127	Orthogonal complement of geometric center subspace	484
128	EDM cone construction by flipping PSD cone	485

<i>LIST OF FIGURES</i>	17
129 Decomposing a member of polar EDM cone	490
130 Ordinary dual EDM cone projected on \mathbb{S}_h^3	496
7 Proximity problems	499
131 Pseudo-Venn diagram	502
132 Elbow placed in path of projection	503
133 Convex envelope	523
A Linear algebra	541
134 Geometrical interpretation of full SVD	572
B Simple matrices	581
135 Four fundamental subspaces of any dyad	583
136 Four fundamental subspaces of a doublet	587
137 Four fundamental subspaces of elementary matrix	588
138 Gimbal	597
D Matrix calculus	613
139 Convex quadratic bowl in $\mathbb{R}^2 \times \mathbb{R}$	624
E Projection	643
140 Action of pseudoinverse	644
141 Nonorthogonal projection of $x \in \mathbb{R}^3$ on $\mathcal{R}(U) = \mathbb{R}^2$	650
142 Biorthogonal expansion of point $x \in \text{aff } \mathcal{K}$	661
143 Dual interpretation of projection on convex set	680
144 Projection product on convex set in subspace	689
145 von Neumann-style projection of point b	692
146 Alternating projection on two halfspaces	693
147 Distance, optimization, feasibility	695
148 Alternating projection on nonnegative orthant and hyperplane	698
149 Geometric convergence of iterates in norm	698
150 Distance between PSD cone and iterate in \mathcal{A}	703
151 Dykstra's alternating projection algorithm	704
152 Polyhedral normal cones	706
153 Normal cone to elliptope	707

List of Tables

2 Convex geometry	
Table 2.9.2.3.1 , rank <i>versus</i> dimension of \mathbb{S}_+^3 faces	115
Table 2.10.0.0.1 , maximum number of c.i. directions	130
Cone Table 1	179
Cone Table S	180
Cone Table A	181
Cone Table 1*	186
4 Semidefinite programming	
Faces of \mathbb{S}_+^3 corresponding to faces of \mathcal{S}_+^3	255
5 Euclidean Distance Matrix	
<i>Précis</i> 5.7.2 : affine dimension in terms of rank	400
B Simple matrices	
Auxiliary V -matrix Table B.4.4	594
D Matrix calculus	
Table D.2.1 , algebraic gradients and derivatives	635
Table D.2.2 , trace Kronecker gradients	636
Table D.2.3 , trace gradients and derivatives	637
Table D.2.4 , logarithmic determinant gradients, derivatives	639
Table D.2.5 , determinant gradients and derivatives	640
Table D.2.6 , logarithmic derivatives	640
Table D.2.7 , exponential gradients and derivatives	641

Chapter 1

Overview

Convex Optimization Euclidean Distance Geometry

People are so afraid of convex analysis.

–[Claude Lemaréchal](#), 2003

In layman’s terms, the mathematical science of Optimization is the study of how to make a good choice when confronted with conflicting requirements. The qualifier *convex* means: when an optimal solution is found, then it is guaranteed to be a best solution; there is no better choice.

Any convex optimization problem has geometric interpretation. If a given optimization problem can be transformed to a convex equivalent, then this interpretive benefit is acquired. That is a powerful attraction: the ability to visualize geometry of an optimization problem. Conversely, recent advances in geometry and in graph theory hold convex optimization within their proofs’ core. [\[353\]](#) [\[283\]](#)

This book is about convex optimization, convex geometry (with particular attention to distance geometry), and nonconvex, combinatorial, and geometrical problems that can be relaxed or transformed into convex problems. A virtual flood of new applications follows by epiphany that many problems, presumed nonconvex, can be so transformed. [\[9\]](#) [\[10\]](#) [\[51\]](#) [\[81\]](#) [\[129\]](#) [\[131\]](#) [\[244\]](#) [\[263\]](#) [\[270\]](#) [\[321\]](#) [\[322\]](#) [\[350\]](#) [\[353\]](#) [\[31\]](#), §4.3, p.316-322



Figure 1: *Orion nebula*. (astrophotography by Massimo Robberto)

Euclidean distance geometry is, fundamentally, a determination of point conformation (configuration, relative position or location) by inference from interpoint distance information. By *inference* we mean: *e.g.*, given only distance information, determine whether there corresponds a *realizable* conformation of points; a list of points in some dimension that attains the given interpoint distances. Each point may represent simply location or, abstractly, any entity expressible as a vector in finite-dimensional Euclidean space; *e.g.*, distance geometry of music [91].

It is a common misconception to presume that some desired point conformation cannot be recovered in the absence of complete interpoint distance information. We might, for example, want to realize a constellation given only interstellar distance (or, equivalently, distance from Earth and relative angular measurement; the Earth as vertex to two stars). At first it may seem $O(N^2)$ data is required, yet there are many circumstances where this can be reduced to $O(N)$.

If we agree that a set of points can have a shape (three points can form a triangle and its interior, for example, four points a tetrahedron), then




Figure 2: Application of trilateration (§5.4.2.2.4) is localization (determining position) of a radio signal source in 2 dimensions; more commonly known by radio engineers as the process “triangulation”. In this scenario, anchors $\check{x}_2, \check{x}_3, \check{x}_4$ are illustrated as fixed antennae. [185] The radio signal source (a sensor $\bullet x_1$) anywhere in affine hull of three antenna bases can be uniquely localized by measuring distance to each (dashed white arrowed line segments). Ambiguity of lone distance measurement to sensor is represented by circle about each antenna. So & Ye proved trilateration is expressible as a semidefinite program; hence, a convex optimization problem. [284]

we can ascribe *shape* of a set of points to their convex hull. It should be apparent: from distance, these shapes can be determined only to within a *rigid transformation* (rotation, reflection, translation).

Absolute position information is generally lost, given only distance information, but we can determine the smallest possible dimension in which an unknown list of points can exist; that attribute is their *affine dimension* (a triangle in any ambient space has affine dimension 2, for example). In circumstances where stationary reference points are also provided, it becomes possible to determine absolute position or location; *e.g.*, Figure 2.




Figure 3: [167] [99] Distance data collected via nuclear magnetic resonance (NMR) helped render this 3-dimensional depiction of a [protein molecule](#). *At the beginning of the 1980s, Kurt Wüthrich [Nobel laureate] developed an idea about how NMR could be extended to cover biological molecules such as proteins. He invented a systematic method of pairing each NMR signal with the right hydrogen nucleus (proton) in the macromolecule. The method is called sequential assignment and is today a cornerstone of all NMR structural investigations. He also showed how it was subsequently possible to determine pairwise distances between a large number of hydrogen nuclei and use this information with a mathematical method based on distance-geometry to calculate a three-dimensional structure for the molecule.* –[247] [337] [162]

Geometric problems involving distance between points can sometimes be reduced to convex optimization problems. Mathematics of this combined study of geometry and optimization is rich and deep. Its application has already proven invaluable discerning organic [molecular conformation](#) by measuring interatomic distance along covalent bonds; *e.g.*, Figure 3. [77] [311] [122] [41] Many disciplines have already benefitted and simplified consequent to this theory; *e.g.*, distance-based [pattern recognition](#) (Figure 4), [localization](#) in wireless sensor networks by measurement of intersensor distance along channels of communication, [42, §5] [348] [40] [wireless location](#) of a radio-signal source such as a cell phone by multiple measurements of signal strength, the [global positioning system](#) (GPS), and [multidimensional scaling](#) which is a numerical representation of qualitative data by finding a low-dimensional scale.




Figure 4: This coarsely discretized triangulated algorithmically flattened human face (made by Kimmel & the Bronsteins [200]) represents a stage in machine recognition of human identity; called *face recognition*. Distance geometry is applied to determine discriminating-features.

Euclidean distance geometry together with convex optimization have also found application in *artificial intelligence*:

- to *machine learning* by discerning naturally occurring manifolds in:
 - Euclidean bodies (Figure 5, §6.4.0.0.1),
 - Fourier spectra of kindred utterances [187],
 - and image sequences [332],
- to *robotics*; *e.g.*, automatic control of vehicles maneuvering in formation. (Figure 8)

by chapter

We study the pervasive convex Euclidean bodies and their various representations. In particular, we make convex polyhedra, cones, and dual cones more visceral through illustration in **chapter 2, Convex geometry**, and we study the geometric relation of polyhedral cones to nonorthogonal




Figure 5: *Swiss roll* from Weinberger & Saul [332]. The problem of manifold learning, illustrated for $N = 800$ data points sampled from a “Swiss roll” ①. A discretized manifold is revealed by connecting each data point and its $k = 6$ nearest neighbors ②. An unsupervised learning algorithm unfolds the Swiss roll while preserving the local geometry of nearby data points ③. Finally, the data points are projected onto the two dimensional subspace that maximizes their variance, yielding a faithful embedding of the original manifold ④.

bases (biorthogonal expansion). We explain conversion between halfspace- and vertex-description of a convex cone, we motivate the dual cone and provide formulae for finding it, and we show how first-order optimality conditions or alternative systems of linear inequality or *linear matrix inequality* [51] can be explained by *generalized inequalities* with respect to convex cones and their duals. The conic analogue to linear independence, called *conic independence*, is introduced as a new tool in the study of cones; the logical next step in the progression: linear, affine, conic.

Any convex optimization problem can be visualized geometrically. Desire to visualize in high dimension is deeply embedded in the [mathematical psyche](#). Chapter 2 provides tools to make visualization easier, and we teach how to visualize in high dimension. The concepts of face, extreme point, and extreme direction of a convex Euclidean body are explained here; crucial to understanding convex optimization. The convex cone of positive semidefinite matrices, in particular, is studied in depth:

- We interpret, for example, inverse image of the positive semidefinite cone under affine transformation. (Example [2.9.1.0.2](#))
- Subsets of the positive semidefinite cone, discriminated by rank exceeding some lower bound, are convex. In other words, high-rank subsets of the positive semidefinite cone boundary united with its interior are convex. (Theorem [2.9.2.6.3](#)) There is a closed form for projection on those convex subsets.
- The positive semidefinite cone is a circular cone in low dimension, while *Geršgorin discs* specify inscription of a polyhedral cone into that positive semidefinite cone. (Figure [44](#))

Chapter 3, Geometry of convex functions, observes analogies between convex sets and functions: We explain, for example, how the real affine function relates to convex functions as the hyperplane relates to convex sets. Partly a cookbook for the most useful of convex functions and optimization problems, methods are drawn from matrix calculus for determining convexity and discerning geometry.

Semidefinite programming is reviewed in **chapter 4** with particular attention to optimality conditions for prototypical primal and dual programs, their interplay, and the *perturbation method* of rank reduction of optimal solutions (extant but not well known). Positive definite Farkas' lemma




Figure 6: About five thousand points along the borders constituting the United States were used to create an exhaustive matrix of interpoint distance for each and every pair of points in the ordered set (a list); called a *Euclidean distance matrix*. From that noiseless distance information, it is easy to reconstruct the map exactly via the Schoenberg criterion (827). (§5.13.1.0.1, confer Figure 113) Map reconstruction is exact (to within a rigid transformation) given any number of interpoint distances; the greater the number of distances, the greater the detail (just as it is for all conventional map preparation).

is derived, and we also show how to determine if a feasible set belongs exclusively to a positive semidefinite cone boundary. A three-dimensional polyhedral analogue for the positive semidefinite cone of 3×3 symmetric matrices is introduced. This analogue is a new tool for visualizing coexistence of low- and high-rank optimal solutions in 6 dimensions. We find a minimum-cardinality Boolean solution to an instance of $Ax = b$:

$$\begin{aligned}
 & \underset{x}{\text{minimize}} && \|x\|_0 \\
 & \text{subject to} && Ax = b \\
 & && x_i \in \{0, 1\}, \quad i = 1 \dots n
 \end{aligned} \tag{622}$$

The *sensor-network localization* problem is solved in any dimension in this chapter. We introduce the method of *convex iteration* for constraining rank in the form $\text{rank } G \leq \rho$ and cardinality in the form $\text{card } x \leq k$.



Figure 7: These bees construct a honeycomb by solving a convex optimization problem. (§5.4.2.2.3) The most dense packing of identical spheres about a central sphere in two dimensions is 6. Packed sphere centers describe a regular lattice.

The EDM is studied in **chapter 5, Euclidean Distance Matrix**, its properties and relationship to both positive semidefinite and Gram matrices. We relate the EDM to the four classical properties of the Euclidean metric; thereby, observing existence of an infinity of properties of the Euclidean metric beyond triangle inequality. We proceed by deriving the fifth Euclidean metric property and then explain why furthering this endeavor is inefficient because the ensuing criteria (while describing polyhedra in angle or area, volume, content, and so on *ad infinitum*) grow linearly in complexity and number with problem size.

Reconstruction methods are explained and applied to a map of the United States; *e.g.*, Figure 6. We also generate a distorted but recognizable isotonic map using only comparative distance information (only ordinal distance data). We demonstrate an elegant method for including dihedral (or *torsion*) angle constraints into a molecular conformation problem. More geometrical problems solvable via EDMs are presented with the best methods for posing them, EDM problems are posed as convex optimizations, and we show how to recover exact relative position given incomplete noiseless interpoint distance information.

The set of all Euclidean distance matrices forms a pointed closed convex cone called the *EDM cone*, \mathbb{EDM}^N . We offer a new proof of Schoenberg’s seminal characterization of EDMs:

$$D \in \mathbb{EDM}^N \Leftrightarrow \begin{cases} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \\ D \in \mathbb{S}_h^N \end{cases} \quad (827)$$

Our proof relies on fundamental geometry; assuming, any EDM must correspond to a list of points contained in some polyhedron (possibly at



Figure 8: Robotic vehicles in concert can move larger objects, guard a perimeter, perform surveillance, find land mines, or localize a plume of gas, liquid, or radio waves. [121]

its vertices) and *vice versa*. It is known, but not obvious, this *Schoenberg criterion* implies nonnegativity of the EDM entries; proved herein.

We characterize the eigenvalue spectrum of an EDM, and then devise a polyhedral spectral cone for determining membership of a given matrix (in Cayley-Menger form) to the convex cone of Euclidean distance matrices; *id est*, a matrix is an EDM if and only if its nonincreasingly ordered vector of eigenvalues belongs to a polyhedral spectral cone for \mathbb{EDM}^N ;

$$D \in \mathbb{EDM}^N \Leftrightarrow \begin{cases} \lambda\left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix}\right) \in \begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \\ D \in \mathbb{S}_h^N \end{cases} \quad (1040)$$

We will see: spectral cones are not unique.

In **chapter 6, Cone of distance matrices**, we explain the geometric relationship between the cone of Euclidean distance matrices, two positive semidefinite cones, and the ellipsope. We illustrate geometric requirements, in particular, for projection of a given matrix on a positive semidefinite cone

that establish its membership to the EDM cone. The faces of the EDM cone are described, but still open is the question whether all its faces are exposed as they are for the positive semidefinite cone. The Schoenberg criterion (827), relating the EDM cone and a positive semidefinite cone, is revealed to be a discretized membership relation (dual generalized inequalities, a new Farkas'-like lemma) between the EDM cone and its ordinary dual, \mathbb{EDM}^{N*} . A matrix criterion for membership to the dual EDM cone is derived that is simpler than the Schoenberg criterion:

$$D^* \in \mathbb{EDM}^{N*} \Leftrightarrow \delta(D^* \mathbf{1}) - D^* \succeq 0 \quad (1189)$$

We derive a concise equality of the EDM cone to two subspaces and a positive semidefinite cone;

$$\mathbb{EDM}^N = \mathbb{S}_h^N \cap (\mathbb{S}_c^{N\perp} - \mathbb{S}_+^N) \quad (1183)$$

In **chapter 7, Proximity problems**, we explore methods of solution to a few fundamental and prevalent Euclidean distance matrix proximity problems; the problem of finding that distance matrix closest, in some sense, to a given matrix H :

$$\begin{array}{ll} \underset{D}{\text{minimize}} & \|-V(D - H)V\|_{\mathbb{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & D \in \mathbb{EDM}^N \end{array} \quad \begin{array}{ll} \underset{\sqrt[D]{D}}{\text{minimize}} & \|\sqrt[D]{D} - H\|_{\mathbb{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & \sqrt[D]{D} \in \sqrt{\mathbb{EDM}^N} \end{array} \quad (1225)$$

$$\begin{array}{ll} \underset{D}{\text{minimize}} & \|D - H\|_{\mathbb{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & D \in \mathbb{EDM}^N \end{array} \quad \begin{array}{ll} \underset{\sqrt[D]{D}}{\text{minimize}} & \|-V(\sqrt[D]{D} - H)V\|_{\mathbb{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & \sqrt[D]{D} \in \sqrt{\mathbb{EDM}^N} \end{array}$$

We apply the new *convex iteration* method for constraining rank. Known heuristics for rank minimization are also explained. We offer a new geometrical proof, of a famous discovery by Eckart & Young in 1936 [112], with particular regard to Euclidean projection of a point on that generally nonconvex subset of the positive semidefinite cone boundary comprising all positive semidefinite matrices having rank not exceeding a prescribed bound ρ . We explain how this problem is transformed to a convex optimization for any rank ρ .

novelty - Convex Optimization & Euclidean Distance Geometry

- p.129 *Conic independence* is introduced as a natural extension to linear and affine independence; a tool in convex analysis for manipulation of cones.
- p.158 Arcane theorems of alternative *generalized inequality* are simply derived from cone *membership relations*; generalizations of *Farkas' lemma* translated to the geometry of convex cones.
- p.164 Smallest face of a proper cone containing \mathcal{C} .
- p.196 Unique *conic coordinates* in a proper conic system.
- p.254 We present an arguably good 3-dimensional polyhedral analogue, to the isomorphically 6-dimensional positive semidefinite cone, as an aid to understand semidefinite programming.
- p.281, p.298 We show how to constrain rank in the form $\text{rank } G \leq \rho$ and cardinality in the form $\text{card } x \leq k$ by *convex iteration*. We show how to transform a rank constrained problem to a rank-1 problem.
- p.329 Lowest image-gradient cardinality for Shepp-Logan phantom in MRI.
- p.365, p.369 *Sphere packing kissing-number* (solved in \mathbb{R}^3 by [Isaac Newton](#)) and *trilateration (localization)* are convex optimization problems.
- p.380 We show how to elegantly include *torsion* or *dihedral* angle constraints into the *molecular conformation problem*.
- p.413 Geometrical proof: *Schoenberg criterion* for Euclidean distance matrix.
- p.431 We experimentally demonstrate a conjecture of Borg & Groenen by reconstructing a map of the USA using only ordinal (comparative) distance information.
- p.6, p.449 There is an equality, relating the convex cone of Euclidean distance matrices to the positive semidefinite cone, apparently overlooked in the literature; an equality between two large convex Euclidean bodies.
- p.493 The *Schoenberg criterion* for a Euclidean distance matrix is revealed to be a discretized *membership relation* (or *dual generalized inequalities*) between the EDM cone and its dual.

appendices

Provided so as to be more self-contained:

- linear algebra (**appendix A** is primarily concerned with proper statements of semidefiniteness for square matrices),
- simple matrices (dyad, doublet, elementary, Householder, Schoenberg, orthogonal, *etcetera*, in **appendix B**),
- a collection of known analytical solutions to some important optimization problems (**appendix C**),
- matrix calculus remains somewhat unsystematized when compared to ordinary calculus (**appendix D** concerns matrix-valued functions, matrix differentiation and directional derivatives, Taylor series, and tables of first- and second-order gradients and matrix derivatives),
- an elaborate exposition offering insight into orthogonal and nonorthogonal projection on convex sets (the connection between projection and positive semidefiniteness, for example, or between projection and a linear objective function in **appendix E**),
- MATLAB code on *Wikimization* to discriminate EDMs, to determine conic independence, to reduce or constrain rank of an optimal solution to a semidefinite program, compressed sensing (compressive sampling) for digital image and audio signal processing, and two distinct methods of reconstructing a map of the United States: one given only distance information, the other given only relative distance data (greater or less than).



Figure 9: Three-dimensional reconstruction of David from distance data.




Figure 10: *Digital Michelangelo Project*, Stanford University. Measuring distance to David by laser rangefinder. Spatial resolution is 0.29mm.

Chapter 2

Convex geometry

Convexity has an immensely rich structure and numerous applications. On the other hand, almost every “convex” idea can be explained by a two-dimensional picture.

–Alexander Barvinok [25, p.vii]

As convex geometry and linear algebra are inextricably bonded by *asymmetry* (linear inequality), we provide much background material on linear algebra (especially in the appendices) although a reader is assumed comfortable with [294] [296] [180] or any other intermediate-level text. The essential references to convex analysis are [177] [271]. The reader is referred to [293] [25] [331] [37] [53] [268] [318] for a comprehensive treatment of convexity. There is relatively less published pertaining to matrix-valued convex functions. [190] [181, §6.6] [260]

2.1 Convex set

A set \mathcal{C} is convex iff for all $Y, Z \in \mathcal{C}$ and $0 \leq \mu \leq 1$

$$\mu Y + (1 - \mu)Z \in \mathcal{C} \quad (1)$$

Under that defining condition on μ , the linear sum in (1) is called a *convex combination* of Y and Z . If Y and Z are points in real finite-dimensional Euclidean vector space \mathbb{R}^n or $\mathbb{R}^{m \times n}$ (matrices), then (1) represents the closed line segment joining them. All line segments are thereby convex sets. Apparent from the definition, a convex set is a connected set. [228, §3.4, §3.5] [37, p.2] A convex set can, but does not necessarily, contain the *origin* $\mathbf{0}$.

An *ellipsoid* centered at $x = a$ (Figure 13, p.39), given matrix $C \in \mathbb{R}^{m \times n}$

$$\{x \in \mathbb{R}^n \mid \|C(x - a)\|^2 = (x - a)^T C^T C (x - a) \leq 1\} \quad (2)$$

is a good icon for a convex set.

2.1.1 subspace

A nonempty subset \mathcal{R} of real Euclidean vector space \mathbb{R}^n is called a *subspace* (§2.5) if every vector^{2.1} of the form $\alpha x + \beta y$, for $\alpha, \beta \in \mathbb{R}$, is in \mathcal{R} whenever vectors x and y are. [220, §2.3] A subspace is a convex set containing the origin, by definition. [271, p.4] Any subspace is therefore open in the sense that it contains no boundary, but closed in the sense [228, §2]

$$\mathcal{R} + \mathcal{R} = \mathcal{R} \quad (3)$$

It is not difficult to show

$$\mathcal{R} = -\mathcal{R} \quad (4)$$

as is true for any subspace \mathcal{R} , because $x \in \mathcal{R} \Leftrightarrow -x \in \mathcal{R}$.

The intersection of an arbitrary collection of subspaces remains a subspace. Any subspace not constituting the entire *ambient vector space* \mathbb{R}^n is a *proper subspace*; e.g.,^{2.2} any line through the origin in two-dimensional Euclidean space \mathbb{R}^2 . The vector space \mathbb{R}^n is itself a conventional subspace, inclusively, [201, §2.1] although not proper.

^{2.1}A *vector* is assumed, throughout, to be a column vector.

^{2.2}We substitute abbreviation *e.g.* in place of the Latin *exempli gratia*.

2.1.2 linear independence

Arbitrary given vectors in Euclidean space $\{\Gamma_i \in \mathbb{R}^n, i=1 \dots N\}$ are *linearly independent* (l.i.) if and only if, for all $\zeta \in \mathbb{R}^N$

$$\Gamma_1 \zeta_1 + \dots + \Gamma_{N-1} \zeta_{N-1} + \Gamma_N \zeta_N = \mathbf{0} \quad (5)$$

has only the *trivial solution* $\zeta = \mathbf{0}$; in other words, iff no vector from the given set can be expressed as a linear combination of those remaining.

2.1.2.1 preservation

(confer §2.4.2.4, §2.10.1) Linear transformation preserves linear dependence. [201, p.86] Conversely, linear independence can be preserved under linear transformation. Given $Y = [y_1 \ y_2 \ \dots \ y_N] \in \mathbb{R}^{N \times N}$, consider the mapping

$$T(\Gamma) : \mathbb{R}^{n \times N} \rightarrow \mathbb{R}^{n \times N} \triangleq \Gamma Y \quad (6)$$

whose domain is the set of all matrices $\Gamma \in \mathbb{R}^{n \times N}$ holding a linearly independent set columnar. Linear independence of $\{\Gamma y_i \in \mathbb{R}^n, i=1 \dots N\}$ demands, by definition, there exist no nontrivial solution $\zeta \in \mathbb{R}^N$ to

$$\Gamma y_1 \zeta_1 + \dots + \Gamma y_{N-1} \zeta_{N-1} + \Gamma y_N \zeta_N = \mathbf{0} \quad (7)$$

By factoring out Γ , we see that triviality is ensured by linear independence of $\{y_i \in \mathbb{R}^N\}$.

2.1.3 Orthant:

name given to a closed convex set that is the higher-dimensional generalization of *quadrant* from the classical Cartesian partition of \mathbb{R}^2 . The most common is the nonnegative orthant \mathbb{R}_+^n or $\mathbb{R}_+^{n \times n}$ (analogue to quadrant I) to which membership denotes nonnegative vector- or matrix-entries respectively; *e.g.*,

$$\mathbb{R}_+^n \triangleq \{x \in \mathbb{R}^n \mid x_i \geq 0 \ \forall i\} \quad (8)$$

The nonpositive orthant \mathbb{R}_-^n or $\mathbb{R}_-^{n \times n}$ (analogue to quadrant III) denotes negative and 0 entries. Orthant convexity^{2.3} is easily verified by definition (1).

^{2.3}All orthants are self-dual simplicial cones. (§2.13.5.1, §2.12.3.1.1)




Figure 11: A *slab* is a convex Euclidean body infinite in extent but not affine. Illustrated in \mathbb{R}^2 , it may be constructed by intersecting two opposing halfspaces whose bounding hyperplanes are parallel but not coincident. Because number of halfspaces used in its construction is finite, slab is a polyhedron. (Cartesian axes and vector inward-normal to each halfspace-boundary are drawn for reference.)

2.1.4 affine set

A nonempty *affine set* (from the word *affinity*) is any subset of \mathbb{R}^n that is a translation of some subspace. Any affine set is convex and open so contains no boundary: *e.g.*, empty set \emptyset , point, line, plane, *hyperplane* (§2.4.2), subspace, *etcetera*. For some *parallel*^{2.4} subspace \mathcal{M} and any point $x \in \mathcal{A}$

$$\begin{aligned} \mathcal{A} \text{ is affine} &\Leftrightarrow \mathcal{A} = x + \mathcal{M} & (9) \\ &= \{y \mid y - x \in \mathcal{M}\} \end{aligned}$$

The intersection of an arbitrary collection of affine sets remains affine. The *affine hull* of a set $\mathcal{C} \subseteq \mathbb{R}^n$ (§2.3.1) is the smallest affine set containing it.

2.1.5 dimension

Dimension of an arbitrary set \mathcal{S} is the dimension of its affine hull; [331, p.14]

$$\dim \mathcal{S} \triangleq \dim \text{aff } \mathcal{S} = \dim \text{aff}(\mathcal{S} - s), \quad s \in \mathcal{S} \quad (10)$$

the same as dimension of the subspace parallel to that affine set $\text{aff } \mathcal{S}$ when nonempty. Hence dimension (of a set) is synonymous with *affine dimension*. [177, A.2.1]

^{2.4}Two affine sets are said to be *parallel* when one is a translation of the other. [271, p.4]

2.1.6 empty set *versus* empty interior

Emptiness \emptyset of a set is handled differently than *interior* in the classical literature. It is common for a nonempty convex set to have empty interior; *e.g.*, paper in the real world:

- An ordinary flat sheet of paper is a nonempty convex set having empty interior in \mathbb{R}^3 but nonempty interior relative to its affine hull.

2.1.6.1 relative interior

Although it is always possible to pass to a smaller ambient Euclidean space where a nonempty set acquires an interior [25, §II.2.3], we prefer the term *relative* which is the conventional fix to this ambiguous terminology.^{2.5} So we distinguish *interior* from *relative interior* throughout: [293] [331] [318]

- Classical interior $\text{int } \mathcal{C}$ is defined as a union of points: x is an interior point of $\mathcal{C} \subseteq \mathbb{R}^n$ if there exists an open ball of dimension n and nonzero radius centered at x that is contained in \mathcal{C} .
- Relative interior $\text{rel int } \mathcal{C}$ of a convex set $\mathcal{C} \subseteq \mathbb{R}^n$ is interior relative to its affine hull.^{2.6}

Thus defined, it is common (though confusing) for $\text{int } \mathcal{C}$ the interior of \mathcal{C} to be empty while its relative interior is not: this happens whenever dimension of its affine hull is less than dimension of the ambient space ($\dim \text{aff } \mathcal{C} < n$; *e.g.*, were \mathcal{C} paper) or in the exception when \mathcal{C} is a single point; [228, §2.2.1]

$$\text{rel int}\{x\} \stackrel{\Delta}{=} \text{aff}\{x\} = \{x\}, \quad \text{int}\{x\} = \emptyset, \quad x \in \mathbb{R}^n \quad (11)$$

In any case, *closure* of the relative interior of a convex set \mathcal{C} always yields the closure of the set itself;

$$\overline{\text{rel int } \mathcal{C}} = \overline{\mathcal{C}} \quad (12)$$

If \mathcal{C} is convex then $\text{rel int } \mathcal{C}$ and $\overline{\mathcal{C}}$ are convex. [177, p.24] If \mathcal{C} has nonempty interior, then

$$\text{rel int } \mathcal{C} = \text{int } \mathcal{C} \quad (13)$$

^{2.5}Superfluous mingling of terms as in *relatively nonempty set* would be an unfortunate consequence. From the opposite perspective, some authors use the term *full* or *full-dimensional* to describe a set having nonempty interior.

^{2.6}Likewise for *relative boundary* (§2.6.1.3), although *relative closure* is superfluous. [177, §A.2.1]




Figure 12: **(a)** Closed convex set. **(b)** Neither open, closed, or convex. Yet the positive semidefinite cone can remain convex in absence of certain boundary components (§2.9.2.6.3). **(c)** Open convex set.

Given the intersection of any convex set \mathcal{C} with an affine set \mathcal{A}

$$\text{rel int}(\mathcal{C} \cap \mathcal{A}) = \text{rel int}(\mathcal{C}) \cap \mathcal{A} \quad (14)$$

2.1.7 classical boundary

(confer §2.6.1.3) *Boundary* of a set \mathcal{C} is the closure of \mathcal{C} less its interior presumed nonempty; [48, §1.1]

$$\partial\mathcal{C} = \bar{\mathcal{C}} \setminus \text{int } \mathcal{C} \quad (15)$$

which follows from the fact

$$\overline{\text{int } \mathcal{C}} = \bar{\mathcal{C}} \quad (16)$$

assuming nonempty interior.^{2.7} Implications are:

^{2.7}Otherwise, for $x \in \mathbb{R}^n$ as in (11), [228, §2.1-§2.3]

$$\overline{\text{int}\{x\}} = \bar{\emptyset} = \emptyset$$

the empty set is both open and closed.




Figure 13: **(a)** Ellipsoid in \mathbb{R} is a line segment whose boundary comprises two points. Intersection of line with ellipsoid in \mathbb{R} , **(b)** in \mathbb{R}^2 , **(c)** in \mathbb{R}^3 . Each ellipsoid illustrated has entire boundary constituted by zero-dimensional faces; in fact, by *vertices* (§2.6.1.0.1). Intersection of line with boundary is a point at entry to interior. These same facts hold in higher dimension.

- $\text{int } \mathcal{C} = \bar{\mathcal{C}} \setminus \partial \mathcal{C}$
- an open set has a boundary defined although not contained in the set
- interior of an open set is equivalent to the set itself

from which an open set is defined: \mathcal{C} is *open* $\Leftrightarrow \text{int } \mathcal{C} = \mathcal{C}$. [228, p.109]

The set illustrated in Figure 12b is not open because it is not equivalent to its interior, it is not closed because it does not contain its boundary, and it is not convex because it does not contain all convex combinations of its boundary points.

2.1.7.1 Line intersection with boundary

A line can intersect the boundary of a convex set in any dimension at a point demarcating the line's entry to the set interior. On one side of that entry-point along the line is the exterior of the set, on the other side is the set interior. In other words,

- starting from any point of a convex set, a move toward the interior is an immediate entry into the interior. [25, §II.2]

When a line intersects the interior of a convex body in any dimension, the boundary appears to the line to be as thin as a point. This is intuitively plausible because, for example, a line intersects the boundary of the ellipsoids in Figure 13 at a point in \mathbb{R} , \mathbb{R}^2 , and \mathbb{R}^3 . Such thinness is a remarkable fact when pondering visualization of convex *polyhedra* (§2.12, §5.14.3) in four dimensions, for example, having boundaries constructed from other three-dimensional convex polyhedra called *faces*.

We formally define *face* in (§2.6). For now, we observe the boundary of a convex body to be entirely constituted by all its faces of dimension lower than the body itself. For example: The ellipsoids in Figure 13 have boundaries composed only of zero-dimensional faces. The two-dimensional slab in Figure 11 is an unbounded *polyhedron* having one-dimensional faces making its boundary. The three-dimensional bounded polyhedron in Figure 18 has zero-, one-, and two-dimensional polygonal faces constituting its boundary.

2.1.7.1.1 Example. Intersection of line with boundary in \mathbb{R}^6 .

The convex cone of positive semidefinite matrices \mathbb{S}_+^3 (§2.9) in the ambient subspace of symmetric matrices \mathbb{S}^3 (§2.2.2.0.1) is a six-dimensional Euclidean body in *isometrically isomorphic* \mathbb{R}^6 (§2.2.1). The boundary of the positive semidefinite cone in this dimension comprises faces having only the dimensions 0, 1, and 3; *id est*, $\{\rho(\rho+1)/2, \rho=0, 1, 2\}$.

Unique minimum-distance projection PX (§E.9) of any point $X \in \mathbb{S}^3$ on that cone \mathbb{S}_+^3 is known in closed form (§7.1.2). Given, for example, $\lambda \in \text{int } \mathbb{R}_+^3$ and *diagonalization* (§A.5.2) of exterior point

$$X = Q\Lambda Q^T \in \mathbb{S}^3, \quad \Lambda \triangleq \begin{bmatrix} \lambda_1 & & \mathbf{0} \\ & \lambda_2 & \\ \mathbf{0} & & -\lambda_3 \end{bmatrix} \quad (17)$$

where $Q \in \mathbb{R}^{3 \times 3}$ is an orthogonal matrix, then the projection on \mathbb{S}_+^3 in \mathbb{R}^6 is

$$PX = Q \begin{bmatrix} \lambda_1 & & \mathbf{0} \\ & \lambda_2 & \\ \mathbf{0} & & 0 \end{bmatrix} Q^T \in \mathbb{S}_+^3 \quad (18)$$

This positive semidefinite matrix PX nearest X thus has rank 2, found by discarding all negative eigenvalues. The line connecting these two points is

$\{X + (PX - X)t \mid t \in \mathbb{R}\}$ where $t=0 \Leftrightarrow X$ and $t=1 \Leftrightarrow PX$. Because this line intersects the boundary of the *positive semidefinite cone* \mathbb{S}_+^3 at point PX and passes through its interior (by assumption), then the matrix corresponding to an infinitesimally positive perturbation of t there should reside interior to the cone (rank 3). Indeed, for ε an arbitrarily small positive constant,

$$X + (PX - X)t|_{t=1+\varepsilon} = Q(\Lambda + (P\Lambda - \Lambda)(1+\varepsilon))Q^T = Q \begin{bmatrix} \lambda_1 & & \mathbf{0} \\ & \lambda_2 & \\ \mathbf{0} & & \varepsilon\lambda_3 \end{bmatrix} Q^T \in \text{int } \mathbb{S}_+^3 \quad (19)$$

□

2.1.7.2 Tangential line intersection with boundary

A higher-dimensional boundary $\partial\mathcal{C}$ of a convex Euclidean body \mathcal{C} is simply a dimensionally larger set through which a line can pass when it does not intersect the body's interior. Still, for example, a line existing in five or more dimensions may pass *tangentially* (intersecting no point interior to \mathcal{C} [194, §15.3]) through a single point relatively interior to a three-dimensional face on $\partial\mathcal{C}$. Let's understand why by inductive reasoning.

Figure 14(a) shows a vertical line-segment whose boundary comprises its two endpoints. For a line to pass through the boundary tangentially (intersecting no point relatively interior to the line-segment), it must exist in an ambient space of at least two dimensions. Otherwise, the line is confined to the same one-dimensional space as the line-segment and must pass along the segment to reach the end points.

Figure 14(b) illustrates a two-dimensional ellipsoid whose boundary is constituted entirely by zero-dimensional faces. Again, a line must exist in at least two dimensions to tangentially pass through any single arbitrarily chosen point on the boundary (without intersecting the ellipsoid interior).

Now let's move to an ambient space of three dimensions. Figure 14(c) shows a polygon rotated into three dimensions. For a line to pass through its zero-dimensional boundary (one of its *vertices*) tangentially, it must exist in at least the two dimensions of the polygon. But for a line to pass tangentially through a single arbitrarily chosen point in the relative interior of a one-dimensional face on the boundary as illustrated, it must exist in at least three dimensions.




Figure 14: Line tangential (a) (b) to relative interior of a zero-dimensional face in \mathbb{R}^2 , (c) (d) to relative interior of a one-dimensional face in \mathbb{R}^3 .

Figure 14(d) illustrates a solid circular pyramid (upside-down) whose one-dimensional faces are line-segments emanating from its pointed end (its *vertex*). This pyramid's boundary is constituted solely by these one-dimensional line-segments. A line may pass through the boundary tangentially, striking only one arbitrarily chosen point relatively interior to a one-dimensional face, if it exists in at least the three-dimensional ambient space of the pyramid.

From these few examples, we deduce a general rule (without proof):

- A line may pass tangentially through a single arbitrarily chosen point relatively interior to a k -dimensional face on the boundary of a convex Euclidean body if the line exists in dimension at least equal to $k + 2$.

Now the interesting part, with regard to Figure 18 showing a bounded polyhedron in \mathbb{R}^3 ; call it \mathcal{P} : A line existing in at least four dimensions is required in order to pass tangentially (without hitting $\text{int } \mathcal{P}$) through a single arbitrary point in the relative interior of any two-dimensional polygonal face on the boundary of polyhedron \mathcal{P} . Now imagine that polyhedron \mathcal{P} is itself a three-dimensional face of some other polyhedron in \mathbb{R}^4 . To pass a line tangentially through polyhedron \mathcal{P} itself, striking only one point from its relative interior $\text{rel int } \mathcal{P}$ as claimed, requires a line existing in at least five dimensions.

This rule can help determine whether there exists unique solution to a convex optimization problem whose *feasible set* is an intersecting line; e.g., the *trilateration* problem (§5.4.2.2.4).

2.1.8 intersection, sum, difference, product

2.1.8.0.1 Theorem. *Intersection.* [53, §2.3.1] [271, §2]

The intersection of an arbitrary collection of convex sets is convex. \diamond

This theorem in converse is implicitly false in so far as a convex set can be formed by the intersection of sets that are not. Unions of convex sets are generally not convex. [177, p.22]

Vector sum of two convex sets \mathcal{C}_1 and \mathcal{C}_2

$$\mathcal{C}_1 + \mathcal{C}_2 = \{x + y \mid x \in \mathcal{C}_1, y \in \mathcal{C}_2\} \quad (20)$$

is convex.

By additive inverse, we can similarly define *vector difference* of two convex sets

$$\mathcal{C}_1 - \mathcal{C}_2 = \{x - y \mid x \in \mathcal{C}_1, y \in \mathcal{C}_2\} \quad (21)$$

which is convex. Applying this definition to nonempty convex set \mathcal{C}_1 , its self-difference $\mathcal{C}_1 - \mathcal{C}_1$ is generally nonempty, nontrivial, and convex; *e.g.*, for any *convex cone* \mathcal{K} , (§2.7.2) the set $\mathcal{K} - \mathcal{K}$ constitutes its affine hull. [271, p.15]

Cartesian product of convex sets

$$\mathcal{C}_1 \times \mathcal{C}_2 = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid x \in \mathcal{C}_1, y \in \mathcal{C}_2 \right\} = \begin{bmatrix} \mathcal{C}_1 \\ \mathcal{C}_2 \end{bmatrix} \quad (22)$$

remains convex. The converse also holds; *id est*, a Cartesian product is convex iff each set is. [177, p.23]

Convex results are also obtained for scaling $\kappa\mathcal{C}$ of a convex set \mathcal{C} , rotation/reflection $Q\mathcal{C}$, or translation $\mathcal{C} + \alpha$; all similarly defined.

Given any operator T and convex set \mathcal{C} , we are prone to write $T(\mathcal{C})$ meaning

$$T(\mathcal{C}) \triangleq \{T(x) \mid x \in \mathcal{C}\} \quad (23)$$

Given linear operator T , it therefore follows from (20),

$$\begin{aligned} T(\mathcal{C}_1 + \mathcal{C}_2) &= \{T(x + y) \mid x \in \mathcal{C}_1, y \in \mathcal{C}_2\} \\ &= \{T(x) + T(y) \mid x \in \mathcal{C}_1, y \in \mathcal{C}_2\} \\ &= T(\mathcal{C}_1) + T(\mathcal{C}_2) \end{aligned} \quad (24)$$

2.1.9 inverse image

While *epigraph* and *sublevel sets* (§3.1.7) of a convex function must be convex, it generally holds that image and inverse image of a convex function are not. Although there are many examples to the contrary, the most prominent are the affine functions (§3.1.6):

2.1.9.0.1 Theorem. *Inverse image.* [271, §3]
Let f be a mapping from $\mathbb{R}^{p \times k}$ to $\mathbb{R}^{m \times n}$.

- The image of a convex set \mathcal{C} under any affine function

$$f(\mathcal{C}) = \{f(X) \mid X \in \mathcal{C}\} \subseteq \mathbb{R}^{m \times n} \quad (25)$$

is convex.




Figure 15:(confer Figure 140) Action of linear map represented by $A \in \mathbb{R}^{m \times n}$: Component of vector x in nullspace $\mathcal{N}(A)$ maps to origin while component in rowspace $\mathcal{R}(A^T)$ maps to range $\mathcal{R}(A)$. For any $A \in \mathbb{R}^{m \times n}$, $AA^\dagger Ax = b$ (§E), and inverse image of $b \in \mathcal{R}(A)$ is a nonempty affine set: $x_p + \mathcal{N}(A)$.

- Inverse image of a convex set \mathcal{F} ,

$$f^{-1}(\mathcal{F}) = \{X \mid f(X) \in \mathcal{F}\} \subseteq \mathbb{R}^{p \times k} \quad (26)$$

a single- or many-valued mapping, under any affine function f is convex. \diamond

In particular, any affine transformation of an affine set remains affine. [271, p.8] Ellipsoids are invariant to any [sic] affine transformation.

Although not precluded, this *inverse image theorem* does not require a uniquely invertible mapping f . Figure 15, for example, mechanizes inverse image under a general linear map. Example 2.9.1.0.2 and Example 3.1.7.0.2 offer further applications.

Each converse of this two-part theorem is generally false; *id est*, given f affine, a convex image $f(\mathcal{C})$ does not imply that set \mathcal{C} is convex, and neither does a convex inverse image $f^{-1}(\mathcal{F})$ imply set \mathcal{F} is convex. A counter-example, invalidating a converse, is easy to visualize when the affine function is an orthogonal projector [294] [220]:

2.1.9.0.2 Corollary. *Projection on subspace.*^{2.8} (1820) [271, §3]
 Orthogonal projection of a convex set on a subspace or nonempty affine set is another convex set. \diamond

Again, the converse is false. Shadows, for example, are umbral projections that can be convex when the body providing the shade is not.

2.2 Vectorized-matrix inner product

Euclidean space \mathbb{R}^n comes equipped with a linear vector inner-product

$$\langle y, z \rangle \triangleq y^T z \quad (27)$$

We prefer those angle brackets to connote a geometric rather than algebraic perspective; *e.g.*, vector y might represent a hyperplane normal (§2.4.2). Two vectors are *orthogonal* (*perpendicular*) to one another if and only if their inner product vanishes;

$$y \perp z \Leftrightarrow \langle y, z \rangle = 0 \quad (28)$$

When orthogonal vectors each have unit *norm*, then they are *orthonormal*. A vector inner-product defines Euclidean norm (vector 2-norm)

$$\|y\|_2 = \|y\| \triangleq \sqrt{y^T y}, \quad \|y\| = 0 \Leftrightarrow y = 0 \quad (29)$$

For linear operation A on a vector, represented by a real matrix, the *adjoint operation* A^T is transposition and defined for matrix A by [201, §3.10]

$$\langle y, A^T z \rangle \triangleq \langle Ay, z \rangle \quad (30)$$

The vector inner-product for matrices is calculated just as it is for vectors; by first transforming a matrix in $\mathbb{R}^{p \times k}$ to a vector in \mathbb{R}^{pk} by concatenating its columns in the *natural order*. For lack of a better term, we shall call that linear *bijective* (one-to-one and onto [201, App.A1.2]) transformation

^{2.8}For hyperplane representations see §2.4.2. For projection of convex sets on hyperplanes see [331, §6.6]. A nonempty affine set is called an *affine subset* (§2.3.1.0.1). Orthogonal projection of points on affine subsets is reviewed in §E.4.

vectorization. For example, the vectorization of $Y = [y_1 \ y_2 \ \cdots \ y_k] \in \mathbb{R}^{p \times k}$ [145] [290] is

$$\text{vec } Y \triangleq \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_k \end{bmatrix} \in \mathbb{R}^{pk} \quad (31)$$

Then the vectorized-matrix inner-product is trace of matrix inner-product; for $Z \in \mathbb{R}^{p \times k}$, [53, §2.6.1] [177, §0.3.1] [341, §8] [325, §2.2]

$$\langle Y, Z \rangle \triangleq \text{tr}(Y^T Z) = \text{vec}(Y)^T \text{vec } Z \quad (32)$$

where (§A.1.1)

$$\text{tr}(Y^T Z) = \text{tr}(Z Y^T) = \text{tr}(Y Z^T) = \text{tr}(Z^T Y) = \mathbf{1}^T (Y \circ Z) \mathbf{1} \quad (33)$$

and where \circ denotes the *Hadamard product*^{2.9} of matrices [139, §1.1.4]. The adjoint operation A^T on a matrix can therefore be defined in like manner:

$$\langle Y, A^T Z \rangle \triangleq \langle AY, Z \rangle \quad (34)$$

Take any element \mathcal{C}_1 from a matrix-valued set in $\mathbb{R}^{p \times k}$, for example, and consider any particular dimensionally compatible real vectors v and w . Then vector inner-product of \mathcal{C}_1 with vw^T is

$$\langle vw^T, \mathcal{C}_1 \rangle = \langle v, \mathcal{C}_1 w \rangle = v^T \mathcal{C}_1 w = \text{tr}(wv^T \mathcal{C}_1) = \mathbf{1}^T ((vw^T) \circ \mathcal{C}_1) \mathbf{1} \quad (35)$$

Further, linear bijective vectorization is *distributive* with respect to Hadamard product; *id est*,

$$\text{vec}(Y \circ Z) = \text{vec}(Y) \circ \text{vec}(Z) \quad (36)$$

2.2.0.0.1 Example. Application of inverse image theorem.

Suppose the set $\mathcal{C} \subseteq \mathbb{R}^{p \times k}$ is convex. Then for any particular vectors $v \in \mathbb{R}^p$ and $w \in \mathbb{R}^k$, the set of vector inner-products

$$\mathcal{Y} \triangleq v^T \mathcal{C} w = \langle vw^T, \mathcal{C} \rangle \subseteq \mathbb{R} \quad (37)$$

^{2.9}The Hadamard product is a simple entrywise product of corresponding entries from two matrices of like size; *id est*, not necessarily square. A commutative operation, the Hadamard product can be extracted from within a Kronecker product. [180, p.475]

is convex. This result is a consequence of the *inverse image theorem*. Yet it is easy to show directly that convex combination of elements from \mathcal{Y} remains an element of \mathcal{Y} .^{2.10} \square

More generally, vw^T in (37) may be replaced with any particular matrix $Z \in \mathbb{R}^{p \times k}$ while convexity of the set $\langle Z, \mathcal{C} \rangle \subseteq \mathbb{R}$ persists. Further, by replacing v and w with any particular respective matrices U and W of dimension compatible with all elements of convex set \mathcal{C} , then set $U^T \mathcal{C} W$ is convex by the *inverse image theorem* because it is a linear mapping of \mathcal{C} .

2.2.1 Frobenius'

2.2.1.0.1 Definition. Isomorphic.

An *isomorphism* of a vector space is a transformation equivalent to a linear bijective mapping. The image and inverse image under the transformation operator are then called isomorphic vector spaces. \triangle

Isomorphic vector spaces are characterized by preservation of *adjacency*; *id est*, if v and w are points connected by a line segment in one vector space, then their images will also be connected by a line segment. Two Euclidean bodies may be considered isomorphic if there exists an isomorphism, of their vector spaces, under which the bodies correspond. [327, §I.1] Projection (§E) is not an isomorphism, for example; hence, perfect reconstruction (inverse projection) is generally impossible without additional information.

When $Z = Y \in \mathbb{R}^{p \times k}$ in (32), *Frobenius' norm* is resultant from vector inner-product; (*confer* (1578))

$$\begin{aligned} \|Y\|_F^2 &= \|\text{vec } Y\|_2^2 = \langle Y, Y \rangle = \text{tr}(Y^T Y) \\ &= \sum_{i,j} Y_{ij}^2 = \sum_i \lambda(Y^T Y)_i = \sum_i \sigma(Y)_i^2 \end{aligned} \quad (38)$$

^{2.10}To verify that, take any two elements \mathcal{C}_1 and \mathcal{C}_2 from the convex matrix-valued set \mathcal{C} , and then form the vector inner-products (37) that are two elements of \mathcal{Y} by definition. Now make a convex combination of those inner products; *videlicet*, for $0 \leq \mu \leq 1$

$$\mu \langle vw^T, \mathcal{C}_1 \rangle + (1 - \mu) \langle vw^T, \mathcal{C}_2 \rangle = \langle vw^T, \mu \mathcal{C}_1 + (1 - \mu) \mathcal{C}_2 \rangle$$

The two sides are equivalent by linearity of inner product. The right-hand side remains a vector inner-product of vw^T with an element $\mu \mathcal{C}_1 + (1 - \mu) \mathcal{C}_2$ from the convex set \mathcal{C} ; hence, it belongs to \mathcal{Y} . Since that holds true for any two elements from \mathcal{Y} , then it must be a convex set. \blacklozenge




Figure 16: Linear injective mapping $Tx = Ax : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ of Euclidean body remains two-dimensional under mapping represented by skinny full-rank matrix $A \in \mathbb{R}^{3 \times 2}$; two bodies are isomorphic by Definition 2.2.1.0.1.

where $\lambda(Y^T Y)_i$ is the i^{th} eigenvalue of $Y^T Y$, and $\sigma(Y)_i$ the i^{th} singular value of Y . Were Y a *normal matrix* (§A.5.2), then $\sigma(Y) = |\lambda(Y)|$ [352, §8.1] thus

$$\|Y\|_F^2 = \sum_i \lambda(Y)_i^2 = \|\lambda(Y)\|_2^2 = \langle \lambda(Y), \lambda(Y) \rangle = \langle Y, Y \rangle \quad (39)$$

The converse (39) \Rightarrow normal matrix Y also holds. [180, §2.5.4]

Frobenius norm is the Euclidean norm of vectorized matrices. Because the metrics are equivalent

$$\|\text{vec } X - \text{vec } Y\|_2 = \|X - Y\|_F \quad (40)$$

and because vectorization (31) is a linear bijective map, then vector space $\mathbb{R}^{p \times k}$ is isometrically isomorphic with vector space \mathbb{R}^{pk} in the Euclidean sense and vec is an isometric isomorphism of $\mathbb{R}^{p \times k}$.^{2.11} Because of this Euclidean structure, all the known results from convex analysis in Euclidean space \mathbb{R}^n carry over directly to the space of real matrices $\mathbb{R}^{p \times k}$.

^{2.11}Given matrix A , its range $\mathcal{R}(A)$ (§2.5) is isometrically isomorphic with its vectorized range $\text{vec } \mathcal{R}(A)$ but not with $\mathcal{R}(\text{vec } A)$.

2.2.1.1 Injective linear operators

Injective mapping (transformation) means one-to-one mapping; synonymous with *uniquely invertible* linear mapping on Euclidean space. Linear injective mappings are fully characterized by lack of a nontrivial nullspace.

2.2.1.1.1 Definition. Isometric isomorphism.

An isometric isomorphism of a vector space having a *metric* defined on it is a linear bijective mapping T that preserves distance; *id est*, for all $x, y \in \text{dom } T$

$$\|Tx - Ty\| = \|x - y\| \quad (41)$$

Then the isometric isomorphism T is a *bijective isometry*. \triangle

Unitary linear operator $Q : \mathbb{R}^k \rightarrow \mathbb{R}^k$, represented by orthogonal matrix $Q \in \mathbb{R}^{k \times k}$ (§B.5), is an isometric isomorphism; *e.g.*, discrete Fourier transform via (755). Suppose $T(X) = UXQ$, for example. Then we say the Frobenius norm is *orthogonally invariant*; meaning, for $X, Y \in \mathbb{R}^{p \times k}$ and dimensionally compatible *orthonormal matrix*^{2.12} U and orthogonal matrix Q

$$\|U(X - Y)Q\|_F = \|X - Y\|_F \quad (42)$$

Yet isometric operator $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, represented by $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$ on \mathbb{R}^2 ,

is injective but not a *surjective* map to \mathbb{R}^3 . [201, §1.6, §2.6] This operator T can therefore be a bijective isometry only with respect to its range.

Any linear injective transformation on Euclidean space is uniquely invertible on its range. In fact, any linear injective transformation has a range whose dimension equals that of its domain. In other words, for any invertible linear transformation T [*ibidem*]

$$\dim \text{dom } T = \dim \mathcal{R}(T) \quad (43)$$

e.g., T represented by skinny-or-square full-rank matrices. (Figure 16) An important consequence of this fact is:

- Affine dimension of image, of any n -dimensional Euclidean body in domain of operator T , is invariant to linear injective transformation.

^{2.12}Any matrix U whose columns are orthonormal with respect to each other ($U^T U = I$); these include the orthogonal matrices.




Figure 17: Linear noninjective mapping $PTx = A^\dagger Ax : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ of three-dimensional Euclidean body \mathcal{B} has affine dimension 2 under projection on rowspace of fat full-rank matrix $A \in \mathbb{R}^{2 \times 3}$. Set of coefficients of orthogonal projection $T\mathcal{B} = \{Ax \mid x \in \mathcal{B}\}$ is isomorphic with projection $P(T\mathcal{B})$ [sic].

2.2.1.2 Noninjective linear operators

Mappings in Euclidean space created by noninjective linear operators can be characterized in terms of an orthogonal projector (§E). Consider noninjective linear operator $Tx = Ax : \mathbb{R}^n \rightarrow \mathbb{R}^m$ represented by fat matrix $A \in \mathbb{R}^{m \times n}$ ($m < n$). What can be said about the nature of this m -dimensional mapping?

Concurrently, consider injective linear operator $Px = A^\dagger x : \mathbb{R}^m \rightarrow \mathbb{R}^n$ where $\mathcal{R}(A^\dagger) = \mathcal{R}(A^T)$. $P(Ax) = PTx$ achieves projection of vector x on the row space $\mathcal{R}(A^T)$. (§E.3.1) This means vector Ax can be succinctly interpreted as coefficients of orthogonal projection.

Pseudoinverse matrix A^\dagger is skinny and full rank, so operator Px is a linear bijection with respect to its range $\mathcal{R}(A^\dagger)$. By Definition 2.2.1.0.1, image $P(T\mathcal{B})$ of projection $PT(\mathcal{B})$ on $\mathcal{R}(A^T)$ in \mathbb{R}^n must therefore be isomorphic with the set of projection coefficients $T\mathcal{B} = \{Ax \mid x \in \mathcal{B}\}$ in \mathbb{R}^m and have the same affine dimension by (43). To illustrate, we present a three-dimensional Euclidean body \mathcal{B} in Figure 17 where any point x in the nullspace $\mathcal{N}(A)$ maps to the origin.

2.2.2 Symmetric matrices

2.2.2.0.1 Definition. *Symmetric matrix subspace.*

Define a subspace of $\mathbb{R}^{M \times M}$: the convex set of all symmetric $M \times M$ matrices;

$$\mathbb{S}^M \triangleq \{A \in \mathbb{R}^{M \times M} \mid A = A^T\} \subseteq \mathbb{R}^{M \times M} \quad (44)$$

This subspace comprising symmetric matrices \mathbb{S}^M is isomorphic with the vector space $\mathbb{R}^{M(M+1)/2}$ whose dimension is the number of free variables in a symmetric $M \times M$ matrix. The *orthogonal complement* [294] [220] of \mathbb{S}^M is

$$\mathbb{S}^{M\perp} \triangleq \{A \in \mathbb{R}^{M \times M} \mid A = -A^T\} \subseteq \mathbb{R}^{M \times M} \quad (45)$$

the subspace of *antisymmetric* matrices in $\mathbb{R}^{M \times M}$; *id est*,

$$\mathbb{S}^M \oplus \mathbb{S}^{M\perp} = \mathbb{R}^{M \times M} \quad (46)$$

where unique vector sum \oplus is defined on page 714. \triangle

All antisymmetric matrices are hollow by definition (have $\mathbf{0}$ main diagonal). Any square matrix $A \in \mathbb{R}^{M \times M}$ can be written as the sum of its symmetric and antisymmetric parts: respectively,

$$A = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T) \quad (47)$$

The symmetric part is orthogonal in \mathbb{R}^{M^2} to the antisymmetric part; *videlicet*,

$$\text{tr}((A^T + A)(A - A^T)) = 0 \quad (48)$$

In the ambient space of real matrices, the antisymmetric matrix subspace can be described

$$\mathbb{S}^{M\perp} \triangleq \left\{ \frac{1}{2}(A - A^T) \mid A \in \mathbb{R}^{M \times M} \right\} \subseteq \mathbb{R}^{M \times M} \quad (49)$$

because any matrix in \mathbb{S}^M is orthogonal to any matrix in $\mathbb{S}^{M\perp}$. Further confined to the ambient subspace of symmetric matrices, because of antisymmetry, $\mathbb{S}^{M\perp}$ would become trivial.

2.2.2.1 Isomorphism of symmetric matrix subspace

When a matrix is symmetric in \mathbb{S}^M , we may still employ the vectorization transformation (31) to \mathbb{R}^{M^2} ; vec , an isometric isomorphism. We might instead choose to realize in the lower-dimensional subspace $\mathbb{R}^{M(M+1)/2}$ by ignoring redundant entries (below the main diagonal) during transformation. Such a realization would remain isomorphic but not isometric. Lack of isometry is a spatial distortion due now to disparity in metric between \mathbb{R}^{M^2} and $\mathbb{R}^{M(M+1)/2}$. To realize isometrically in $\mathbb{R}^{M(M+1)/2}$, we must make a correction: For $Y = [Y_{ij}] \in \mathbb{S}^M$ we introduce the symmetric vectorization

$$\text{svec } Y \triangleq \begin{bmatrix} Y_{11} \\ \sqrt{2}Y_{12} \\ Y_{22} \\ \sqrt{2}Y_{13} \\ \sqrt{2}Y_{23} \\ Y_{33} \\ \vdots \\ Y_{MM} \end{bmatrix} \in \mathbb{R}^{M(M+1)/2} \quad (50)$$

where all entries off the main diagonal have been scaled. Now for $Z \in \mathbb{S}^M$

$$\langle Y, Z \rangle \triangleq \text{tr}(Y^T Z) = \text{vec}(Y)^T \text{vec } Z = \text{svec}(Y)^T \text{svec } Z \quad (51)$$

Then because the metrics become equivalent, for $X \in \mathbb{S}^M$

$$\|\text{svec } X - \text{svec } Y\|_2 = \|X - Y\|_F \quad (52)$$

and because symmetric vectorization (50) is a linear bijective mapping, then svec is an isometric isomorphism of the symmetric matrix subspace. In other words, \mathbb{S}^M is isometrically isomorphic with $\mathbb{R}^{M(M+1)/2}$ in the Euclidean sense under transformation svec .

The set of all symmetric matrices \mathbb{S}^M forms a proper subspace in $\mathbb{R}^{M \times M}$, so for it there exists a standard orthonormal *basis* in isometrically isomorphic $\mathbb{R}^{M(M+1)/2}$

$$\{E_{ij} \in \mathbb{S}^M\} = \left\{ \begin{array}{ll} e_i e_i^T, & i = j = 1 \dots M \\ \frac{1}{\sqrt{2}}(e_i e_j^T + e_j e_i^T), & 1 \leq i < j \leq M \end{array} \right\} \quad (53)$$

where $M(M+1)/2$ standard basis matrices E_{ij} are formed from the standard basis vectors

$$e_i = \left[\begin{array}{c} \left\{ \begin{array}{l} 1, \quad i = j \\ 0, \quad i \neq j \end{array} \right. , \quad j = 1 \dots M \end{array} \right] \in \mathbb{R}^M \quad (54)$$

Thus we have a basic *orthogonal expansion* for $Y \in \mathbb{S}^M$

$$Y = \sum_{j=1}^M \sum_{i=1}^j \langle E_{ij}, Y \rangle E_{ij} \quad (55)$$

whose coefficients

$$\langle E_{ij}, Y \rangle = \begin{cases} Y_{ii}, & i = 1 \dots M \\ Y_{ij}\sqrt{2}, & 1 \leq i < j \leq M \end{cases} \quad (56)$$

correspond to entries of the symmetric vectorization (50).

2.2.3 Symmetric hollow subspace

2.2.3.0.1 Definition. *Hollow subspaces.* [312]

Define a subspace of $\mathbb{R}^{M \times M}$: the convex set of all (real) symmetric $M \times M$ matrices having $\mathbf{0}$ main diagonal;

$$\mathbb{R}_h^{M \times M} \triangleq \{A \in \mathbb{R}^{M \times M} \mid A = A^T, \delta(A) = \mathbf{0}\} \subset \mathbb{R}^{M \times M} \quad (57)$$

where the main diagonal of $A \in \mathbb{R}^{M \times M}$ is denoted (§A.1)

$$\delta(A) \in \mathbb{R}^M \quad (1328)$$

Operating on a vector, linear operator δ naturally returns a diagonal matrix; $\delta(\delta(A))$ is a diagonal matrix. Operating recursively on a vector $\Lambda \in \mathbb{R}^N$ or diagonal matrix $\Lambda \in \mathbb{S}^N$, operator $\delta(\delta(\Lambda))$ returns Λ itself;

$$\delta^2(\Lambda) \equiv \delta(\delta(\Lambda)) \triangleq \Lambda \quad (1330)$$

The subspace $\mathbb{R}_h^{M \times M}$ (57) comprising (real) symmetric hollow matrices is isomorphic with subspace $\mathbb{R}^{M(M-1)/2}$. The orthogonal complement of $\mathbb{R}_h^{M \times M}$ is

$$\mathbb{R}_h^{M \times M \perp} \triangleq \{A \in \mathbb{R}^{M \times M} \mid A = -A^T + 2\delta^2(A)\} \subseteq \mathbb{R}^{M \times M} \quad (58)$$

the subspace of *antisymmetric antihollow* matrices in $\mathbb{R}^{M \times M}$; *id est*,

$$\mathbb{R}_h^{M \times M} \oplus \mathbb{R}_h^{M \times M \perp} = \mathbb{R}^{M \times M} \quad (59)$$

Yet defined instead as a proper subspace of \mathbb{S}^M

$$\mathbb{S}_h^M \triangleq \{A \in \mathbb{S}^M \mid \delta(A) = \mathbf{0}\} \subset \mathbb{S}^M \quad (60)$$

the orthogonal complement $\mathbb{S}_h^{M \perp}$ of \mathbb{S}_h^M in ambient \mathbb{S}^M

$$\mathbb{S}_h^{M \perp} \triangleq \{A \in \mathbb{S}^M \mid A = \delta^2(A)\} \subseteq \mathbb{S}^M \quad (61)$$

is simply the subspace of diagonal matrices; *id est*,

$$\mathbb{S}_h^M \oplus \mathbb{S}_h^{M \perp} = \mathbb{S}^M \quad (62)$$

△

Any matrix $A \in \mathbb{R}^{M \times M}$ can be written as the sum of its symmetric hollow and antisymmetric antihollow parts: respectively,

$$A = \left(\frac{1}{2}(A + A^T) - \delta^2(A) \right) + \left(\frac{1}{2}(A - A^T) + \delta^2(A) \right) \quad (63)$$

The symmetric hollow part is orthogonal in \mathbb{R}^{M^2} to the antisymmetric antihollow part; *videlicet*,

$$\text{tr} \left(\left(\frac{1}{2}(A + A^T) - \delta^2(A) \right) \left(\frac{1}{2}(A - A^T) + \delta^2(A) \right) \right) = 0 \quad (64)$$

In the ambient space of real matrices, the antisymmetric antihollow subspace is described

$$\mathbb{S}_h^{M \perp} \triangleq \left\{ \frac{1}{2}(A - A^T) + \delta^2(A) \mid A \in \mathbb{R}^{M \times M} \right\} \subseteq \mathbb{R}^{M \times M} \quad (65)$$

because any matrix in \mathbb{S}_h^M is orthogonal to any matrix in $\mathbb{S}_h^{M \perp}$. Yet in the ambient space of symmetric matrices \mathbb{S}^M , the antihollow subspace is nontrivial;

$$\mathbb{S}_h^{M \perp} \triangleq \{\delta^2(A) \mid A \in \mathbb{S}^M\} = \{\delta(u) \mid u \in \mathbb{R}^M\} \subseteq \mathbb{S}^M \quad (66)$$

In anticipation of their utility with Euclidean distance matrices (EDMs) in §5, for symmetric hollow matrices we introduce the linear bijective vectorization dvec that is the natural analogue to symmetric matrix vectorization svec (50): for $Y = [Y_{ij}] \in \mathbb{S}_h^M$

$$\text{dvec } Y \triangleq \sqrt{2} \begin{bmatrix} Y_{12} \\ Y_{13} \\ Y_{23} \\ Y_{14} \\ Y_{24} \\ Y_{34} \\ \vdots \\ Y_{M-1,M} \end{bmatrix} \in \mathbb{R}^{M(M-1)/2} \quad (67)$$

Like svec (50), dvec is an isometric isomorphism on the symmetric hollow subspace.

The set of all symmetric hollow matrices \mathbb{S}_h^M forms a proper subspace in $\mathbb{R}^{M \times M}$, so for it there must be a standard orthonormal basis in isometrically isomorphic $\mathbb{R}^{M(M-1)/2}$

$$\{E_{ij} \in \mathbb{S}_h^M\} = \left\{ \frac{1}{\sqrt{2}}(e_i e_j^T + e_j e_i^T), \quad 1 \leq i < j \leq M \right\} \quad (68)$$

where $M(M-1)/2$ standard basis matrices E_{ij} are formed from the standard basis vectors $e_i \in \mathbb{R}^M$.

The *symmetric hollow majorization corollary* on page 545 characterizes eigenvalues of symmetric hollow matrices.

2.3 Hulls

We focus on the affine, convex, and conic hulls: convex sets that may be regarded as kinds of Euclidean container or vessel united with its interior.

2.3.1 Affine hull, affine dimension

Affine dimension of any set in \mathbb{R}^n is the dimension of the smallest affine set (empty set, point, line, plane, hyperplane (§2.4.2), translated subspace, \mathbb{R}^n) that contains it. For nonempty sets, affine dimension is the same as dimension of the subspace parallel to that affine set. [271, §1] [177, §A.2.1]




Figure 18: Convex hull of a random list of points in \mathbb{R}^3 . Some points from that generating list reside interior to this *convex polyhedron*. [333, *Convex Polyhedron*] (Avis-Fukuda-Mizukoshi)

Ascribe the points in a list $\{x_\ell \in \mathbb{R}^n, \ell=1 \dots N\}$ to the columns of matrix X :

$$X = [x_1 \cdots x_N] \in \mathbb{R}^{n \times N} \quad (69)$$

In particular, we define *affine dimension* r of the N -point list X as dimension of the smallest affine set in Euclidean space \mathbb{R}^n that contains X ;

$$r \triangleq \dim \text{aff } X \quad (70)$$

Affine dimension r is a lower bound sometimes called *embedding dimension*. [312] [164] That affine set \mathcal{A} in which those points are embedded is unique and called the *affine hull* [293, §2.1];

$$\begin{aligned} \mathcal{A} &\triangleq \text{aff } \{x_\ell \in \mathbb{R}^n, \ell=1 \dots N\} &= \text{aff } X \\ &= x_1 + \mathcal{R}\{x_\ell - x_1, \ell=2 \dots N\} &= \{Xa \mid a^T \mathbf{1} = 1\} \subseteq \mathbb{R}^n \end{aligned} \quad (71)$$

parallel to subspace

$$\mathcal{R}\{x_\ell - x_1, \ell=2 \dots N\} = \mathcal{R}(X - x_1 \mathbf{1}^T) \subseteq \mathbb{R}^n \quad (72)$$

where

$$\mathcal{R}(A) = \{Ax \mid \forall x\} \quad (133)$$

Given some arbitrary set \mathcal{C} and any $x \in \mathcal{C}$

$$\text{aff } \mathcal{C} = x + \text{aff}(\mathcal{C} - x) \quad (73)$$

where $\text{aff}(\mathcal{C} - x)$ is a subspace.

2.3.1.0.1 Definition. *Affine subset.*

We analogize *affine subset* to subspace,^{2.13} defining it to be any nonempty affine set (§2.1.4). △

$$\text{aff } \emptyset \triangleq \emptyset \tag{74}$$

The affine hull of a point x is that point itself;

$$\text{aff } \{x\} = \{x\} \tag{75}$$

The affine hull of two distinct points is the unique line through them. (Figure 19) The affine hull of three noncollinear points in any dimension is that unique plane containing the points, and so on. The subspace of symmetric matrices \mathbb{S}^m is the affine hull of the cone of positive semidefinite matrices; (§2.9)

$$\text{aff } \mathbb{S}_+^m = \mathbb{S}^m \tag{76}$$

2.3.1.0.2 Example. *Affine hull of rank-1 correlation matrices.* [193]

The set of all $m \times m$ *rank-1 correlation matrices* is defined by all the binary vectors y in \mathbb{R}^m (confer §5.9.1.0.1)

$$\{yy^T \in \mathbb{S}_+^m \mid \delta(yy^T) = \mathbf{1}\} \tag{77}$$

Affine hull of the rank-1 correlation matrices is equal to the set of normalized symmetric matrices; *id est*,

$$\text{aff}\{yy^T \in \mathbb{S}_+^m \mid \delta(yy^T) = \mathbf{1}\} = \{A \in \mathbb{S}^m \mid \delta(A) = \mathbf{1}\} \tag{78}$$

□

2.3.1.0.3 Exercise. *Affine hull of correlation matrices.*

Prove (78) via definition of affine hull. Find the convex hull instead. ▼

^{2.13}The popular term *affine subspace* is an oxymoron.




Figure 19: Given two points in Euclidean vector space of any dimension, their various hulls are illustrated. Each hull is a subset of range; generally, $\mathcal{A}, \mathcal{C}, \mathcal{K} \subseteq \mathcal{R} \ni \mathbf{0}$. (Cartesian axes drawn for reference.)

2.3.1.1 Partial order induced by \mathbb{R}_+^N and \mathbb{S}_+^M

Notation $a \succeq 0$ means vector a belongs to the nonnegative orthant \mathbb{R}_+^N , while $a \succ 0$ means vector a belongs to the nonnegative orthant's interior $\text{int } \mathbb{R}_+^N$, whereas $a \succeq b$ denotes comparison of vector a to vector b on \mathbb{R}^N with respect to the nonnegative orthant; *id est*, $a \succeq b$ means $a - b$ belongs to the nonnegative orthant, but neither a or b necessarily belongs to that orthant. With particular respect to the nonnegative orthant, $a \succeq b \Leftrightarrow a_i \geq b_i \ \forall i$ (335). More generally, $a \succeq_{\mathcal{K}} b$ or $a \succ_{\mathcal{K}} b$ denotes comparison with respect to pointed closed convex cone \mathcal{K} , but equivalence with entrywise comparison does not hold. (§2.7.2.2)

The symbol \geq is reserved for scalar comparison on the real line \mathbb{R} with respect to the nonnegative real line \mathbb{R}_+ as in $a^T y \geq b$. Comparison of matrices with respect to the positive semidefinite cone \mathbb{S}_+^M , like $I \succeq A \succeq 0$ in Example 2.3.2.0.1, is explained in §2.9.0.1.

2.3.2 Convex hull

The *convex hull* [177, §A.1.4] [271] of any *bounded*^{2.14} list or set of N points $X \in \mathbb{R}^{n \times N}$ forms a unique bounded convex polyhedron (*confer* §2.12.0.0.1) whose vertices constitute some subset of that list;

$$\mathcal{P} \triangleq \text{conv} \{x_\ell, \ell=1 \dots N\} = \text{conv } X = \{Xa \mid a^T \mathbf{1} = 1, a \succeq 0\} \subseteq \mathbb{R}^n \quad (79)$$

Union of relative interior and relative boundary (§2.6.1.3) of the polyhedron comprise its convex hull \mathcal{P} , the smallest closed convex set that contains the list X ; *e.g.*, Figure 18. Given \mathcal{P} , the *generating list* $\{x_\ell\}$ is not unique. But because every bounded polyhedron is the convex hull of its vertices, [293, §2.12.2] the vertices of \mathcal{P} comprise a *minimal set* of generators.

Given some arbitrary set $\mathcal{C} \subseteq \mathbb{R}^n$, its convex hull $\text{conv } \mathcal{C}$ is equivalent to the smallest closed convex set containing it. (*confer* §2.4.1.1.1) The convex hull is a subset of the affine hull;

$$\text{conv } \mathcal{C} \subseteq \text{aff } \mathcal{C} = \text{aff } \overline{\mathcal{C}} = \overline{\text{aff } \mathcal{C}} = \text{aff } \text{conv } \mathcal{C} \quad (80)$$

Any closed bounded convex set \mathcal{C} is equal to the convex hull of its boundary;

$$\mathcal{C} = \text{conv } \partial \mathcal{C} \quad (81)$$

^{2.14}A set in \mathbb{R}^n is bounded iff it can be contained in a Euclidean ball having finite radius. [96, §2.2] (*confer* §5.7.3.0.1)

$$\text{conv } \emptyset \stackrel{\Delta}{=} \emptyset \quad (82)$$

2.3.2.0.1 Example. *Hull of outer product.* [124] [254] [10, §4.1] [258, §3] [211, §2.4] [212] Convex hull of the set comprising outer product of orthonormal matrices has equivalent expression: for $1 \leq k \leq N$ (§2.9.0.1)

$$\text{conv}\{UU^T \mid U \in \mathbb{R}^{N \times k}, U^T U = I\} = \{A \in \mathbb{S}^N \mid I \succeq A \succeq 0, \langle I, A \rangle = k\} \subset \mathbb{S}_+^N \quad (83)$$

This important convex body we call *Fantope* (after mathematician Ky Fan). In case $k = 1$, there is slight simplification: ((1516), Example 2.9.2.4.1)

$$\text{conv}\{UU^T \mid U \in \mathbb{R}^N, U^T U = 1\} = \{A \in \mathbb{S}^N \mid A \succeq 0, \langle I, A \rangle = 1\} \quad (84)$$

In case $k = N$, the Fantope is identity matrix I . More generally, the set

$$\{UU^T \mid U \in \mathbb{R}^{N \times k}, U^T U = I\} \quad (85)$$

comprises the extreme points (§2.6.0.0.1) of its convex hull. By (1376), each and every extreme point UU^T has only k nonzero eigenvalues λ and they all equal 1; *id est*, $\lambda(UU^T)_{1:k} = \lambda(U^T U) = \mathbf{1}$. So the Frobenius norm of each and every extreme point equals the same constant

$$\|UU^T\|_{\text{F}}^2 = k \quad (86)$$

Each extreme point simultaneously lies on the boundary of the positive semidefinite cone (when $k < N$, §2.9) and on the boundary of a *hypersphere* of dimension $k(N - \frac{k}{2} + \frac{1}{2})$ and radius $\sqrt{k(1 - \frac{k}{N})}$ centered at $\frac{k}{N}I$ along the ray (base $\mathbf{0}$) through the identity matrix I in isomorphic vector space $\mathbb{R}^{N(N+1)/2}$ (§2.2.2.1).

Figure 20 illustrates extreme points (85) comprising the boundary of a Fantope, the boundary of a *disc* corresponding to $k = 1$, $N = 2$; but that circumscription does not hold in higher dimension. (§2.9.2.5) \square

2.3.2.0.2 Example. *Nuclear norm ball: convex hull of rank-1 matrices.* From (84), in Example 2.3.2.0.1, we learn that the convex hull of normalized symmetric rank-1 matrices is a slice of the positive semidefinite cone. In §2.9.2.4 we find the convex hull of all symmetric rank-1 matrices to be the entire positive semidefinite cone.




Figure 20: Two Fantopes. Circle, (radius $\frac{1}{\sqrt{2}}$) shown here on boundary of positive semidefinite cone \mathbb{S}_+^2 in isometrically isomorphic \mathbb{R}^3 from Figure 39, comprises boundary of a Fantope (83) in this dimension ($k=1$, $N=2$). Lone point illustrated is identity matrix I and is that Fantope corresponding to $k=2$, $N=2$. (View is from inside PSD cone looking toward origin.)

In the present example we abandon symmetry; instead posing, what is the convex hull of bounded nonsymmetric rank-1 matrices:

$$\text{conv}\{uv^T \mid \|uv^T\| \leq 1, u \in \mathbb{R}^m, v \in \mathbb{R}^n\} = \{X \in \mathbb{R}^{m \times n} \mid \sum_i \sigma(X)_i \leq 1\} \quad (87)$$

where $\sigma(X)$ is a vector of singular values. (Since $\|uv^T\| = \|u\|\|v\|$ (1508), norm of each vector constituting a *dyad* uv^T (§B.1) in the hull is effectively bounded above by 1.)

Proof. (\Leftarrow) Suppose $\sum \sigma(X)_i \leq 1$. As in §A.6, define a singular value decomposition: $X = U\Sigma V^T$ where $U = [u_1 \dots u_{\min\{m,n\}}] \in \mathbb{R}^{m \times \min\{m,n\}}$, $V = [v_1 \dots v_{\min\{m,n\}}] \in \mathbb{R}^{n \times \min\{m,n\}}$, and whose sum of singular values is $\sum \sigma(X)_i = \text{tr } \Sigma = \kappa \leq 1$. Then we may write $X = \sum \frac{\sigma_i}{\kappa} \sqrt{\kappa} u_i \sqrt{\kappa} v_i^T$ which is a convex combination of dyads each of whose norm does not exceed 1. (Srebro)

(\Rightarrow) Now suppose we are given a convex combination of dyads $X = \sum \alpha_i u_i v_i^T$ such that $\sum \alpha_i = 1$, $\alpha_i \geq 0 \forall i$, and $\|u_i v_i^T\| \leq 1 \forall i$. Then by triangle inequality for singular values [181, cor.3.4.3] $\sum \sigma(X)_i \leq \sum \sigma(\alpha_i u_i v_i^T) = \sum \alpha_i \|u_i v_i^T\| \leq \sum \alpha_i$. \blacklozenge

Given any particular dyad uv_p^T in the convex hull, because its polar $-uv_p^T$ and every convex combination of the two belong to that hull, then the unique line containing those two points $\pm uv_p^T$ (their affine combination (71)) must intersect the hull's boundary at the normalized dyads $\{\pm uv^T \mid \|uv^T\| = 1\}$. Any point formed by convex combination of dyads in the hull must therefore be expressible as a convex combination of dyads on the boundary: Figure 21.

$$\text{conv}\{uv^T \mid \|uv^T\| \leq 1, u \in \mathbb{R}^m, v \in \mathbb{R}^n\} \equiv \text{conv}\{uv^T \mid \|uv^T\| = 1, u \in \mathbb{R}^m, v \in \mathbb{R}^n\} \quad (88)$$

id est, dyads may be normalized and the hull's boundary contains them;

$$\partial\{X \in \mathbb{R}^{m \times n} \mid \sum_i \sigma(X)_i \leq 1\} \supseteq \{uv^T \mid \|uv^T\| = 1, u \in \mathbb{R}^m, v \in \mathbb{R}^n\} \quad (89)$$

Normalized dyads constitute the set of extreme points (§2.6.0.0.1) of this nuclear norm ball which is, therefore, their convex hull. \square

2.3.2.0.3 Exercise. Convex hull of factorized matrices.

Find the convex hull of nonorthogonal projection matrices (§E.1.1):

$$\{UV^T \mid U \in \mathbb{R}^{N \times k}, V \in \mathbb{R}^{N \times k}, V^T U = I\} \quad (90)$$




Figure 21: uv_p^T is a convex combination of normalized dyads $\|\pm uv^T\|=1$; similarly for xy_p^T . Any point in line segment joining xy_p^T to uv_p^T is expressible as a convex combination of two to four points indicated on boundary. Affine dimension of vectorized nuclear norm ball in $X \in \mathbb{R}^{2 \times 2}$ is 2 because (1508) $\sigma_1 = \|u\| \|v\| = 1$ by (88).

Find the convex hull of nonsymmetric matrices bounded under some norm:

$$\{UV^T \mid U \in \mathbb{R}^{m \times k}, V \in \mathbb{R}^{n \times k}, \|UV^T\| \leq 1\} \quad (91)$$



2.3.2.0.4 Example. *Convex hull of permutation matrices.* [179] [281] [231] A permutation matrix Ξ is formed by interchanging rows and columns of identity matrix I . Since Ξ is square and $\Xi^T \Xi = I$, the set of all permutation matrices is a proper subset of the nonconvex manifold of orthogonal matrices (§B.5). In fact, the only orthogonal matrices having all nonnegative entries are permutations of the identity.

Regarding the permutation matrices as a set of points in Euclidean space, its convex hull is a bounded polyhedron (§2.12) described (Birkhoff, 1946)

$$\text{conv}\{\Xi = \Pi_i(I \in \mathbb{S}^n) \in \mathbb{R}^{n \times n}, i=1 \dots n!\} = \{X \in \mathbb{R}^{n \times n} \mid X^T \mathbf{1} = \mathbf{1}, X \mathbf{1} = \mathbf{1}, X \geq \mathbf{0}\} \quad (92)$$

where Π_i is a linear operator representing the i^{th} permutation. This polyhedral hull, whose $n!$ vertices are the permutation matrices, is known as the set of *doubly stochastic matrices*. The permutation matrices are the minimum cardinality (fewest nonzero entries) doubly stochastic matrices. The only orthogonal matrices belonging to this polyhedron are the permutation matrices.

It is remarkable that $n!$ permutation matrices can be described as the extreme points (§2.6.0.0.1) of a bounded polyhedron, of affine dimension $(n-1)^2$, that is itself described by only $2n$ equalities ($2n-1$ linearly independent equality constraints in nonnegative variables). By *Carathéodory's theorem*, conversely, any doubly stochastic matrix can be described as a convex combination of at most $(n-1)^2+1$ permutation matrices. [180, §8.7] This polyhedron, then, can be a device for relaxing an integer, combinatorial, or Boolean optimization problem.^{2.15} [58] [249, §3.1]

□

2.3.2.0.5 Example. *Convex hull of orthonormal matrices.* [26, §1.2] Consider rank- k matrices $U \in \mathbb{R}^{n \times k}$ such that $U^T U = I$. These are the orthonormal matrices; a closed bounded submanifold, of all orthogonal

^{2.15}Dantzig first showed in 1951 that, by this device, the so-called *assignment problem* can be formulated as a linear program. [280] [25, §II.5]

matrices, having dimension $nk - \frac{1}{2}k(k+1)$ [45]. Their convex hull is expressed, for $1 \leq k \leq n$

$$\begin{aligned} \text{conv}\{U \in \mathbb{R}^{n \times k} \mid U^T U = I\} &= \{X \in \mathbb{R}^{n \times k} \mid \|X\|_2 \leq 1\} \\ &= \{X \in \mathbb{R}^{n \times k} \mid \|X^T a\| \leq \|a\| \quad \forall a \in \mathbb{R}^n\} \end{aligned} \quad (93)$$

When $k=n$, matrices U are orthogonal and the convex hull is called the *spectral norm ball* which is the set of all *contractions*. [181, p.158] [292, p.313] The orthogonal matrices then constitute the extreme points (§2.6.0.0.1) of this hull.

By Schur complement (§A.4), the spectral norm $\|X\|_2$ constraining largest singular value $\sigma_1(X)$ can be expressed as a semidefinite constraint

$$\begin{bmatrix} I & X \\ X^T & I \end{bmatrix} \succeq 0 \quad (94)$$

because of equivalence $X^T X \preceq I \Leftrightarrow \sigma(X) \leq \mathbf{1}$ with singular values. (1465) (1363) (1364) \square

2.3.3 Conic hull

In terms of a finite-length point list (or set) arranged columnar in $X \in \mathbb{R}^{n \times N}$ (69), its conic hull is expressed

$$\mathcal{K} \triangleq \text{cone}\{x_\ell, \ell=1 \dots N\} = \text{cone } X = \{Xa \mid a \succeq 0\} \subseteq \mathbb{R}^n \quad (95)$$

id est, every nonnegative combination of points from the list. The conic hull of any finite-length list forms a *polyhedral cone* [177, §A.4.3] (§2.12.1.0.1; e.g., Figure 22); the smallest closed convex cone that contains the list.

By convention, the aberration [293, §2.1]

$$\text{cone } \emptyset \triangleq \{\mathbf{0}\} \quad (96)$$

Given some arbitrary set \mathcal{C} , it is apparent

$$\text{conv } \mathcal{C} \subseteq \text{cone } \mathcal{C} \quad (97)$$




Figure 22: A simplicial cone (§2.12.3.1.1) in \mathbb{R}^3 whose boundary is drawn truncated; constructed using $A \in \mathbb{R}^{3 \times 3}$ and $C = \mathbf{0}$ in (260). By the most fundamental definition of a cone (§2.7.1), entire boundary can be constructed from an aggregate of rays emanating exclusively from the origin. Each of three extreme directions corresponds to an edge (§2.6.0.0.3); they are conically, affinely, and linearly independent for this cone. Because this set is polyhedral, exposed directions are in one-to-one correspondence with extreme directions; there are only three. Its extreme directions give rise to what is called a *vertex-description* of this polyhedral cone; simply, the conic hull of extreme directions. Obviously this cone can also be constructed by intersection of three halfspaces; hence the equivalent *halfspace-description*.

2.3.4 Vertex-description

The conditions in (71), (79), and (95) respectively define an *affine combination*, *convex combination*, and *conic combination* of elements from the set or list. Whenever a Euclidean body can be described as some hull or span of a set of points, then that representation is loosely called a *vertex-description*.

2.4 Halfspace, Hyperplane

A two-dimensional affine subset is called a *plane*. An $(n-1)$ -dimensional affine subset in \mathbb{R}^n is called a *hyperplane*. [271] [177] Every hyperplane partially bounds a *halfspace* (which is convex but not affine).

2.4.1 Halfspaces \mathcal{H}_+ and \mathcal{H}_-

Euclidean space \mathbb{R}^n is partitioned in two by any hyperplane $\partial\mathcal{H}$; *id est*, $\mathcal{H}_- + \mathcal{H}_+ = \mathbb{R}^n$. The resulting (closed convex) halfspaces, both partially bounded by $\partial\mathcal{H}$, may be described as an asymmetry

$$\mathcal{H}_- = \{y \mid a^T y \leq b\} = \{y \mid a^T(y - y_p) \leq 0\} \subset \mathbb{R}^n \quad (98)$$

$$\mathcal{H}_+ = \{y \mid a^T y \geq b\} = \{y \mid a^T(y - y_p) \geq 0\} \subset \mathbb{R}^n \quad (99)$$

where nonzero vector $a \in \mathbb{R}^n$ is an *outward-normal* to the hyperplane partially bounding \mathcal{H}_- while an *inward-normal* with respect to \mathcal{H}_+ . For any vector $y - y_p$ that makes an obtuse angle with normal a , vector y will lie in the halfspace \mathcal{H}_- on one side (shaded in Figure 23) of the hyperplane while acute angles denote y in \mathcal{H}_+ on the other side.

An equivalent more intuitive representation of a halfspace comes about when we consider all the points in \mathbb{R}^n closer to point d than to point c or equidistant, in the Euclidean sense; from Figure 23,

$$\mathcal{H}_- = \{y \mid \|y - d\| \leq \|y - c\|\} \quad (100)$$

This representation, in terms of proximity, is resolved with the more conventional representation of a halfspace (98) by squaring both sides of the inequality in (100);

$$\mathcal{H}_- = \left\{ y \mid (c - d)^T y \leq \frac{\|c\|^2 - \|d\|^2}{2} \right\} = \left\{ y \mid (c - d)^T \left(y - \frac{c + d}{2} \right) \leq 0 \right\} \quad (101)$$




Figure 23: Hyperplane illustrated $\partial\mathcal{H}$ is a line partially bounding halfspaces $\mathcal{H}_- = \{y \mid a^T(y - y_p) \leq 0\}$ and $\mathcal{H}_+ = \{y \mid a^T(y - y_p) \geq 0\}$ in \mathbb{R}^2 . Shaded is a rectangular piece of semi-infinite \mathcal{H}_- with respect to which vector a is outward-normal to bounding hyperplane; vector a is inward-normal with respect to \mathcal{H}_+ . Halfspace \mathcal{H}_- contains nullspace $\mathcal{N}(a^T)$ (dashed line through origin) because $a^T y_p > 0$. Hyperplane, halfspace, and nullspace are each drawn truncated. Points c and d are equidistant from hyperplane, and vector $c - d$ is normal to it. Δ is distance from origin to hyperplane.

2.4.1.1 PRINCIPLE 1: Halfspace-description of convex sets

The most fundamental principle in convex geometry follows from the *geometric Hahn-Banach theorem* [220, §5.12] [17, §1] [115, §I.1.2] which guarantees any closed convex set to be an intersection of halfspaces.

2.4.1.1.1 Theorem. *Halfspaces.* [177, §A.4.2(b)] [37, §2.4]

A closed convex set in \mathbb{R}^n is equivalent to the intersection of all halfspaces that contain it. \diamond

Intersection of multiple halfspaces in \mathbb{R}^n may be represented using a matrix constant A

$$\bigcap_i \mathcal{H}_{i-} = \{y \mid A^T y \preceq b\} = \{y \mid A^T(y - y_p) \preceq 0\} \quad (102)$$

$$\bigcap_i \mathcal{H}_{i+} = \{y \mid A^T y \succeq b\} = \{y \mid A^T(y - y_p) \succeq 0\} \quad (103)$$

where b is now a vector, and the i^{th} column of A is normal to a hyperplane $\partial\mathcal{H}_i$ partially bounding \mathcal{H}_i . By the *halfspaces theorem*, intersections like this can describe interesting convex Euclidean bodies such as polyhedra and cones, giving rise to the term *halfspace-description*.

2.4.2 Hyperplane $\partial\mathcal{H}$ representations

Every hyperplane $\partial\mathcal{H}$ is an affine set parallel to an $(n-1)$ -dimensional subspace of \mathbb{R}^n ; it is itself a subspace if and only if it contains the origin.

$$\dim \partial\mathcal{H} = n - 1 \quad (104)$$

so a hyperplane is a point in \mathbb{R} , a line in \mathbb{R}^2 , a plane in \mathbb{R}^3 , and so on. Every hyperplane can be described as the intersection of complementary halfspaces; [271, §19]

$$\partial\mathcal{H} = \mathcal{H}_- \cap \mathcal{H}_+ = \{y \mid a^T y \leq b, a^T y \geq b\} = \{y \mid a^T y = b\} \quad (105)$$

a halfspace-description. Assuming *normal* $a \in \mathbb{R}^n$ to be nonzero, then any hyperplane in \mathbb{R}^n can be described as the solution set to vector equation $a^T y = b$ (illustrated in Figure 23 and Figure 24 for \mathbb{R}^2);

$$\partial\mathcal{H} \triangleq \{y \mid a^T y = b\} = \{y \mid a^T(y - y_p) = 0\} = \{Z\xi + y_p \mid \xi \in \mathbb{R}^{n-1}\} \subset \mathbb{R}^n \quad (106)$$




Figure 24: **(a)-(d)** Hyperplanes in \mathbb{R}^2 (truncated). Movement in normal direction increases vector inner-product. This visual concept is exploited to attain analytical solution of linear programs; *e.g.*, Example 2.4.2.6.2, Exercise 2.5.1.2.2, Example 3.1.6.0.2, [53, exer.4.8-exer.4.20]. Each graph is also interpretable as a contour plot of a real affine function of two variables as in Figure 66. **(e)** Ratio $|\beta|/\|\alpha\|$ from $\{x \mid \alpha^T x = \beta\}$ represents radius of hypersphere about $\mathbf{0}$ supported by hyperplane whose normal is α .

All solutions y constituting the hyperplane are *offset* from the nullspace of a^T by the same constant vector $y_p \in \mathbb{R}^n$ that is any particular solution to $a^T y = b$; *id est*,

$$y = Z\xi + y_p \quad (107)$$

where the columns of $Z \in \mathbb{R}^{n \times n-1}$ constitute a basis for the nullspace $\mathcal{N}(a^T) = \{x \in \mathbb{R}^n \mid a^T x = \mathbf{0}\}$. ^{2.16}

Conversely, given any point y_p in \mathbb{R}^n , the unique hyperplane containing it having normal a is the affine set $\partial\mathcal{H}$ (106) where b equals $a^T y_p$ and where a basis for $\mathcal{N}(a^T)$ is arranged in Z columnar. Hyperplane dimension is apparent from the dimensions of Z ; that hyperplane is parallel to the span of its columns.

2.4.2.0.1 Exercise. *Hyperplane scaling.*

Given normal y , draw a hyperplane $\{x \in \mathbb{R}^2 \mid x^T y = 1\}$. Suppose $z = \frac{1}{2}y$. On the same plot, draw the hyperplane $\{x \in \mathbb{R}^2 \mid x^T z = 1\}$. Now suppose $z = 2y$, then draw the last hyperplane again with this new z . What is the apparent effect of scaling normal y ? \blacktriangledown

2.4.2.0.2 Example. *Distance from origin to hyperplane.*

Given the (shortest) distance $\Delta \in \mathbb{R}_+$ from the origin to a hyperplane having normal vector a , we can find its representation $\partial\mathcal{H}$ by dropping a perpendicular. The point thus found is the orthogonal projection of the origin on $\partial\mathcal{H}$ (§E.5.0.0.5), equal to $a\Delta/\|a\|$ if the origin is known *a priori* to belong to halfspace \mathcal{H}_- (Figure 23), or equal to $-a\Delta/\|a\|$ if the origin belongs to halfspace \mathcal{H}_+ ; *id est*, when $\mathcal{H}_- \ni \mathbf{0}$

$$\partial\mathcal{H} = \{y \mid a^T(y - a\Delta/\|a\|) = 0\} = \{y \mid a^T y = \|a\|\Delta\} \quad (108)$$

or when $\mathcal{H}_+ \ni \mathbf{0}$

$$\partial\mathcal{H} = \{y \mid a^T(y + a\Delta/\|a\|) = 0\} = \{y \mid a^T y = -\|a\|\Delta\} \quad (109)$$

Knowledge of only distance Δ and normal a thus introduces ambiguity into the hyperplane representation. \square

^{2.16}We will later find this expression for y in terms of nullspace of a^T (more generally, of matrix A^T (135)) to be a useful device for eliminating affine equality constraints, much as we did here.

2.4.2.1 Matrix variable

Any halfspace in \mathbb{R}^{mn} may be represented using a matrix variable. For variable $Y \in \mathbb{R}^{m \times n}$, given constants $A \in \mathbb{R}^{m \times n}$ and $b = \langle A, Y_p \rangle \in \mathbb{R}$

$$\mathcal{H}_- = \{Y \in \mathbb{R}^{mn} \mid \langle A, Y \rangle \leq b\} = \{Y \in \mathbb{R}^{mn} \mid \langle A, Y - Y_p \rangle \leq 0\} \quad (110)$$

$$\mathcal{H}_+ = \{Y \in \mathbb{R}^{mn} \mid \langle A, Y \rangle \geq b\} = \{Y \in \mathbb{R}^{mn} \mid \langle A, Y - Y_p \rangle \geq 0\} \quad (111)$$

Recall vector inner-product from §2.2, $\langle A, Y \rangle = \text{tr}(A^T Y)$.

Hyperplanes in \mathbb{R}^{mn} may, of course, also be represented using matrix variables.

$$\partial\mathcal{H} = \{Y \mid \langle A, Y \rangle = b\} = \{Y \mid \langle A, Y - Y_p \rangle = 0\} \subset \mathbb{R}^{mn} \quad (112)$$

Vector a from Figure 23 is normal to the hyperplane illustrated. Likewise, nonzero vectorized matrix A is normal to hyperplane $\partial\mathcal{H}$;

$$A \perp \partial\mathcal{H} \text{ in } \mathbb{R}^{mn} \quad (113)$$

2.4.2.2 Vertex-description of hyperplane

Any hyperplane in \mathbb{R}^n may be described as the affine hull of a minimal set of points $\{x_\ell \in \mathbb{R}^n, \ell = 1 \dots n\}$ arranged columnar in a matrix $X \in \mathbb{R}^{n \times n}$ (69):

$$\begin{aligned} \partial\mathcal{H} &= \text{aff}\{x_\ell \in \mathbb{R}^n, \ell = 1 \dots n\}, & \dim \text{aff}\{x_\ell \forall \ell\} &= n-1 \\ &= \text{aff } X, & \dim \text{aff } X &= n-1 \\ &= x_1 + \mathcal{R}\{x_\ell - x_1, \ell = 2 \dots n\}, & \dim \mathcal{R}\{x_\ell - x_1, \ell = 2 \dots n\} &= n-1 \\ &= x_1 + \mathcal{R}(X - x_1 \mathbf{1}^T), & \dim \mathcal{R}(X - x_1 \mathbf{1}^T) &= n-1 \end{aligned} \quad (114)$$

where

$$\mathcal{R}(A) = \{Ax \mid \forall x\} \quad (133)$$

2.4.2.3 Affine independence, minimal set

For any particular affine set, a *minimal set* of points constituting its vertex-description is an affinely independent generating set and *vice versa*.

Arbitrary given points $\{x_i \in \mathbb{R}^n, i = 1 \dots N\}$ are *affinely independent* (a.i.) if and only if, over all $\zeta \in \mathbb{R}^N \ni \zeta^T \mathbf{1} = 1, \zeta_k = 0$ (confer §2.1.2)

$$x_i \zeta_i + \dots + x_j \zeta_j - x_k = \mathbf{0}, \quad i \neq \dots \neq j \neq k = 1 \dots N \quad (115)$$




Figure 25: Any one particular point of three points illustrated does not belong to affine hull \mathcal{A}_i ($i \in 1, 2, 3$, each drawn truncated) of points remaining. Three corresponding vectors in \mathbb{R}^2 are, therefore, affinely independent (but neither linearly or conically independent).

has no solution ζ ; in words, iff no point from the given set can be expressed as an affine combination of those remaining. We deduce

$$\text{l.i.} \Rightarrow \text{a.i.} \quad (116)$$

Consequently, $\{x_i, i = 1 \dots N\}$ is an affinely independent set if and only if $\{x_i - x_1, i = 2 \dots N\}$ is a linearly independent (l.i.) set. [183, §3] (Figure 25)

This is equivalent to the property that the columns of $\begin{bmatrix} X \\ \mathbf{1}^T \end{bmatrix}$ (for $X \in \mathbb{R}^{n \times N}$ as in (69)) form a linearly independent set. [177, §A.1.3]

2.4.2.4 Preservation of affine independence

Independence in the linear (§2.1.2.1), affine, and conic (§2.10.1) senses can be preserved under linear transformation. Suppose a matrix $X \in \mathbb{R}^{n \times N}$ (69) holds an affinely independent set in its columns. Consider a transformation on the domain of such matrices

$$T(X) : \mathbb{R}^{n \times N} \rightarrow \mathbb{R}^{n \times N} \triangleq XY \quad (117)$$

where fixed matrix $Y \triangleq [y_1 \ y_2 \ \dots \ y_N] \in \mathbb{R}^{N \times N}$ represents linear operator T . Affine independence of $\{Xy_i \in \mathbb{R}^n, i = 1 \dots N\}$ demands (by definition (115))

there exist no solution $\zeta \in \mathbb{R}^N$ \ni $\zeta^T \mathbf{1} = 1$, $\zeta_k = 0$, to

$$Xy_i \zeta_i + \cdots + Xy_j \zeta_j - Xy_k = \mathbf{0}, \quad i \neq \cdots \neq j \neq k = 1 \dots N \quad (118)$$

By factoring out X , we see that is ensured by affine independence of $\{y_i \in \mathbb{R}^N\}$ and by $\mathcal{R}(Y) \cap \mathcal{N}(X) = \mathbf{0}$ where

$$\mathcal{N}(A) = \{x \mid Ax = \mathbf{0}\} \quad (134)$$

2.4.2.5 affine maps

Affine transformations preserve affine hulls. Given any affine mapping T of vector spaces and some arbitrary set \mathcal{C} [271, p.8]

$$\text{aff}(T\mathcal{C}) = T(\text{aff } \mathcal{C}) \quad (119)$$

2.4.2.6 PRINCIPLE 2: Supporting hyperplane

The second most fundamental principle of convex geometry also follows from the *geometric Hahn-Banach theorem* [220, §5.12] [17, §1] that guarantees existence of at least one hyperplane in \mathbb{R}^n supporting a convex set (having nonempty interior)^{2.17} at each point on its boundary.

The partial boundary $\partial\mathcal{H}$ of a halfspace that contains arbitrary set \mathcal{Y} is called a *supporting hyperplane* $\underline{\partial\mathcal{H}}$ to \mathcal{Y} when the hyperplane contains at least one point of $\overline{\mathcal{Y}}$. [271, §11]

2.4.2.6.1 Definition. Supporting hyperplane $\underline{\partial\mathcal{H}}$.

Assuming set \mathcal{Y} and some normal $a \neq \mathbf{0}$ reside in opposite halfspaces^{2.18} (Figure 27(a)), then a hyperplane supporting \mathcal{Y} at $y_p \in \partial\mathcal{Y}$ is

$$\underline{\partial\mathcal{H}}_- = \{y \mid a^T(y - y_p) = 0, \quad y_p \in \overline{\mathcal{Y}}, \quad a^T(z - y_p) \leq 0 \quad \forall z \in \overline{\mathcal{Y}}\} \quad (120)$$

Given normal a , instead, the supporting hyperplane is

$$\underline{\partial\mathcal{H}}_- = \{y \mid a^T y = \sup\{a^T z \mid z \in \mathcal{Y}\}\} \quad (121)$$

^{2.17}It is conventional to speak of a hyperplane supporting set \mathcal{C} but not containing \mathcal{C} ; called *nontrivial support*. [271, p.100] Hyperplanes in support of lower-dimensional bodies are admitted.

^{2.18}Normal a belongs to \mathcal{H}_+ by definition.




Figure 26: Each shaded line segment $\{z \in \mathcal{C} \mid a^T z = \kappa_i\}$ belonging to set $\mathcal{C} \subset \mathbb{R}^2$ shows intersection with hyperplane parametrized by scalar κ_i ; each shows a (linear) contour in vector z of equal inner product with normal vector a . Cartesian axes drawn for reference. (*confer* Figure 66)




Figure 27: (a) Hyperplane $\underline{\partial\mathcal{H}}_-$ (120) supporting closed set $\mathcal{Y} \in \mathbb{R}^2$. Vector a is inward-normal to hyperplane with respect to halfspace \mathcal{H}_+ , but outward-normal with respect to set \mathcal{Y} . A supporting hyperplane can be considered the limit of an increasing sequence in the normal-direction like that in Figure 26. (b) Hyperplane $\underline{\partial\mathcal{H}}_+$ nontraditionally supporting \mathcal{Y} . Vector \tilde{a} is inward-normal to hyperplane now with respect to both halfspace \mathcal{H}_+ and set \mathcal{Y} . Tradition [177] [271] recognizes only positive normal polarity in support function $\sigma_{\mathcal{Y}}$ as in (121); *id est*, normal a , figure (a). But both interpretations of supporting hyperplane are useful.

where real function

$$\sigma_{\mathcal{Y}}(a) \triangleq \sup\{a^T z \mid z \in \mathcal{Y}\} \quad (499)$$

is called the *support function* for \mathcal{Y} .

An equivalent but nontraditional representation^{2.19} for a supporting hyperplane is obtained by reversing polarity of normal a ; (1572)

$$\begin{aligned} \underline{\partial\mathcal{H}}_+ &= \{y \mid \tilde{a}^T(y - y_p) = 0, \quad y_p \in \overline{\mathcal{Y}}, \quad \tilde{a}^T(z - y_p) \geq 0 \quad \forall z \in \overline{\mathcal{Y}}\} \\ &= \{y \mid \tilde{a}^T y = -\inf\{\tilde{a}^T z \mid z \in \mathcal{Y}\} = \sup\{-\tilde{a}^T z \mid z \in \mathcal{Y}\}\} \end{aligned} \quad (122)$$

where normal \tilde{a} and set \mathcal{Y} both now reside in \mathcal{H}_+ . (Figure 27(b))

When a supporting hyperplane contains only a single point of $\overline{\mathcal{Y}}$, that hyperplane is termed *strictly supporting*.^{2.20} \triangle

A set of nonempty interior that has a supporting hyperplane at every point on its boundary, conversely, is convex. A convex set $\mathcal{C} \subset \mathbb{R}^n$, for example, can be expressed as the intersection of all halfspaces partially bounded by hyperplanes supporting it; *videlicet*, [220, p.135]

$$\overline{\mathcal{C}} = \bigcap_{a \in \mathbb{R}^n} \{y \mid a^T y \leq \sigma_{\mathcal{C}}(a)\} \quad (123)$$

by the *halfspaces theorem* (§2.4.1.1.1).

There is no geometric difference between supporting hyperplane $\underline{\partial\mathcal{H}}_+$ or $\underline{\partial\mathcal{H}}_-$ or $\underline{\partial\mathcal{H}}$ and^{2.21} an ordinary hyperplane $\partial\mathcal{H}$ coincident with them.

2.4.2.6.2 Example. Minimization on a hypercube.

Consider minimization of a linear function on a *hypercube*, given vector c

$$\begin{aligned} &\underset{x}{\text{minimize}} && c^T x \\ &\text{subject to} && -\mathbf{1} \preceq x \preceq \mathbf{1} \end{aligned} \quad (124)$$

^{2.19} useful for constructing the *dual cone*; e.g., Figure 51(b). Tradition would instead have us construct the *polar cone*; which is, the negative dual cone.

^{2.20}Rockafellar terms a strictly supporting hyperplane *tangent* to \mathcal{Y} if it is unique there; [271, §18, p.169] a definition we do not adopt because our only criterion for tangency is intersection exclusively with a relative boundary. Hiriart-Urruty & Lemaréchal [177, p.44] (*confer* [271, p.100]) do not demand any tangency of a supporting hyperplane.

^{2.21}If vector-normal polarity is unimportant, we may instead signify a supporting hyperplane by $\underline{\partial\mathcal{H}}$.

This convex optimization problem is called a *linear program*^{2.22} because the objective of minimization is linear and the constraints describe a polyhedron (intersection of a finite number of halfspaces and hyperplanes). Applying graphical concepts from Figure 24, Figure 26, and Figure 27, an *optimal solution* can be shown to be $x^* = -\text{sgn}(c)$ but is not necessarily unique. Because an optimal solution always exists at a hypercube vertex (§2.6.1.0.1) regardless of the value of nonzero vector c [82], mathematicians see this geometry as a means to *relax* a discrete problem (whose desired solution is integer or combinatorial, *confer* Example 4.2.3.1.1). [211, §3.1] [212] □

2.4.2.6.3 Exercise. Unbounded below.

Suppose instead we minimize over the unit hypersphere in Example 2.4.2.6.2; $\|x\| \leq 1$. What is an expression for optimal solution now? Is that program still linear?

Now suppose minimization of absolute value in (124). Are the following programs equivalent for some arbitrary real convex set \mathcal{C} ? (*confer* (467))

$$\begin{array}{ll} \underset{x \in \mathbb{R}}{\text{minimize}} & |x| \\ \text{subject to} & -1 \leq x \leq 1 \\ & x \in \mathcal{C} \end{array} \quad \equiv \quad \begin{array}{ll} \underset{\alpha, \beta}{\text{minimize}} & \alpha + \beta \\ \text{subject to} & 1 \geq \beta \geq 0 \\ & 1 \geq \alpha \geq 0 \\ & \alpha - \beta \in \mathcal{C} \end{array} \quad (125)$$

Many optimization problems of interest and some methods of solution require nonnegative variables. The method illustrated below splits a variable into parts; $x = \alpha - \beta$ (extensible to vectors). Under what conditions on vector a and scalar b is an optimal solution x^* negative infinity?

$$\begin{array}{ll} \underset{\alpha \in \mathbb{R}, \beta \in \mathbb{R}}{\text{minimize}} & \alpha - \beta \\ \text{subject to} & \beta \geq 0 \\ & \alpha \geq 0 \\ & a^T \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = b \end{array} \quad (126)$$

Minimization of the *objective function*^{2.23} entails maximization of β . ▼

^{2.22}The term *program* has its roots in economics. It was originally meant with regard to a plan or to efficient organization of some industrial process. [82, §2]

^{2.23}The *objective* is the function that is argument to minimization or maximization.

2.4.2.7 PRINCIPLE 3: Separating hyperplane

The third most fundamental principle of convex geometry again follows from the *geometric Hahn-Banach theorem* [220, §5.12] [17, §1] [115, §I.1.2] that guarantees existence of a hyperplane separating two nonempty convex sets in \mathbb{R}^n whose relative interiors are nonintersecting. *Separation* intuitively means each set belongs to a halfspace on an opposing side of the hyperplane. There are two cases of interest:

- 1) If the two sets intersect only at their relative boundaries (§2.6.1.3), then there exists a separating hyperplane $\partial\mathcal{H}$ containing the intersection but containing no points relatively interior to either set. If at least one of the two sets is open, conversely, then the existence of a separating hyperplane implies the two sets are nonintersecting. [53, §2.5.1]
- 2) A *strictly separating hyperplane* $\partial\mathcal{H}$ intersects the closure of neither set; its existence is guaranteed when the intersection of the closures is empty and at least one set is bounded. [177, §A.4.1]

2.4.3 Angle between hyperspaces

Given halfspace-descriptions, the *dihedral* angle between hyperplanes and halfspaces is defined as the angle between their defining normals. Given normals a and b respectively describing $\partial\mathcal{H}_a$ and $\partial\mathcal{H}_b$, for example

$$\sphericalangle(\partial\mathcal{H}_a, \partial\mathcal{H}_b) \triangleq \arccos\left(\frac{\langle a, b \rangle}{\|a\| \|b\|}\right) \text{ radians} \quad (127)$$

2.5 Subspace representations

There are two common forms of expression for Euclidean subspaces, both coming from elementary linear algebra: *range form* \mathcal{R} and *nullspace form* \mathcal{N} ; a.k.a, vertex-description and halfspace-description respectively.

The fundamental vector subspaces associated with a matrix $A \in \mathbb{R}^{m \times n}$ [294, §3.1] are ordinarily related

$$\mathcal{R}(A^T) \perp \mathcal{N}(A), \quad \mathcal{N}(A^T) \perp \mathcal{R}(A) \quad (128)$$

$$\mathcal{R}(A^T) \oplus \mathcal{N}(A) = \mathbb{R}^n, \quad \mathcal{N}(A^T) \oplus \mathcal{R}(A) = \mathbb{R}^m \quad (129)$$

and of dimension:

$$\dim \mathcal{R}(A^T) = \dim \mathcal{R}(A) = \text{rank } A \leq \min\{m, n\} \quad (130)$$

with complementarity

$$\dim \mathcal{N}(A) = n - \text{rank } A, \quad \dim \mathcal{N}(A^T) = m - \text{rank } A \quad (131)$$

These equations (128)-(131) comprise the *fundamental theorem of linear algebra*. [294, p.95, p.138]

From these four fundamental subspaces, the rowspace and range identify one form of subspace description (range form or vertex-description (§2.3.4))

$$\mathcal{R}(A^T) \triangleq \text{span } A^T = \{A^T y \mid y \in \mathbb{R}^m\} = \{x \in \mathbb{R}^n \mid A^T y = x, \ y \in \mathcal{R}(A)\} \quad (132)$$

$$\mathcal{R}(A) \triangleq \text{span } A = \{Ax \mid x \in \mathbb{R}^n\} = \{y \in \mathbb{R}^m \mid Ax = y, \ x \in \mathcal{R}(A^T)\} \quad (133)$$

while the nullspaces identify the second common form (nullspace form or halfspace-description (105))

$$\mathcal{N}(A) \triangleq \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}\} \quad (134)$$

$$\mathcal{N}(A^T) \triangleq \{y \in \mathbb{R}^m \mid A^T y = \mathbf{0}\} \quad (135)$$

Range forms (132) (133) are realized as the respective span of the column vectors in matrices A^T and A , whereas nullspace form (134) or (135) is the solution set to a linear equation similar to hyperplane definition (106). Yet because matrix A generally has multiple rows, halfspace-description $\mathcal{N}(A)$ is actually the intersection of as many hyperplanes through the origin; for (134), each row of A is normal to a hyperplane while each row of A^T is a normal for (135).

2.5.1 Subspace or affine subset ...

Any particular vector subspace \mathcal{R}_p can be described as $\mathcal{N}(A)$ the nullspace of some matrix A or as $\mathcal{R}(B)$ the range of some matrix B .

More generally, we have the choice of expressing an $n - m$ -dimensional affine subset in \mathbb{R}^n as the intersection of m hyperplanes, or as the offset span of $n - m$ vectors:

2.5.1.1 ... as hyperplane intersection

Any affine subset \mathcal{A} of dimension $n - m$ can be described as an intersection of m hyperplanes in \mathbb{R}^n ; given *fat* ($m \leq n$) *full-rank* ($\text{rank} = \min\{m, n\}$) matrix

$$A \triangleq \begin{bmatrix} a_1^T \\ \vdots \\ a_m^T \end{bmatrix} \in \mathbb{R}^{m \times n} \quad (136)$$

and vector $b \in \mathbb{R}^m$,

$$\mathcal{A} \triangleq \{x \in \mathbb{R}^n \mid Ax = b\} = \bigcap_{i=1}^m \{x \mid a_i^T x = b_i\} \quad (137)$$

a halfspace-description. (105)

For example: The intersection of any two independent^{2.24} hyperplanes in \mathbb{R}^3 is a line, whereas three independent hyperplanes intersect at a point. In \mathbb{R}^4 , the intersection of two independent hyperplanes is a plane (Example 2.5.1.2.1), whereas three hyperplanes intersect at a line, four at a point, and so on. \mathcal{A} describes a subspace whenever $b = 0$ in (137).

For $n > k$

$$\mathcal{A} \cap \mathbb{R}^k = \{x \in \mathbb{R}^n \mid Ax = b\} \cap \mathbb{R}^k = \bigcap_{i=1}^m \{x \in \mathbb{R}^k \mid a_i(1:k)^T x = b_i\} \quad (138)$$

The result in §2.4.2.2 is extensible; *id est*, any affine subset \mathcal{A} also has a vertex-description:

2.5.1.2 ... as span of nullspace basis

Alternatively, we may compute a basis for nullspace of matrix A in (136) and then equivalently express the affine subset \mathcal{A} as its range plus an offset: Define

$$Z \triangleq \text{basis } \mathcal{N}(A) \in \mathbb{R}^{n \times n-m} \quad (139)$$

so $AZ = \mathbf{0}$. Then we have the vertex-description in Z ,

$$\mathcal{A} = \{x \in \mathbb{R}^n \mid Ax = b\} = \{Z\xi + x_p \mid \xi \in \mathbb{R}^{n-m}\} \subseteq \mathbb{R}^n \quad (140)$$

the offset span of $n - m$ column vectors, where x_p is any particular solution to $Ax = b$. For example, \mathcal{A} describes a subspace whenever $x_p = 0$.

^{2.24}Hyperplanes are said to be *independent* iff the normals defining them are linearly independent.

2.5.1.2.1 Example. *Intersecting planes in 4-space.*

Two planes can intersect at a point in four-dimensional Euclidean vector space. It is easy to visualize intersection of two planes in three dimensions; a line can be formed. In four dimensions it is harder to visualize. So let's resort to the tools acquired.

Suppose an intersection of two hyperplanes in four dimensions is specified by a fat full-rank matrix $A_1 \in \mathbb{R}^{2 \times 4}$ ($m = 2, n = 4$) as in (137):

$$\mathcal{A}_1 \triangleq \left\{ x \in \mathbb{R}^4 \mid \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \end{bmatrix} x = b_1 \right\} \quad (141)$$

The nullspace of A_1 is two dimensional (from Z in (140)), so \mathcal{A}_1 represents a plane in four dimensions. Similarly define a second plane in terms of $A_2 \in \mathbb{R}^{2 \times 4}$:

$$\mathcal{A}_2 \triangleq \left\{ x \in \mathbb{R}^4 \mid \begin{bmatrix} a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix} x = b_2 \right\} \quad (142)$$

If the two planes are independent (meaning any line in one is linearly independent of any line from the other), they will intersect at a point because then $\begin{bmatrix} A_1 \\ A_2 \end{bmatrix}$ is invertible;

$$\mathcal{A}_1 \cap \mathcal{A}_2 = \left\{ x \in \mathbb{R}^4 \mid \begin{bmatrix} A_1 \\ A_2 \end{bmatrix} x = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} \right\} \quad (143)$$

□

2.5.1.2.2 Exercise. *Linear program.*

Minimize a hyperplane over affine set \mathcal{A} in the nonnegative orthant

$$\begin{aligned} & \underset{x}{\text{minimize}} && c^T x \\ & \text{subject to} && Ax = b \\ & && x \succeq 0 \end{aligned} \quad (144)$$

where $\mathcal{A} = \{x \mid Ax = b\}$. Two cases of interest are drawn in Figure 28. Graphically illustrate and explain optimal solutions indicated in the caption. Why is α^* negative in both cases? Is there solution on the vertical axis? ▼




Figure 28: Minimizing hyperplane over affine set \mathcal{A} in nonnegative orthant \mathbb{R}_+^2 . (a) Optimal solution is \bullet . (b) Optimal objective $\alpha^* = -\infty$.

2.5.2 Intersection of subspaces

The intersection of nullspaces associated with two matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{k \times n}$ can be expressed most simply as

$$\mathcal{N}(A) \cap \mathcal{N}(B) = \mathcal{N}\left(\begin{bmatrix} A \\ B \end{bmatrix}\right) \triangleq \{x \in \mathbb{R}^n \mid \begin{bmatrix} A \\ B \end{bmatrix} x = \mathbf{0}\} \quad (145)$$

the nullspace of their rowwise concatenation.

Suppose the columns of a matrix Z constitute a basis for $\mathcal{N}(A)$ while the columns of a matrix W constitute a basis for $\mathcal{N}(BZ)$. Then [139, §12.4.2]

$$\mathcal{N}(A) \cap \mathcal{N}(B) = \mathcal{R}(ZW) \quad (146)$$

If each basis is orthonormal, then the columns of ZW constitute an orthonormal basis for the intersection.

In the particular circumstance A and B are each positive semidefinite [20, §6], or in the circumstance A and B are two linearly independent dyads (§B.1.1), then

$$\mathcal{N}(A) \cap \mathcal{N}(B) = \mathcal{N}(A + B), \quad \begin{cases} A, B \in \mathbb{S}_+^M \\ \text{or} \\ A + B = u_1 v_1^T + u_2 v_2^T \quad (\text{l.i.}) \end{cases} \quad (147)$$

2.6 Extreme, Exposed

2.6.0.0.1 Definition. *Extreme point.*

An extreme point x_ε of a convex set \mathcal{C} is a point, belonging to its closure $\bar{\mathcal{C}}$ [37, §3.3], that is not expressible as a convex combination of points in $\bar{\mathcal{C}}$ distinct from x_ε ; *id est*, for $x_\varepsilon \in \bar{\mathcal{C}}$ and all $x_1, x_2 \in \bar{\mathcal{C}} \setminus x_\varepsilon$

$$\mu x_1 + (1 - \mu)x_2 \neq x_\varepsilon, \quad \mu \in [0, 1] \quad (148)$$

△

In other words, x_ε is an extreme point of \mathcal{C} if and only if x_ε is not a point relatively interior to any line segment in $\bar{\mathcal{C}}$. [318, §2.10]

Borwein & Lewis offer: [48, §4.1.6] An extreme point of a convex set \mathcal{C} is a point x_ε in $\bar{\mathcal{C}}$ whose *relative complement* $\bar{\mathcal{C}} \setminus x_\varepsilon$ is convex.

The set consisting of a single point $\mathcal{C} = \{x_\varepsilon\}$ is itself an extreme point.

2.6.0.0.2 Theorem. *Extreme existence.* [271, §18.5.3] [25, §II.3.5]
 A nonempty closed convex set containing no lines has at least one extreme point. \diamond

2.6.0.0.3 Definition. *Face, edge.* [177, §A.2.3]

- A *face* \mathcal{F} of convex set \mathcal{C} is a convex subset $\mathcal{F} \subseteq \bar{\mathcal{C}}$ such that every closed line segment $\overline{x_1x_2}$ in $\bar{\mathcal{C}}$, having a relatively interior point ($x \in \text{rel int } \overline{x_1x_2}$) in \mathcal{F} , has both endpoints in \mathcal{F} . The zero-dimensional faces of \mathcal{C} constitute its extreme points. The empty set and $\bar{\mathcal{C}}$ itself are conventional faces of \mathcal{C} . [271, §18]
- All faces \mathcal{F} are extreme sets by definition; *id est*, for $\mathcal{F} \subseteq \bar{\mathcal{C}}$ and all $x_1, x_2 \in \bar{\mathcal{C}} \setminus \mathcal{F}$

$$\mu x_1 + (1 - \mu)x_2 \notin \mathcal{F}, \quad \mu \in [0, 1] \quad (149)$$
- A one-dimensional face of a convex set is called an *edge*. \triangle

Dimension of a face is the penultimate number of affinely independent points (§2.4.2.3) belonging to it;

$$\dim \mathcal{F} = \sup_{\rho} \dim \{x_2 - x_1, x_3 - x_1, \dots, x_{\rho} - x_1 \mid x_i \in \mathcal{F}, i = 1 \dots \rho\} \quad (150)$$

The point of intersection in $\bar{\mathcal{C}}$ with a strictly supporting hyperplane identifies an extreme point, but not *vice versa*. The nonempty intersection of any supporting hyperplane with $\bar{\mathcal{C}}$ identifies a face, in general, but not *vice versa*. To acquire a converse, the concept *exposed face* requires introduction:

2.6.1 Exposure

2.6.1.0.1 Definition. *Exposed face, exposed point, vertex, facet.*
 [177, §A.2.3, A.2.4]

- \mathcal{F} is an *exposed face* of an n -dimensional convex set \mathcal{C} iff there is a supporting hyperplane $\partial\mathcal{H}$ to $\bar{\mathcal{C}}$ such that

$$\mathcal{F} = \bar{\mathcal{C}} \cap \partial\mathcal{H} \quad (151)$$

Only faces of dimension -1 through $n-1$ can be exposed by a hyperplane.




Figure 29: Closed convex set in \mathbb{R}^2 . Point A is exposed hence extreme; a classical vertex. Point B is extreme but not an exposed point. Point C is exposed and extreme; zero-dimensional exposure makes it a vertex. Point D is neither an exposed or extreme point although it belongs to a one-dimensional exposed face. [177, §A.2.4] [293, §3.6] Closed face \overline{AB} is exposed; a facet. The arc is not a conventional face, yet it is composed entirely of extreme points. Union of all rotations of this entire set about its vertical edge produces another convex set in three dimensions having no edges; but that convex set produced by rotation about horizontal edge containing D has edges.

- An *exposed point*, the definition of *vertex*, is equivalent to a zero-dimensional exposed face; the point of intersection with a strictly supporting hyperplane.
- A *facet* is an $(n - 1)$ -dimensional exposed face of an n -dimensional convex set \mathcal{C} ; facets exist in one-to-one correspondence with the $(n - 1)$ -dimensional faces.^{2.25}
- $\overline{\{\text{exposed points}\}} = \{\text{extreme points}\}$
 $\{\text{exposed faces}\} \subseteq \{\text{faces}\}$ △

2.6.1.1 Density of exposed points

For any closed convex set \mathcal{C} , its exposed points constitute a *dense* subset of its extreme points; [271, §18] [298] [293, §3.6, p.115] dense in the sense [333] that closure of that subset yields the set of extreme points.

For the convex set illustrated in Figure 29, point B cannot be exposed because it relatively bounds both the facet \overline{AB} and the closed quarter circle, each bounding the set. Since B is not relatively interior to any line segment in the set, then B is an extreme point by definition. Point B may be regarded as the limit of some sequence of exposed points beginning at vertex C.

2.6.1.2 Face transitivity and algebra

Faces of a convex set enjoy transitive relation. If \mathcal{F}_1 is a face (an extreme set) of \mathcal{F}_2 which in turn is a face of \mathcal{F}_3 , then it is always true that \mathcal{F}_1 is a face of \mathcal{F}_3 . (The parallel statement for exposed faces is false. [271, §18]) For example, any extreme point of \mathcal{F}_2 is an extreme point of \mathcal{F}_3 ; in this example, \mathcal{F}_2 could be a face exposed by a hyperplane supporting polyhedron \mathcal{F}_3 . [198, def.115/6, p.358] Yet it is erroneous to presume that a face, of dimension 1 or more, consists entirely of extreme points, nor is a face of dimension 2 or more entirely composed of edges, and so on.

For the polyhedron in \mathbb{R}^3 from Figure 18, for example, the nonempty faces exposed by a hyperplane are the vertices, edges, and facets; there are no more. The zero-, one-, and two-dimensional faces are in one-to-one correspondence with the exposed faces in that example.

^{2.25}This coincidence occurs simply because the facet's dimension is the same as the dimension of the supporting hyperplane exposing it.

Define the smallest face \mathcal{F} that contains some element G of a convex set \mathcal{C} :

$$\mathcal{F}(\mathcal{C} \ni G) \quad (152)$$

videlicet, $\mathcal{C} \supseteq \mathcal{F}(\mathcal{C} \ni G) \ni G$. An affine set has no faces except itself and the empty set. The smallest face that contains G of the intersection of convex set \mathcal{C} with an affine set \mathcal{A} [211, §2.4] [212]

$$\mathcal{F}((\mathcal{C} \cap \mathcal{A}) \ni G) = \mathcal{F}(\mathcal{C} \ni G) \cap \mathcal{A} \quad (153)$$

equals the intersection of \mathcal{A} with the smallest face that contains G of set \mathcal{C} .

2.6.1.3 Boundary

(confer §2.1.7) The classical definition of *boundary* of a set \mathcal{C} presumes nonempty interior:

$$\partial\mathcal{C} = \bar{\mathcal{C}} \setminus \text{int } \mathcal{C} \quad (15)$$

More suitable for the study of convex sets is the *relative boundary*; defined [177, §A.2.1.2]

$$\text{rel } \partial\mathcal{C} = \bar{\mathcal{C}} \setminus \text{rel int } \mathcal{C} \quad (154)$$

the boundary relative to the affine hull of \mathcal{C} , conventionally equivalent to:

2.6.1.3.1 Definition. *Conventional boundary of convex set.* [177, §C.3.1] The relative boundary $\partial\mathcal{C}$ of a nonempty convex set \mathcal{C} is the union of all exposed faces of $\bar{\mathcal{C}}$. △

Equivalence of this definition to (154) comes about because it is conventionally presumed that any supporting hyperplane, central to the definition of exposure, does not contain \mathcal{C} . [271, p.100]

Any face \mathcal{F} of convex set \mathcal{C} (that is not \mathcal{C} itself) belongs to $\text{rel } \partial\mathcal{C}$. (§2.8.2.1) In the exception when \mathcal{C} is a single point $\{x\}$, (11)

$$\text{rel } \partial\{x\} = \overline{\{x\}} \setminus \{x\} = \emptyset, \quad x \in \mathbb{R}^n \quad (155)$$

A bounded convex polyhedron (§2.3.2, §2.12.0.0.1) having nonempty interior, for example, in \mathbb{R} has a boundary constructed from two points, in \mathbb{R}^2 from at least three line segments, in \mathbb{R}^3 from convex polygons, while a convex

polychoron (a bounded polyhedron in \mathbb{R}^4 [333]) has a boundary constructed from three-dimensional convex polyhedra. A halfspace is partially bounded by a hyperplane; its interior therefore excludes that hyperplane. By Definition 2.6.1.3.1, an affine set has no relative boundary.

2.7 Cones

In optimization, convex cones achieve prominence because they generalize subspaces. Most compelling is the projection analogy: Projection on a subspace can be ascertained from projection on its orthogonal complement (§E), whereas projection on a closed convex cone can be determined from projection instead on its *algebraic complement* (§2.13, §E.9.2.1); called the *polar cone*.

2.7.0.0.1 Definition. Ray.

The one-dimensional set

$$\{\zeta\Gamma + B \mid \zeta \geq 0, \Gamma \neq \mathbf{0}\} \subset \mathbb{R}^n \quad (156)$$

defines a *halfline* called a *ray* in nonzero *direction* $\Gamma \in \mathbb{R}^n$ having *base* $B \in \mathbb{R}^n$. When $B = \mathbf{0}$, a ray is the conic hull of direction Γ ; hence a convex cone. \triangle

The relative boundary of a single ray, base $\mathbf{0}$, in any dimension is the origin because that is the union of all exposed faces not containing the entire set. Its relative interior is the ray itself excluding the origin.

2.7.1 Cone defined

A set \mathcal{X} is called, simply, *cone* if and only if

$$\Gamma \in \mathcal{X} \Rightarrow \zeta\Gamma \in \overline{\mathcal{X}} \text{ for all } \zeta \geq 0 \quad (157)$$

where $\overline{\mathcal{X}}$ denotes closure of cone \mathcal{X} . An example of such a cone is the union of two opposing quadrants; *e.g.*, $\mathcal{X} = \{x \in \mathbb{R}^2 \mid x_1 x_2 \geq 0\}$ which is not convex. [331, §2.5] Similar examples are shown in Figure 30 and Figure 34.

All cones can be defined by an aggregate of rays emanating exclusively from the origin (but not all cones are convex). Hence all closed cones contain




Figure 30: **(a)** Two-dimensional nonconvex cone drawn truncated. Boundary of this cone is itself a cone. Each polar half is itself a convex cone. **(b)** This convex cone (drawn truncated) is a line through the origin in any dimension. It has no relative boundary, while its relative interior comprises entire line.




Figure 31: This nonconvex cone in \mathbb{R}^2 is a pair of lines through the origin. [220, §2.4]




Figure 32: Boundary of a convex cone in \mathbb{R}^2 is a nonconvex cone; a pair of rays emanating from the origin.




Figure 33: Nonconvex cone \mathcal{X} drawn truncated in \mathbb{R}^2 . Boundary is also a cone. [220, §2.4] Cone exterior is convex cone.




Figure 34: Truncated nonconvex cone $\mathcal{X} = \{x \in \mathbb{R}^2 \mid x_1 \geq x_2, x_1 x_2 \geq 0\}$. Boundary is also a cone. [220, §2.4] Cartesian axes drawn for reference. Each half (about the origin) is itself a convex cone.

the origin $\mathbf{0}$ and are unbounded, excepting the simplest cone $\{\mathbf{0}\}$. The empty set \emptyset is not a cone, but its conic hull is;

$$\text{cone } \emptyset \triangleq \{\mathbf{0}\} \quad (96)$$

2.7.2 Convex cone

We call the set $\mathcal{K} \subseteq \mathbb{R}^M$ a *convex cone* iff

$$\Gamma_1, \Gamma_2 \in \mathcal{K} \Rightarrow \zeta \Gamma_1 + \xi \Gamma_2 \in \overline{\mathcal{K}} \quad \text{for all } \zeta, \xi \geq 0 \quad (158)$$

Apparent from this definition, $\zeta \Gamma_1 \in \overline{\mathcal{K}}$ and $\xi \Gamma_2 \in \overline{\mathcal{K}} \quad \forall \zeta, \xi \geq 0$; meaning, \mathcal{K} is a cone. Set \mathcal{K} is convex since, for any particular $\zeta, \xi \geq 0$

$$\mu \zeta \Gamma_1 + (1 - \mu) \xi \Gamma_2 \in \overline{\mathcal{K}} \quad \forall \mu \in [0, 1] \quad (159)$$

because $\mu \zeta, (1 - \mu) \xi \geq 0$. Obviously,

$$\{\mathcal{X}\} \supset \{\mathcal{K}\} \quad (160)$$

the set of all convex cones is a proper subset of all cones. The set of convex cones is a narrower but more familiar class of cone, any member

of which can be equivalently described as the intersection of a possibly (but not necessarily) infinite number of hyperplanes (through the origin) and halfspaces whose bounding hyperplanes pass through the origin; a halfspace-description (§2.4). Convex cones need not be full-dimensional.




Figure 35: Not a cone; ironically, the three-dimensional *flared horn* (with or without its interior) resembling the mathematical symbol \succ denoting cone membership and partial order.

Familiar examples of convex cones include an unbounded *ice-cream cone* united with its interior (a.k.a: *second-order cone*, *quadratic cone*, *circular cone* (§2.9.2.5.1), *Lorentz cone* (confer Figure 42) [53, exmps.2.3 & 2.25])

$$\mathcal{K}_\ell = \left\{ \begin{bmatrix} x \\ t \end{bmatrix} \in \mathbb{R}^n \times \mathbb{R} \mid \|x\|_\ell \leq t \right\}, \quad \ell=2 \quad (161)$$

and any polyhedral cone (§2.12.1.0.1); e.g., any orthant generated by Cartesian half-axes (§2.1.3). Esoteric examples of convex cones include the point at the origin, any line through the origin, any ray having the origin as base such as the nonnegative real line \mathbb{R}_+ in subspace \mathbb{R} , any halfspace partially bounded by a hyperplane through the origin, the positive semidefinite cone \mathbb{S}_+^M (174), the cone of Euclidean distance matrices EDM^N (810) (§6), any subspace, and Euclidean vector space \mathbb{R}^n .

2.7.2.1 cone invariance

More Euclidean bodies are cones, it seems, than are not.^{2.26} This class of convex body, the convex cone, is invariant to nonnegative scaling, linear and single- or many-valued inverse linear transformation, their vector summation, Cartesian product, and intersection, [271, p.22] but is not invariant to translation or projection: *e.g.*, Figure 41.

2.7.2.1.1 Theorem. Cone intersection. [271, §2, §19]

The intersection of an arbitrary collection of convex cones is a convex cone. Intersection of an arbitrary collection of closed convex cones is a closed convex cone. [228, §2.3] Intersection of a finite number of polyhedral cones (§2.12.1.0.1, Figure 46 p.131) is polyhedral. \diamond

The property *pointedness* is associated with a convex cone.

2.7.2.1.2 Definition. Pointed convex cone. (confer §2.12.2.2)

A convex cone \mathcal{K} is *pointed* iff it contains no line. Equivalently, \mathcal{K} is not pointed iff there exists any nonzero direction $\Gamma \in \overline{\mathcal{K}}$ such that $-\Gamma \in \overline{\mathcal{K}}$. [53, §2.4.1] If the origin is an extreme point of $\overline{\mathcal{K}}$ or, equivalently, if

$$\overline{\mathcal{K}} \cap -\overline{\mathcal{K}} = \{\mathbf{0}\} \quad (162)$$

then \mathcal{K} is pointed, and *vice versa*. [293, §2.10] A convex cone is pointed iff the origin is the smallest nonempty face of its closure. \triangle

Then a pointed closed convex cone, by principle of separating hyperplane (§2.4.2.7), has a strictly supporting hyperplane at the origin. The simplest and only bounded [331, p.75] convex cone $\mathcal{K} = \{\mathbf{0}\} \subseteq \mathbb{R}^n$ is pointed by convention, but has empty interior. Its relative boundary is the empty set (155) while its relative interior is the point itself (11). The pointed convex cone that is a halfline emanating from the origin in \mathbb{R}^n has the origin as relative boundary while its relative interior is the halfline itself, excluding the origin.

^{2.26} *confer* Figures: 22 30 31 32 33 34 35 37 39 46 50 53 55 56 57 58 59 60 61 117 130 153

2.7.2.1.3 Theorem. *Pointed cones.* [48, §3.3.15, exer.20]

A closed convex cone $\mathcal{K} \subset \mathbb{R}^n$ is pointed if and only if there exists a normal α such that the set

$$\mathcal{C} \triangleq \{x \in \mathcal{K} \mid \langle x, \alpha \rangle = 1\} \quad (163)$$

is closed, bounded, and $\mathcal{K} = \text{cone } \mathcal{C}$. Equivalently, \mathcal{K} is pointed if and only if there exists a vector β normal to a hyperplane strictly supporting \mathcal{K} ; *id est*, for some positive scalar ϵ

$$\langle x, \beta \rangle \geq \epsilon \|x\| \quad \forall x \in \mathcal{K} \quad (164)$$

◇

If closed convex cone \mathcal{K} is not pointed, then it has no extreme point.^{2.27} Yet a pointed closed convex cone has only one extreme point [37, §3.3]: the exposed point residing at the origin; its vertex. And from the *cone intersection theorem* it follows that an intersection of convex cones is pointed if at least one of the cones is.

2.7.2.2 Pointed closed convex cone and partial order

Relation \preceq is a *partial order* on some set if the relation possesses^{2.28}

reflexivity ($x \preceq x$)

antisymmetry ($x \preceq z, z \preceq x \Rightarrow x = z$)

transitivity ($x \preceq y, y \preceq z \Rightarrow x \preceq z$), ($x \prec y, y \prec z \Rightarrow x \prec z$)

A pointed closed convex cone \mathcal{K} induces partial order on \mathbb{R}^n or $\mathbb{R}^{m \times n}$, [20, §1] [287, p.7] essentially defined by vector or matrix inequality;

$$x \preceq_{\mathcal{K}} z \Leftrightarrow z - x \in \mathcal{K} \quad (165)$$

$$x \prec_{\mathcal{K}} z \Leftrightarrow z - x \in \text{rel int } \mathcal{K} \quad (166)$$

^{2.27} nor does it have extreme directions (§2.8.1).

^{2.28} A set is *totally ordered* if it further obeys a *comparability* property of the relation (a.k.a, *trichotomy law*): for each and every x and y from the set, $x \preceq y$ or $y \preceq x$.




Figure 36: **(a)** Point x is the minimum element of set \mathcal{C}_1 with respect to cone \mathcal{K} because cone translated to $x \in \mathcal{C}_1$ contains set. **(b)** Point y is a minimal element of set \mathcal{C}_2 with respect to cone \mathcal{K} because negative cone translated to $y \in \mathcal{C}_2$ contains only y . (Cones drawn truncated in \mathbb{R}^2 .) These concepts, minimum/minimal, become equivalent under a *total order*.

Neither x or z is necessarily a member of \mathcal{K} for these relations to hold. Only when \mathcal{K} is the nonnegative orthant do these inequalities reduce to ordinary entrywise comparison. (§2.13.4.2.3) Inclusive of that special case, we ascribe nomenclature *generalized inequality* to comparison with respect to a pointed closed convex cone.

Visceral mechanics of actually comparing points when cone \mathcal{K} is not an orthant are well illustrated in the example of Figure 58 which relies on the equivalent membership-interpretation in definition (165) or (166). *Comparable points* and the *minimum element* of some vector- or matrix-valued partially ordered set are thus well defined, so nonincreasing sequences with respect to cone \mathcal{K} can therefore converge in this sense: Point $x \in \mathcal{C}$ is the (unique) minimum element of set \mathcal{C} with respect to cone \mathcal{K} iff for each and every $z \in \mathcal{C}$ we have $x \preceq z$; equivalently, iff $\mathcal{C} \subseteq x + \mathcal{K}$.^{2.29}

Further properties of partial order with respect to pointed closed convex cone \mathcal{K} are:

$$\text{homogeneity } (x \preceq y, \lambda \geq 0 \Rightarrow \lambda x \preceq \lambda z), \quad (x \prec y, \lambda > 0 \Rightarrow \lambda x \prec \lambda z)$$

$$\text{additivity } (x \preceq z, u \preceq v \Rightarrow x + u \preceq z + v), \quad (x \prec z, u \preceq v \Rightarrow x + u \prec z + v)$$

A closely related concept, *minimal element*, is useful for partially ordered sets having no minimum element: Point $x \in \mathcal{C}$ is a minimal element of set \mathcal{C} with respect to \mathcal{K} if and only if $(x - \mathcal{K}) \cap \mathcal{C} = x$. (Figure 36) No uniqueness is implied here, although implicit is the assumption: $\dim \mathcal{K} \geq \dim \text{aff } \mathcal{C}$.

2.7.2.2.1 Definition. *Proper cone:* a cone that is

- pointed
- closed
- convex
- has nonempty interior (is full-dimensional). △

A proper cone remains proper under injective linear transformation. [79, §5.1] Examples of proper cones are the positive semidefinite cone \mathbb{S}_+^M in the ambient space of symmetric matrices (§2.9), the nonnegative real line \mathbb{R}_+ in vector space \mathbb{R} , or any orthant in \mathbb{R}^n .

^{2.29}Borwein & Lewis [48, §3.3, exer.21] ignore possibility of equality to $x + \mathcal{K}$ in this condition, and require a second condition: ... and $\mathcal{C} \subset y + \mathcal{K}$ for some y in \mathbb{R}^n implies $x \in y + \mathcal{K}$.

2.8 Cone boundary

Every hyperplane supporting a convex cone contains the origin. [177, §A.4.2] Because any supporting hyperplane to a convex cone must therefore itself be a cone, then from the *cone intersection theorem* it follows:

2.8.0.0.1 Lemma. *Cone faces.* [25, §II.8]
Each nonempty exposed face of a convex cone is a convex cone. \diamond

2.8.0.0.2 Theorem. *Proper-cone boundary.*
Suppose a nonzero point Γ lies on the boundary $\partial\mathcal{K}$ of proper cone \mathcal{K} in \mathbb{R}^n . Then it follows that the ray $\{\zeta\Gamma \mid \zeta \geq 0\}$ also belongs to $\partial\mathcal{K}$. \diamond

Proof. By virtue of its propriety, a proper cone guarantees existence of a strictly supporting hyperplane at the origin. [271, cor.11.7.3]^{2.30} Hence the origin belongs to the boundary of \mathcal{K} because it is the zero-dimensional exposed face. The origin belongs to the ray through Γ , and the ray belongs to \mathcal{K} by definition (157). By the *cone faces lemma*, each and every nonempty exposed face must include the origin. Hence the closed line segment $\overline{\mathbf{0}\Gamma}$ must lie in an exposed face of \mathcal{K} because both endpoints do by Definition 2.6.1.3.1. That means there exists a supporting hyperplane $\underline{\partial\mathcal{H}}$ to \mathcal{K} containing $\overline{\mathbf{0}\Gamma}$. So the ray through Γ belongs both to \mathcal{K} and to $\underline{\partial\mathcal{H}}$. $\underline{\partial\mathcal{H}}$ must therefore expose a face of \mathcal{K} that contains the ray; *id est*,

$$\{\zeta\Gamma \mid \zeta \geq 0\} \subseteq \mathcal{K} \cap \underline{\partial\mathcal{H}} \subset \partial\mathcal{K} \quad (167)$$

\blacklozenge

Proper cone $\{\mathbf{0}\}$ in \mathbb{R}^0 has no boundary (154) because (11)

$$\text{rel int}\{\mathbf{0}\} = \{\mathbf{0}\} \quad (168)$$

The boundary of any proper cone in \mathbb{R} is the origin.

The boundary of any convex cone whose dimension exceeds 1 can be constructed entirely from an aggregate of rays emanating exclusively from the origin.

^{2.30}Rockafellar's corollary yields a supporting hyperplane at the origin to any convex cone in \mathbb{R}^n not equal to \mathbb{R}^n .




Figure 37: \mathcal{K} is a pointed polyhedral cone having empty interior in \mathbb{R}^3 (drawn truncated and in a plane parallel to the floor upon which you stand). \mathcal{K}^* is a *wedge* whose truncated boundary is illustrated (drawn perpendicular to the floor). In this particular instance, $\mathcal{K} \subset \text{int } \mathcal{K}^*$ (excepting the origin). Cartesian coordinate axes drawn for reference.

2.8.1 Extreme direction

The property *extreme direction* arises naturally in connection with the pointed closed convex cone $\mathcal{K} \subset \mathbb{R}^n$, being analogous to extreme point. [271, §18, p.162]^{2.31} An extreme direction Γ_ε of pointed \mathcal{K} is a vector corresponding to an edge that is a ray emanating from the origin.^{2.32} Nonzero direction Γ_ε in pointed \mathcal{K} is extreme if and only if

$$\zeta_1 \Gamma_1 + \zeta_2 \Gamma_2 \neq \Gamma_\varepsilon \quad \forall \zeta_1, \zeta_2 \geq 0, \quad \forall \Gamma_1, \Gamma_2 \in \mathcal{K} \setminus \{\zeta \Gamma_\varepsilon \in \mathcal{K} \mid \zeta \geq 0\} \quad (169)$$

In words, an extreme direction in a pointed closed convex cone is the direction of a ray, called an *extreme ray*, that cannot be expressed as a conic combination of directions of any rays in the cone distinct from it.

An extreme ray is a one-dimensional face of \mathcal{K} . By (97), extreme direction Γ_ε is not a point relatively interior to any line segment in $\mathcal{K} \setminus \{\zeta \Gamma_\varepsilon \in \mathcal{K} \mid \zeta \geq 0\}$. Thus, by analogy, the corresponding extreme ray $\{\zeta \Gamma_\varepsilon \in \mathcal{K} \mid \zeta \geq 0\}$ is not a ray relatively interior to any *plane segment*^{2.33} in \mathcal{K} .

2.8.1.1 extreme distinction, uniqueness

An extreme *direction* is unique, but its vector representation Γ_ε is not because any positive scaling of it produces another vector in the same (extreme) direction. Hence an extreme direction is *unique* to within a positive scaling. When we say extreme directions are *distinct*, we are referring to distinctness of rays containing them. Nonzero vectors of various length in the same extreme direction are therefore interpreted to be identical extreme directions.^{2.34}

The extreme directions of the polyhedral cone in Figure 22 (p.67), for example, correspond to its three edges. For any pointed polyhedral cone, there is a one-to-one correspondence of one-dimensional faces with extreme directions.

The extreme directions of the positive semidefinite cone (§2.9) comprise the infinite set of all symmetric rank-one matrices. [20, §6] [174, §III] It

^{2.31}We diverge from Rockafellar's extreme direction: "extreme point at infinity".

^{2.32}An edge (§2.6.0.0.3) of a convex cone is not necessarily a ray. A convex cone may contain an edge that is a line; *e.g.*, a wedge-shaped polyhedral cone (\mathcal{K}^* in Figure 37).

^{2.33}A planar fragment; in this context, a planar cone.

^{2.34}Like vectors, an extreme direction can be identified by the Cartesian point at the vector's head with respect to the origin.

is sometimes prudent to instead consider the less infinite but complete normalized set, for $M > 0$ (*confer* (207))

$$\{zz^T \in \mathbb{S}^M \mid \|z\| = 1\} \quad (170)$$

The positive semidefinite cone in one dimension $M = 1$, \mathbb{S}_+ the nonnegative real line, has one extreme direction belonging to its relative interior; an idiosyncrasy of dimension 1.

Pointed closed convex cone $\mathcal{K} = \{\mathbf{0}\}$ has no extreme direction because extreme directions are nonzero by definition.

- If closed convex cone \mathcal{K} is not pointed, then it has no extreme directions and no vertex. [20, §1]

Conversely, pointed closed convex cone \mathcal{K} is equivalent to the convex hull of its vertex and all its extreme directions. [271, §18, p.167] That is the practical utility of extreme direction; to facilitate construction of polyhedral sets, apparent from the *extremes theorem*:

2.8.1.1.1 Theorem. (Klee) *Extremes.* [293, §3.6] [271, §18, p.166] (*confer* §2.3.2, §2.12.2.0.1) Any closed convex set containing no lines can be expressed as the convex hull of its extreme points and extreme rays. \diamond

It follows that any element of a convex set containing no lines may be expressed as a linear combination of its extreme elements; *e.g.*, Example 2.9.2.4.1.

2.8.1.2 Generators

In the narrowest sense, generators for a convex set comprise any collection of points and directions whose convex hull constructs the set.

When the *extremes theorem* applies, the extreme points and directions are called generators of a convex set. An arbitrary collection of generators for a convex set includes its extreme elements as a subset; the set of extreme elements of a convex set is a minimal set of generators for that convex set. Any polyhedral set has a minimal set of generators whose *cardinality* is finite.

When the convex set under scrutiny is a closed convex cone, conic combination of generators during construction is implicit as shown in Example 2.8.1.2.1 and Example 2.10.2.0.1. So, a vertex at the origin (if it exists) becomes benign.

We can, of course, generate affine sets by taking the affine hull of any collection of points and directions. We broaden, thereby, the meaning of generator to be inclusive of all kinds of hulls.

Any hull of generators is loosely called a vertex-description. (§2.3.4) Hulls encompass subspaces, so any basis constitutes generators for a vertex-description; span basis $\mathcal{R}(A)$.

2.8.1.2.1 Example. Application of extremes theorem.

Given an extreme point at the origin and N extreme rays, denoting the i^{th} extreme direction by $\Gamma_i \in \mathbb{R}^n$, then the convex hull is (79)

$$\begin{aligned} \mathcal{P} &= \{[\mathbf{0} \ \Gamma_1 \ \Gamma_2 \cdots \Gamma_N] a \zeta \mid a^T \mathbf{1} = 1, a \succeq 0, \zeta \geq 0\} \\ &= \{[\Gamma_1 \ \Gamma_2 \cdots \Gamma_N] a \zeta \mid a^T \mathbf{1} \leq 1, a \succeq 0, \zeta \geq 0\} \\ &= \{[\Gamma_1 \ \Gamma_2 \cdots \Gamma_N] b \mid b \succeq 0\} \subset \mathbb{R}^n \end{aligned} \quad (171)$$

a closed convex set that is simply a conic hull like (95). \square

2.8.2 Exposed direction

2.8.2.0.1 Definition. Exposed point & direction of pointed convex cone. [271, §18] (confer §2.6.1.0.1)

- When a convex cone has a vertex, an exposed point, it resides at the origin; there can be only one.
- In the closure of a pointed convex cone, an *exposed direction* is the direction of a one-dimensional exposed face that is a ray emanating from the origin.
- $\{\text{exposed directions}\} \subseteq \{\text{extreme directions}\} \quad \triangle$

For a proper cone in vector space \mathbb{R}^n with $n \geq 2$, we can say more:

$$\overline{\{\text{exposed directions}\}} = \{\text{extreme directions}\} \quad (172)$$

It follows from Lemma 2.8.0.0.1 for any pointed closed convex cone, there is one-to-one correspondence of one-dimensional exposed faces with exposed directions; *id est*, there is no one-dimensional exposed face that is not a ray base $\mathbf{0}$.

The pointed closed convex cone \mathbb{EDM}^2 , for example, is a ray in isomorphic subspace \mathbb{R} whose relative boundary (§2.6.1.3.1) is the origin. The conventionally exposed directions of \mathbb{EDM}^2 constitute the empty set $\emptyset \subset \{\text{extreme direction}\}$. This cone has one extreme direction belonging to its relative interior; an idiosyncrasy of dimension 1.

2.8.2.1 Connection between boundary and extremes

2.8.2.1.1 Theorem. *Exposed.* [271, §18.7] (confer §2.8.1.1.1)

Any closed convex set \mathcal{C} containing no lines (and whose dimension is at least 2) can be expressed as the closure of the convex hull of its exposed points and exposed rays. \diamond

From Theorem 2.8.1.1.1,

$$\left. \begin{aligned} \text{rel } \partial \mathcal{C} &= \bar{\mathcal{C}} \setminus \text{rel int } \mathcal{C} && (154) \\ &= \overline{\text{conv}\{\text{exposed points and exposed rays}\}} \setminus \text{rel int } \mathcal{C} \\ &= \text{conv}\{\text{extreme points and extreme rays}\} \setminus \text{rel int } \mathcal{C} \end{aligned} \right\} (173)$$

Thus each and every extreme point of a convex set (that is not a point) resides on its relative boundary, while each and every extreme direction of a convex set (that is not a halfline and contains no line) resides on its relative boundary because extreme points and directions of such respective sets do not belong to relative interior by definition.

The relationship between extreme sets and the relative boundary actually goes deeper: Any face \mathcal{F} of convex set \mathcal{C} (that is not \mathcal{C} itself) belongs to $\text{rel } \partial \mathcal{C}$, so $\dim \mathcal{F} < \dim \mathcal{C}$. [271, §18.1.3]

2.8.2.2 Converse caveat

It is inconsequent to presume that each and every extreme point and direction is necessarily exposed, as might be erroneously inferred from the *conventional boundary definition* (§2.6.1.3.1); although it can correctly be inferred: each and every extreme point and direction belongs to some exposed face.

Arbitrary points residing on the relative boundary of a convex set are not necessarily exposed or extreme points. Similarly, the direction of an arbitrary ray, base $\mathbf{0}$, on the boundary of a convex cone is not necessarily an exposed or extreme direction. For the polyhedral cone illustrated in Figure 22, for




Figure 38: Properties of extreme points carry over to extreme directions. [271, §18] Four rays (drawn truncated) on boundary of conic hull of two-dimensional closed convex set from Figure 29 lifted to \mathbb{R}^3 . Ray through point A is exposed hence extreme. Extreme direction B on cone boundary is not an exposed direction, although it belongs to the exposed face $\text{cone}\{A, B\}$. Extreme ray through C is exposed. Point D is neither an exposed or extreme direction although it belongs to a two-dimensional exposed face of the conic hull.

example, there are three two-dimensional exposed faces constituting the entire boundary, each composed of an infinity of rays. Yet there are only three exposed directions.

Neither is an extreme direction on the boundary of a pointed convex cone necessarily an exposed direction. Lift the two-dimensional set in Figure 29, for example, into three dimensions such that no two points in the set are collinear with the origin. Then its conic hull can have an extreme direction B on the boundary that is not an exposed direction, illustrated in Figure 38.

2.9 Positive semidefinite (PSD) cone

The cone of positive semidefinite matrices studied in this section is arguably the most important of all non-polyhedral cones whose facial structure we completely understand.

–Alexander Barvinok [25, p.78]

2.9.0.0.1 Definition. Positive semidefinite cone.

The set of all symmetric positive semidefinite matrices of particular dimension M is called the *positive semidefinite cone*:

$$\begin{aligned} \mathbb{S}_+^M &\triangleq \{A \in \mathbb{S}^M \mid A \succeq 0\} \\ &= \{A \in \mathbb{S}^M \mid y^T A y \geq 0 \quad \forall \|y\| = 1\} \\ &= \bigcap_{\|y\|=1} \{A \in \mathbb{S}^M \mid \langle yy^T, A \rangle \geq 0\} \end{aligned} \tag{174}$$

formed by the intersection of an infinite number of halfspaces (§2.4.1.1) in vectorized variable A ,^{2.35} each halfspace having partial boundary containing the origin in isomorphic $\mathbb{R}^{M(M+1)/2}$. It is a unique immutable proper cone in the ambient space of symmetric matrices \mathbb{S}^M .

^{2.35} infinite in number when $M > 1$. Because $y^T A y = y^T A^T y$, matrix A is almost always assumed symmetric. (§A.2.1)

The positive definite (full-rank) matrices comprise the cone interior

$$\begin{aligned}
\text{int } \mathbb{S}_+^M &= \{A \in \mathbb{S}^M \mid A \succ 0\} \\
&= \{A \in \mathbb{S}^M \mid y^T A y > 0 \quad \forall \|y\| = 1\} \\
&= \bigcap_{\|y\|=1} \{A \in \mathbb{S}^M \mid \langle yy^T, A \rangle > 0\} \\
&= \{A \in \mathbb{S}_+^M \mid \text{rank } A = M\}
\end{aligned} \tag{175}$$

while all singular positive semidefinite matrices (having at least one 0 eigenvalue) reside on the cone boundary (Figure 39); (§A.7.5)

$$\begin{aligned}
\partial \mathbb{S}_+^M &= \{A \in \mathbb{S}^M \mid \min\{\lambda(A)_i, i=1 \dots M\} = 0\} \\
&= \{A \in \mathbb{S}_+^M \mid \langle yy^T, A \rangle = 0 \text{ for some } \|y\| = 1\} \\
&= \{A \in \mathbb{S}_+^M \mid \text{rank } A < M\}
\end{aligned} \tag{176}$$


where $\lambda(A) \in \mathbb{R}^M$ holds the eigenvalues of A . △

The only symmetric positive semidefinite matrix in \mathbb{S}_+^M having M 0-eigenvalues resides at the origin. (§A.7.3.0.1)

2.9.0.1 Membership

Observe the notation $A \succeq 0$; meaning (*confer* §2.3.1.1), matrix A belongs to the positive semidefinite cone in the subspace of symmetric matrices, whereas $A \succ 0$ denotes membership to that cone's interior. (§2.13.2) Notation $A \succ 0$ can be read: *symmetric matrix A is greater than zero with respect to the positive semidefinite cone*. This notation further implies that coordinates [*sic*] for orthogonal expansion of a positive (semi)definite matrix must be its (nonnegative) positive eigenvalues (§2.13.7.1.1, §E.6.4.1.1) when expanded in its *eigenmatrices* (§A.5.1).

Generalizing comparison on the real line, the notation $A \succeq B$ denotes comparison with respect to the positive semidefinite cone; (§A.3.1) *id est*, $A \succeq B \Leftrightarrow A - B \in \mathbb{S}_+^M$ but neither matrix A or B necessarily belongs to the positive semidefinite cone. Yet, (1396) $A \succeq B, B \succeq 0 \Rightarrow A \succeq 0$; *id est*, $A \in \mathbb{S}_+^M$. (*confer* Figure 58)



Minimal set of generators are the extreme directions: $\text{svec}\{yy^T \mid y \in \mathbb{R}^M\}$

Figure 39: (d'Aspremont) Truncated boundary of PSD cone in \mathbb{S}^2 plotted in isometrically isomorphic \mathbb{R}^3 via svec (50); 0-contour of smallest eigenvalue (176). Lightest shading is closest, darkest shading is farthest and inside shell. Entire boundary can be constructed from an aggregate of rays (§2.7.0.0.1) emanating exclusively from the origin: $\{\kappa^2[z_1^2 \ \sqrt{2}z_1z_2 \ z_2^2]^T \mid \kappa \in \mathbb{R}\}$. A circular cone in this dimension (§2.9.2.5), each and every ray on boundary corresponds to an extreme direction but such is not the case in any higher dimension (*confer* Figure 22). PSD cone geometry is not as simple in higher dimensions [25, §II.12] although PSD cone is self-dual (341) in ambient real space of symmetric matrices. [174, §II] PSD cone has no two-dimensional face in any dimension, its only extreme point residing at $\mathbf{0}$.

2.9.0.1.1 Example. *Equality constraints in semidefinite program (592).* Employing properties of partial order (§2.7.2.2) for the pointed closed convex positive semidefinite cone, it is easy to show, given $A + S = C$

$$S \succeq 0 \Leftrightarrow A \preceq C \quad (177)$$

□

2.9.1 Positive semidefinite cone is convex

The set of all positive semidefinite matrices forms a convex cone in the ambient space of symmetric matrices because any pair satisfies definition (158); [180, §7.1] *videlicet*, for all $\zeta_1, \zeta_2 \geq 0$ and each and every $A_1, A_2 \in \mathbb{S}^M$

$$\zeta_1 A_1 + \zeta_2 A_2 \succeq 0 \Leftrightarrow A_1 \succeq 0, A_2 \succeq 0 \quad (178)$$

a fact easily verified by the definitive test for positive semidefiniteness of a symmetric matrix (§A):

$$A \succeq 0 \Leftrightarrow x^T A x \geq 0 \text{ for each and every } \|x\| = 1 \quad (179)$$

id est, for $A_1, A_2 \succeq 0$ and each and every $\zeta_1, \zeta_2 \geq 0$

$$\zeta_1 x^T A_1 x + \zeta_2 x^T A_2 x \geq 0 \text{ for each and every normalized } x \in \mathbb{R}^M \quad (180)$$

The convex cone \mathbb{S}_+^M is more easily visualized in the isomorphic vector space $\mathbb{R}^{M(M+1)/2}$ whose dimension is the number of free variables in a symmetric $M \times M$ matrix. When $M=2$ the PSD cone is semi-infinite in expanse in \mathbb{R}^3 , having boundary illustrated in Figure 39. When $M=3$ the PSD cone is six-dimensional, and so on.

2.9.1.0.1 Example. *Sets from maps of positive semidefinite cone.*

The set

$$\mathcal{C} = \{X \in \mathbb{S}^n \times x \in \mathbb{R}^n \mid X \succeq xx^T\} \quad (181)$$

is convex because it has *Schur-form*; (§A.4)

$$X - xx^T \succeq 0 \Leftrightarrow f(X, x) \triangleq \begin{bmatrix} X & x \\ x^T & 1 \end{bmatrix} \succeq 0 \quad (182)$$




Figure 40: Convex set $\mathcal{C} = \{X \in \mathbb{S} \times x \in \mathbb{R} \mid X \succeq xx^T\}$ drawn truncated.

e.g., Figure 40. Set \mathcal{C} is the inverse image (§2.1.9.0.1) of \mathbb{S}_+^{n+1} under affine mapping f . The set $\{X \in \mathbb{S}^n \times x \in \mathbb{R}^n \mid X \preceq xx^T\}$ is not convex, in contrast, having no Schur-form. Yet for fixed $x = x_p$, the set

$$\{X \in \mathbb{S}^n \mid X \preceq x_p x_p^T\} \quad (183)$$

is simply the negative semidefinite cone shifted to $x_p x_p^T$. \square

2.9.1.0.2 Example. Inverse image of positive semidefinite cone.

Now consider finding the set of all matrices $X \in \mathbb{S}^N$ satisfying

$$AX + B \succeq 0 \quad (184)$$

given $A, B \in \mathbb{S}^N$. Define the set

$$\mathcal{X} \triangleq \{X \mid AX + B \succeq 0\} \subseteq \mathbb{S}^N \quad (185)$$

which is the inverse image of the positive semidefinite cone under affine transformation $g(X) \triangleq AX + B$. Set \mathcal{X} must therefore be convex by Theorem 2.1.9.0.1.

Yet we would like a less amorphous characterization of this set, so instead we consider its vectorization (31) which is easier to visualize:

$$\text{vec } g(X) = \text{vec}(AX) + \text{vec } B = (I \otimes A) \text{vec } X + \text{vec } B \quad (186)$$

where

$$I \otimes A \triangleq Q\Lambda Q^T \in \mathbb{S}^{N^2} \quad (187)$$

is a block-diagonal matrix formed by *Kronecker product* (§A.1.1 no.31, §D.1.2.1). Assign

$$\begin{aligned} x &\triangleq \text{vec } X \in \mathbb{R}^{N^2} \\ b &\triangleq \text{vec } B \in \mathbb{R}^{N^2} \end{aligned} \quad (188)$$

then make the equivalent problem: Find

$$\text{vec } \mathcal{X} = \{x \in \mathbb{R}^{N^2} \mid (I \otimes A)x + b \in \mathcal{K}\} \quad (189)$$

where

$$\mathcal{K} \triangleq \text{vec } \mathbb{S}_+^N \quad (190)$$

is a proper cone isometrically isomorphic with the positive semidefinite cone in the subspace of symmetric matrices; the vectorization of every element of \mathbb{S}_+^N . Utilizing the diagonalization (187),

$$\begin{aligned} \text{vec } \mathcal{X} &= \{x \mid \Lambda Q^T x \in Q^T(\mathcal{K} - b)\} \\ &= \{x \mid \Phi Q^T x \in \Lambda^\dagger Q^T(\mathcal{K} - b)\} \subseteq \mathbb{R}^{N^2} \end{aligned} \quad (191)$$

where \dagger denotes matrix *pseudoinverse* (§E) and

$$\Phi \triangleq \Lambda^\dagger \Lambda \quad (192)$$

is a diagonal projection matrix whose entries are either 1 or 0 (§E.3). We have the complementary sum

$$\Phi Q^T x + (I - \Phi)Q^T x = Q^T x \quad (193)$$

So, adding $(I - \Phi)Q^T x$ to both sides of the membership within (191) admits

$$\begin{aligned} \text{vec } \mathcal{X} &= \{x \in \mathbb{R}^{N^2} \mid Q^T x \in \Lambda^\dagger Q^T(\mathcal{K} - b) + (I - \Phi)Q^T x\} \\ &= \{x \mid Q^T x \in \Phi(\Lambda^\dagger Q^T(\mathcal{K} - b)) \oplus (I - \Phi)\mathbb{R}^{N^2}\} \\ &= \{x \in Q\Lambda^\dagger Q^T(\mathcal{K} - b) \oplus Q(I - \Phi)\mathbb{R}^{N^2}\} \\ &= (I \otimes A)^\dagger(\mathcal{K} - b) \oplus \mathcal{N}(I \otimes A) \end{aligned} \quad (194)$$

where we used the facts: linear function $Q^T x$ in x on \mathbb{R}^{N^2} is a *bijection*, and $\Phi\Lambda^\dagger = \Lambda^\dagger$.

$$\text{vec } \mathcal{X} = (I \otimes A)^\dagger \text{vec}(\mathbb{S}_+^N - B) \oplus \mathcal{N}(I \otimes A) \quad (195)$$

In words, set $\text{vec } \mathcal{X}$ is the vector sum of the translated PSD cone (linearly mapped onto the rowspace of $I \otimes A$ (§E)) and the nullspace of $I \otimes A$ (synthesis of fact from §A.6.3 and §A.7.3.0.1). Should $I \otimes A$ have no nullspace, then $\text{vec } \mathcal{X} = (I \otimes A)^{-1} \text{vec}(\mathbb{S}_+^N - B)$ which is the expected result. \square

2.9.2 Positive semidefinite cone boundary

For any symmetric positive semidefinite matrix A of rank ρ , there must exist a rank ρ matrix Y such that A be expressible as an outer product in Y ; [294, §6.3]

$$A = YY^T \in \mathbb{S}_+^M, \quad \text{rank } A = \rho, \quad Y \in \mathbb{R}^{M \times \rho} \quad (196)$$

Then the boundary of the positive semidefinite cone may be expressed

$$\partial \mathbb{S}_+^M = \{A \in \mathbb{S}_+^M \mid \text{rank } A < M\} = \{YY^T \mid Y \in \mathbb{R}^{M \times M-1}\} \quad (197)$$

Because the boundary of any convex body is obtained with closure of its relative interior (§2.1.7, §2.6.1.3), from (175) we must also have

$$\begin{aligned} \mathbb{S}_+^M &= \overline{\{A \in \mathbb{S}_+^M \mid \text{rank } A < M\}} = \overline{\{YY^T \mid Y \in \mathbb{R}^{M \times M}, \text{rank } Y = M\}} \\ &= \{YY^T \mid Y \in \mathbb{R}^{M \times M}\} \end{aligned} \quad (198)$$

2.9.2.1 rank ρ subset of the positive semidefinite cone

For the same reason (closure), this applies more generally; for $0 \leq \rho \leq M$

$$\overline{\{A \in \mathbb{S}_+^M \mid \text{rank } A = \rho\}} = \{A \in \mathbb{S}_+^M \mid \text{rank } A \leq \rho\} \quad (199)$$

For easy reference, we give such generally nonconvex sets a name: *rank ρ subset of a positive semidefinite cone*. For $\rho < M$ this subset, nonconvex for $M > 1$, resides on the positive semidefinite cone boundary.

2.9.2.1.1 Exercise. Closure and rank ρ subset.

Prove equality in (199). \blacktriangledown

For example,

$$\partial \mathbb{S}_+^M = \overline{\{A \in \mathbb{S}_+^M \mid \text{rank } A = M-1\}} = \{A \in \mathbb{S}_+^M \mid \text{rank } A \leq M-1\} \quad (200)$$




Figure 41: **(a)** Projection of the PSD cone \mathbb{S}_+^2 , truncated above $\gamma=1$, on $\alpha\beta$ -plane in isometrically isomorphic \mathbb{R}^3 . View is from above with respect to Figure 39. **(b)** Truncated above $\gamma=2$. From these plots we may infer, for example, the line $\{[0 \ 1/\sqrt{2} \ \gamma]^T \mid \gamma \in \mathbb{R}\}$ intercepts the PSD cone at some large value of γ ; in fact, $\gamma=\infty$.

In \mathbb{S}^2 , each and every ray on the boundary of the positive semidefinite cone in isomorphic \mathbb{R}^3 corresponds to a symmetric rank-1 matrix (Figure 39), but that does not hold in any higher dimension.

2.9.2.2 Subspace tangent to open rank- ρ subset

When the positive semidefinite cone subset in (199) is left unclosed as in

$$\mathcal{M}(\rho) \triangleq \{A \in \mathbb{S}_+^M \mid \text{rank } A = \rho\} \quad (201)$$

then we can specify a subspace tangent to the positive semidefinite cone at a particular member of manifold $\mathcal{M}(\rho)$. Specifically, the subspace $\mathcal{R}_{\mathcal{M}}$ tangent to manifold $\mathcal{M}(\rho)$ at $B \in \mathcal{M}(\rho)$ [165, §5, prop.1.1]

$$\mathcal{R}_{\mathcal{M}}(B) \triangleq \{XB + BX^T \mid X \in \mathbb{R}^{M \times M}\} \subseteq \mathbb{S}^M \quad (202)$$

has dimension

$$\dim \text{svec } \mathcal{R}_{\mathcal{M}}(B) = \rho \left(M - \frac{\rho - 1}{2} \right) = \rho(M - \rho) + \frac{\rho(\rho + 1)}{2} \quad (203)$$

Tangent subspace $\mathcal{R}_{\mathcal{M}}$ contains no member of the positive semidefinite cone \mathbb{S}_+^M whose rank exceeds ρ .

Subspace $\mathcal{R}_{\mathcal{M}}(B)$ is a supporting hyperplane when $B \in \mathcal{M}(M-1)$. Another good example of tangent subspace is given in §E.7.2.0.2 by (1887); $\mathcal{R}_{\mathcal{M}}(\mathbf{1}\mathbf{1}^T) = \mathbb{S}_c^{M\perp}$, orthogonal complement to the *geometric center subspace*. (Figure 127, p.484)

2.9.2.3 Faces of PSD cone, their dimension *versus* rank

Each and every face of the positive semidefinite cone, having dimension less than that of the cone, is exposed. [217, §6] [190, §2.3.4] Because each and every face of the positive semidefinite cone contains the origin (§2.8.0.0.1), each face belongs to a subspace of the same dimension.

Given positive semidefinite matrix $A \in \mathbb{S}_+^M$, define $\mathcal{F}(\mathbb{S}_+^M \ni A)$ (152) as the smallest face that contains A of the positive semidefinite cone \mathbb{S}_+^M . Then matrix A , having ordered diagonalization $Q\Lambda Q^T$ (§A.5.2), is relatively interior to [25, §II.12] [96, §31.5.3] [211, §2.4] [212] (§A.7.4.0.1)

$$\begin{aligned} \mathcal{F}(\mathbb{S}_+^M \ni A) &= \{X \in \mathbb{S}_+^M \mid \mathcal{N}(X) \supseteq \mathcal{N}(A)\} \\ &= \{X \in \mathbb{S}_+^M \mid \langle Q(I - \Lambda\Lambda^\dagger)Q^T, X \rangle = 0\} \\ &\simeq \mathbb{S}_+^{\text{rank } A} \end{aligned} \quad (204)$$

which is isomorphic with the convex cone $\mathbb{S}_+^{\text{rank } A}$. Thus dimension of the smallest face containing given matrix A is

$$\dim \mathcal{F}(\mathbb{S}_+^M \ni A) = \text{rank}(A)(\text{rank}(A) + 1)/2 \quad (205)$$

in isomorphic $\mathbb{R}^{M(M+1)/2}$, and each and every face of \mathbb{S}_+^M is isomorphic with a positive semidefinite cone having dimension the same as the face. Observe: not all dimensions are represented, and the only zero-dimensional face is the origin. The positive semidefinite cone has no facets, for example.

2.9.2.3.1 Table: Rank k versus dimension of \mathbb{S}_+^3 faces

	k	$\dim \mathcal{F}(\mathbb{S}_+^3 \ni \text{rank-}k \text{ matrix})$
boundary	0	0
	1	1
	2	3
interior	3	6

For the positive semidefinite cone \mathbb{S}_+^2 in isometrically isomorphic \mathbb{R}^3 depicted in Figure 39, for example, rank-2 matrices belong to the interior of that face having dimension 3 (the entire closed cone), rank-1 matrices belong to relative interior of a face having dimension 1, ^{2.36} and the only rank-0 matrix is the point at the origin (the zero-dimensional face).

Any *simultaneously diagonalizable* positive semidefinite rank- k matrices belong to the same face (204). That observation leads to the following hyperplane characterization of PSD cone faces: Any rank- $k < M$ positive semidefinite matrix A belongs to a face, of the positive semidefinite cone, described by an intersection with a hyperplane: for $A = Q\Lambda Q^T$ and $0 \leq k < M$

$$\begin{aligned} \mathcal{F}(\mathbb{S}_+^M \ni A \ni \text{rank}(A) = k) &= \{X \in \mathbb{S}_+^M \mid \langle Q(I - \Lambda\Lambda^\dagger)Q^T, X \rangle = 0\} \\ &= \left\{ X \in \mathbb{S}_+^M \mid \left\langle Q \left(I - \begin{bmatrix} I \in \mathbb{S}^k & \mathbf{0} \\ \mathbf{0}^T & \mathbf{0} \end{bmatrix} \right) Q^T, X \right\rangle = 0 \right\} \\ &= \mathbb{S}_+^M \cap \underline{\partial\mathcal{H}}_+ \end{aligned} \quad (206)$$

Faces are doubly indexed: continuously indexed by orthogonal matrix Q , and discretely indexed by rank k . Each and every orthogonal matrix Q

^{2.36}The boundary constitutes all the one-dimensional faces, in \mathbb{R}^3 , which are rays emanating from the origin.

makes projectors $Q(:, k+1:M)Q(:, k+1:M)^T$ indexed by k , in other words, each projector describing a normal $\text{svec}(Q(:, k+1:M)Q(:, k+1:M)^T)$ to a supporting hyperplane $\underline{\partial\mathcal{H}}_+$ (containing the origin) exposing a face (§2.11) of the positive semidefinite cone containing only rank- k matrices.

2.9.2.4 Extreme directions of positive semidefinite cone

Because the positive semidefinite cone is pointed (§2.7.2.1.2), there is a one-to-one correspondence of one-dimensional faces with extreme directions in any dimension M ; *id est*, because of the *cone faces lemma* (§2.8.0.0.1) and the direct correspondence of exposed faces to faces of \mathbb{S}_+^M , it follows there is no one-dimensional face of the positive semidefinite cone that is not a ray emanating from the origin.

Symmetric dyads constitute the set of all extreme directions: For $M > 1$

$$\{yy^T \in \mathbb{S}^M \mid y \in \mathbb{R}^M\} \subset \partial\mathbb{S}_+^M \quad (207)$$

this superset of extreme directions (infinite in number, *confer* (170)) for the positive semidefinite cone is, generally, a subset of the boundary. By the *extremes theorem* (§2.8.1.1.1), the convex hull of extreme rays and the origin is the PSD cone:

$$\text{conv}\{yy^T \in \mathbb{S}^M \mid y \in \mathbb{R}^M\} = \mathbb{S}_+^M \quad (208)$$

For two-dimensional matrices ($M=2$, Figure 39)

$$\{yy^T \in \mathbb{S}^2 \mid y \in \mathbb{R}^2\} = \partial\mathbb{S}_+^2 \quad (209)$$

while for one-dimensional matrices, in exception, ($M=1$, §2.7)

$$\{yy \in \mathbb{S} \mid y \neq \mathbf{0}\} = \text{int } \mathbb{S}_+ \quad (210)$$

Each and every extreme direction yy^T makes the same angle with the identity matrix in isomorphic $\mathbb{R}^{M(M+1)/2}$, dependent only on dimension; *videlicet*,^{2.37}

$$\sphericalangle(yy^T, I) = \arccos \frac{\langle yy^T, I \rangle}{\|yy^T\|_F \|I\|_F} = \arccos \left(\frac{1}{\sqrt{M}} \right) \quad \forall y \in \mathbb{R}^M \quad (211)$$

^{2.37}Analogy with respect to the *EDM cone* is considered in [164, p.162] where it is found: angle is not constant. Extreme directions of the EDM cone can be found in §6.5.3.2. The cone's axis is $-E = \mathbf{1}\mathbf{1}^T - I$ (1010).

2.9.2.4.1 Example. *Positive semidefinite matrix from extreme directions.* Diagonalizability (§A.5) of symmetric matrices yields the following results:

Any symmetric positive semidefinite matrix (1363) can be written in the form

$$A = \sum_i \lambda_i z_i z_i^T = \hat{A} \hat{A}^T = \sum_i \hat{a}_i \hat{a}_i^T \succeq 0, \quad \lambda \succeq 0 \quad (212)$$

a conic combination of linearly independent extreme directions ($\hat{a}_i \hat{a}_i^T$ or $z_i z_i^T$ where $\|z_i\|=1$), where λ is a vector of eigenvalues.

If we limit consideration to all symmetric positive semidefinite matrices bounded via unity trace

$$\mathcal{C} \triangleq \{A \succeq 0 \mid \text{tr } A = 1\} \quad (213)$$

then any matrix A from that set may be expressed as a convex combination of linearly independent extreme directions;

$$A = \sum_i \lambda_i z_i z_i^T \in \mathcal{C}, \quad \mathbf{1}^T \lambda = 1, \quad \lambda \succeq 0 \quad (214)$$

Implications are:

1. set \mathcal{C} is convex (an intersection of PSD cone with hyperplane),
2. because the set of eigenvalues corresponding to a given square matrix A is unique (§A.5.0.1), no single eigenvalue can exceed 1; *id est*, $I \succeq A$
3. the converse holds: set \mathcal{C} is an instance of Fantope (84). \square

2.9.2.5 Positive semidefinite cone is generally not circular

Extreme angle equation (211) suggests that the positive semidefinite cone might be invariant to rotation about its axis of revolution; *id est*, a circular cone. We investigate this now:

2.9.2.5.1 Definition. *Circular cone:*^{2.38}

a pointed closed convex cone having hyperspherical sections orthogonal to its *axis of revolution* about which the cone is invariant to rotation. \triangle

^{2.38}A circular cone is assumed convex throughout, although not so by other authors. We also assume a *right* circular cone.




Figure 42: This circular cone continues upward infinitely. Axis of revolution is illustrated as vertical line segment through origin. R represents radius: distance measured from an extreme direction to axis of revolution. Were this a Lorentz cone, any plane slice containing axis of revolution would make a right angle.

A *conic section* is the intersection of a cone with any hyperplane. In three dimensions, an intersecting plane perpendicular to a circular cone's axis of revolution produces a section bounded by a circle. (Figure 42) A prominent example of a circular cone in convex analysis is the Lorentz cone (161). We also find that the positive semidefinite cone and cone of Euclidean distance matrices are circular cones, but only in low dimension.

The positive semidefinite cone has axis of revolution that is the ray (base $\mathbf{0}$) through the identity matrix I . Consider a set of normalized extreme directions of the positive semidefinite cone: for some arbitrary positive constant $a \in \mathbb{R}_+$

$$\{yy^T \in \mathbb{S}^M \mid \|y\| = \sqrt{a}\} \subset \partial\mathbb{S}_+^M \quad (215)$$

The distance from each extreme direction to the axis of revolution is radius

$$R \triangleq \inf_c \|yy^T - cI\|_F = a\sqrt{1 - \frac{1}{M}} \quad (216)$$

which is the distance from yy^T to $\frac{a}{M}I$; the length of vector $yy^T - \frac{a}{M}I$.

Because distance R (in a particular dimension) from the axis of revolution to each and every normalized extreme direction is identical, the extreme directions lie on the boundary of a hypersphere in isometrically isomorphic $\mathbb{R}^{M(M+1)/2}$. From Example 2.9.2.4.1, the convex hull (excluding the vertex at the origin) of the normalized extreme directions is a conic section

$$\mathcal{C} \triangleq \text{conv}\{yy^T \mid y \in \mathbb{R}^M, y^T y = a\} = \mathbb{S}_+^M \cap \{A \in \mathbb{S}^M \mid \langle I, A \rangle = a\} \quad (217)$$

orthogonal to identity matrix I ;

$$\langle \mathcal{C} - \frac{a}{M}I, I \rangle = \text{tr}(\mathcal{C} - \frac{a}{M}I) = 0 \quad (218)$$

Proof. Although the positive semidefinite cone possesses some characteristics of a circular cone, we can show it is not by demonstrating shortage of extreme directions; *id est*, some extreme directions corresponding to each and every angle of rotation about the axis of revolution are nonexistent: Referring to Figure 43, [339, §1-7]

$$\cos \theta = \frac{\langle \frac{a}{M}\mathbf{1}\mathbf{1}^T - \frac{a}{M}I, yy^T - \frac{a}{M}I \rangle}{a^2(1 - \frac{1}{M})} \quad (219)$$




Figure 43: Illustrated is a section, perpendicular to axis of revolution, of circular cone from Figure 42. Radius R is distance from any extreme direction to axis at $\frac{a}{M}I$. Vector $\frac{a}{M}\mathbf{1}\mathbf{1}^T$ is an arbitrary reference by which to measure angle θ .

Solving for vector y we get

$$a(1 + (M - 1) \cos \theta) = (\mathbf{1}^T y)^2 \quad (220)$$

which does not have real solution $\forall 0 \leq \theta \leq 2\pi$ in every matrix dimension M .



From the foregoing proof we can conclude that the positive semidefinite cone might be circular but only in matrix dimensions 1 and 2.^{2.39} Because of a shortage of extreme directions, conic section (217) cannot be hyperspherical by the *extremes theorem* (§2.8.1.1.1, Figure 38).

^{2.39}In fact, the positive semidefinite cone is circular in matrix dimensions 1 and 2 while it is a rotation of the Lorentz cone in matrix dimension 2.




Figure 44: Proper polyhedral cone \mathcal{K} , created by intersection of halfspaces, inscribes PSD cone in isometrically isomorphic \mathbb{R}^3 as predicted by *Geršgorin discs theorem* for $A=[A_{ij}] \in \mathbb{S}^2$. Hyperplanes supporting \mathcal{K} intersect along boundary of PSD cone. Four extreme directions of \mathcal{K} coincide with extreme directions of PSD cone.

2.9.2.5.2 Example. *PSD cone inscription in three dimensions.*

Theorem. *Geršgorin discs.* [180, §6.1] [324] [225, p.140]

For $p \in \mathbb{R}_+^m$ given $A = [A_{ij}] \in \mathbb{S}^m$, then all eigenvalues of A belong to the union of m closed intervals on the real line;

$$\lambda(A) \in \bigcup_{i=1}^m \left\{ \xi \in \mathbb{R} \mid |\xi - A_{ii}| \leq \varrho_i \triangleq \frac{1}{p_i} \sum_{\substack{j=1 \\ j \neq i}}^m p_j |A_{ij}| \right\} = \bigcup_{i=1}^m [A_{ii} - \varrho_i, A_{ii} + \varrho_i] \quad (221)$$

Furthermore, if a union of k of these m [intervals] forms a connected region that is disjoint from all the remaining $n - k$ [intervals], then there are precisely k eigenvalues of A in this region. \diamond

To apply the theorem to determine positive semidefiniteness of symmetric matrix A , we observe that for each i we must have

$$A_{ii} \geq \varrho_i \quad (222)$$

Suppose

$$m = 2 \quad (223)$$

so $A \in \mathbb{S}^2$. Vectorizing A as in (50), $\text{svec } A$ belongs to isometrically isomorphic \mathbb{R}^3 . Then we have $m2^{m-1} = 4$ inequalities, in the matrix entries A_{ij} with Geršgorin parameters $p = [p_i] \in \mathbb{R}_+^2$,

$$\begin{aligned} p_1 A_{11} &\geq \pm p_2 A_{12} \\ p_2 A_{22} &\geq \pm p_1 A_{12} \end{aligned} \quad (224)$$

which describe an intersection of four halfspaces in $\mathbb{R}^{m(m+1)/2}$. That intersection creates the proper polyhedral cone \mathcal{K} (§2.12.1) whose construction is illustrated in Figure 44. Drawn truncated is the boundary of the positive semidefinite cone $\text{svec } \mathbb{S}_+^2$ and the bounding hyperplanes supporting \mathcal{K} .

Created by means of Geršgorin discs, \mathcal{K} always belongs to the positive semidefinite cone for any nonnegative value of $p \in \mathbb{R}_+^m$. Hence any point in \mathcal{K} corresponds to some positive semidefinite matrix A . Only the extreme directions of \mathcal{K} intersect the positive semidefinite cone boundary in this dimension; the four extreme directions of \mathcal{K} are extreme directions of the

positive semidefinite cone. As p_1/p_2 increases in value from 0, two extreme directions of \mathcal{K} sweep the entire boundary of this positive semidefinite cone. Because the entire positive semidefinite cone can be swept by \mathcal{K} , the system of linear inequalities

$$Y^T \text{svec } A \triangleq \begin{bmatrix} p_1 & \pm p_2/\sqrt{2} & 0 \\ 0 & \pm p_1/\sqrt{2} & p_2 \end{bmatrix} \text{svec } A \succeq 0 \quad (225)$$

when made dynamic can replace a semidefinite constraint $A \succeq 0$; *id est*, for

$$\mathcal{K} = \{z \mid Y^T z \succeq 0\} \subset \text{svec } \mathbb{S}_+^m \quad (226)$$

given p where $Y \in \mathbb{R}^{m(m+1)/2 \times m 2^{m-1}}$

$$\text{svec } A \in \mathcal{K} \Rightarrow A \in \mathbb{S}_+^m \quad (227)$$

but

$$\exists p \ni Y^T \text{svec } A \succeq 0 \Leftrightarrow A \succeq 0 \quad (228)$$

In other words, *diagonal dominance* [180, p.349, §7.2.3]

$$A_{ii} \geq \sum_{\substack{j=1 \\ j \neq i}}^m |A_{ij}|, \quad \forall i = 1 \dots m \quad (229)$$

is only a sufficient condition for membership to the PSD cone; but by dynamic weighting p in this dimension, it was made necessary and sufficient. \square

In higher dimension ($m > 2$), boundary of the positive semidefinite cone is no longer constituted completely by its extreme directions (symmetric rank-one matrices); the geometry becomes complicated. How all the extreme directions can be swept by an inscribed polyhedral cone,^{2.40} similarly to the foregoing example, remains an open question.

2.9.2.5.3 Exercise. *Dual inscription.*

Find dual proper polyhedral cone \mathcal{K}^* from Figure 44. \blacktriangledown

^{2.40}It is not necessary to sweep the entire boundary in higher dimension.

2.9.2.6 Boundary constituents of the positive semidefinite cone

2.9.2.6.1 Lemma. *Sum of positive semidefinite matrices.*

For $A, B \in \mathbb{S}_+^M$

$$\text{rank}(A + B) = \text{rank}(\mu A + (1 - \mu)B) \quad (230)$$

over the open interval $(0, 1)$ of μ . \diamond

Proof. Any positive semidefinite matrix belonging to the PSD cone has an eigen decomposition that is a positively scaled sum of linearly independent symmetric dyads. By the *linearly independent dyads definition* in §B.1.1.0.1, rank of the sum $A + B$ is equivalent to the number of linearly independent dyads constituting it. Linear independence is insensitive to further positive scaling by μ . The assumption of positive semidefiniteness prevents annihilation of any dyad from the sum $A + B$. \blacklozenge

2.9.2.6.2 Example. *Rank function quasiconcavity.* (confer §3.3)

For $A, B \in \mathbb{R}^{m \times n}$ [180, §0.4]

$$\text{rank } A + \text{rank } B \geq \text{rank}(A + B) \quad (231)$$

that follows from the fact [294, §3.6]

$$\dim \mathcal{R}(A) + \dim \mathcal{R}(B) = \dim \mathcal{R}(A + B) + \dim(\mathcal{R}(A) \cap \mathcal{R}(B)) \quad (232)$$

For $A, B \in \mathbb{S}_+^M$

$$\text{rank } A + \text{rank } B \geq \text{rank}(A + B) \geq \min\{\text{rank } A, \text{rank } B\} \quad (1379)$$

that follows from the fact

$$\mathcal{N}(A + B) = \mathcal{N}(A) \cap \mathcal{N}(B), \quad A, B \in \mathbb{S}_+^M \quad (147)$$

Rank is a *quasiconcave* function on \mathbb{S}_+^M because the right-hand inequality in (1379) has the concave form (586); *videlicet*, Lemma 2.9.2.6.1. \square

From this example we see, unlike convex functions, *quasiconvex* functions are not necessarily continuous. (§3.3) We also glean:

2.9.2.6.3 Theorem. *Convex subsets of positive semidefinite cone.*
Subsets of the positive semidefinite cone \mathbb{S}_+^M , for $0 \leq \rho \leq M$

$$\mathbb{S}_+^M(\rho) \triangleq \{X \in \mathbb{S}_+^M \mid \text{rank } X \geq \rho\} \quad (233)$$

are pointed convex cones, but not closed unless $\rho = 0$; *id est*, $\mathbb{S}_+^M(0) = \mathbb{S}_+^M$.
◇

Proof. Given ρ , a subset $\mathbb{S}_+^M(\rho)$ is convex if and only if convex combination of any two members has rank at least ρ . That is confirmed applying identity (230) from Lemma 2.9.2.6.1 to (1379); *id est*, for $A, B \in \mathbb{S}_+^M(\rho)$ on the closed interval $\mu \in [0, 1]$

$$\text{rank}(\mu A + (1 - \mu)B) \geq \min\{\text{rank } A, \text{rank } B\} \quad (234)$$

It can similarly be shown, almost identically to proof of the lemma, any conic combination of A, B in subset $\mathbb{S}_+^M(\rho)$ remains a member; *id est*, $\forall \zeta, \xi \geq 0$

$$\text{rank}(\zeta A + \xi B) \geq \min\{\text{rank}(\zeta A), \text{rank}(\xi B)\} \quad (235)$$

Therefore, $\mathbb{S}_+^M(\rho)$ is a convex cone. ◆

Another proof of convexity can be made by projection arguments:

2.9.2.7 Projection on $\mathbb{S}_+^M(\rho)$

Because these cones $\mathbb{S}_+^M(\rho)$ indexed by ρ (233) are convex, projection on them is straightforward. Given a symmetric matrix H having diagonalization $H \triangleq Q\Lambda Q^T \in \mathbb{S}^M$ (§A.5.2) with eigenvalues Λ arranged in nonincreasing order, then its *Euclidean projection* (minimum-distance projection) on $\mathbb{S}_+^M(\rho)$

$$P_{\mathbb{S}_+^M(\rho)} H = Q\Upsilon^* Q^T \quad (236)$$

corresponds to a map of its eigenvalues:

$$\Upsilon_{ii}^* = \begin{cases} \max\{\epsilon, \Lambda_{ii}\}, & i = 1 \dots \rho \\ \max\{0, \Lambda_{ii}\}, & i = \rho + 1 \dots M \end{cases} \quad (237)$$

where ϵ is positive but arbitrarily close to 0.

2.9.2.7.1 Exercise. *Projection on open convex cones.*

Prove (237) using Theorem E.9.2.0.1. ▼

Because each $H \in \mathbb{S}^M$ has unique projection on $\mathbb{S}_+^M(\rho)$ (despite possibility of repeated eigenvalues in Λ), we may conclude it is a convex set by the *Bunt-Motzkin theorem* (§E.9.0.0.1).

Compare (237) to the well-known result regarding Euclidean projection on a rank ρ subset of the positive semidefinite cone (§2.9.2.1)

$$\mathbb{S}_+^M \setminus \mathbb{S}_+^M(\rho+1) = \{X \in \mathbb{S}_+^M \mid \text{rank } X \leq \rho\} \quad (238)$$

$$P_{\mathbb{S}_+^M \setminus \mathbb{S}_+^M(\rho+1)} H = Q \Upsilon^* Q^T \quad (239)$$

As proved in §7.1.4, this projection of H corresponds to the eigenvalue map

$$\Upsilon_{ii}^* = \begin{cases} \max\{0, \Lambda_{ii}\}, & i=1 \dots \rho \\ 0, & i=\rho+1 \dots M \end{cases} \quad (1256)$$

Together these two results (237) and (1256) mean: A higher-rank solution to projection on the positive semidefinite cone lies arbitrarily close to any given lower-rank projection, but not *vice versa*. Were the number of nonnegative eigenvalues in Λ known *a priori* not to exceed ρ , then these two different projections would produce identical results in the limit $\epsilon \rightarrow 0$.

2.9.2.8 Uniting constituents

Interior of the PSD cone $\text{int } \mathbb{S}_+^M$ is convex by Theorem 2.9.2.6.3, for example, because all positive semidefinite matrices having rank M constitute the cone interior.

All positive semidefinite matrices of rank less than M constitute the cone boundary; an amalgam of positive semidefinite matrices of different rank. Thus each nonconvex subset of positive semidefinite matrices, for $0 < \rho < M$

$$\{Y \in \mathbb{S}_+^M \mid \text{rank } Y = \rho\} \quad (240)$$

having rank ρ successively 1 lower than M , appends a nonconvex constituent to the cone boundary; but only in their union is the boundary complete: (*confer* §2.9.2)

$$\partial \mathbb{S}_+^M = \bigcup_{\rho=0}^{M-1} \{Y \in \mathbb{S}_+^M \mid \text{rank } Y = \rho\} \quad (241)$$

The composite sequence, the cone interior in union with each successive constituent, remains convex at each step; *id est*, for $0 \leq k \leq M$

$$\bigcup_{\rho=k}^M \{Y \in \mathbb{S}_+^M \mid \text{rank } Y = \rho\} \quad (242)$$

is convex for each k by Theorem 2.9.2.6.3.

2.9.2.9 Peeling constituents

Proceeding the other way: To peel constituents off the complete positive semidefinite cone boundary, one starts by removing the origin; the only rank-0 positive semidefinite matrix. What remains is convex. Next, the extreme directions are removed because they constitute all the rank-1 positive semidefinite matrices. What remains is again convex, and so on. Proceeding in this manner eventually removes the entire boundary leaving, at last, the convex interior of the PSD cone; all the positive definite matrices.

2.9.2.9.1 Exercise. *Difference* $A - B$.

What about the difference of matrices A, B belonging to the positive semidefinite cone? Show:

- The difference of any two points on the boundary belongs to the boundary or exterior.
- The difference $A - B$, where A belongs to the boundary while B is interior, belongs to the exterior. ▼

2.9.3 Barvinok's proposition

Barvinok posits existence and quantifies an upper bound on rank of a positive semidefinite matrix belonging to the intersection of the PSD cone with an affine subset:

2.9.3.0.1 Proposition. (Barvinok) *Affine intersection with PSD cone.* [25, §II.13] [23, §2.2] Consider finding a matrix $X \in \mathbb{S}^N$ satisfying

$$X \succeq 0, \quad \langle A_j, X \rangle = b_j, \quad j = 1 \dots m \quad (243)$$

given nonzero linearly independent $A_j \in \mathbb{S}^N$ and real b_j . Define the affine subset

$$\mathcal{A} \triangleq \{X \mid \langle A_j, X \rangle = b_j, j=1 \dots m\} \subseteq \mathbb{S}^N \quad (244)$$

If the intersection $\mathcal{A} \cap \mathbb{S}_+^N$ is nonempty given a number m of equalities, then there exists a matrix $X \in \mathcal{A} \cap \mathbb{S}_+^N$ such that

$$\text{rank } X (\text{rank } X + 1)/2 \leq m \quad (245)$$

whence the upper bound^{2.41}

$$\text{rank } X \leq \left\lfloor \frac{\sqrt{8m+1} - 1}{2} \right\rfloor \quad (246)$$

Given desired rank instead, equivalently,

$$m < (\text{rank } X + 1)(\text{rank } X + 2)/2 \quad (247)$$

An extreme point of $\mathcal{A} \cap \mathbb{S}_+^N$ satisfies (246) and (247). (confer §4.1.1.3)

A matrix $X \triangleq R^T R$ is an extreme point if and only if the smallest face that contains X of $\mathcal{A} \cap \mathbb{S}_+^N$ has dimension 0; [211, §2.4] [212] *id est*, iff (152)

$$\dim \mathcal{F}((\mathcal{A} \cap \mathbb{S}_+^N) \ni X) \quad (248)$$

$$= \text{rank}(X)(\text{rank}(X) + 1)/2 - \text{rank}[\text{svec } RA_1 R^T \text{ svec } RA_2 R^T \dots \text{svec } RA_m R^T]$$

equals 0 in isomorphic $\mathbb{R}^{N(N+1)/2}$.

Now the intersection $\mathcal{A} \cap \mathbb{S}_+^N$ is assumed bounded: Assume a given nonzero upper bound ρ on rank, a number of equalities

$$m = (\rho + 1)(\rho + 2)/2 \quad (249)$$

and matrix dimension $N \geq \rho + 2 \geq 3$. If the intersection is nonempty and bounded, then there exists a matrix $X \in \mathcal{A} \cap \mathbb{S}_+^N$ such that

$$\text{rank } X \leq \rho \quad (250)$$

This represents a *tightening* of the upper bound; a reduction by exactly 1 of the bound provided by (246) given the same specified number m (249) of equalities; *id est*,

$$\text{rank } X \leq \frac{\sqrt{8m+1} - 1}{2} - 1 \quad (251)$$

◇

^{2.41} §4.1.1.3 contains an intuitive explanation. This bound is itself limited above, of course, by N ; a *tight* limit corresponding to an interior point of \mathbb{S}_+^N .

When the intersection $\mathcal{A} \cap \mathbb{S}_+^N$ is known *a priori* to consist only of a single point, then Barvinok's proposition provides the greatest upper bound on its rank not exceeding N . The intersection can be a single nonzero point only if the number of linearly independent hyperplanes m constituting \mathcal{A} satisfies^{2.42}

$$N(N+1)/2 - 1 \leq m \leq N(N+1)/2 \quad (252)$$

2.10 Conic independence (c.i.)

In contrast to extreme direction, the property *conically independent direction* is more generally applicable; inclusive of all closed convex cones (not only pointed closed convex cones). Arbitrary given directions $\{\Gamma_i \in \mathbb{R}^n, i=1 \dots N\}$ are *conically independent* if and only if, for all $\zeta \in \mathbb{R}_+^N$ (confer §2.1.2, §2.4.2.3)

$$\Gamma_i \zeta_i + \dots + \Gamma_j \zeta_j - \Gamma_\ell = \mathbf{0}, \quad i \neq \dots \neq j \neq \ell = 1 \dots N \quad (253)$$

has no solution ζ ; in words, iff no direction from the given set can be expressed as a conic combination of those remaining. (Figure 45, for example. MATLAB implementation of test (253) is given on *Wikimization*.) It is evident that linear independence (l.i.) of N directions implies their conic independence;

- l.i. \Rightarrow c.i.

Arranging any set of generators for a particular convex cone in a matrix columnar,

$$X \triangleq [\Gamma_1 \ \Gamma_2 \ \dots \ \Gamma_N] \in \mathbb{R}^{n \times N} \quad (254)$$

then the relationship l.i. \Rightarrow c.i. suggests: the number of l.i. generators in the columns of X cannot exceed the number of c.i. generators. Denoting by \mathbf{k} the number of conically independent generators contained in X , we have the most fundamental rank inequality for convex cones

$$\dim \text{aff } \mathcal{K} = \dim \text{aff}[\mathbf{0} \ X] = \text{rank } X \leq \mathbf{k} \leq N \quad (255)$$

Whereas N directions in n dimensions can no longer be linearly independent once N exceeds n , conic independence remains possible:

^{2.42}For $N > 1$, $N(N+1)/2 - 1$ independent hyperplanes in $\mathbb{R}^{N(N+1)/2}$ can make a line




Figure 45: Vectors in \mathbb{R}^2 : (a) affinely and conically independent, (b) affinely independent but not conically independent, (c) conically independent but not affinely independent. None of the examples exhibits linear independence. (In general, a.i. $\not\Rightarrow$ c.i.)

2.10.0.0.1 Table: Maximum number of c.i. directions

dimension n	sup \mathbf{k} (pointed)	sup \mathbf{k} (not pointed)
0	0	0
1	1	2
2	2	4
3	∞	∞
\vdots	\vdots	\vdots

Assuming veracity of this table, there is an apparent vastness between two and three dimensions. The finite numbers of conically independent directions indicate:

- Convex cones in dimensions 0, 1, and 2 must be polyhedral. (§2.12.1)

Conic independence is certainly one convex idea that cannot be completely explained by a two-dimensional picture as Barvinok suggests [25, p.vii].

From this table it is also evident that dimension of Euclidean space cannot exceed the number of conically independent directions possible;

- $n \leq \text{sup } \mathbf{k}$

tangent to $\text{svec } \partial \mathbb{S}_+^N$ at a point because all one-dimensional faces of \mathbb{S}_+^N are exposed. Because a pointed convex cone has only one vertex, the origin, there can be no intersection of $\text{svec } \partial \mathbb{S}_+^N$ with any higher-dimensional affine subset \mathcal{A} that will make a nonzero point.




Figure 46: **(a)** A pointed polyhedral cone (drawn truncated) in \mathbb{R}^3 having six facets. The extreme directions, corresponding to six edges emanating from the origin, are generators for this cone; not linearly independent but they must be conically independent. **(b)** The boundary of dual cone \mathcal{K}^* (drawn truncated) is now added to the drawing of same \mathcal{K} . \mathcal{K}^* is polyhedral, proper, and has the same number of extreme directions as \mathcal{K} has facets.

2.10.1 Preservation of conic independence

Independence in the linear (§2.1.2.1), affine (§2.4.2.4), and conic senses can be preserved under linear transformation. Suppose a matrix $X \in \mathbb{R}^{n \times N}$ (254) holds a conically independent set columnar. Consider a transformation on the domain of such matrices

$$T(X) : \mathbb{R}^{n \times N} \rightarrow \mathbb{R}^{n \times N} \triangleq XY \quad (256)$$

where fixed matrix $Y \triangleq [y_1 \ y_2 \ \cdots \ y_N] \in \mathbb{R}^{N \times N}$ represents linear operator T . Conic independence of $\{Xy_i \in \mathbb{R}^n, i=1 \dots N\}$ demands, by definition (253),

$$Xy_i \zeta_i + \cdots + Xy_j \zeta_j - Xy_\ell = \mathbf{0}, \quad i \neq \cdots \neq j \neq \ell = 1 \dots N \quad (257)$$

have no solution $\zeta \in \mathbb{R}_+^N$. That is ensured by conic independence of $\{y_i \in \mathbb{R}^N\}$ and by $\mathcal{R}(Y) \cap \mathcal{N}(X) = \mathbf{0}$; seen by factoring out X .

2.10.1.1 linear maps of cones

[20, §7] If \mathcal{K} is a convex cone in Euclidean space \mathcal{R} and T is any linear mapping from \mathcal{R} to Euclidean space \mathcal{M} , then $T(\mathcal{K})$ is a convex cone in \mathcal{M} and $x \preceq y$ with respect to \mathcal{K} implies $T(x) \preceq T(y)$ with respect to $T(\mathcal{K})$. If \mathcal{K} is closed or has nonempty interior in \mathcal{R} , then so is $T(\mathcal{K})$ in \mathcal{M} .

If T is a linear bijection, then $x \preceq y \Leftrightarrow T(x) \preceq T(y)$. Further, if \mathcal{F} is a face of \mathcal{K} , then $T(\mathcal{F})$ is a face of $T(\mathcal{K})$.

2.10.2 Pointed closed convex \mathcal{K} & conic independence

The following bullets can be derived from definitions (169) and (253) in conjunction with the *extremes theorem* (§2.8.1.1.1):

The set of all extreme directions from a pointed closed convex cone $\mathcal{K} \subset \mathbb{R}^n$ is not necessarily a linearly independent set, yet it must be a conically independent set; (compare Figure 22 on page 67 with Figure 46(a))

- $\{\text{extreme directions}\} \Rightarrow \{\text{c.i.}\}$

Conversely, when a conically independent set of directions from pointed closed convex cone \mathcal{K} is known to comprise generators, then all directions from that set must be extreme directions of the cone;

- $\{\text{extreme directions}\} \Leftrightarrow \{\text{c.i. generators of pointed closed convex } \mathcal{K}\}$

Barker & Carlson [20, §1] call the extreme directions a *minimal generating set* for a pointed closed convex cone. A minimal set of generators is therefore a conically independent set of generators, and *vice versa*,^{2.43} for a pointed closed convex cone.

An arbitrary collection of n or fewer distinct extreme directions from pointed closed convex cone $\mathcal{K} \subset \mathbb{R}^n$ is not necessarily a linearly independent set; *e.g.*, dual extreme directions (437) from Example 2.13.11.0.3.

- $\{\leq n \text{ extreme directions in } \mathbb{R}^n\} \not\Rightarrow \{\text{l.i.}\}$

Linear dependence of few extreme directions is another convex idea that cannot be explained by a two-dimensional picture as Barvinok suggests [25, p.vii]; indeed, it only first comes to light in four dimensions! But there is a converse: [293, §2.10.9]

- $\{\text{extreme directions}\} \Leftarrow \{\text{l.i. generators of closed convex } \mathcal{K}\}$

2.10.2.0.1 Example. Vertex-description of halfspace \mathcal{H} about origin.

From $n + 1$ points in \mathbb{R}^n we can make a vertex-description of a convex cone that is a halfspace \mathcal{H} , where $\{x_\ell \in \mathbb{R}^n, \ell = 1 \dots n\}$ constitutes a minimal set of generators for a hyperplane $\partial\mathcal{H}$ through the origin. An example is illustrated in Figure 47. By demanding the augmented set $\{x_\ell \in \mathbb{R}^n, \ell = 1 \dots n + 1\}$ be affinely independent (we want vector x_{n+1} not parallel to $\partial\mathcal{H}$), then

$$\begin{aligned} \mathcal{H} &= \bigcup_{\zeta \geq 0} (\zeta x_{n+1} + \partial\mathcal{H}) \\ &= \{\zeta x_{n+1} + \text{cone}\{x_\ell \in \mathbb{R}^n, \ell = 1 \dots n\} \mid \zeta \geq 0\} \\ &= \text{cone}\{x_\ell \in \mathbb{R}^n, \ell = 1 \dots n + 1\} \end{aligned} \quad (258)$$

a union of parallel hyperplanes. Cardinality is one step beyond dimension of the ambient space, but $\{x_\ell \forall \ell\}$ is a minimal set of generators for this convex cone \mathcal{H} which has no extreme elements. \square

2.10.2.0.2 Exercise. Enumerating conically independent directions.

Describe a nonpointed polyhedral cone in three dimensions having more than 8 conically independent generators. (*confer* Table 2.10.0.0.1) \blacktriangledown

^{2.43}This converse does not hold for nonpointed closed convex cones as Table 2.10.0.0.1 implies; *e.g.*, ponder four conically independent generators for a plane ($n=2$, Figure 45).




Figure 47: Minimal set of generators $X = [x_1 \ x_2 \ x_3] \in \mathbb{R}^{2 \times 3}$ (not extreme directions) for halfspace about origin; affinely and conically independent.

2.10.3 Utility of conic independence

Perhaps the most useful application of conic independence is determination of the intersection of closed convex cones from their halfspace-descriptions, or representation of the sum of closed convex cones from their vertex-descriptions.

- $\bigcap \mathcal{K}_i$ A halfspace-description for the intersection of any number of closed convex cones \mathcal{K}_i can be acquired by pruning normals; specifically, only the conically independent normals from the aggregate of all the halfspace-descriptions need be retained.
- $\sum \mathcal{K}_i$ Generators for the sum of any number of closed convex cones \mathcal{K}_i can be determined by retaining only the conically independent generators from the aggregate of all the vertex-descriptions.

Such conically independent sets are not necessarily unique or minimal.

2.11 When extreme means exposed

For any convex polyhedral set in \mathbb{R}^n having nonempty interior, distinction between the terms *extreme* and *exposed* vanishes [293, §2.4] [96, §2.2] for faces of all dimensions except n ; their meanings become equivalent as we saw in Figure 18 (discussed in §2.6.1.2). In other words, each and every face of any polyhedral set (except the set itself) can be exposed by a hyperplane, and *vice versa*; *e.g.*, Figure 22.

Lewis [217, §6] [190, §2.3.4] claims nonempty extreme proper subsets and the exposed subsets coincide for \mathbb{S}_+^n ; *id est*, each and every face of the positive semidefinite cone, whose dimension is less than dimension of the cone, is exposed. A more general discussion of cones having this property can be found in [303]; *e.g.*, Lorentz cone (161) [19, §II.A].

2.12 Convex polyhedra

Every polyhedron, such as the convex hull (79) of a bounded list X , can be expressed as the solution set of a finite system of linear equalities and inequalities, and *vice versa*. [96, §2.2]

2.12.0.0.1 Definition. Convex polyhedra, halfspace-description.

A convex polyhedron is the intersection of a finite number of halfspaces and hyperplanes;

$$\mathcal{P} = \{y \mid Ay \succeq b, Cy = d\} \subseteq \mathbb{R}^n \quad (259)$$

where coefficients A and C generally denote matrices. Each row of C is a vector normal to a hyperplane, while each row of A is a vector inward-normal to a hyperplane partially bounding a halfspace. \triangle

By the *halfspaces theorem* in §2.4.1.1.1, a polyhedron thus described is a closed convex set having possibly empty interior; *e.g.*, Figure 18. Convex polyhedra^{2.44} are finite-dimensional comprising all affine sets (§2.3.1), polyhedral cones, line segments, rays, halfspaces, convex polygons, *solids* [198, def.104/6, p.343], polychora, *polytopes*,^{2.45} *etcetera*.

^{2.44}We consider only convex polyhedra throughout, but acknowledge the existence of concave polyhedra. [333, *Kepler-Poinsot Solid*]

^{2.45}Some authors distinguish bounded polyhedra via the designation *polytope*. [96, §2.2]

It follows from definition (259) by exposure that each face of a convex polyhedron is a convex polyhedron.

The projection of any polyhedron on a subspace remains a polyhedron. More generally, the image of a polyhedron under any linear transformation is a polyhedron. [25, §I.9]

When b and d in (259) are $\mathbf{0}$, the resultant is a polyhedral cone. The set of all polyhedral cones is a subset of convex cones:

2.12.1 Polyhedral cone

From our study of cones, we see: the number of intersecting hyperplanes and halfspaces constituting a convex cone is possibly but not necessarily infinite. When the number is finite, the convex cone is termed *polyhedral*. That is the primary distinguishing feature between the set of all convex cones and polyhedra; all polyhedra, including polyhedral cones, are *finitely generated* [271, §19]. (Figure 48) We distinguish polyhedral cones in the set of all convex cones for this reason, although all convex cones of dimension 2 or less are polyhedral.

2.12.1.0.1 Definition. *Polyhedral cone, halfspace-description.*^{2.46} (*confer* (95)) A polyhedral cone is the intersection of a finite number of halfspaces and hyperplanes about the origin;

$$\begin{aligned} \mathcal{K} &= \{y \mid Ay \succeq 0, Cy = \mathbf{0}\} \subseteq \mathbb{R}^n & (a) \\ &= \{y \mid Ay \succeq 0, Cy \succeq 0, Cy \preceq 0\} & (b) \\ &= \left\{ y \mid \begin{bmatrix} A \\ C \\ -C \end{bmatrix} y \succeq 0 \right\} & (c) \end{aligned} \tag{260}$$

where coefficients A and C generally denote matrices of finite dimension. Each row of C is a vector normal to a hyperplane containing the origin, while each row of A is a vector inward-normal to a hyperplane containing the origin and partially bounding a halfspace. \triangle

^{2.46}Rockafellar [271, §19] proposes affine sets be handled via complementary pairs of affine inequalities; *e.g.*, $Cy \succeq d$ and $Cy \preceq d$ which, when taken together, can present severe difficulties to some interior-point methods of numerical solution.




Figure 48: Polyhedral cones are finitely generated, unbounded, and convex.

A polyhedral cone thus defined is closed, convex, possibly has empty interior, and only a finite number of generators (§2.8.1.2), and *vice versa*. (Minkowski/Weyl) [293, §2.8] [271, thm.19.1]

From the definition it follows that any single hyperplane through the origin, or any halfspace partially bounded by a hyperplane through the origin is a polyhedral cone. The most familiar example of polyhedral cone is any quadrant (or orthant, §2.1.3) generated by Cartesian half-axes. Esoteric examples of polyhedral cone include the point at the origin, any line through the origin, any ray having the origin as base such as the nonnegative real line \mathbb{R}_+ in subspace \mathbb{R} , polyhedral flavors of the (proper) Lorentz cone (287), any subspace, and \mathbb{R}^n . More polyhedral cones are illustrated in Figure 46 and Figure 22.

2.12.2 Vertices of convex polyhedra

By definition, a vertex (§2.6.1.0.1) always lies on the relative boundary of a convex polyhedron. [198, def.115/6, p.358] In Figure 18, each vertex of the polyhedron is located at the intersection of three or more facets, and every edge belongs to precisely two facets [25, §VI.1, p.252]. In Figure 22, the only vertex of that polyhedral cone lies at the origin.

The set of all polyhedral cones is clearly a subset of convex polyhedra and a subset of convex cones (Figure 48). Not all convex polyhedra are bounded; evidently, neither can they all be described by the convex hull of a bounded set of points as defined in (79). Hence a universal vertex-description of polyhedra in terms of that same finite-length list X (69):

2.12.2.0.1 Definition. *Convex polyhedra, vertex-description.*
(confer §2.8.1.1.1) Denote the truncated a -vector

$$a_{i:\ell} = \begin{bmatrix} a_i \\ \vdots \\ a_\ell \end{bmatrix} \quad (261)$$

By discriminating a suitable finite-length *generating list* (or set) arranged columnar in $X \in \mathbb{R}^{n \times N}$, then any particular polyhedron may be described

$$\mathcal{P} = \{Xa \mid a_{1:k}^T \mathbf{1} = 1, a_{m:N} \succeq 0, \{1 \dots k\} \cup \{m \dots N\} = \{1 \dots N\}\} \quad (262)$$

where $0 \leq k \leq N$ and $1 \leq m \leq N+1$. Setting $k=0$ removes the affine equality condition. Setting $m=N+1$ removes the inequality. \triangle

Coefficient indices in (262) may or may not be overlapping, but all the coefficients are assumed constrained. From (71), (79), and (95), we summarize how the coefficient conditions may be applied;

$$\left. \begin{array}{l} \text{affine sets} \quad \longrightarrow \quad a_{1:k}^T \mathbf{1} = 1 \\ \text{polyhedral cones} \quad \longrightarrow \quad a_{m:N} \succeq 0 \end{array} \right\} \longleftarrow \text{convex hull } (m \leq k) \quad (263)$$

It is always possible to describe a convex hull in the region of overlapping indices because, for $1 \leq m \leq k \leq N$

$$\{a_{m:k} \mid a_{m:k}^T \mathbf{1} = 1, a_{m:k} \succeq 0\} \subseteq \{a_{m:k} \mid a_{1:k}^T \mathbf{1} = 1, a_{m:N} \succeq 0\} \quad (264)$$

Members of a generating list are not necessarily vertices of the corresponding polyhedron; certainly true for (79) and (262), some subset of list members reside in the polyhedron's relative interior. Conversely, when boundedness (79) applies, the convex hull of the vertices is a polyhedron identical to the convex hull of the generating list.

2.12.2.1 Vertex-description of polyhedral cone

Given closed convex cone \mathcal{K} in a subspace of \mathbb{R}^n having any set of generators for it arranged in a matrix $X \in \mathbb{R}^{n \times N}$ as in (254), then that cone is described setting $m=1$ and $k=0$ in vertex-description (262):

$$\mathcal{K} = \text{cone } X = \{Xa \mid a \succeq 0\} \subseteq \mathbb{R}^n \quad (95)$$

a conic hull of N generators.^{2.47}

^{2.47}This vertex-description is extensible to an infinite number of generators; which follows, for a pointed cone, from the *extremes theorem* (§2.8.1.1.1) and Example 2.8.1.2.1.

2.12.2.2 Pointedness

[293, §2.10] Assuming all generators constituting the columns of $X \in \mathbb{R}^{n \times N}$ are nonzero, polyhedral cone \mathcal{K} is pointed (§2.7.2.1.2) if and only if there is no nonzero $a \succeq 0$ that solves $Xa = \mathbf{0}$; *id est*, iff

$$\begin{aligned} & \text{find} && a \\ & \text{subject to} && \begin{aligned} Xa &= \mathbf{0} \\ \mathbf{1}^T a &= 1 \\ a &\succeq 0 \end{aligned} \end{aligned} \tag{265}$$

is infeasible or iff $\mathcal{N}(X) \cap \mathbb{R}_+^N = \mathbf{0}$.^{2.48}

A proper polyhedral cone in \mathbb{R}^n must have at least n linearly independent generators, or be the intersection of at least n halfspaces whose partial boundaries have normals that are linearly independent. Otherwise, the cone will contain at least one line and there can be no vertex; *id est*, the cone cannot otherwise be pointed.

Examples of pointed closed convex cones \mathcal{K} are not limited to polyhedral cones: the origin, any $\mathbf{0}$ -based ray in a subspace, any two-dimensional V-shaped cone in a subspace, the Lorentz (ice-cream) cone and its polyhedral flavors, the cone of Euclidean distance matrices EDM^N in \mathbb{S}_h^N , the proper cones: \mathbb{S}_+^M in ambient \mathbb{S}^M , any orthant in \mathbb{R}^n or $\mathbb{R}^{m \times n}$; *e.g.*, the nonnegative real line \mathbb{R}_+ in vector space \mathbb{R} .

2.12.3 Unit simplex

A peculiar subset of the nonnegative orthant with halfspace-description

$$\mathcal{S} \triangleq \{s \mid s \succeq 0, \mathbf{1}^T s \leq 1\} \subseteq \mathbb{R}_+^n \tag{266}$$

is a unique bounded convex polyhedron called *unit simplex* (Figure 49) having nonempty interior, $n + 1$ vertices, and dimension [53, §2.2.4]

$$\dim \mathcal{S} = n \tag{267}$$

The origin supplies one vertex while heads of the *standard basis* [180] [294] $\{e_i, i=1 \dots n\}$ in \mathbb{R}^n constitute those remaining;^{2.49} thus its

^{2.48}If $\text{rank } X = n$, then the dual cone \mathcal{K}^* (§2.13.1) is pointed. (281)

^{2.49}In \mathbb{R}^0 the unit simplex is the point at the origin, in \mathbb{R} the unit simplex is the line segment $[0, 1]$, in \mathbb{R}^2 it is a triangle and its relative interior, in \mathbb{R}^3 it is the convex hull of a tetrahedron (Figure 49), in \mathbb{R}^4 it is the convex hull of a pentatope [333], and so on.




Figure 49: Unit simplex \mathcal{S} in \mathbb{R}^3 is a unique solid tetrahedron but not *regular*.

vertex-description:

$$\begin{aligned} \mathcal{S} &= \text{conv} \{ \mathbf{0}, \{e_i, i=1 \dots n\} \} \\ &= \{ [\mathbf{0} \ e_1 \ e_2 \ \dots \ e_n] a \mid a^T \mathbf{1} = 1, a \succeq 0 \} \end{aligned} \quad (268)$$

2.12.3.1 Simplex

The unit simplex comes from a class of general polyhedra called *simplex*, having vertex-description: [82] [271] [331] [96]

$$\text{conv} \{ x_\ell \in \mathbb{R}^n \mid \ell = 1 \dots k+1, \dim \text{aff} \{ x_\ell \} = k, n \geq k \} \quad (269)$$

So defined, a simplex is a closed bounded convex set having possibly empty interior. Examples of simplices, by increasing affine dimension, are: a point, any line segment, any triangle and its relative interior, a general tetrahedron, polychoron, and so on.

2.12.3.1.1 Definition. *Simplicial cone.*

A proper polyhedral (§2.7.2.2.1) cone \mathcal{K} in \mathbb{R}^n is called *simplicial* iff \mathcal{K} has exactly n extreme directions; [19, §II.A] equivalently, iff proper \mathcal{K} has exactly n linearly independent generators contained in any given set of generators.

△

- simplicial cone \Rightarrow proper polyhedral cone

There are an infinite variety of simplicial cones in \mathbb{R}^n ; *e.g.*, Figure 22, Figure 50, Figure 59. Any orthant is simplicial, as is any rotation thereof.

2.12.4 Converting between descriptions

Conversion between halfspace-descriptions (259) (260) and equivalent vertex-descriptions (79) (262) is nontrivial, in general, [14] [96, §2.2] but the conversion is easy for simplices. [53, §2.2] Nonetheless, we tacitly assume the two descriptions to be equivalent. [271, §19, thm.19.1] We explore conversions in §2.13.4, §2.13.9, and §2.13.11:





Figure 50: Two views of a simplicial cone and its dual in \mathbb{R}^3 (second view on next page). Semi-infinite boundary of each cone is truncated for illustration. Each cone has three facets (*confer* §2.13.11.0.3). Cartesian axes are drawn for reference.



2.13 Dual cone & generalized inequality & biorthogonal expansion

These three concepts, dual cone, generalized inequality, and biorthogonal expansion, are inextricably melded; meaning, it is difficult to completely discuss one without mentioning the others. The dual cone is critical in tests for convergence by contemporary primal/dual methods for numerical solution of conic problems. [350] [244, §4.5] For unique minimum-distance projection on a closed convex cone \mathcal{K} , the negative dual cone $-\mathcal{K}^*$ plays the role that orthogonal complement plays for subspace projection.^{2.50} (§E.9.2.1) Indeed, $-\mathcal{K}^*$ is the *algebraic complement* in \mathbb{R}^n ;

$$\mathcal{K} \boxplus -\mathcal{K}^* = \mathbb{R}^n \quad (1911)$$

where \boxplus denotes unique orthogonal vector sum.

One way to think of a pointed closed convex cone is as a new kind of coordinate system whose basis is generally nonorthogonal; a conic system, very much like the familiar Cartesian system whose analogous cone is the first quadrant (the nonnegative orthant). Generalized inequality $\succeq_{\mathcal{K}}$ is a formalized means to determine membership to any pointed closed convex cone \mathcal{K} (§2.7.2.2) whereas *biorthogonal expansion* is, fundamentally, an expression of coordinates in a pointed conic system whose axes are linearly independent but not necessarily orthogonal. When cone \mathcal{K} is the nonnegative orthant, then these three concepts come into alignment with the Cartesian prototype: biorthogonal expansion becomes orthogonal expansion, the dual cone becomes identical to the orthant, and generalized inequality obeys a total order entrywise.

2.13.1 Dual cone

For any set \mathcal{K} (convex or not), the *dual cone* [92, §4.2]

$$\mathcal{K}^* \triangleq \{y \in \mathbb{R}^n \mid \langle y, x \rangle \geq 0 \text{ for all } x \in \mathcal{K}\} \quad (270)$$

is a unique cone^{2.51} that is always closed and convex because it is an intersection of halfspaces (*halfspaces theorem* (§2.4.1.1.1)) whose partial

^{2.50}Namely, projection on a subspace is ascertainable from its projection on the orthogonal complement.

^{2.51}The dual cone is the negative polar cone defined by many authors; $\mathcal{K}^* = -\mathcal{K}^\circ$. [177, §A.3.2] [271, §14] [36] [25] [293, §2.7]

boundaries each contain the origin, each halfspace having inward-normal x belonging to \mathcal{K} ; *e.g.*, Figure 51(a).

When cone \mathcal{K} is convex, there is a second and equivalent construction: Dual cone \mathcal{K}^* is the union of each and every vector y inward-normal to a hyperplane supporting \mathcal{K} or bounding a halfspace containing \mathcal{K} ; *e.g.*, Figure 51(b). When \mathcal{K} is represented by a halfspace-description such as (260), for example, where

$$A \triangleq \begin{bmatrix} a_1^T \\ \vdots \\ a_m^T \end{bmatrix} \in \mathbb{R}^{m \times n}, \quad C \triangleq \begin{bmatrix} c_1^T \\ \vdots \\ c_p^T \end{bmatrix} \in \mathbb{R}^{p \times n} \quad (271)$$

then the dual cone can be represented as the conic hull

$$\mathcal{K}^* = \text{cone}\{a_1, \dots, a_m, \pm c_1, \dots, \pm c_p\} \quad (272)$$

a vertex-description, because each and every conic combination of normals from the halfspace-description of \mathcal{K} yields another inward-normal to a hyperplane supporting or bounding a halfspace containing \mathcal{K} .

\mathcal{K}^* can also be constructed pointwise using projection theory from §E.9.2: for $P_{\mathcal{K}}x$ the Euclidean projection of point x on closed convex cone \mathcal{K}

$$-\mathcal{K}^* = \{x - P_{\mathcal{K}}x \mid x \in \mathbb{R}^n\} = \{x \in \mathbb{R}^n \mid P_{\mathcal{K}}x = \mathbf{0}\} \quad (1912)$$

2.13.1.0.1 Exercise. Dual cone manual-construction.

Perhaps the most instructive graphical method of dual cone construction is cut-and-try. Find the dual of each polyhedral cone from Figure 52 by using dual cone equation (270). ▼

2.13.1.0.2 Exercise. Dual cone definitions.

What is $\{x \in \mathbb{R}^n \mid x^T z \geq 0 \quad \forall z \in \mathbb{R}^n\}$?

What is $\{x \in \mathbb{R}^n \mid x^T z \geq 1 \quad \forall z \in \mathbb{R}^n\}$?

What is $\{x \in \mathbb{R}^n \mid x^T z \geq 1 \quad \forall z \in \mathbb{R}_+^n\}$? ▼

As defined, dual cone \mathcal{K}^* exists even when the affine hull of the original cone is a proper subspace; *id est*, even when the original cone has empty interior. Rockafellar formulates dimension of \mathcal{K} and \mathcal{K}^* . [271, §14]^{2.52}

^{2.52}His monumental work *Convex Analysis* has not one figure or illustration. See [25, §II.16] for a good illustration of Rockafellar's *recession cone* [37].





Figure 51: Two equivalent constructions of dual cone \mathcal{K}^* in \mathbb{R}^2 : (a) Showing construction by intersection of halfspaces about $\mathbf{0}$ (drawn truncated). Only those two halfspaces whose bounding hyperplanes have inward-normal corresponding to an extreme direction of this pointed closed convex cone $\mathcal{K} \subset \mathbb{R}^2$ need be drawn; by (334). (b) Suggesting construction by union of inward-normals y to each and every hyperplane $\partial\mathcal{H}_+$ supporting \mathcal{K} . This interpretation is valid when \mathcal{K} is convex because existence of a supporting hyperplane is then guaranteed (§2.4.2.6).



$$x \in \mathcal{K} \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } y \in \mathcal{G}(\mathcal{K}^*) \quad (331)$$

Figure 52: Dual cone construction by right angle. Each extreme direction of a proper polyhedral cone is orthogonal to a facet of its dual cone, and *vice versa*, in any dimension. (§2.13.6.1) (a) This characteristic guides graphical construction of dual cone in two dimensions: It suggests finding dual-cone boundary ∂ by making right angles with extreme directions of polyhedral cone. The construction is then pruned so that each dual boundary vector does not exceed $\pi/2$ radians in angle with each and every vector from polyhedral cone. Were dual cone in \mathbb{R}^2 to narrow, Figure 53 would be reached in limit. (b) Same polyhedral cone and its dual continued into three dimensions. (*confer* Figure 59)




Figure 53: \mathcal{K} is a halfspace about the origin in \mathbb{R}^2 . \mathcal{K}^* is a ray base $\mathbf{0}$, hence has empty interior in \mathbb{R}^2 ; so \mathcal{K} cannot be pointed. (Both convex cones appear truncated.)

To further motivate our understanding of the dual cone, consider the ease with which convergence can be observed in the following optimization problem (p):

2.13.1.0.3 Example. *Dual problem.* (confer §4.1)

Duality is a powerful and widely employed tool in applied mathematics for a number of reasons. First, the dual program is always convex even if the primal is not. Second, the number of variables in the dual is equal to the number of constraints in the primal which is often less than the number of variables in the primal program. Third, the maximum value achieved by the dual problem is often equal to the minimum of the primal. [264, §2.1.3]

Essentially, duality theory concerns representation of a given optimization problem as half a *minimax problem*. [271, §36] [53, §5.4] Given any real function $f(x, z)$

$$\underset{x}{\text{minimize}} \underset{z}{\text{maximize}} f(x, z) \geq \underset{z}{\text{maximize}} \underset{x}{\text{minimize}} f(x, z) \quad (273)$$

always holds. When

$$\underset{x}{\text{minimize}} \underset{z}{\text{maximize}} f(x, z) = \underset{z}{\text{maximize}} \underset{x}{\text{minimize}} f(x, z) \quad (274)$$




Figure 54: Although objective functions from conic problems (275p) and (275d) are linear, this is a mnemonic icon for primal and dual problems. When problems are *strong duals*, duality gap is 0; meaning, functions $f(x)$ and $g(z)$ (dotted) *kiss* at saddle value, as depicted at center. Otherwise, dual functions never meet ($f(x) > g(z)$) by (273). Drawing by Kieff.

we have *strong duality* and then a *saddle value* [133] exists. (Figure 54) [268, p.3] Consider primal conic problem (p) (over cone \mathcal{K}) and its corresponding *dual problem* (d): [259, §3.3.1] [211, §2.1] [212] given vectors α, β and matrix constant C

$$\begin{array}{ll}
 \text{minimize}_{x} & \alpha^T x \\
 \text{subject to} & x \in \mathcal{K} \\
 & Cx = \beta
 \end{array}
 \quad
 \begin{array}{ll}
 \text{maximize}_{y, z} & \beta^T z \\
 \text{subject to} & y \in \mathcal{K}^* \\
 & C^T z + y = \alpha
 \end{array}
 \quad (d) \quad (275)$$

Observe the dual problem is also conic, and its objective function value never exceeds that of the primal;

$$\begin{aligned}
 \alpha^T x &\geq \beta^T z \\
 x^T(C^T z + y) &\geq (Cx)^T z \\
 x^T y &\geq 0
 \end{aligned} \quad (276)$$

which holds by definition (270). Under the sufficient condition: (275p) is a *convex problem* and satisfies *Slater's condition*,^{2.53} then each problem (p) and (d) attains the same optimal value of its objective and each problem is called a *strong dual* to the other because the *duality gap* (primal–dual optimal objective difference) is 0. Then (p) and (d) are together equivalent to the minimax problem

$$\begin{array}{ll}
 \text{minimize}_{x, y, z} & \alpha^T x - \beta^T z \\
 \text{subject to} & x \in \mathcal{K}, \quad y \in \mathcal{K}^* \\
 & Cx = \beta, \quad C^T z + y = \alpha
 \end{array}
 \quad (p)-(d) \quad (277)$$

whose optimal objective always has the saddle value 0 (regardless of the particular convex cone \mathcal{K} and other problem parameters). [319, §3.2] Thus determination of convergence for either primal or dual problem is facilitated.

Were convex cone \mathcal{K} polyhedral (§2.12.1), then problems (p) and (d) would be linear programs.^{2.54} Were \mathcal{K} a positive semidefinite cone, then

^{2.53}A convex problem, essentially, has convex objective function optimized over a convex set. (§4) In this context, (p) is convex if \mathcal{K} is a convex cone. Slater's condition is satisfied whenever any primal strictly feasible point exists. (p.260)

^{2.54}The self-dual nonnegative orthant yields the primal prototypical linear program and its dual.

problem (p) has the form of prototypical *semidefinite program*^{2.55} (592) with (d) its dual. It is sometimes possible to solve a primal problem by way of its dual; advantageous *when the dual problem is easier to solve than the primal problem, for example, because it can be solved analytically, or has some special structure that can be exploited.* [53, §5.5.5] (§4.2.3.1) \square

2.13.1.1 Key properties of dual cone

- For any cone, $(-\mathcal{K})^* = -\mathcal{K}^*$
- For any cones \mathcal{K}_1 and \mathcal{K}_2 , $\mathcal{K}_1 \subseteq \mathcal{K}_2 \Rightarrow \mathcal{K}_1^* \supseteq \mathcal{K}_2^*$ [293, §2.7]
- (Cartesian product) For closed convex cones \mathcal{K}_1 and \mathcal{K}_2 , their Cartesian product $\mathcal{K} = \mathcal{K}_1 \times \mathcal{K}_2$ is a closed convex cone, and

$$\mathcal{K}^* = \mathcal{K}_1^* \times \mathcal{K}_2^* \quad (278)$$

- (conjugation) [271, §14] [92, §4.5] When \mathcal{K} is any convex cone, the dual of the dual cone is the closure of the original cone; $\mathcal{K}^{**} = \overline{\mathcal{K}}$. Because $\mathcal{K}^{***} = \mathcal{K}^*$

$$\mathcal{K}^* = (\overline{\mathcal{K}})^* \quad (279)$$

When \mathcal{K} is closed and convex, then the dual of the dual cone is the original cone; $\mathcal{K}^{**} = \mathcal{K} \Leftrightarrow \mathcal{K}$ is a closed convex cone: [293, p.95]

$$\mathcal{K} = \left\{ x \in \mathbb{R}^n \mid \langle y, x \rangle \geq 0 \text{ for all } y \in \mathcal{K}^* \right\} \quad (280)$$

- If any cone \mathcal{K} has nonempty interior, then \mathcal{K}^* is pointed;

$$\mathcal{K} \text{ nonempty interior} \Rightarrow \mathcal{K}^* \text{ pointed} \quad (281)$$

Conversely, if the closure of any convex cone \mathcal{K} is pointed, then \mathcal{K}^* has nonempty interior;

$$\overline{\mathcal{K}} \text{ pointed} \Rightarrow \mathcal{K}^* \text{ nonempty interior} \quad (282)$$

Given that a cone $\mathcal{K} \subset \mathbb{R}^n$ is closed and convex, \mathcal{K} is pointed if and only if $\mathcal{K}^* - \mathcal{K}^* = \mathbb{R}^n$; *id est*, iff \mathcal{K}^* has nonempty interior. [48, §3.3, exer.20]

^{2.55}A *semidefinite program* is any convex program constraining a variable to any subset of a positive semidefinite cone. The qualifier *prototypical* conventionally means: a program having linear objective, affine equality constraints, but no inequality constraints except for cone membership.

- (vector sum) [271, thm.3.8] For convex cones \mathcal{K}_1 and \mathcal{K}_2

$$\mathcal{K}_1 + \mathcal{K}_2 = \text{conv}(\mathcal{K}_1 \cup \mathcal{K}_2) \quad (283)$$

- (dual vector-sum) [271, §16.4.2] [92, §4.6] For convex cones \mathcal{K}_1 and \mathcal{K}_2

$$\mathcal{K}_1^* \cap \mathcal{K}_2^* = (\mathcal{K}_1 + \mathcal{K}_2)^* = (\mathcal{K}_1 \cup \mathcal{K}_2)^* \quad (284)$$

- (closure of vector sum of duals)^{2.56} For closed convex cones \mathcal{K}_1 and \mathcal{K}_2

$$(\mathcal{K}_1 \cap \mathcal{K}_2)^* = \overline{\mathcal{K}_1^* + \mathcal{K}_2^*} = \overline{\text{conv}(\mathcal{K}_1^* \cup \mathcal{K}_2^*)} \quad (285)$$

[293, p.96] where operation closure becomes superfluous under the sufficient condition $\mathcal{K}_1 \cap \text{int } \mathcal{K}_2 \neq \emptyset$ [48, §3.3 exer.16, §4.1 exer.7].

- (Krein-Rutman) Given closed convex cones $\mathcal{K}_1 \subseteq \mathbb{R}^m$ and $\mathcal{K}_2 \subseteq \mathbb{R}^n$ and any linear map $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then provided $\text{int } \mathcal{K}_1 \cap A\mathcal{K}_2 \neq \emptyset$ [48, §3.3.13, confer §4.1 exer.9]

$$(A^{-1}\mathcal{K}_1 \cap \mathcal{K}_2)^* = A^T\mathcal{K}_1^* + \mathcal{K}_2^* \quad (286)$$

where dual of cone \mathcal{K}_1 is with respect to its ambient space \mathbb{R}^m and dual of cone \mathcal{K}_2 is with respect to \mathbb{R}^n , where $A^{-1}\mathcal{K}_1$ denotes inverse image (§2.1.9.0.1) of \mathcal{K}_1 under mapping A , and where A^T denotes the adjoint operation.

- \mathcal{K} is proper if and only if \mathcal{K}^* is proper.
- \mathcal{K} is polyhedral if and only if \mathcal{K}^* is polyhedral. [293, §2.8]
- \mathcal{K} is simplicial if and only if \mathcal{K}^* is simplicial. (§2.13.9.2) A simplicial cone and its dual are proper polyhedral cones (Figure 59, Figure 50), but not the converse.
- $\mathcal{K} \boxplus -\mathcal{K}^* = \mathbb{R}^n \Leftrightarrow \mathcal{K}$ is closed and convex. (1911) (p.714)
- Any direction in a proper cone \mathcal{K} is normal to a hyperplane separating \mathcal{K} from $-\mathcal{K}^*$.

^{2.56}These parallel analogous results for subspaces $\mathcal{R}_1, \mathcal{R}_2 \subseteq \mathbb{R}^n$; [92, §4.6]

$$(\mathcal{R}_1 + \mathcal{R}_2)^\perp = \mathcal{R}_1^\perp \cap \mathcal{R}_2^\perp$$

$$(\mathcal{R}_1 \cap \mathcal{R}_2)^\perp = \overline{\mathcal{R}_1^\perp + \mathcal{R}_2^\perp}$$

$\mathcal{R}^{\perp\perp} = \mathcal{R}$ for any subspace \mathcal{R} .




Figure 55: When convex cone \mathcal{K} is any one Cartesian axis, its dual cone is the convex hull of all axes remaining. In \mathbb{R}^3 , dual cone \mathcal{K}^* (drawn tiled and truncated) is a hyperplane through origin; its normal belongs to line \mathcal{K} . In \mathbb{R}^2 , dual cone \mathcal{K}^* is a line through origin while convex cone \mathcal{K} is that line orthogonal.

2.13.1.2 Examples of dual cone

When \mathcal{K} is \mathbb{R}^n , \mathcal{K}^* is the point at the origin, and *vice versa*.

When \mathcal{K} is a subspace, \mathcal{K}^* is its orthogonal complement, and *vice versa*. (§E.9.2.1, Figure 55)

When cone \mathcal{K} is a halfspace in \mathbb{R}^n with $n > 0$ (Figure 53 for example), the dual cone \mathcal{K}^* is a ray (base $\mathbf{0}$) belonging to that halfspace but orthogonal to its bounding hyperplane (that contains the origin), and *vice versa*.

When convex cone \mathcal{K} is a closed halfplane in \mathbb{R}^3 (Figure 56), it is neither pointed or of nonempty interior; hence, the dual cone \mathcal{K}^* can be neither of nonempty interior or pointed.

When \mathcal{K} is any particular orthant in \mathbb{R}^n , the dual cone is identical; *id est*, $\mathcal{K} = \mathcal{K}^*$.

When \mathcal{K} is any quadrant in subspace \mathbb{R}^2 , \mathcal{K}^* is a *wedge*-shaped polyhedral cone in \mathbb{R}^3 ; *e.g.*, for \mathcal{K} equal to quadrant I, $\mathcal{K}^* = \begin{bmatrix} \mathbb{R}_+^2 \\ \mathbb{R} \end{bmatrix}$.

When \mathcal{K} is a polyhedral flavor of the Lorentz cone (*confer*(161))

$$\mathcal{K}_\ell = \left\{ \begin{bmatrix} x \\ t \end{bmatrix} \in \mathbb{R}^n \times \mathbb{R} \mid \|x\|_\ell \leq t \right\}, \quad \ell \in \{1, \infty\} \quad (287)$$

its dual is the proper cone

$$\mathcal{K}_q = \mathcal{K}_\ell^* = \left\{ \begin{bmatrix} x \\ t \end{bmatrix} \in \mathbb{R}^n \times \mathbb{R} \mid \|x\|_q \leq t \right\}, \quad \ell \in \{1, 2, \infty\} \quad (288)$$

where $\|x\|_q$ is the *dual norm* [53, §A.1.6] determined via solution to $1/\ell + 1/q = 1$.

2.13.2 Abstractions of *Farkas' lemma*

2.13.2.0.1 Corollary. *Generalized inequality and membership relation.*

[177, §A.4.2] Let \mathcal{K} be any closed convex cone and \mathcal{K}^* its dual, and let x and y belong to a vector space \mathbb{R}^n . Then

$$y \in \mathcal{K}^* \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } x \in \mathcal{K} \quad (289)$$

which is, merely, a statement of fact by definition of dual cone (270). By closure we have conjugation:

$$x \in \mathcal{K} \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } y \in \mathcal{K}^* \quad (290)$$




Figure 56: \mathcal{K} and \mathcal{K}^* are halfplanes in \mathbb{R}^3 ; *blades*. Both semi-infinite convex cones appear truncated. Each cone is like \mathcal{K} in Figure 53, but embedded in a two-dimensional subspace of \mathbb{R}^3 . Cartesian coordinate axes drawn for reference.

which may be regarded as a simple translation of the *Farkas lemma* [116] as in [271, §22] to the language of convex cones, and a generalization of the well-known Cartesian fact

$$x \succeq 0 \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } y \succeq 0 \quad (291)$$

for which implicitly $\mathcal{K} = \mathcal{K}^* = \mathbb{R}_+^n$ the nonnegative orthant.

Membership relation (290) is often written instead as *dual generalized inequalities*, when \mathcal{K} and \mathcal{K}^* are pointed closed convex cones,

$$x \underset{\mathcal{K}}{\succeq} 0 \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } y \underset{\mathcal{K}^*}{\succeq} 0 \quad (292)$$

meaning, coordinates for biorthogonal expansion of x (§2.13.8) [325] must be nonnegative when x belongs to \mathcal{K} . By conjugation [271, thm.14.1]

$$y \underset{\mathcal{K}^*}{\succeq} 0 \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } x \underset{\mathcal{K}}{\succeq} 0 \quad (293)$$

◇

When pointed closed convex cone \mathcal{K} is not polyhedral, coordinate axes for biorthogonal expansion asserted by the corollary are taken from extreme directions of \mathcal{K} ; expansion is assured by *Carathéodory's theorem* (§E.6.4.1.1).

We presume, throughout, the obvious:

$$\begin{aligned} x \in \mathcal{K} &\Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } y \in \mathcal{K}^* & (290) \\ &\Leftrightarrow & (294) \\ x \in \mathcal{K} &\Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } y \in \mathcal{K}^*, \|y\|=1 \end{aligned}$$

2.13.2.0.2 Exercise. Test of dual generalized inequalities.

Test Corollary 2.13.2.0.1 and (294) graphically on the two-dimensional polyhedral cone and its dual in Figure 52. ▼

When pointed closed convex cone \mathcal{K} is implicit from context: (confer §2.7.2.2)

$$\begin{aligned} x \succeq 0 &\Leftrightarrow x \in \mathcal{K} \\ x \succ 0 &\Leftrightarrow x \in \text{rel int } \mathcal{K} \end{aligned} \quad (295)$$

Strict inequality $x \succ 0$ means coordinates for biorthogonal expansion of x must be positive when x belongs to $\text{rel int } \mathcal{K}$. Strict membership relations are useful; *e.g.*, for any proper cone \mathcal{K} and its dual \mathcal{K}^*

$$x \in \text{int } \mathcal{K} \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } y \in \mathcal{K}^*, y \neq \mathbf{0} \quad (296)$$

$$x \in \mathcal{K}, x \neq \mathbf{0} \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } y \in \text{int } \mathcal{K}^* \quad (297)$$

By conjugation, we also have the dual relations:

$$y \in \text{int } \mathcal{K}^* \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } x \in \mathcal{K}, x \neq \mathbf{0} \quad (298)$$

$$y \in \mathcal{K}^*, y \neq \mathbf{0} \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } x \in \text{int } \mathcal{K} \quad (299)$$

Boundary-membership relations for proper cones are also useful:

$$x \in \partial \mathcal{K} \Leftrightarrow \exists y \neq \mathbf{0} \ni \langle y, x \rangle = 0, y \in \mathcal{K}^*, x \in \mathcal{K} \quad (300)$$

$$y \in \partial \mathcal{K}^* \Leftrightarrow \exists x \neq \mathbf{0} \ni \langle y, x \rangle = 0, x \in \mathcal{K}, y \in \mathcal{K}^* \quad (301)$$

2.13.2.0.3 Example. *Linear inequality.* [299, §4]
(confer §2.13.5.1.1) Consider a given matrix A and closed convex cone \mathcal{K} .
By membership relation we have

$$\begin{aligned} Ay \in \mathcal{K}^* &\Leftrightarrow x^T A y \geq 0 \quad \forall x \in \mathcal{K} \\ &\Leftrightarrow y^T z \geq 0 \quad \forall z \in \{A^T x \mid x \in \mathcal{K}\} \\ &\Leftrightarrow y \in \{A^T x \mid x \in \mathcal{K}\}^* \end{aligned} \quad (302)$$

This implies

$$\{y \mid Ay \in \mathcal{K}^*\} = \{A^T x \mid x \in \mathcal{K}\}^* \quad (303)$$

If we regard A as a linear operator, then A^T is its adjoint. When \mathcal{K} is the *self-dual* nonnegative orthant (§2.13.5.1), for example, then

$$\{y \mid Ay \succeq 0\} = \{A^T x \mid x \succeq 0\}^* \quad (304)$$

□

2.13.2.1 Null certificate, Theorem of the alternative

If in particular $x_p \notin \mathcal{K}$ a closed convex cone, then the construction in Figure 51(b) suggests there exists a supporting hyperplane (having inward-normal belonging to dual cone \mathcal{K}^*) separating x_p from \mathcal{K} ; indeed, (290)

$$x_p \notin \mathcal{K} \Leftrightarrow \exists y \in \mathcal{K}^* \ni \langle y, x_p \rangle < 0 \quad (305)$$

The existence of any one such y is a certificate of null membership. From a different perspective,

$$\begin{aligned} x_p &\in \mathcal{K} \\ \text{or in the alternative} & \\ \exists y \in \mathcal{K}^* \ni \langle y, x_p \rangle &< 0 \end{aligned} \quad (306)$$

By *alternative* is meant: these two systems are incompatible; one system is feasible while the other is not.

2.13.2.1.1 Example. *Theorem of the alternative for linear inequality.*

Myriad alternative systems of linear inequality can be explained in terms of pointed closed convex cones and their duals.

Beginning from the simplest Cartesian dual generalized inequalities (291) (with respect to the nonnegative orthant \mathbb{R}_+^m),

$$y \succeq 0 \Leftrightarrow x^T y \geq 0 \text{ for all } x \succeq 0 \quad (307)$$

Given $A \in \mathbb{R}^{n \times m}$, we make vector substitution $y \leftarrow A^T y$

$$A^T y \succeq 0 \Leftrightarrow x^T A^T y \geq 0 \text{ for all } x \succeq 0 \quad (308)$$

Introducing a new vector by calculating $b \triangleq Ax$ we get

$$A^T y \succeq 0 \Leftrightarrow b^T y \geq 0, \quad b = Ax \text{ for all } x \succeq 0 \quad (309)$$

By complementing sense of the scalar inequality:

$$\begin{aligned} A^T y &\succeq 0 \\ \text{or in the alternative} & \\ b^T y &< 0, \quad \exists b = Ax, \quad x \succeq 0 \end{aligned} \quad (310)$$

If one system has a solution, then the other does not; define a convex cone $\mathcal{K} = \{y \mid A^T y \succeq 0\}$, then $y \in \mathcal{K}$ or in the alternative $y \notin \mathcal{K}$. Scalar inequality $b^T y < 0$ can be moved to the other side of the alternative, but that requires some explanation: From the results of Example 2.13.2.0.3, the dual cone is $\mathcal{K}^* = \{Ax \mid x \succeq 0\}$. From (290) we have

$$\begin{aligned} y \in \mathcal{K} &\Leftrightarrow b^T y \geq 0 \text{ for all } b \in \mathcal{K}^* \\ A^T y \succeq 0 &\Leftrightarrow b^T y \geq 0 \text{ for all } b \in \{Ax \mid x \succeq 0\} \end{aligned} \quad (311)$$

Given some b vector and $y \in \mathcal{K}$, then $b^T y < 0$ can only mean $b \notin \mathcal{K}^*$. An alternative system is therefore simply $b \in \mathcal{K}^*$: [177, p.59] (Farkas/Tucker)

$$\begin{aligned} A^T y &\succeq 0, \quad b^T y < 0 \\ \text{or in the alternative} & \\ b &= Ax, \quad x \succeq 0 \end{aligned} \tag{312}$$

Geometrically this means, either vector b belongs to convex cone \mathcal{K}^* or it does not. When $b \notin \mathcal{K}^*$, then there is a vector $y \in \mathcal{K}$ normal to a hyperplane separating point b from cone \mathcal{K}^* .

For another example, from membership relation (289) with affine transformation of dual variable we may write: Given $A \in \mathbb{R}^{n \times m}$ and $b \in \mathbb{R}^n$

$$b - Ay \in \mathcal{K}^* \Leftrightarrow x^T(b - Ay) \geq 0 \quad \forall x \in \mathcal{K} \tag{313}$$

$$A^T x = \mathbf{0}, \quad b - Ay \in \mathcal{K}^* \Rightarrow x^T b \geq 0 \quad \forall x \in \mathcal{K} \tag{314}$$

From membership relation (313), conversely, suppose we allow any $y \in \mathbb{R}^m$. Then because $-x^T A y$ is unbounded below, $x^T(b - Ay) \geq 0$ implies $A^T x = \mathbf{0}$: for $y \in \mathbb{R}^m$

$$A^T x = \mathbf{0}, \quad b - Ay \in \mathcal{K}^* \Leftrightarrow x^T(b - Ay) \geq 0 \quad \forall x \in \mathcal{K} \tag{315}$$

In toto,

$$b - Ay \in \mathcal{K}^* \Leftrightarrow x^T b \geq 0, \quad A^T x = \mathbf{0} \quad \forall x \in \mathcal{K} \tag{316}$$

Vector x belongs to cone \mathcal{K} but is also constrained to lie in a subspace of \mathbb{R}^n specified by an intersection of hyperplanes through the origin $\{x \in \mathbb{R}^n \mid A^T x = \mathbf{0}\}$. From this, alternative systems of generalized inequality with respect to pointed closed convex cones \mathcal{K} and \mathcal{K}^*

$$Ay \preceq_{\mathcal{K}^*} b$$

$$\text{or in the alternative} \tag{317}$$

$$x^T b < 0, \quad A^T x = \mathbf{0}, \quad x \succeq_{\mathcal{K}} 0$$

derived from (316) simply by taking the complementary sense of the inequality in $x^T b$. These two systems are alternatives; if one system has a solution, then the other does not.^{2.57} [271, p.201]

By invoking a strict membership relation between proper cones (296), we can construct a more exotic alternative strengthened by demand for an interior point;

$$b - Ay \succ_{\kappa^*} 0 \Leftrightarrow x^T b > 0, \quad A^T x = \mathbf{0} \quad \forall x \succeq_{\kappa} 0, \quad x \neq \mathbf{0} \quad (318)$$

From this, alternative systems of generalized inequality [53, pages:50,54,262]

$$Ay \prec_{\kappa^*} b$$

$$\text{or in the alternative} \quad (319)$$

$$x^T b \leq 0, \quad A^T x = \mathbf{0}, \quad x \succeq_{\kappa} 0, \quad x \neq \mathbf{0}$$

derived from (318) by taking the complementary sense of the inequality in $x^T b$.

And from this, alternative systems with respect to the nonnegative orthant attributed to Gordan in 1873: [140] [48, §2.2] substituting $A \leftarrow A^T$ and setting $b = \mathbf{0}$

$$A^T y \prec 0$$

$$\text{or in the alternative} \quad (320)$$

$$Ax = \mathbf{0}, \quad x \succeq 0, \quad \|x\|_1 = 1$$

□

^{2.57}If solutions at $\pm\infty$ are disallowed, then the alternative systems become instead *mutually exclusive* with respect to nonpolyhedral cones. Simultaneous infeasibility of the two systems is not precluded by mutual exclusivity; called a *weak alternative*. Ye provides an example illustrating simultaneous infeasibility with respect to the positive semidefinite cone: $x \in \mathbb{S}^2$, $y \in \mathbb{R}$, $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, and $b = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ where $x^T b$ means $\langle x, b \rangle$. A better strategy than disallowing solutions at $\pm\infty$ is to demand an interior point as in (319) or Lemma 4.2.1.1.2. Then question of simultaneous infeasibility is moot.

2.13.3 Optimality condition

(confer §2.13.10.1) The general first-order necessary and sufficient condition for optimality of solution x^* to a minimization problem ((275p) for example) with real differentiable convex objective function $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ is [270, §3]

$$\nabla f(x^*)^T(x - x^*) \geq 0 \quad \forall x \in \mathcal{C}, \quad x^* \in \mathcal{C} \quad (321)$$

where \mathcal{C} is a convex *feasible set*,^{2.58} and where $\nabla f(x^*)$ is the *gradient* of f (§3.1.8) with respect to x evaluated at x^* .^{2.59} In words, the negative gradient is normal to a hyperplane supporting the feasible set at a point of optimality. (Figure 62)

In the unconstrained case, optimality condition (321) reduces to the classical rule

$$\nabla f(x^*) = \mathbf{0}, \quad x^* \in \text{dom } f \quad (322)$$

which can be inferred from the following application:

2.13.3.0.1 Example. Equality constrained problem.

Given a real differentiable convex function $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ defined on domain \mathbb{R}^n , a fat full-rank matrix $C \in \mathbb{R}^{p \times n}$, and vector $d \in \mathbb{R}^p$, the convex optimization problem

$$\begin{aligned} & \underset{x}{\text{minimize}} && f(x) \\ & \text{subject to} && Cx = d \end{aligned} \quad (323)$$

is characterized by the well-known necessary and sufficient optimality condition [53, §4.2.3]

$$\nabla f(x^*) + C^T \nu = \mathbf{0} \quad (324)$$

where $\nu \in \mathbb{R}^p$ is the eminent *Lagrange multiplier*. [269] [220, p.188] In other words, feasible solution x^* is optimal if and only if $\nabla f(x^*)$ belongs to $\mathcal{R}(C^T)$.

Via membership relation, we now derive condition (324) from the general first-order condition for optimality (321): The feasible set for problem (323) is

$$\mathcal{C} \triangleq \{x \in \mathbb{R}^n \mid Cx = d\} = \{Z\xi + x_p \mid \xi \in \mathbb{R}^{n-\text{rank } C}\} \quad (325)$$

^{2.58} a set of all variable values satisfying the problem constraints.

^{2.59} Direct solution to (321), instead of the minimization, is known as a *variation inequality* from the *calculus of variations*. [220, p.178] [115, p.37]

where $Z \in \mathbb{R}^{n \times n - \text{rank } C}$ holds basis $\mathcal{N}(C)$ columnar, and x_p is any particular solution to $Cx = d$. Since $x^* \in \mathcal{C}$, we arbitrarily choose $x_p = x^*$ which yields the equivalent optimality condition

$$\nabla f(x^*)^T Z \xi \geq 0 \quad \forall \xi \in \mathbb{R}^{n - \text{rank } C} \quad (326)$$

But this is simply half of a membership relation, and the cone dual to $\mathbb{R}^{n - \text{rank } C}$ is the origin in $\mathbb{R}^{n - \text{rank } C}$. We must therefore have

$$Z^T \nabla f(x^*) = \mathbf{0} \Leftrightarrow \nabla f(x^*)^T Z \xi \geq 0 \quad \forall \xi \in \mathbb{R}^{n - \text{rank } C} \quad (327)$$

meaning, $\nabla f(x^*)$ must be orthogonal to $\mathcal{N}(C)$. This condition

$$Z^T \nabla f(x^*) = \mathbf{0}, \quad x^* \in \mathcal{C} \quad (328)$$

is necessary and sufficient for optimality of x^* . \square

2.13.4 Discretization of membership relation

2.13.4.1 Dual halfspace-description

Halfspace-description of the dual cone is equally simple (and extensible to an infinite number of generators) as vertex-description (95) for the corresponding closed convex cone: By definition (270), for $X \in \mathbb{R}^{n \times N}$ as in (254), (*confer*(260))

$$\begin{aligned} \mathcal{K}^* &= \{y \in \mathbb{R}^n \mid z^T y \geq 0 \text{ for all } z \in \mathcal{K}\} \\ &= \{y \in \mathbb{R}^n \mid z^T y \geq 0 \text{ for all } z = Xa, a \succeq 0\} \\ &= \{y \in \mathbb{R}^n \mid a^T X^T y \geq 0, a \succeq 0\} \\ &= \{y \in \mathbb{R}^n \mid X^T y \succeq 0\} \end{aligned} \quad (329)$$

that follows from the *generalized inequality and membership corollary* (291). The semi-infinity of tests specified by all $z \in \mathcal{K}$ has been reduced to a set of generators for \mathcal{K} constituting the columns of X ; *id est*, the test has been discretized.

Whenever \mathcal{K} is known to be closed and convex, then the converse must also hold; *id est*, given any set of generators for \mathcal{K}^* arranged columnar in Y , then the consequent vertex-description of the dual cone connotes a halfspace-description for \mathcal{K} : [293, §2.8]

$$\mathcal{K}^* = \{Ya \mid a \succeq 0\} \Leftrightarrow \mathcal{K}^{**} = \mathcal{K} = \{z \mid Y^T z \succeq 0\} \quad (330)$$

2.13.4.2 First dual-cone formula

From these two results (329) and (330) we deduce a general principle:

- From any *[sic]* given vertex-description of a convex cone \mathcal{K} , a halfspace-description of the dual cone \mathcal{K}^* is immediate by matrix transposition (329); conversely, from any given halfspace-description of \mathcal{K} , a dual vertex-description is immediate (330).

Various other converses are just a little trickier. (§2.13.9, §2.13.11)

We deduce further: For any polyhedral cone \mathcal{K} , the dual cone \mathcal{K}^* is also polyhedral and $\mathcal{K}^{**} = \mathcal{K}$. [293, §2.8]

The *generalized inequality and membership corollary* is discretized in the following theorem [20, §1]^{2.60} that follows directly from (329) and (330):

2.13.4.2.1 Theorem. *Discretized membership.* (confer §2.13.2.0.1)

Given any set of generators (§2.8.1.2) denoted by $\mathcal{G}(\mathcal{K})$ for closed convex cone $\mathcal{K} \subseteq \mathbb{R}^n$ and any set of generators denoted $\mathcal{G}(\mathcal{K}^*)$ for its dual, then discretization of the *generalized inequality and membership corollary* (p.154) is necessary and sufficient for certifying cone membership: for x and y in vector space \mathbb{R}^n

$$x \in \mathcal{K} \Leftrightarrow \langle \gamma^*, x \rangle \geq 0 \text{ for all } \gamma^* \in \mathcal{G}(\mathcal{K}^*) \quad (331)$$

$$y \in \mathcal{K}^* \Leftrightarrow \langle \gamma, y \rangle \geq 0 \text{ for all } \gamma \in \mathcal{G}(\mathcal{K}) \quad (332)$$

◇

2.13.4.2.2 Exercise. *Test of discretized dual generalized inequalities.*

Test Theorem 2.13.4.2.1 on Figure 52(a) using the extreme directions as generators. ▼

From the *discretized membership theorem* we may further deduce a more surgical description of closed convex cone that prescribes only a finite number of halfspaces for its construction when polyhedral: (Figure 51(a))

$$\mathcal{K} = \{x \in \mathbb{R}^n \mid \langle \gamma^*, x \rangle \geq 0 \text{ for all } \gamma^* \in \mathcal{G}(\mathcal{K}^*)\} \quad (333)$$

$$\mathcal{K}^* = \{y \in \mathbb{R}^n \mid \langle \gamma, y \rangle \geq 0 \text{ for all } \gamma \in \mathcal{G}(\mathcal{K})\} \quad (334)$$

^{2.60}Barker & Carlson state the theorem only for the pointed closed convex case.

2.13.4.2.3 Exercise. *Partial order induced by orthant.*

When comparison is with respect to the nonnegative orthant $\mathcal{K} = \mathbb{R}_+^n$, then from the *discretized membership theorem* it directly follows:

$$x \preceq z \Leftrightarrow x_i \leq z_i \quad \forall i \quad (335)$$

Generate simple counterexamples demonstrating that this equivalence with entrywise inequality holds only when the underlying cone inducing partial order is the nonnegative orthant. \blacktriangledown

2.13.4.2.4 Example. *Boundary membership to proper polyhedral cone.*

For a polyhedral cone, test (300) of boundary membership can be formulated as a linear program. Say proper polyhedral cone \mathcal{K} is specified completely by its extreme directions that are arranged columnar in

$$X = [\Gamma_1 \ \cdots \ \Gamma_N] \in \mathbb{R}^{n \times N} \quad (254)$$

is est, $\mathcal{K} = \{Xa \mid a \succeq 0\}$. Then the membership relation

$$c \in \partial\mathcal{K} \Leftrightarrow \exists y \neq \mathbf{0} \ni \langle y, c \rangle = 0, \quad y \in \mathcal{K}^*, \quad c \in \mathcal{K} \quad (300)$$

may be expressed

$$\begin{aligned} & \underset{a, y}{\text{find}} && y \neq \mathbf{0} \\ & \text{subject to} && c^T y = 0 \\ & && X^T y \succeq 0 \\ & && Xa = c \\ & && a \succeq 0 \end{aligned} \quad (336)$$

This linear feasibility problem has a solution iff $c \in \partial\mathcal{K}$. \square

2.13.4.3 smallest face of proper cone

The *smallest face* containing \mathcal{C} , a given nonempty convex subset of a convex set \mathcal{K} , is equivalent to the intersection of all faces of \mathcal{K} containing \mathcal{C} . [271, p.164] By (280), membership relation (300) implies that each and every point on boundary $\partial\mathcal{K}$ of proper cone \mathcal{K} belongs to a hyperplane supporting \mathcal{K} whose normal y belongs to dual cone \mathcal{K}^* . It follows that the smallest face \mathcal{F} , containing $\mathcal{C} \subset \partial\mathcal{K} \subset \mathbb{R}^n$ on boundary of proper cone \mathcal{K} , is the intersection of all hyperplanes containing \mathcal{C} whose normals are in \mathcal{K}^* ;

$$\mathcal{F}(\mathcal{K} \supset \mathcal{C}) = \{x \in \mathcal{K} \mid x \perp \mathcal{K}^* \cap \mathcal{C}^\perp\} \quad (337)$$

where

$$\mathcal{C}^\perp \triangleq \{y \in \mathbb{R}^n \mid \langle z, y \rangle = 0 \quad \forall z \in \mathcal{C}\} \quad (338)$$

2.13.4.3.1 Example. *Finding smallest face.*

Suppose proper polyhedral cone \mathcal{K} is specified by extreme directions arranged columnar in

$$X = [\Gamma_1 \ \cdots \ \Gamma_N] \in \mathbb{R}^{n \times N} \quad (254)$$

Then it is easy to find its smallest face $\mathcal{F}(\mathcal{K} \ni c)$ containing a given point $c \in \partial\mathcal{K}$. By the *discretized membership theorem*, it is necessary and sufficient to find generators for the smallest face by solving a linear *feasibility problem*: *videlicet*, for each $i \in \{1 \dots N\}$

$$\begin{aligned} & \text{find } z \\ & \text{subject to } \begin{aligned} c^\text{T}z &= 0 \\ X^\text{T}z &\succeq 0 \\ \Gamma_i^\text{T}z &= 1 \end{aligned} \end{aligned} \quad (339)$$

If this problem is infeasible for extreme direction $\Gamma_i \in \mathcal{K}$, then $\Gamma_i \in \mathcal{F}(\mathcal{K} \ni c)$ by (337) and (329) because $\mathcal{K}^* \cap c^\perp = \{z \in \mathbb{R}^n \mid X^\text{T}z \succeq 0, c^\text{T}z = 0\}$; in that case, Γ_i is a generator of $\mathcal{F}(\mathcal{K} \ni c)$. If there exists a feasible vector z for which $\Gamma_i^\text{T}z = 1$, conversely, then $\Gamma_i \notin \mathcal{K}^* \cap c^\perp$ so $\Gamma_i \notin \mathcal{F}(\mathcal{K} \ni c)$; vector z is a certificate of null membership.

Since the constant in constraint $\Gamma_i^\text{T}z = 1$ is arbitrary positively, then by *theorem of the alternative* there is correspondence between (339) and (317) admitting the alternative linear problem: for a given point $c \in \partial\mathcal{K}$

$$\begin{aligned} & \text{find } a, t \\ & \text{subject to } \begin{aligned} a \in \mathbb{R}^N, t \in \mathbb{R} \\ tc - \Gamma_i &= Xa \\ a &\succeq 0 \end{aligned} \end{aligned} \quad (340)$$

Now if this problem is feasible (bounded) for extreme direction $\Gamma_i \in \mathcal{K}$, then (339) is not feasible and $\Gamma_i \in \mathcal{F}(\mathcal{K} \ni c)$ is a generator of the smallest face containing c . \square

2.13.4.3.2 Exercise. *Smallest face of positive semidefinite cone.*

Derive (204) from (337). \blacktriangledown

2.13.5 Dual PSD cone and generalized inequality

The *dual positive semidefinite cone* \mathcal{K}^* is confined to \mathbb{S}^M by convention;

$$\mathbb{S}_+^{M*} \triangleq \{Y \in \mathbb{S}^M \mid \langle Y, X \rangle \geq 0 \text{ for all } X \in \mathbb{S}_+^M\} = \mathbb{S}_+^M \quad (341)$$

The positive semidefinite cone is self-dual in the ambient space of symmetric matrices [53, exmp.2.24] [35] [174, §II]; $\mathcal{K} = \mathcal{K}^*$.

Dual generalized inequalities with respect to the positive semidefinite cone in the ambient space of symmetric matrices can therefore be simply stated: (Fejér)

$$X \succeq 0 \Leftrightarrow \text{tr}(Y^T X) \geq 0 \text{ for all } Y \succeq 0 \quad (342)$$

Membership to this cone can be determined in the isometrically isomorphic Euclidean space \mathbb{R}^{M^2} via (32). (§2.2.1) By the two interpretations in §2.13.1, positive semidefinite matrix Y can be interpreted as inward-normal to a hyperplane supporting the positive semidefinite cone.

The fundamental statement of positive semidefiniteness, $y^T X y \geq 0 \forall y$ (§A.3.0.0.1), evokes a particular instance of these dual generalized inequalities (342):

$$X \succeq 0 \Leftrightarrow \langle yy^T, X \rangle \geq 0 \forall yy^T (\succeq 0) \quad (1355)$$

Discretization (§2.13.4.2.1) allows replacement of positive semidefinite matrices Y with this minimal set of generators comprising the extreme directions of the positive semidefinite cone (§2.9.2.4).

2.13.5.1 self-dual cones

From (123) (a consequence of the *halfspaces theorem*, §2.4.1.1.1), where the only finite value of the support function for a convex cone is 0 [177, §C.2.3.1], or from discretized definition (334) of the dual cone we get a rather self-evident characterization of self-duality:

$$\mathcal{K} = \mathcal{K}^* \Leftrightarrow \mathcal{K} = \bigcap_{\gamma \in \mathcal{G}(\mathcal{K})} \{y \mid \gamma^T y \geq 0\} \quad (343)$$

In words: Cone \mathcal{K} is *self-dual* iff its own extreme directions are inward-normals to a (minimal) set of hyperplanes bounding halfspaces whose intersection constructs it. This means each extreme direction of \mathcal{K} is normal




Figure 57: Two (truncated) views of a polyhedral cone \mathcal{K} and its dual in \mathbb{R}^3 . Each of four extreme directions from \mathcal{K} belongs to a face of dual cone \mathcal{K}^* . Cone \mathcal{K} , shrouded by its dual, is symmetrical about its axis of revolution. Each pair of diametrically opposed extreme directions from \mathcal{K} makes a right angle. An orthant (or any rotation thereof; a simplicial cone) is not the only self-dual polyhedral cone in three or more dimensions; [19, §2.A.21] *e.g.*, consider an *equilateral* having five extreme directions. In fact, every self-dual polyhedral cone in \mathbb{R}^3 has an odd number of extreme directions. [21, thm.3]

to a hyperplane exposing one of its own faces; a necessary but insufficient condition for self-duality (Figure 57, for example).

Self-dual cones are of necessarily nonempty interior [29, §I] and invariant to rotation about the origin. Their most prominent representatives are the orthants, the positive semidefinite cone \mathbb{S}_+^M in the ambient space of symmetric matrices (341), and the Lorentz cone (161) [19, §II.A] [53, exmp.2.25]. In three dimensions, a plane containing the axis of revolution of a self-dual cone (and the origin) will produce a *slice* whose boundary makes a right angle.

2.13.5.1.1 Example. *Linear matrix inequality.* (confer §2.13.2.0.3)

Consider a peculiar vertex-description for a closed convex cone \mathcal{K} defined over a positive semidefinite cone (instead of a nonnegative orthant as in definition (95)): for $X \in \mathbb{S}_+^n$ given $A_j \in \mathbb{S}^n$, $j = 1 \dots m$

$$\begin{aligned} \mathcal{K} &= \left\{ \left[\begin{array}{c} \langle A_1, X \rangle \\ \vdots \\ \langle A_m, X \rangle \end{array} \right] \mid X \succeq 0 \right\} \subseteq \mathbb{R}^m \\ &= \left\{ \left[\begin{array}{c} \text{svec}(A_1)^\text{T} \\ \vdots \\ \text{svec}(A_m)^\text{T} \end{array} \right] \text{svec } X \mid X \succeq 0 \right\} \\ &\triangleq \{A \text{ svec } X \mid X \succeq 0\} \end{aligned} \quad (344)$$

where $A \in \mathbb{R}^{m \times n(n+1)/2}$, and where symmetric vectorization svec is defined in (50). Cone \mathcal{K} is indeed convex because, by (158)

$$A \text{ svec } X_{p_1}, A \text{ svec } X_{p_2} \in \mathcal{K} \Rightarrow A(\zeta \text{ svec } X_{p_1} + \xi \text{ svec } X_{p_2}) \in \mathcal{K} \text{ for all } \zeta, \xi \geq 0 \quad (345)$$

since a nonnegatively weighted sum of positive semidefinite matrices must be positive semidefinite. (§A.3.1.0.2) Although matrix A is finite-dimensional, \mathcal{K} is generally not a polyhedral cone (unless m equals 1 or 2) simply because $X \in \mathbb{S}_+^n$. Provided the A_j matrices are linearly independent, then

$$\text{rel int } \mathcal{K} = \text{int } \mathcal{K} \quad (13)$$

meaning, cone \mathcal{K} interior is nonempty; implying, dual cone \mathcal{K}^* is pointed by (281). If matrix A has no nullspace, then $A \text{ svec } X$ is an isomorphism

in X between positive semidefinite cone \mathbb{S}_+^n and $\mathcal{R}(A)$. (§2.2.1.0.1, §2.10.1.1)
In that case, convex cone \mathcal{K} has relative interior

$$\text{rel int } \mathcal{K} = \{A \text{ svec } X \mid X \succ 0\} \quad (346)$$

and boundary

$$\text{rel } \partial \mathcal{K} = \{A \text{ svec } X \mid X \succeq 0, X \neq 0\} \quad (347)$$

Now consider the (closed convex) dual cone:

$$\begin{aligned} \mathcal{K}^* &= \{y \mid \langle z, y \rangle \geq 0 \text{ for all } z \in \mathcal{K}\} \subseteq \mathbb{R}^m \\ &= \{y \mid \langle z, y \rangle \geq 0 \text{ for all } z = A \text{ svec } X, X \succeq 0\} \\ &= \{y \mid \langle A \text{ svec } X, y \rangle \geq 0 \text{ for all } X \succeq 0\} \\ &= \{y \mid \langle \text{svec } X, A^T y \rangle \geq 0 \text{ for all } X \succeq 0\} \\ &= \{y \mid \text{svec}^{-1}(A^T y) \succeq 0\} \end{aligned} \quad (348)$$

that follows from (342) and leads to an equally peculiar halfspace-description

$$\mathcal{K}^* = \{y \in \mathbb{R}^m \mid \sum_{j=1}^m y_j A_j \succeq 0\} \quad (349)$$

The summation inequality with respect to positive semidefinite cone \mathbb{S}_+^n is known as a *linear matrix inequality*. [51] [131] [232] [322] Although we already know that the dual cone is convex (§2.13.1), the *inverse image theorem* certifies convexity of \mathcal{K}^* which is the inverse image of positive semidefinite cone \mathbb{S}_+^n under linear transformation $g(y) \triangleq \sum y_j A_j$. Function $g(y)$ on \mathbb{R}^m is an isomorphism when the A_j matrices are linearly independent; hence, uniquely invertible. Inverse image \mathcal{K}^* must therefore have dimension equal to $\dim(\mathcal{R}(A^T) \cap \text{svec } \mathbb{S}_+^n)$ by (43). When this dimension is m , then dual cone \mathcal{K}^* interior is nonempty

$$\text{int } \mathcal{K}^* = \{y \in \mathbb{R}^m \mid \sum_{j=1}^m y_j A_j \succ 0\} \quad (350)$$

having boundary

$$\partial \mathcal{K}^* = \{y \in \mathbb{R}^m \mid \sum_{j=1}^m y_j A_j \succeq 0, \sum_{j=1}^m y_j A_j \neq 0\} \quad (351)$$

and then cone \mathcal{K} is pointed (§2.13.1.1). \square

2.13.6 Dual of pointed polyhedral cone

In a subspace of \mathbb{R}^n , now we consider a pointed polyhedral cone \mathcal{K} given in terms of its extreme directions Γ_i arranged columnar in

$$X = [\Gamma_1 \ \Gamma_2 \ \cdots \ \Gamma_N] \in \mathbb{R}^{n \times N} \quad (254)$$

The *extremes theorem* (§2.8.1.1.1) provides the vertex-description of a pointed polyhedral cone in terms of its finite number of extreme directions and its lone vertex at the origin:

2.13.6.0.1 Definition. *Pointed polyhedral cone, vertex-description.*

Given pointed polyhedral cone \mathcal{K} in a subspace of \mathbb{R}^n , denoting its i^{th} extreme direction by $\Gamma_i \in \mathbb{R}^n$ arranged in a matrix X as in (254), then that cone may be described: (79) (confer (171) (266))

$$\begin{aligned} \mathcal{K} &= \{[\mathbf{0} \ X] a \zeta \mid a^T \mathbf{1} = 1, a \succeq 0, \zeta \geq 0\} \\ &= \{X a \zeta \mid a^T \mathbf{1} \leq 1, a \succeq 0, \zeta \geq 0\} \\ &= \{X b \mid b \succeq 0\} \subseteq \mathbb{R}^n \end{aligned} \quad (352)$$

that is simply a conic hull (like (95)) of a finite number N of directions.

△

Whenever cone \mathcal{K} is pointed closed and convex (not only polyhedral), then dual cone \mathcal{K}^* has a halfspace-description in terms of the extreme directions Γ_i of \mathcal{K} :

$$\mathcal{K}^* = \{y \mid \gamma^T y \geq 0 \text{ for all } \gamma \in \{\Gamma_i, i=1 \dots N\} \subseteq \text{rel } \partial \mathcal{K}\} \quad (353)$$

because when $\{\Gamma_i\}$ constitutes any set of generators for \mathcal{K} , the discretization result in §2.13.4.1 allows relaxation of the requirement $\forall x \in \mathcal{K}$ in (270) to $\forall \gamma \in \{\Gamma_i\}$ directly.^{2.61} That dual cone so defined is unique, identical to (270), polyhedral whenever the number of generators N is finite

$$\mathcal{K}^* = \{y \mid X^T y \succeq 0\} \subseteq \mathbb{R}^n \quad (329)$$

and has nonempty interior because \mathcal{K} is assumed pointed (but \mathcal{K}^* is not necessarily pointed unless \mathcal{K} has nonempty interior (§2.13.1.1)).

^{2.61}The extreme directions of \mathcal{K} constitute a minimal set of generators. Formulae and conversions to vertex-descriptions of the dual cone are in §2.13.9 and §2.13.11.

2.13.6.1 Facet normal & extreme direction

We see from (329) that the conically independent generators of cone \mathcal{K} (namely, the extreme directions of pointed closed convex cone \mathcal{K} constituting the N columns of X) each define an inward-normal to a hyperplane supporting dual cone \mathcal{K}^* (§2.4.2.6.1) and exposing a dual facet when N is finite. Were \mathcal{K}^* pointed and finitely generated, then by conjugation the dual statement would also hold; *id est*, the extreme directions of pointed \mathcal{K}^* each define an inward-normal to a hyperplane supporting \mathcal{K} and exposing a facet when N is finite. Examine Figure 52 or Figure 57, for example.

We may conclude, the extreme directions of proper polyhedral \mathcal{K} are respectively orthogonal to the facets of \mathcal{K}^* ; likewise, the extreme directions of proper polyhedral \mathcal{K}^* are respectively orthogonal to the facets of \mathcal{K} .

2.13.7 Biorthogonal expansion by example

2.13.7.0.1 Example. Relationship to dual polyhedral cone.

Simplicial cone \mathcal{K} illustrated in Figure 58 induces a partial order on \mathbb{R}^2 . All points greater than x with respect to \mathcal{K} , for example, are contained in the translated cone $x + \mathcal{K}$. The extreme directions Γ_1 and Γ_2 of \mathcal{K} do not make an orthogonal set; neither do extreme directions Γ_3 and Γ_4 of dual cone \mathcal{K}^* ; rather, we have the *biorthogonality condition*, [325]

$$\begin{aligned} \Gamma_4^T \Gamma_1 &= \Gamma_3^T \Gamma_2 = 0 \\ \Gamma_3^T \Gamma_1 &\neq 0, \quad \Gamma_4^T \Gamma_2 \neq 0 \end{aligned} \quad (354)$$

Biorthogonal expansion of $x \in \mathcal{K}$ is then

$$x = \Gamma_1 \frac{\Gamma_3^T x}{\Gamma_3^T \Gamma_1} + \Gamma_2 \frac{\Gamma_4^T x}{\Gamma_4^T \Gamma_2} \quad (355)$$

where $\Gamma_3^T x / (\Gamma_3^T \Gamma_1)$ is the nonnegative coefficient of nonorthogonal projection (§E.6.1) of x on Γ_1 in the direction orthogonal to Γ_3 (y in Figure 142, p.661), and where $\Gamma_4^T x / (\Gamma_4^T \Gamma_2)$ is the nonnegative coefficient of nonorthogonal projection of x on Γ_2 in the direction orthogonal to Γ_4 (z in Figure 142); they are coordinates in this nonorthogonal system. Those coefficients must be nonnegative $x \succeq_{\mathcal{K}} 0$ because $x \in \mathcal{K}$ (295) and \mathcal{K} is simplicial.

If we ascribe the extreme directions of \mathcal{K} to the columns of a matrix

$$X \triangleq [\Gamma_1 \quad \Gamma_2] \quad (356)$$




Figure 58: (*confer* Figure 142) Simplicial cone \mathcal{K} in \mathbb{R}^2 and its dual \mathcal{K}^* drawn truncated. Conically independent generators Γ_1 and Γ_2 constitute extreme directions of \mathcal{K} while Γ_3 and Γ_4 constitute extreme directions of \mathcal{K}^* . Dotted ray-pairs bound translated cones \mathcal{K} . Point x is comparable to point z (and *vice versa*) but not to y ; $z \succeq_{\mathcal{K}} x \Leftrightarrow z - x \in \mathcal{K} \Leftrightarrow z - x \succeq_{\mathcal{K}} 0$ iff \exists nonnegative coordinates for biorthogonal expansion of $z - x$. Point y is not comparable to z because z does not belong to $y \pm \mathcal{K}$. Translating a negated cone is quite helpful for visualization: $u \preceq_{\mathcal{K}} w \Leftrightarrow u \in w - \mathcal{K} \Leftrightarrow u - w \preceq_{\mathcal{K}} 0$. Points need not belong to \mathcal{K} to be comparable; *e.g.*, all points less than w (with respect to \mathcal{K}) belong to $w - \mathcal{K}$.

then we find that the pseudoinverse transpose matrix

$$X^{\dagger T} = \begin{bmatrix} \Gamma_3 \frac{1}{\Gamma_3^T \Gamma_1} & \Gamma_4 \frac{1}{\Gamma_4^T \Gamma_2} \end{bmatrix} \quad (357)$$

holds the extreme directions of the dual cone. Therefore,

$$x = XX^{\dagger}x \quad (363)$$

is the biorthogonal expansion (355) (§E.0.1), and the biorthogonality condition (354) can be expressed succinctly (§E.1.1)^{2.62}

$$X^{\dagger}X = I \quad (364)$$

Expansion $w = XX^{\dagger}w$ for any $w \in \mathbb{R}^2$ is unique if and only if the extreme directions of \mathcal{K} are linearly independent; *id est*, iff X has no nullspace. \square

2.13.7.1 Pointed cones and biorthogonality

Biorthogonality condition $X^{\dagger}X = I$ from Example 2.13.7.0.1 means Γ_1 and Γ_2 are linearly independent generators of \mathcal{K} (§B.1.1.1); generators because every $x \in \mathcal{K}$ is their conic combination. From §2.10.2 we know that means Γ_1 and Γ_2 must be extreme directions of \mathcal{K} .

A biorthogonal expansion is necessarily associated with a pointed closed convex cone; pointed, otherwise there can be no extreme directions (§2.8.1). We will address biorthogonal expansion with respect to a pointed polyhedral cone having empty interior in §2.13.8.

2.13.7.1.1 Example. *Expansions implied by diagonalization.* (confer §6.5.3.2.1) When matrix $X \in \mathbb{R}^{M \times M}$ is diagonalizable (§A.5),

$$X = S\Lambda S^{-1} = [s_1 \cdots s_M] \Lambda \begin{bmatrix} w_1^T \\ \vdots \\ w_M^T \end{bmatrix} = \sum_{i=1}^M \lambda_i s_i w_i^T \quad (1449)$$

^{2.62}Possibly confusing is the fact that formula $XX^{\dagger}x$ is simultaneously the orthogonal projection of x on $\mathcal{R}(X)$ (1797), and a sum of nonorthogonal projections of $x \in \mathcal{R}(X)$ on the range of each and every column of full-rank X skinny-or-square (§E.5.0.0.2).

coordinates for biorthogonal expansion are its eigenvalues λ_i (contained in diagonal matrix Λ) when expanded in S ;

$$X = SS^{-1}X = [s_1 \cdots s_M] \begin{bmatrix} w_1^T X \\ \vdots \\ w_M^T X \end{bmatrix} = \sum_{i=1}^M \lambda_i s_i w_i^T \quad (358)$$

Coordinate value depend upon the geometric relationship of X to its linearly independent eigenmatrices $s_i w_i^T$. (§A.5.1, §B.1.1)

- Eigenmatrices $s_i w_i^T$ are linearly independent dyads constituted by right and left eigenvectors of diagonalizable X and are generators of some pointed polyhedral cone \mathcal{K} in a subspace of $\mathbb{R}^{M \times M}$.

When S is real and X belongs to that polyhedral cone \mathcal{K} , for example, then coordinates of expansion (the eigenvalues λ_i) must be nonnegative.

When $X = Q\Lambda Q^T$ is symmetric, coordinates for biorthogonal expansion are its eigenvalues when expanded in Q ; *id est*, for $X \in \mathbb{S}^M$

$$X = QQ^T X = \sum_{i=1}^M q_i q_i^T X = \sum_{i=1}^M \lambda_i q_i q_i^T \in \mathbb{S}^M \quad (359)$$

becomes an orthogonal expansion with *orthonormality condition* $Q^T Q = I$ where λ_i is the i^{th} eigenvalue of X , q_i is the corresponding i^{th} eigenvector arranged columnar in orthogonal matrix

$$Q = [q_1 \ q_2 \ \cdots \ q_M] \in \mathbb{R}^{M \times M} \quad (360)$$

and where eigenmatrix $q_i q_i^T$ is an extreme direction of some pointed polyhedral cone $\mathcal{K} \subset \mathbb{S}^M$ and an extreme direction of the positive semidefinite cone \mathbb{S}_+^M .

- Orthogonal expansion is a special case of biorthogonal expansion of $X \in \text{aff } \mathcal{K}$ occurring when polyhedral cone \mathcal{K} is any rotation about the origin of an orthant belonging to a subspace.

Similarly, when $X = Q\Lambda Q^T$ belongs to the positive semidefinite cone in the subspace of symmetric matrices, coordinates for orthogonal expansion

must be its nonnegative eigenvalues (1363) when expanded in Q ; *id est*, for $X \in \mathbb{S}_+^M$

$$X = QQ^T X = \sum_{i=1}^M q_i q_i^T X = \sum_{i=1}^M \lambda_i q_i q_i^T \in \mathbb{S}_+^M \quad (361)$$

where $\lambda_i \geq 0$ is the i^{th} eigenvalue of X . This means matrix X simultaneously belongs to the positive semidefinite cone and to the pointed polyhedral cone \mathcal{K} formed by the conic hull of its eigenmatrices. \square

2.13.7.1.2 Example. Expansion respecting nonpositive orthant.

Suppose $x \in \mathcal{K}$ any orthant in \mathbb{R}^n .^{2.63} Then coordinates for biorthogonal expansion of x must be nonnegative; in fact, absolute value of the Cartesian coordinates.

Suppose, in particular, x belongs to the nonpositive orthant $\mathcal{K} = \mathbb{R}_-^n$. Then the biorthogonal expansion becomes an orthogonal expansion

$$x = XX^T x = \sum_{i=1}^n -e_i (-e_i^T x) = \sum_{i=1}^n -e_i |e_i^T x| \in \mathbb{R}_-^n \quad (362)$$

and the coordinates of expansion are nonnegative. For this orthant \mathcal{K} we have orthonormality condition $X^T X = I$ where $X = -I$, $e_i \in \mathbb{R}^n$ is a standard basis vector, and $-e_i$ is an extreme direction (§2.8.1) of \mathcal{K} .

Of course, this expansion $x = XX^T x$ applies more broadly to domain \mathbb{R}^n , but then the coordinates each belong to all of \mathbb{R} . \square

2.13.8 Biorthogonal expansion, derivation

Biorthogonal expansion is a means for determining coordinates in a pointed conic coordinate system characterized by a nonorthogonal basis. Study of nonorthogonal bases invokes pointed polyhedral cones and their duals; extreme directions of a cone \mathcal{K} are assumed to constitute the *basis* while those of the dual cone \mathcal{K}^* determine coordinates.

Unique biorthogonal expansion with respect to \mathcal{K} depends upon existence of its linearly independent extreme directions: Polyhedral cone \mathcal{K} must be pointed; then it possesses extreme directions. Those extreme directions must be linearly independent to uniquely represent any point in their span.

^{2.63}An orthant is simplicial and self-dual.

We consider nonempty pointed polyhedral cone \mathcal{K} having possibly empty interior; *id est*, we consider a basis spanning a subspace. Then we need only observe that section of dual cone \mathcal{K}^* in the affine hull of \mathcal{K} because, by *expansion* of x , membership $x \in \text{aff } \mathcal{K}$ is implicit and because any breach of the ordinary dual cone into ambient space becomes irrelevant (§2.13.9.3). *Biorthogonal expansion*

$$x = XX^\dagger x \in \text{aff } \mathcal{K} = \text{aff cone}(X) \quad (363)$$

is expressed in the extreme directions $\{\Gamma_i\}$ of \mathcal{K} arranged columnar in

$$X = [\Gamma_1 \ \Gamma_2 \ \cdots \ \Gamma_N] \in \mathbb{R}^{n \times N} \quad (254)$$

under assumption of *biorthogonality*

$$X^\dagger X = I \quad (364)$$

where † denotes matrix pseudoinverse (§E). We therefore seek, in this section, a vertex-description for $\mathcal{K}^* \cap \text{aff } \mathcal{K}$ in terms of linearly independent dual generators $\{\Gamma_i^*\} \subset \text{aff } \mathcal{K}$ in the same finite quantity^{2.64} as the extreme directions $\{\Gamma_i\}$ of

$$\mathcal{K} = \text{cone}(X) = \{Xa \mid a \succeq 0\} \subseteq \mathbb{R}^n \quad (95)$$

We assume the quantity of extreme directions N does not exceed the dimension n of ambient vector space because, otherwise, the expansion could not be unique; *id est*, assume N linearly independent extreme directions hence $N \leq n$ (X *skinny*^{2.65}-or-square full-rank). In other words, fat full-rank matrix X is prohibited by uniqueness because of the existence of an infinity of right-inverses;

- polyhedral cones whose extreme directions number in excess of the ambient space dimension are precluded in biorthogonal expansion.

^{2.64}When \mathcal{K} is contained in a proper subspace of \mathbb{R}^n , the ordinary dual cone \mathcal{K}^* will have more generators in any minimal set than \mathcal{K} has extreme directions.

^{2.65}“Skinny” meaning thin; more rows than columns.

2.13.8.1 $x \in \mathcal{K}$

Suppose x belongs to $\mathcal{K} \subseteq \mathbb{R}^n$. Then $x = Xa$ for some $a \succeq 0$. Vector a is unique only when $\{\Gamma_i\}$ is a linearly independent set.^{2.66} Vector $a \in \mathbb{R}^N$ can take the form $a = Bx$ if $\mathcal{R}(B) = \mathbb{R}^N$. Then we require $Xa = XBx = x$ and $Bx = BXa = a$. The pseudoinverse $B = X^\dagger \in \mathbb{R}^{N \times n}$ (§E) is suitable when X is skinny-or-square and full-rank. In that case $\text{rank } X = N$, and for all $c \succeq 0$ and $i = 1 \dots N$

$$a \succeq 0 \Leftrightarrow X^\dagger X a \succeq 0 \Leftrightarrow a^\top X^\top X^\dagger c \geq 0 \Leftrightarrow \Gamma_i^\top X^\dagger c \geq 0 \quad (365)$$

The penultimate inequality follows from the *generalized inequality and membership corollary*, while the last inequality is a consequence of that corollary's discretization (§2.13.4.2.1).^{2.67} From (365) and (353) we deduce

$$\mathcal{K}^* \cap \text{aff } \mathcal{K} = \text{cone}(X^{\dagger\top}) = \{X^{\dagger\top} c \mid c \succeq 0\} \subseteq \mathbb{R}^n \quad (366)$$

is the vertex-description for that section of \mathcal{K}^* in the affine hull of \mathcal{K} because $\mathcal{R}(X^{\dagger\top}) = \mathcal{R}(X)$ by definition of the pseudoinverse. From (281), we know $\mathcal{K}^* \cap \text{aff } \mathcal{K}$ must be pointed if $\text{relint } \mathcal{K}$ is logically assumed nonempty with respect to $\text{aff } \mathcal{K}$.

Conversely, suppose full-rank skinny-or-square matrix ($N \leq n$)

$$X^{\dagger\top} \triangleq \begin{bmatrix} \Gamma_1^* & \Gamma_2^* & \dots & \Gamma_N^* \end{bmatrix} \in \mathbb{R}^{n \times N} \quad (367)$$

comprises the extreme directions $\{\Gamma_i^*\} \subset \text{aff } \mathcal{K}$ of the dual-cone intersection with the affine hull of \mathcal{K} .^{2.68} From the *discretized membership theorem* and

^{2.66}Conic independence alone (§2.10) is insufficient to guarantee uniqueness.
^{2.67}

$$a \succeq 0 \Leftrightarrow a^\top X^\top X^\dagger c \geq 0 \quad \forall (c \succeq 0 \Leftrightarrow a^\top X^\top X^\dagger c \geq 0 \quad \forall a \succeq 0) \\ \forall (c \succeq 0 \Leftrightarrow \Gamma_i^\top X^\dagger c \geq 0 \quad \forall i) \quad \blacklozenge$$

Intuitively, any nonnegative vector a is a conic combination of the standard basis $\{e_i \in \mathbb{R}^N\}$; $a \succeq 0 \Leftrightarrow a_i e_i \succeq 0$ for all i . The last inequality in (365) is a consequence of the fact that $x = Xa$ may be any extreme direction of \mathcal{K} , in which case a is a standard basis vector; $a = e_i \succeq 0$. Theoretically, because $c \succeq 0$ defines a pointed polyhedral cone (in fact, the nonnegative orthant in \mathbb{R}^N), we can take (365) one step further by discretizing c :

$$a \succeq 0 \Leftrightarrow \Gamma_i^\top \Gamma_j^* \geq 0 \text{ for } i, j = 1 \dots N \Leftrightarrow X^\dagger X \geq \mathbf{0}$$

In words, $X^\dagger X$ must be a matrix whose entries are each nonnegative.

^{2.68}When closed convex cone \mathcal{K} has empty interior, \mathcal{K}^* has no extreme directions.

(285) we get a partial dual to (353); *id est*, assuming $x \in \text{aff cone } X$

$$x \in \mathcal{K} \Leftrightarrow \gamma^{*\text{T}}x \geq 0 \text{ for all } \gamma^* \in \left\{ \Gamma_i^*, i=1 \dots N \right\} \subset \partial \mathcal{K}^* \cap \text{aff } \mathcal{K} \quad (368)$$

$$\Leftrightarrow X^\dagger x \succeq 0 \quad (369)$$

that leads to a partial halfspace-description,

$$\mathcal{K} = \{x \in \text{aff cone } X \mid X^\dagger x \succeq 0\} \quad (370)$$

For $\gamma^* = X^{\dagger\text{T}}e_i$, any $x = Xa$, and for all i we have $e_i^\text{T}X^\dagger Xa = e_i^\text{T}a \geq 0$ only when $a \succeq 0$. Hence $x \in \mathcal{K}$.

When X is full-rank, then the unique biorthogonal expansion of $x \in \mathcal{K}$ becomes (363)

$$x = XX^\dagger x = \sum_{i=1}^N \Gamma_i \Gamma_i^{*\text{T}} x \quad (371)$$

whose *coordinates* $\Gamma_i^{*\text{T}}x$ must be nonnegative because \mathcal{K} is assumed pointed closed and convex. Whenever X is full-rank, so is its pseudoinverse X^\dagger . (§E) In the present case, the columns of $X^{\dagger\text{T}}$ are linearly independent and generators of the dual cone $\mathcal{K}^* \cap \text{aff } \mathcal{K}$; hence, the columns constitute its extreme directions. (§2.10.2) That section of the dual cone is itself a polyhedral cone (by (260) or the *cone intersection theorem*, §2.7.2.1.1) having the same number of extreme directions as \mathcal{K} .

2.13.8.2 $x \in \text{aff } \mathcal{K}$

The extreme directions of \mathcal{K} and $\mathcal{K}^* \cap \text{aff } \mathcal{K}$ have a distinct relationship; because $X^\dagger X = I$, then for $i, j = 1 \dots N$, $\Gamma_i^\text{T} \Gamma_i^* = 1$, while for $i \neq j$, $\Gamma_i^\text{T} \Gamma_j^* = 0$. Yet neither set of extreme directions, $\{\Gamma_i\}$ nor $\{\Gamma_i^*\}$, is necessarily orthogonal. This is a biorthogonality condition, precisely, [325, §2.2.4] [180] implying each set of extreme directions is linearly independent. (§B.1.1.1)

The biorthogonal expansion therefore applies more broadly; meaning, for any $x \in \text{aff } \mathcal{K}$, vector x can be uniquely expressed $x = Xb$ where $b \in \mathbb{R}^N$ because $\text{aff } \mathcal{K}$ contains the origin. Thus, for any such $x \in \mathcal{R}(X)$ (*confer* §E.1.1), biorthogonal expansion (371) becomes $x = XX^\dagger Xb = Xb$.

2.13.9 Formulae finding dual cone

2.13.9.1 Pointed \mathcal{K} , dual, X skinny-or-square full-rank

We wish to derive expressions for a convex cone and its ordinary dual under the general assumptions: pointed polyhedral \mathcal{K} denoted by its linearly independent extreme directions arranged columnar in matrix X such that

$$\text{rank}(X \in \mathbb{R}^{n \times N}) = N \stackrel{\Delta}{=} \dim \text{aff } \mathcal{K} \leq n \quad (372)$$

The vertex-description is given:

$$\mathcal{K} = \{Xa \mid a \succeq 0\} \subseteq \mathbb{R}^n \quad (373)$$

from which a halfspace-description for the dual cone follows directly:

$$\mathcal{K}^* = \{y \in \mathbb{R}^n \mid X^T y \succeq 0\} \quad (374)$$

By defining a matrix

$$X^\perp \stackrel{\Delta}{=} \text{basis } \mathcal{N}(X^T) \quad (375)$$

(a columnar basis for the orthogonal complement of $\mathcal{R}(X)$), we can say

$$\text{aff cone } X = \text{aff } \mathcal{K} = \{x \mid X^{\perp T} x = \mathbf{0}\} \quad (376)$$

meaning \mathcal{K} lies in a subspace, perhaps \mathbb{R}^n . Thus a halfspace-description

$$\mathcal{K} = \{x \in \mathbb{R}^n \mid X^\dagger x \succeq 0, X^{\perp T} x = \mathbf{0}\} \quad (377)$$

and a vertex-description^{2.69} from (285)

$$\mathcal{K}^* = \{[X^{\dagger T} \quad X^\perp \quad -X^\perp]b \mid b \succeq 0\} \subseteq \mathbb{R}^n \quad (378)$$

These results are summarized for a pointed polyhedral cone, having linearly independent generators, and its ordinary dual:

Cone Table 1	\mathcal{K}	\mathcal{K}^*
vertex-description	X	$X^{\dagger T}, \pm X^\perp$
halfspace-description	$X^\dagger, X^{\perp T}$	X^T

^{2.69}These descriptions are not unique. A vertex-description of the dual cone, for example, might use four conically independent generators for a plane (§2.10.0.0.1, Figure 45) when only three would suffice.

2.13.9.2 Simplicial case

When a convex cone is simplicial (§2.12.3), Cone Table 1 simplifies because then $\text{aff cone } X = \mathbb{R}^n$: For square X and assuming simplicial \mathcal{K} such that

$$\text{rank}(X \in \mathbb{R}^{n \times n}) = n \stackrel{\Delta}{=} \dim \text{aff } \mathcal{K} = n \quad (379)$$

we have

Cone Table S	\mathcal{K}	\mathcal{K}^*
vertex-description	X	$X^{\dagger T}$
halfspace-description	X^\dagger	X^T

For example, vertex-description (378) simplifies to

$$\mathcal{K}^* = \{X^{\dagger T} b \mid b \succeq 0\} \subset \mathbb{R}^n \quad (380)$$

Now, because $\dim \mathcal{R}(X) = \dim \mathcal{R}(X^{\dagger T})$, (§E) the dual cone \mathcal{K}^* is simplicial whenever \mathcal{K} is.

2.13.9.3 Cone membership relations in a subspace

It is obvious, by definition (270) of the ordinary dual cone \mathcal{K}^* in ambient vector space \mathcal{R} , that its determination instead in subspace $\mathcal{M} \subseteq \mathcal{R}$ is identical to its intersection with \mathcal{M} ; *id est*, assuming closed convex cone $\mathcal{K} \subseteq \mathcal{M}$ and $\mathcal{K}^* \subseteq \mathcal{R}$

$$(\mathcal{K}^* \text{ were ambient } \mathcal{M}) \equiv (\mathcal{K}^* \text{ in ambient } \mathcal{R}) \cap \mathcal{M} \quad (381)$$

because

$$\{y \in \mathcal{M} \mid \langle y, x \rangle \geq 0 \text{ for all } x \in \mathcal{K}\} = \{y \in \mathcal{R} \mid \langle y, x \rangle \geq 0 \text{ for all } x \in \mathcal{K}\} \cap \mathcal{M} \quad (382)$$

From this, a constrained membership relation for the ordinary dual cone $\mathcal{K}^* \subseteq \mathcal{R}$, assuming $x, y \in \mathcal{M}$ and closed convex cone $\mathcal{K} \subseteq \mathcal{M}$

$$y \in \mathcal{K}^* \cap \mathcal{M} \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } x \in \mathcal{K} \quad (383)$$

By closure in subspace \mathcal{M} we have conjugation (§2.13.1.1):

$$x \in \mathcal{K} \Leftrightarrow \langle y, x \rangle \geq 0 \text{ for all } y \in \mathcal{K}^* \cap \mathcal{M} \quad (384)$$

This means membership determination in subspace \mathcal{M} requires knowledge of the dual cone only in \mathcal{M} . For sake of completeness, for proper cone \mathcal{K} with respect to subspace \mathcal{M} (*confer*(296))

$$x \in \text{int } \mathcal{K} \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } y \in \mathcal{K}^* \cap \mathcal{M}, y \neq \mathbf{0} \quad (385)$$

$$x \in \mathcal{K}, x \neq \mathbf{0} \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } y \in \text{int } \mathcal{K}^* \cap \mathcal{M} \quad (386)$$

(By conjugation, we also have the dual relations.) Yet when \mathcal{M} equals $\text{aff } \mathcal{K}$ for \mathcal{K} a closed convex cone

$$x \in \text{rel int } \mathcal{K} \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } y \in \mathcal{K}^* \cap \text{aff } \mathcal{K}, y \neq \mathbf{0} \quad (387)$$

$$x \in \mathcal{K}, x \neq \mathbf{0} \Leftrightarrow \langle y, x \rangle > 0 \text{ for all } y \in \text{rel int}(\mathcal{K}^* \cap \text{aff } \mathcal{K}) \quad (388)$$

2.13.9.4 Subspace $\mathcal{M} = \text{aff } \mathcal{K}$

Assume now a subspace \mathcal{M} that is the affine hull of cone \mathcal{K} : Consider again a pointed polyhedral cone \mathcal{K} denoted by its extreme directions arranged columnar in matrix X such that

$$\text{rank}(X \in \mathbb{R}^{n \times N}) = N \stackrel{\Delta}{=} \dim \text{aff } \mathcal{K} \leq n \quad (372)$$

We want expressions for the convex cone and its dual in subspace $\mathcal{M} = \text{aff } \mathcal{K}$:

Cone Table A	\mathcal{K}	$\mathcal{K}^* \cap \text{aff } \mathcal{K}$
vertex-description	X	$X^{\dagger T}$
halfspace-description	$X^{\dagger}, X^{\perp T}$	$X^T, X^{\perp T}$

When $\dim \text{aff } \mathcal{K} = n$, this table reduces to Cone Table **S**. These descriptions facilitate work in a proper subspace. The subspace of symmetric matrices \mathbb{S}^N , for example, often serves as ambient space. [2.70](#)

2.13.9.4.1 Exercise. *Conically independent columns and rows.*

We suspect the number of conically independent columns (rows) of X to be the same for $X^{\dagger T}$, where \dagger denotes matrix pseudoinverse (§E). Prove whether it holds that the columns (rows) of X are c.i. \Leftrightarrow the columns (rows) of $X^{\dagger T}$ are c.i. ▼

^{2.70}The dual cone of positive semidefinite matrices $\mathbb{S}_+^{N*} = \mathbb{S}_+^N$ remains in \mathbb{S}^N by convention, whereas the ordinary dual cone would venture into $\mathbb{R}^{N \times N}$.




Figure 59: Simplicial cones. **(a)** Monotone nonnegative cone $\mathcal{K}_{\mathcal{M}+}$ and its dual $\mathcal{K}_{\mathcal{M}+}^*$ (drawn truncated) in \mathbb{R}^2 . **(b)** Monotone nonnegative cone and boundary of its dual (both drawn truncated) in \mathbb{R}^3 . Extreme directions of $\mathcal{K}_{\mathcal{M}+}^*$ are indicated.

2.13.9.4.2 Example. *Monotone nonnegative cone.*

[53, exer.2.33] [313, §2] Simplicial cone (§2.12.3.1.1) $\mathcal{K}_{\mathcal{M}+}$ is the cone of all nonnegative vectors having their entries sorted in nonincreasing order:

$$\begin{aligned}\mathcal{K}_{\mathcal{M}+} &\triangleq \{x \mid x_1 \geq x_2 \geq \cdots \geq x_n \geq 0\} \subseteq \mathbb{R}_+^n \\ &= \{x \mid (e_i - e_{i+1})^T x \geq 0, i = 1 \dots n-1, e_n^T x \geq 0\} \\ &= \{x \mid X^\dagger x \succeq 0\}\end{aligned}\quad (389)$$

a halfspace-description where e_i is the i^{th} standard basis vector, and where^{2.71}

$$X^{\dagger T} \triangleq [e_1 - e_2 \quad e_2 - e_3 \quad \cdots \quad e_n] \in \mathbb{R}^{n \times n} \quad (390)$$

For any vectors x and y , simple algebra demands

$$\begin{aligned}x^T y &= \sum_{i=1}^n x_i y_i = (x_1 - x_2)y_1 + (x_2 - x_3)(y_1 + y_2) + (x_3 - x_4)(y_1 + y_2 + y_3) + \cdots \\ &\quad + (x_{n-1} - x_n)(y_1 + \cdots + y_{n-1}) + x_n(y_1 + \cdots + y_n)\end{aligned}\quad (391)$$

Because $x_i - x_{i+1} \geq 0 \forall i$ by assumption whenever $x \in \mathcal{K}_{\mathcal{M}+}$, we can employ dual generalized inequalities (293) with respect to the self-dual nonnegative orthant \mathbb{R}_+^n to find the halfspace-description of the dual cone $\mathcal{K}_{\mathcal{M}+}^*$. We can say $x^T y \geq 0$ for all $X^\dagger x \succeq 0$ [*sic*] if and only if

$$y_1 \geq 0, \quad y_1 + y_2 \geq 0, \quad \dots, \quad y_1 + y_2 + \cdots + y_n \geq 0 \quad (392)$$

id est,

$$x^T y \geq 0 \quad \forall X^\dagger x \succeq 0 \quad \Leftrightarrow \quad X^T y \succeq 0 \quad (393)$$

where

$$X = [e_1 \quad e_1 + e_2 \quad e_1 + e_2 + e_3 \quad \cdots \quad \mathbf{1}] \in \mathbb{R}^{n \times n} \quad (394)$$

Because $X^\dagger x \succeq 0$ connotes membership of x to pointed $\mathcal{K}_{\mathcal{M}+}$, then by (270) the dual cone we seek comprises all y for which (393) holds; thus its halfspace-description

$$\mathcal{K}_{\mathcal{M}+}^* = \{y \succeq 0\} = \{y \mid \sum_{i=1}^k y_i \geq 0, k = 1 \dots n\} = \{y \mid X^T y \succeq 0\} \subset \mathbb{R}^n \quad (395)$$

^{2.71}With X^\dagger in hand, we might concisely scribe the remaining vertex- and halfspace-descriptions from the tables for $\mathcal{K}_{\mathcal{M}+}$ and its dual. Instead we use dual generalized inequalities in their derivation.




Figure 60: Monotone cone $\mathcal{K}_{\mathcal{M}}$ and its dual $\mathcal{K}_{\mathcal{M}}^*$ (drawn truncated) in \mathbb{R}^2 .

The monotone nonnegative cone and its dual are simplicial, illustrated for two Euclidean spaces in Figure 59.

From §2.13.6.1, the extreme directions of proper $\mathcal{K}_{\mathcal{M}+}$ are respectively orthogonal to the facets of $\mathcal{K}_{\mathcal{M}+}^*$. Because $\mathcal{K}_{\mathcal{M}+}^*$ is simplicial, the inward-normals to its facets constitute the linearly independent rows of X^T by (395). Hence the vertex-description for $\mathcal{K}_{\mathcal{M}+}$ employs the columns of X in agreement with Cone Table S because $X^\dagger = X^{-1}$. Likewise, the extreme directions of proper $\mathcal{K}_{\mathcal{M}+}^*$ are respectively orthogonal to the facets of $\mathcal{K}_{\mathcal{M}+}$ whose inward-normals are contained in the rows of X^\dagger by (389). So the vertex-description for $\mathcal{K}_{\mathcal{M}+}^*$ employs the columns of $X^{\dagger T}$. \square

2.13.9.4.3 Example. Monotone cone.

(Figure 60, Figure 61) Of nonempty interior but not pointed, the monotone cone is polyhedral and defined by the halfspace-description

$$\mathcal{K}_{\mathcal{M}} \triangleq \{x \in \mathbb{R}^n \mid x_1 \geq x_2 \geq \cdots \geq x_n\} = \{x \in \mathbb{R}^n \mid X^{*T}x \succeq 0\} \quad (396)$$

Its dual is therefore pointed but of empty interior, having vertex-description

$$\mathcal{K}_{\mathcal{M}}^* = \{X^*b \triangleq [e_1 - e_2 \ e_2 - e_3 \ \cdots \ e_{n-1} - e_n]b \mid b \succeq 0\} \subset \mathbb{R}^n \quad (397)$$

where the columns of X^* comprise the extreme directions of $\mathcal{K}_{\mathcal{M}}^*$. Because $\mathcal{K}_{\mathcal{M}}^*$ is pointed and satisfies

$$\text{rank}(X^* \in \mathbb{R}^{n \times N}) = N \triangleq \dim \text{aff } \mathcal{K}^* \leq n \quad (398)$$




Figure 61: Two views of monotone cone $\mathcal{K}_{\mathcal{M}}$ and its dual $\mathcal{K}_{\mathcal{M}}^*$ (drawn truncated) in \mathbb{R}^3 . Monotone cone is not pointed. Dual monotone cone has empty interior. Cartesian coordinate axes are drawn for reference.

where $N = n - 1$, and because $\mathcal{K}_{\mathcal{M}}$ is closed and convex, we may adapt Cone Table 1 (p.179) as follows:

Cone Table 1*	\mathcal{K}^*	$\mathcal{K}^{**} = \mathcal{K}$
vertex-description	X^*	$X^{*\dagger T}, \pm X^{*\perp}$
halfspace-description	$X^{*\dagger}, X^{*\perp T}$	X^{*T}

The vertex-description for $\mathcal{K}_{\mathcal{M}}$ is therefore

$$\mathcal{K}_{\mathcal{M}} = \{[X^{*\dagger T} \quad X^{*\perp} \quad -X^{*\perp}]a \mid a \succeq 0\} \subset \mathbb{R}^n \quad (399)$$

where $X^{*\perp} = \mathbf{1}$ and

$$X^{*\dagger} = \frac{1}{n} \begin{bmatrix} n-1 & -1 & -1 & \cdots & -1 & -1 & -1 \\ n-2 & n-2 & -2 & \ddots & \cdots & -2 & -2 \\ \vdots & n-3 & n-3 & \ddots & -(n-4) & \vdots & -3 \\ 3 & \vdots & n-4 & \ddots & -(n-3) & -(n-3) & \vdots \\ 2 & 2 & \cdots & \ddots & 2 & -(n-2) & -(n-2) \\ 1 & 1 & 1 & \cdots & 1 & 1 & -(n-1) \end{bmatrix} \in \mathbb{R}^{n-1 \times n} \quad (400)$$

while

$$\mathcal{K}_{\mathcal{M}}^* = \{y \in \mathbb{R}^n \mid X^{*\dagger}y \succeq 0, X^{*\perp T}y = \mathbf{0}\} \quad (401)$$

is the dual monotone cone halfspace-description. \square

2.13.9.4.4 Exercise. Inside the monotone cones.

Mathematically describe the respective interior of the monotone nonnegative cone and monotone cone. In three dimensions, also describe the relative interior of each face. \blacktriangledown

2.13.9.5 More pointed cone descriptions with equality condition

Consider pointed polyhedral cone \mathcal{K} having a linearly independent set of generators and whose subspace membership is explicit; *id est*, we are given the ordinary halfspace-description

$$\mathcal{K} = \{x \mid Ax \succeq 0, Cx = \mathbf{0}\} \subseteq \mathbb{R}^n \quad (260a)$$

where $A \in \mathbb{R}^{m \times n}$ and $C \in \mathbb{R}^{p \times n}$. This can be equivalently written in terms of nullspace of C and vector ξ :

$$\mathcal{K} = \{Z\xi \in \mathbb{R}^n \mid AZ\xi \succeq 0\} \quad (402)$$

where $\mathcal{R}(Z \in \mathbb{R}^{n \times n - \text{rank } C}) \triangleq \mathcal{N}(C)$. Assuming (372) is satisfied

$$\text{rank } X \triangleq \text{rank}((AZ)^\dagger \in \mathbb{R}^{n - \text{rank } C \times m}) = m - \ell = \dim \text{aff } \mathcal{K} \leq n - \text{rank } C \quad (403)$$

where ℓ is the number of conically dependent rows in AZ which must be removed to make $\hat{A}Z$ before the cone tables become applicable.^{2.72} Then the results collected in the cone tables admit the assignment $\hat{X} \triangleq (\hat{A}Z)^\dagger \in \mathbb{R}^{n - \text{rank } C \times m - \ell}$, where $\hat{A} \in \mathbb{R}^{m - \ell \times n}$, followed with linear transformation by Z . So we get the vertex-description, for $(\hat{A}Z)^\dagger$ skinny-or-square full-rank,

$$\mathcal{K} = \{Z(\hat{A}Z)^\dagger b \mid b \succeq 0\} \quad (404)$$

From this and (329) we get a halfspace-description of the dual cone

$$\mathcal{K}^* = \{y \in \mathbb{R}^n \mid (Z^T \hat{A}^T)^\dagger Z^T y \succeq 0\} \quad (405)$$

From this and Cone Table 1 (p.179) we get a vertex-description, (1760)

$$\mathcal{K}^* = \{[Z^{\dagger T} (\hat{A}Z)^T \quad C^T \quad -C^T]c \mid c \succeq 0\} \quad (406)$$

Yet because

$$\mathcal{K} = \{x \mid Ax \succeq 0\} \cap \{x \mid Cx = \mathbf{0}\} \quad (407)$$

then, by (285), we get an equivalent vertex-description for the dual cone

$$\begin{aligned} \mathcal{K}^* &= \overline{\{x \mid Ax \succeq 0\}^* + \{x \mid Cx = \mathbf{0}\}^*} \\ &= \{[A^T \quad C^T \quad -C^T]b \mid b \succeq 0\} \end{aligned} \quad (408)$$

from which the conically dependent columns may, of course, be removed.

^{2.72}When the conically dependent rows (§2.10) are removed, the rows remaining must be linearly independent for the cone tables to apply.

2.13.10 Dual cone-translate

First-order optimality condition (321) inspires a dual-cone variant: For any set \mathcal{K} , the negative dual of its translation by any $a \in \mathbb{R}^n$ is

$$\begin{aligned} -(\mathcal{K} - a)^* &\stackrel{\Delta}{=} \{y \in \mathbb{R}^n \mid \langle y, x - a \rangle \leq 0 \text{ for all } x \in \mathcal{K}\} \\ &= \{y \in \mathbb{R}^n \mid \langle y, x \rangle \leq 0 \text{ for all } x \in \mathcal{K} - a\} \end{aligned} \quad (409)$$

a closed convex cone called the *normal cone* to \mathcal{K} at point a . (§E.10.3.2.1) From this, a new membership relation like (290) for closed convex cone \mathcal{K} :

$$y \in -(\mathcal{K} - a)^* \Leftrightarrow \langle y, x - a \rangle \leq 0 \text{ for all } x \in \mathcal{K} \quad (410)$$

2.13.10.1 first-order optimality condition - restatement

(confer §2.13.3) The general first-order necessary and sufficient condition for optimality of solution x^* to a minimization problem with real differentiable convex objective function $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ over convex feasible set \mathcal{C} is [270, §3]

$$-\nabla f(x^*) \in -(\mathcal{C} - x^*)^*, \quad x^* \in \mathcal{C} \quad (411)$$

id est, the negative gradient (§3.1.8) belongs to the normal cone at x^* as in Figure 62.

2.13.10.1.1 Example. Normal cone to orthant.

Consider proper cone $\mathcal{K} = \mathbb{R}_+^n$, the self-dual nonnegative orthant in \mathbb{R}^n . The normal cone to \mathbb{R}_+^n at $a \in \mathcal{K}$ is (1979)

$$\mathcal{K}_{\mathbb{R}_+^n}^\perp(a \in \mathbb{R}_+^n) = -(\mathbb{R}_+^n - a)^* = -\mathbb{R}_+^n \cap a^\perp, \quad a \in \mathbb{R}_+^n \quad (412)$$

where $-\mathbb{R}_+^n = -\mathcal{K}^*$ is the algebraic complement of \mathbb{R}_+^n , and a^\perp is the orthogonal complement of point a . This means: When point a is interior to \mathbb{R}_+^n , the normal cone is the origin. If n_p represents the number of nonzero entries in point $a \in \partial \mathbb{R}_+^n$, then $\dim(-\mathbb{R}_+^n \cap a^\perp) = n - n_p$ and there is a complementary relationship between the nonzero entries in point a and the nonzero entries in any vector $x \in -\mathbb{R}_+^n \cap a^\perp$. \square




Figure 62: Shown is a plausible contour plot in \mathbb{R}^2 of some arbitrary differentiable real convex function $f(x)$ at selected levels α , β , and γ ; *id est*, contours of equal level f (*level sets*) drawn (dashed) in function's domain. From results in §3.1.9 (p.235), gradient $\nabla f(x^*)$ is normal to γ -sublevel set by Definition E.9.1.0.1. Function is minimized over convex set \mathcal{C} at point x^* iff negative gradient $-\nabla f(x^*)$ belongs to normal cone to \mathcal{C} there. In circumstance depicted, normal cone is a ray whose direction is coincident with negative gradient. So, gradient is normal to a hyperplane supporting both \mathcal{C} and the γ -sublevel set.

2.13.10.1.2 Example. *Optimality conditions for conic problem.*

Consider a convex optimization problem having real differentiable convex objective function $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ defined on domain \mathbb{R}^n

$$\begin{aligned} & \underset{x}{\text{minimize}} && f(x) \\ & \text{subject to} && x \in \mathcal{K} \end{aligned} \quad (413)$$

The feasible set is a pointed polyhedral cone \mathcal{K} possessing a linearly independent set of generators and whose subspace membership is made explicit by fat full-rank matrix $C \in \mathbb{R}^{p \times n}$; *id est*, we are given the halfspace-description

$$\mathcal{K} = \{x \mid Ax \succeq 0, Cx = \mathbf{0}\} \subseteq \mathbb{R}^n \quad (260a)$$

where $A \in \mathbb{R}^{m \times n}$. The vertex-description of this cone, assuming $(\hat{A}Z)^\dagger$ skinny-or-square full-rank, is

$$\mathcal{K} = \{Z(\hat{A}Z)^\dagger b \mid b \succeq 0\} \quad (404)$$

where $\hat{A} \in \mathbb{R}^{m-\ell \times n}$, ℓ is the number of conically dependent rows in AZ (§2.10) which must be removed, and $Z \in \mathbb{R}^{n \times n - \text{rank } C}$ holds basis $\mathcal{N}(C)$ columnar.

From optimality condition (321),

$$\nabla f(x^*)^\top (Z(\hat{A}Z)^\dagger b - x^*) \geq 0 \quad \forall b \succeq 0 \quad (414)$$

$$-\nabla f(x^*)^\top Z(\hat{A}Z)^\dagger (b - b^*) \leq 0 \quad \forall b \succeq 0 \quad (415)$$

because

$$x^* \triangleq Z(\hat{A}Z)^\dagger b^* \in \mathcal{K} \quad (416)$$

From membership relation (410) and Example 2.13.10.1.1

$$\begin{aligned} \langle -(Z^\top \hat{A}^\top)^\dagger Z^\top \nabla f(x^*), b - b^* \rangle &\leq 0 \quad \text{for all } b \in \mathbb{R}_+^{m-\ell} \\ &\Leftrightarrow \\ -(Z^\top \hat{A}^\top)^\dagger Z^\top \nabla f(x^*) &\in -\mathbb{R}_+^{m-\ell} \cap b^{\star\perp} \end{aligned} \quad (417)$$

Then the equivalent necessary and sufficient conditions for optimality of the conic problem (413) with pointed polyhedral feasible set \mathcal{K} are: (*confer* (328))

$$(Z^\top \hat{A}^\top)^\dagger Z^\top \nabla f(x^*) \succeq_{\mathbb{R}_+^{m-\ell}} 0, \quad b^* \succeq_{\mathbb{R}_+^{m-\ell}} 0, \quad \nabla f(x^*)^\top Z(\hat{A}Z)^\dagger b^* = 0 \quad (418)$$

When $\mathcal{K} = \mathbb{R}_+^n$, in particular, then $C = \mathbf{0}$, $A = Z = I \in \mathbb{S}^n$; *id est*,

$$\begin{aligned} & \underset{x}{\text{minimize}} && f(x) \\ & \text{subject to} && x \succeq_{\mathbb{R}_+^n} 0 \end{aligned} \quad (419)$$

The necessary and sufficient conditions become (*confer* [53, §4.2.3])

$$\begin{aligned} \nabla f(x^*) \succeq_{\mathbb{R}_+^n} 0, & \quad x^* \succeq_{\mathbb{R}_+^n} 0, & \quad \nabla f(x^*)^T x^* = 0 \end{aligned} \quad (420)$$

□

2.13.10.1.3 Example. *Linear complementarity.* [241] [275]

Given matrix $A \in \mathbb{R}^{n \times n}$ and vector $q \in \mathbb{R}^n$, the *complementarity problem* is a feasibility problem:

$$\begin{aligned} & \text{find} && w, z \\ & \text{subject to} && w \succeq 0 \\ & && z \succeq 0 \\ & && w^T z = 0 \\ & && w = q + Az \end{aligned} \quad (421)$$

Volumes have been written about this problem, most notably by Cottle [76]. The problem is not convex if both vectors w and z are variable. But if one of them is fixed, then the problem becomes convex with a very simple geometric interpretation: Define the affine subset

$$\mathcal{A} \triangleq \{y \in \mathbb{R}^n \mid Ay = w - q\} \quad (422)$$

For $w^T z$ to vanish, there must be a complementary relationship between the nonzero entries of vectors w and z ; *id est*, $w_i z_i = 0 \forall i$. Given $w \succeq 0$, then z belongs to the convex set of feasible solutions:

$$z \in -\mathcal{K}_{\mathbb{R}_+^n}^\perp(w \in \mathbb{R}_+^n) \cap \mathcal{A} = \mathbb{R}_+^n \cap w^\perp \cap \mathcal{A} \quad (423)$$

where $\mathcal{K}_{\mathbb{R}_+^n}^\perp(w)$ is the normal cone to \mathbb{R}_+^n at w (412). If this intersection is nonempty, then the problem is solvable. □

2.13.11 Proper nonsimplicial \mathcal{K} , dual, X fat full-rank

Since conically dependent columns can always be removed from X to construct \mathcal{K} or to determine \mathcal{K}^* (*Wikimization*), then assume we are given a set of N conically independent generators (§2.10) of an arbitrary proper polyhedral cone \mathcal{K} in \mathbb{R}^n arranged columnar in $X \in \mathbb{R}^{n \times N}$ such that $N > n$ (fat) and $\text{rank } X = n$. Having found formula (380) to determine the dual of a simplicial cone, the easiest way to find a vertex-description of proper dual cone \mathcal{K}^* is to first decompose \mathcal{K} into simplicial parts \mathcal{K}_i so that $\mathcal{K} = \bigcup \mathcal{K}_i$.^{2.73} Each component simplicial cone in \mathcal{K} corresponds to some subset of n linearly independent columns from X . The key idea, here, is how the extreme directions of the simplicial parts must remain extreme directions of \mathcal{K} . Finding the dual of \mathcal{K} amounts to finding the dual of each simplicial part:

2.13.11.0.1 Theorem. *Dual cone intersection.* [293, §2.7]

Suppose proper cone $\mathcal{K} \subset \mathbb{R}^n$ equals the union of M simplicial cones \mathcal{K}_i whose extreme directions all coincide with those of \mathcal{K} . Then proper dual cone \mathcal{K}^* is the intersection of M dual simplicial cones \mathcal{K}_i^* ; *id est*,

$$\mathcal{K} = \bigcup_{i=1}^M \mathcal{K}_i \quad \Rightarrow \quad \mathcal{K}^* = \bigcap_{i=1}^M \mathcal{K}_i^* \quad (424)$$

◇

Proof. For $X_i \in \mathbb{R}^{n \times n}$, a complete matrix of linearly independent extreme directions (p.133) arranged columnar, corresponding simplicial \mathcal{K}_i (§2.12.3.1.1) has vertex-description

$$\mathcal{K}_i = \{X_i c \mid c \succeq 0\} \quad (425)$$

Now suppose,

$$\mathcal{K} = \bigcup_{i=1}^M \mathcal{K}_i = \bigcup_{i=1}^M \{X_i c \mid c \succeq 0\} \quad (426)$$

^{2.73}That proposition presupposes, of course, that we know how to perform simplicial decomposition efficiently; also called “triangulation”. [267] [152, §3.1] [153, §3.1] Existence of multiple simplicial parts means expansion of $x \in \mathcal{K}$ like (371) can no longer be unique because the number N of extreme directions in \mathcal{K} exceeds dimension n of the space.

The union of all \mathcal{K}_i can be equivalently expressed

$$\mathcal{K} = \left\{ [X_1 \ X_2 \ \cdots \ X_M] \begin{bmatrix} a \\ b \\ \vdots \\ c \end{bmatrix} \mid a, b, \dots, c \succeq 0 \right\} \quad (427)$$

Because extreme directions of the simplices \mathcal{K}_i are extreme directions of \mathcal{K} by assumption, then by the *extremes theorem* (§2.8.1.1.1),

$$\mathcal{K} = \{ [X_1 \ X_2 \ \cdots \ X_M] d \mid d \succeq 0 \} \quad (428)$$

Defining $X \triangleq [X_1 \ X_2 \ \cdots \ X_M]$ (with any redundant [*sic*] columns optionally removed from X), then \mathcal{K}^* can be expressed, (329) (Cone Table **S**, p.180)

$$\mathcal{K}^* = \{y \mid X^T y \succeq 0\} = \bigcap_{i=1}^M \{y \mid X_i^T y \succeq 0\} = \bigcap_{i=1}^M \mathcal{K}_i^* \quad (429)$$

◆

To find the extreme directions of the dual cone, first we observe that some facets of each simplicial part \mathcal{K}_i are common to facets of \mathcal{K} by assumption, and the union of all those common facets comprises the set of all facets of \mathcal{K} by design. For any particular proper polyhedral cone \mathcal{K} , the extreme directions of dual cone \mathcal{K}^* are respectively orthogonal to the facets of \mathcal{K} . (§2.13.6.1) Then the extreme directions of the dual cone can be found among inward-normals to facets of the component simplicial cones \mathcal{K}_i ; those normals are extreme directions of the dual simplicial cones \mathcal{K}_i^* . From the theorem and Cone Table **S** (p.180),

$$\mathcal{K}^* = \bigcap_{i=1}^M \mathcal{K}_i^* = \bigcap_{i=1}^M \{X_i^{\dagger T} c \mid c \succeq 0\} \quad (430)$$

The set of extreme directions $\{\Gamma_i^*\}$ for proper dual cone \mathcal{K}^* is therefore constituted by those conically independent generators, from the columns of all the dual simplicial matrices $\{X_i^{\dagger T}\}$, that do not violate discrete definition (329) of \mathcal{K}^* ;

$$\{\Gamma_1^*, \Gamma_2^*, \dots, \Gamma_N^*\} = \text{c.i.} \left\{ X_i^{\dagger T}(:, j), \ i=1 \dots M, \ j=1 \dots n \mid X_i^{\dagger}(j, :) \Gamma_\ell \geq 0, \ \ell=1 \dots N \right\} \quad (431)$$

where c.i. denotes selection of only the conically independent vectors from the argument set, argument $(:,j)$ denotes the j^{th} column while $(j,:)$ denotes the j^{th} row, and $\{\Gamma_\ell\}$ constitutes the extreme directions of \mathcal{K} . Figure 46(b) (p.131) shows a cone and its dual found via this algorithm.

2.13.11.0.2 Example. *Dual of \mathcal{K} nonsimplicial in subspace $\text{aff } \mathcal{K}$.*

Given conically independent generators for pointed closed convex cone \mathcal{K} in \mathbb{R}^4 arranged columnar in

$$X = [\Gamma_1 \ \Gamma_2 \ \Gamma_3 \ \Gamma_4] = \begin{bmatrix} 1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & -1 \end{bmatrix} \quad (432)$$

having $\dim \text{aff } \mathcal{K} = \text{rank } X = 3$, (255) then performing the most inefficient simplicial decomposition in $\text{aff } \mathcal{K}$ we find

$$\begin{aligned} X_1 &= \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, & X_2 &= \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & -1 \end{bmatrix} \\ X_3 &= \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & -1 \end{bmatrix}, & X_4 &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & -1 \end{bmatrix} \end{aligned} \quad (433)$$

The corresponding dual simplicial cones in $\text{aff } \mathcal{K}$ have generators respectively columnar in

$$\begin{aligned} 4X_1^{\dagger T} &= \begin{bmatrix} 2 & 1 & 1 \\ -2 & 1 & 1 \\ 2 & -3 & 1 \\ -2 & 1 & -3 \end{bmatrix}, & 4X_2^{\dagger T} &= \begin{bmatrix} 1 & 2 & 1 \\ -3 & 2 & 1 \\ 1 & -2 & 1 \\ 1 & -2 & -3 \end{bmatrix} \\ 4X_3^{\dagger T} &= \begin{bmatrix} 3 & 2 & -1 \\ -1 & 2 & -1 \\ -1 & -2 & 3 \\ -1 & -2 & -1 \end{bmatrix}, & 4X_4^{\dagger T} &= \begin{bmatrix} 3 & -1 & 2 \\ -1 & 3 & -2 \\ -1 & -1 & 2 \\ -1 & -1 & -2 \end{bmatrix} \end{aligned} \quad (434)$$

Applying algorithm (431) we get

$$\begin{bmatrix} \Gamma_1^* & \Gamma_2^* & \Gamma_3^* & \Gamma_4^* \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 & 2 & 3 & 2 \\ 1 & 2 & -1 & -2 \\ 1 & -2 & -1 & 2 \\ -3 & -2 & -1 & -2 \end{bmatrix} \quad (435)$$

whose rank is 3, and is the known result,^{2.74} a conically independent set of generators for that pointed section of the dual cone \mathcal{K}^* in $\text{aff } \mathcal{K}$; *id est*, $\mathcal{K}^* \cap \text{aff } \mathcal{K}$. \square

2.13.11.0.3 Example. *Dual of proper polyhedral \mathcal{K} in \mathbb{R}^4 .*

Given conically independent generators for a full-dimensional pointed closed convex cone \mathcal{K}

$$X = [\Gamma_1 \ \Gamma_2 \ \Gamma_3 \ \Gamma_4 \ \Gamma_5] = \begin{bmatrix} 1 & 1 & 0 & 1 & 0 \\ -1 & 0 & 1 & 0 & 1 \\ 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & -1 & -1 & 0 \end{bmatrix} \quad (436)$$

we count $5!/((5-4)!4!) = 5$ component simplices.^{2.75} Applying algorithm (431), we find the six extreme directions of dual cone \mathcal{K}^* (with $\Gamma_2 = \Gamma_5^*$)

$$X^* = [\Gamma_1^* \ \Gamma_2^* \ \Gamma_3^* \ \Gamma_4^* \ \Gamma_5^* \ \Gamma_6^*] = \begin{bmatrix} 1 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & -1 & 1 \\ 1 & -1 & -1 & 1 & 0 & 0 \end{bmatrix} \quad (437)$$

which means, (§2.13.6.1) this proper polyhedral $\mathcal{K} = \text{cone}(X)$ has six (three-dimensional) facets generated \mathcal{G} by its {extreme directions}:

$$\mathcal{G} \left\{ \begin{array}{l} \mathcal{F}_1 \\ \mathcal{F}_2 \\ \mathcal{F}_3 \\ \mathcal{F}_4 \\ \mathcal{F}_5 \\ \mathcal{F}_6 \end{array} \right\} = \left\{ \begin{array}{l} \Gamma_1 \ \Gamma_2 \ \Gamma_3 \\ \Gamma_1 \ \Gamma_2 \ \Gamma_5 \\ \Gamma_1 \ \Gamma_4 \ \Gamma_5 \\ \Gamma_1 \ \Gamma_3 \ \Gamma_4 \\ \Gamma_3 \ \Gamma_4 \ \Gamma_5 \\ \Gamma_2 \ \Gamma_3 \ \Gamma_5 \end{array} \right\} \quad (438)$$

^{2.74}These calculations proceed so as to be consistent with [97, §6]; as if the ambient vector space were proper subspace $\text{aff } \mathcal{K}$ whose dimension is 3. In that ambient space, \mathcal{K} may be regarded as a proper cone. Yet that author (from the citation) erroneously states dimension of the ordinary dual cone to be 3; it is, in fact, 4.

^{2.75}There are no linearly dependent combinations of three or four extreme directions in the primal cone.

whereas dual proper polyhedral cone \mathcal{K}^* has only five:

$$\mathcal{G} \begin{pmatrix} \mathcal{F}_1^* \\ \mathcal{F}_2^* \\ \mathcal{F}_3^* \\ \mathcal{F}_4^* \\ \mathcal{F}_5^* \end{pmatrix} = \begin{pmatrix} \Gamma_1^* & \Gamma_2^* & \Gamma_3^* & \Gamma_4^* & & \\ \Gamma_1^* & \Gamma_2^* & & & & \Gamma_6^* \\ \Gamma_1^* & & & \Gamma_4^* & \Gamma_5^* & \Gamma_6^* \\ & & \Gamma_3^* & \Gamma_4^* & \Gamma_5^* & \\ & \Gamma_2^* & \Gamma_3^* & & \Gamma_5^* & \Gamma_6^* \end{pmatrix} \quad (439)$$

Six two-dimensional cones, having generators respectively $\{\Gamma_1^* \ \Gamma_3^*\}$ $\{\Gamma_2^* \ \Gamma_4^*\}$ $\{\Gamma_1^* \ \Gamma_5^*\}$ $\{\Gamma_4^* \ \Gamma_6^*\}$ $\{\Gamma_2^* \ \Gamma_5^*\}$ $\{\Gamma_3^* \ \Gamma_6^*\}$, are relatively interior to dual facets; so cannot be two-dimensional faces of \mathcal{K}^* (by Definition 2.6.0.0.3).

We can check this result (437) by reversing the process; we find $6!/((6-4)!4!) - 3 = 12$ component simplices in the dual cone.^{2.76} Applying algorithm (431) to those simplices returns a conically independent set of generators for \mathcal{K} equivalent to (436). \square

2.13.11.0.4 Exercise. *Reaching proper polyhedral cone interior.*

Name two extreme directions Γ_i of cone \mathcal{K} from Example 2.13.11.0.3 whose convex hull passes through that cone's interior. Explain why. Are there two such extreme directions of dual cone \mathcal{K}^* ? \blacktriangledown

A natural question pertains to whether a theory of unique coordinates, like biorthogonal expansion, is extensible to proper cones whose extreme directions number in excess of ambient spatial dimensionality.

2.13.11.0.5 Theorem. *Conic coordinates.*

With respect to vector v in some finite-dimensional Euclidean space \mathbb{R}^n , define a coordinate t_v^* of point x in full-dimensional pointed closed convex cone \mathcal{K}

$$t_v^*(x) \triangleq \sup\{t \in \mathbb{R} \mid x - tv \in \mathcal{K}\} \quad (440)$$

Given points x and y in cone \mathcal{K} , if $t_v^*(x) = t_v^*(y)$ for each and every extreme direction v of \mathcal{K} then $x = y$. \diamond

^{2.76}Three combinations of four dual extreme directions are linearly dependent; they belong to the dual facets. But there are no linearly dependent combinations of three dual extreme directions.

2.13.11.0.6 Proof. Vector $x - t^*v$ must belong to the cone boundary $\partial\mathcal{K}$ by definition (440). So there must exist a nonzero vector λ that is inward-normal to a hyperplane supporting cone \mathcal{K} and containing $x - t^*v$; *id est*, by boundary membership relation for full-dimensional pointed closed convex cones (§2.13.2)

$$x - t^*v \in \partial\mathcal{K} \Leftrightarrow \exists \lambda \neq \mathbf{0} \ni \langle \lambda, x - t^*v \rangle = 0, \quad \lambda \in \mathcal{K}^*, \quad x - t^*v \in \mathcal{K} \quad (300)$$

where

$$\mathcal{K}^* = \{w \in \mathbb{R}^n \mid \langle v, w \rangle \geq 0 \text{ for all } v \in \mathcal{G}(\mathcal{K})\} \quad (334)$$

is the full-dimensional pointed closed convex dual cone. The set $\mathcal{G}(\mathcal{K})$, of possibly infinite cardinality N , comprises generators for cone \mathcal{K} ; *e.g.*, its extreme directions which constitute a minimal generating set. If $x - t^*v$ is nonzero, any such vector λ must belong to the dual cone boundary by conjugate boundary membership relation

$$\lambda \in \partial\mathcal{K}^* \Leftrightarrow \exists x - t^*v \neq \mathbf{0} \ni \langle \lambda, x - t^*v \rangle = 0, \quad x - t^*v \in \mathcal{K}, \quad \lambda \in \mathcal{K}^* \quad (301)$$

where

$$\mathcal{K} = \{z \in \mathbb{R}^n \mid \langle \lambda, z \rangle \geq 0 \text{ for all } \lambda \in \mathcal{G}(\mathcal{K}^*)\} \quad (333)$$

This description of \mathcal{K} means: cone \mathcal{K} is an intersection of halfspaces whose inward-normals are generators of the dual cone. Each and every face of cone \mathcal{K} (except the cone itself) belongs to a hyperplane supporting \mathcal{K} . Each and every vector $x - t^*v$ on the cone boundary must therefore be orthogonal to an extreme direction constituting generators $\mathcal{G}(\mathcal{K}^*)$ of the dual cone.

To the i^{th} extreme direction $v = \Gamma_i \in \mathbb{R}^n$ of cone \mathcal{K} , ascribe a coordinate $t_i^*(x) \in \mathbb{R}$ of x from definition (440). On domain \mathcal{K} , the mapping

$$t^*(x) = \begin{bmatrix} t_1^*(x) \\ \vdots \\ t_N^*(x) \end{bmatrix} : \mathbb{R}^n \rightarrow \mathbb{R}^N \quad (441)$$

has no nontrivial nullspace. Because $x - t^*v$ must belong to $\partial\mathcal{K}$ by definition, the mapping $t^*(x)$ is equivalent to a convex problem (separable in index i) whose objective (by (300)) is tightly bounded below by 0:

$$\begin{aligned} t^*(x) \equiv \arg \underset{t \in \mathbb{R}^N}{\text{minimize}} & \sum_{i=1}^N \Gamma_{j(i)}^* \Gamma_i (x - t_i \Gamma_i) \\ \text{subject to} & \quad x - t_i \Gamma_i \in \mathcal{K}, \quad i = 1 \dots N \end{aligned} \quad (442)$$

where index $j \in \mathcal{I}$ is dependent on i and where (by (333)) $\lambda = \Gamma_j^* \in \mathbb{R}^n$ is an extreme direction of dual cone \mathcal{K}^* that is normal to a hyperplane supporting \mathcal{K} and containing $x - t_i^* \Gamma_i$. Because extreme-direction cardinality N for cone \mathcal{K} is not necessarily the same as for dual cone \mathcal{K}^* , index j must be judiciously selected from a set \mathcal{I} .

To prove injectivity when extreme-direction cardinality $N > n$ exceeds spatial dimension, we need only show mapping $t^*(x)$ to be invertible; [111, thm.9.2.3] *id est*, x is recoverable given $t^*(x)$:

$$\begin{aligned} x = \arg \underset{\tilde{x} \in \mathbb{R}^n}{\text{minimize}} \quad & \sum_{i=1}^N \Gamma_{j(i)}^{*\text{T}} (\tilde{x} - t_i^* \Gamma_i) \\ \text{subject to} \quad & \tilde{x} - t_i^* \Gamma_i \in \mathcal{K}, \quad i = 1 \dots N \end{aligned} \quad (443)$$

The feasible set of this nonseparable convex problem is an intersection of translated full-dimensional pointed closed convex cones $\bigcap_i \mathcal{K} + t_i^* \Gamma_i$. The objective function's linear part describes movement in normal-direction $-\Gamma_j^*$ for each of N hyperplanes. The optimal point of hyperplane intersection is the unique solution x when $\{\Gamma_j^*\}$ comprises n linearly independent normals that come from the dual cone and make the objective vanish. Because the dual cone \mathcal{K}^* is full-dimensional, pointed, closed, and convex by assumption, there exist N extreme directions $\{\Gamma_j^*\}$ from $\mathcal{K}^* \subset \mathbb{R}^n$ that span \mathbb{R}^n . So we need simply choose N spanning dual extreme directions that make the optimal objective vanish. Because such dual extreme directions preexist by (300), $t^*(x)$ is invertible.

Otherwise, in the case $N \leq n$, $t^*(x)$ holds coordinates of biorthogonal expansion. Reconstruction of x is therefore unique. \blacklozenge

2.13.11.1 conic coordinate computation

The foregoing proof of the *conic coordinates theorem* is not constructive; it establishes existence of dual extreme directions $\{\Gamma_j^*\}$ that will reconstruct a point x from its coordinates $t^*(x)$ via (443), but does not prescribe the index set \mathcal{I} . There are at least two computational methods for specifying $\{\Gamma_{j(i)}^*\}$: one is combinatorial but sure to succeed, the other is a geometric method that searches for a minimum of a nonconvex function. We describe the latter:

Convex problem (P)

$$(P) \quad \begin{array}{ll} \text{maximize} & t \\ \text{subject to} & x - tv \in \mathcal{K} \end{array} \quad \begin{array}{ll} \text{minimize} & \lambda^T x \\ \text{subject to} & \lambda^T v = 1 \\ & \lambda \in \mathcal{K}^* \end{array} \quad (D) \quad (444)$$

is equivalent to definition (440) whereas convex problem (D) is its dual,^{2.77} meaning, primal and dual optimal objectives are equal $t^* = \lambda^{*T} x$ assuming Slater's condition (p.260) is satisfied. Under this assumption of strong duality, $\lambda^{*T}(x - t^*v) = t^*(1 - \lambda^{*T}v) = 0$; which implies, the primal problem is equivalent to

$$\begin{array}{ll} \text{minimize} & \lambda^{*T}(x - tv) \\ \text{subject to} & x - tv \in \mathcal{K} \end{array} \quad (P) \quad (445)$$

while the dual problem is equivalent to

$$\begin{array}{ll} \text{minimize} & \lambda^T(x - t^*v) \\ \text{subject to} & \lambda^T v = 1 \\ & \lambda \in \mathcal{K}^* \end{array} \quad (D) \quad (446)$$

Instead given coordinates $t^*(x)$ and a description of cone \mathcal{K} , we propose inversion by alternating solution of primal and dual problems

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^N \Gamma_i^{*T}(x - t_i^* \Gamma_i) \\ \text{subject to} & x - t_i^* \Gamma_i \in \mathcal{K}, \quad i=1 \dots N \end{array} \quad (447)$$

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^N \Gamma_i^{*T}(x^* - t_i^* \Gamma_i) \\ \text{subject to} & \Gamma_i^{*T} \Gamma_i = 1, \quad i=1 \dots N \\ & \Gamma_i^* \in \mathcal{K}^*, \quad i=1 \dots N \end{array} \quad (448)$$

^{2.77}Form the Lagrangian associated with primal problem (P):

$$\begin{aligned} \mathfrak{L}(t, \lambda) &= t + \lambda^T(x - tv) = \lambda^T x + t(1 - \lambda^T v), & \lambda &\succeq 0 \\ & & & \mathcal{K}^* \\ \sup_t \mathfrak{L}(t, \lambda) &= \lambda^T x, & 1 - \lambda^T v &= 0 \end{aligned}$$

Dual variable (Lagrange multiplier [220, p.216]) λ generally has a nonnegative sense for primal maximization with any cone membership constraint, whereas λ would have a nonpositive sense were the primal instead a minimization problem having a membership constraint.

where dual extreme directions Γ_i^* are initialized arbitrarily and ultimately ascertained by the alternation. Convex problems (447) and (448) are iterated until convergence which is guaranteed by virtue of a monotonically nonincreasing real sequence of objective values. Convergence can be fast. The mapping $t^*(x)$ is uniquely inverted when the necessarily nonnegative objective vanishes; *id est*, when $\Gamma_i^{*T}(x^* - t_i^* \Gamma_i) = 0 \quad \forall i$. Here, a zero objective can occur only at the true solution x . But this global optimality condition cannot be guaranteed by the alternation because the common objective function, when regarded in both primal x and dual Γ_i^* variables simultaneously, is generally neither quasiconvex or monotonic.

Conversely, a nonzero objective at convergence is a certificate that inversion was not performed properly. A nonzero objective indicates that the global minimum of a multimodal objective function could not be found by this alternation. That is a flaw in this particular iterative algorithm for inversion; not in theory.^{2.78} A numerical remedy is to reinitialize the Γ_i^* to different values.

^{2.78}The Proof 2.13.11.0.6 that suitable dual extreme directions $\{\Gamma_j^*\}$ always exist means that a global optimization algorithm would always find the zero objective of alternation (447) (448); hence, the unique inversion x . But such an algorithm can be combinatorial.

Chapter 3

Geometry of convex functions

The link between convex sets and convex functions is via the epigraph: A function is convex if and only if its epigraph is a convex set.

–Werner Fenchel

We limit our treatment of *multidimensional functions*^{3.1} to finite-dimensional Euclidean space. Then an icon for a one-dimensional (real) *convex function* is bowl-shaped (Figure 70), whereas the *concave* icon is the inverted bowl; respectively characterized by a unique global minimum and maximum whose existence is assumed. Because of this simple relationship, usage of the term *convexity* is often implicitly inclusive of *concavity* in the literature. Despite the iconic imagery, the reader is reminded that the set of all convex, concave, quasiconvex, and quasiconcave functions contains the *monotonic functions* [181] [190, §2.3.5]; *e.g.*, [53, §3.6, exer.3.46].

^{3.1} vector- or matrix-valued functions including the real functions. Appendix D, with its tables of first- and second-order gradients, is the practical adjunct to this chapter.

3.1 Convex function

3.1.1 real and vector-valued function

Vector-valued function

$$f(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}^M = \begin{bmatrix} f_1(X) \\ \vdots \\ f_M(X) \end{bmatrix} \quad (449)$$

assigns each X in its domain $\text{dom } f$ (a subset of ambient vector space $\mathbb{R}^{p \times k}$) to a specific *element* [228, p.3] of its range (a subset of \mathbb{R}^M). Function $f(X)$ is *linear* in X on its domain if and only if, for each and every $Y, Z \in \text{dom } f$ and $\alpha, \beta \in \mathbb{R}$

$$f(\alpha Y + \beta Z) = \alpha f(Y) + \beta f(Z) \quad (450)$$

A vector-valued function $f(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}^M$ is convex in X if and only if $\text{dom } f$ is a convex set and, for each and every $Y, Z \in \text{dom } f$ and $0 \leq \mu \leq 1$

$$f(\mu Y + (1 - \mu)Z) \preceq_{\mathbb{R}_+^M} \mu f(Y) + (1 - \mu)f(Z) \quad (451)$$

As defined, continuity is implied but not differentiability (nor *smoothness*).^{3.2} Apparently some, but not all, nonlinear functions are convex. Reversing sense of the inequality flips this definition to concavity. Linear (and affine §3.1.6) functions attain equality in this definition. Linear functions are therefore simultaneously convex and concave.

Vector-valued functions are most often compared (165) as in (451) with respect to the M -dimensional self-dual nonnegative orthant \mathbb{R}_+^M , a proper cone.^{3.3} In this case, the test prescribed by (451) is simply a comparison on \mathbb{R} of each *entry* f_i of a vector-valued function f . (§2.13.4.2.3) The vector-valued function case is therefore a straightforward generalization of conventional convexity theory for a real function. This conclusion follows from theory of dual generalized inequalities (§2.13.2.0.1) which asserts

^{3.2}Differentiability is not a requirement for optimization of constrained convex functions by modern numerical methods.

^{3.3}Definition of convexity can be broadened to other (not necessarily proper) cones; referred to in the literature as *K-convexity*. [260]




Figure 63: Each convex real function has a unique minimizer x^* but, for $x \in \mathbb{R}$, $f_1(x) = x^2$ is strictly convex whereas $f_2(x) = \sqrt{x^2} = |x|$ is not. Strict convexity of a real function is therefore only a sufficient condition for minimizer uniqueness.

$$f \text{ convex} \Leftrightarrow w^T f \text{ convex} \quad \forall w \in \mathcal{G}(\mathbb{R}_+^M) \quad (452)$$

shown by substitution of the defining inequality (451). Discretization (§2.13.4.2.1) allows relaxation of the semi-infinite number of conditions $\forall w \succeq 0$ to:

$$\forall w \in \mathcal{G}(\mathbb{R}_+^M) = \{e_i, i = 1 \dots M\} \quad (453)$$

(the standard basis for \mathbb{R}^M and a minimal set of generators (§2.8.1.2) for \mathbb{R}_+^M) from which the stated conclusion follows; *id est*, the test for convexity of a vector-valued function is a comparison on \mathbb{R} of each entry.

3.1.1.0.1 Exercise. Cone of convex functions.

Prove that relation (452) implies: the set of all vector-valued convex functions in \mathbb{R}^M is a convex cone. Indeed, any nonnegatively weighted sum of (strictly) convex functions remains (strictly) convex.^{3.4} Interior to the cone are the strictly convex functions: ▼

^{3.4}Strict case excludes cone's point at origin. By these definitions (452) (455), positively weighted sums mixing convex and strictly convex real functions are not strictly convex because each summand is considered to be an entry from a vector-valued function.

3.1.2 strict convexity

When $f(X)$ instead satisfies, for each and every distinct Y and Z in $\text{dom } f$ and all $0 < \mu < 1$

$$f(\mu Y + (1 - \mu)Z) \underset{\mathbb{R}_+^M}{<} \mu f(Y) + (1 - \mu)f(Z) \quad (454)$$

then we shall say f is a *strictly convex function*.

Similarly to (452)

$$f \text{ strictly convex} \Leftrightarrow w^T f \text{ strictly convex} \quad \forall w \in \mathcal{G}(\mathbb{R}_+^M) \quad (455)$$

discretization allows relaxation of the semi-infinite number of conditions $\forall w \succeq 0, w \neq \mathbf{0}$ (296) to a finite number (453). More tests for strict convexity are given in §3.1.8.1.2, §3.1.11, and §3.2.3.0.2.

Any convex real function $f(X)$ has unique minimum value over any convex subset of its domain. Yet solution to some convex optimization problem is, in general, not unique; *e.g.*, given minimization of a convex real function over some convex *feasible solution* set \mathcal{C}

$$\begin{aligned} & \underset{X}{\text{minimize}} && f(X) \\ & \text{subject to} && X \in \mathcal{C} \end{aligned} \quad (456)$$

any *optimal solution* X^* comes from a convex set of optimal solutions

$$X^* \in \{X \mid f(X) = \inf_{Y \in \mathcal{C}} f(Y)\} \subseteq \mathcal{C} \quad (457)$$

But a strictly convex real function has a unique minimizer X^* ; *id est*, for the optimal solution set in (457) to be a single point, it is sufficient (Figure 63) that $f(X)$ be a strictly convex real^{3.5} function and set \mathcal{C} convex.

Quadratic real functions $x^T A x + b^T x + c$ are convex in x iff $A \succeq 0$. Quadratics characterized by positive definite matrix $A \succ 0$ are strictly convex. The vector 2-norm square $\|x\|^2$ (Euclidean norm square) and Frobenius norm square $\|X\|_F^2$, for example, are strictly convex functions of their respective argument (each absolute norm is convex but not strictly convex). Figure 63(a) illustrates a strictly convex real function.

^{3.5}It is more customary to consider only a real function for the objective of a convex optimization problem because vector- or matrix-valued functions can introduce ambiguity into the optimal objective value. (§2.7.2.2) Study of multidimensional objective functions is called *multicriteria-* [287] or *multiobjective-* or *vector-optimization*.

3.1.3 norm functions, absolute value

A vector *norm* on \mathbb{R}^n is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfying: for $x, y \in \mathbb{R}^n$, $\alpha \in \mathbb{R}$ [201, p.59] [139, p.52]

1. $f(x) \geq 0$ ($f(x) = 0 \Leftrightarrow x = 0$) (nonnegativity)
2. $f(x + y) \leq f(x) + f(y)$ (triangle inequality)
3. $f(\alpha x) = |\alpha|f(x)$ (nonnegative homogeneity)

Most useful of the convex norms are 1-, 2-, and infinity-norm:

$$\begin{aligned} \|x\|_1 = \underset{t \in \mathbb{R}^n}{\text{minimize}} \quad & \mathbf{1}^T t \\ \text{subject to} \quad & -t \preceq x \preceq t \end{aligned} \quad (458)$$

where $|x| = t^*$ (entrywise absolute value equals optimal t).^{3.6}

$$\begin{aligned} \|x\|_2 = \underset{t \in \mathbb{R}}{\text{minimize}} \quad & t \\ \text{subject to} \quad & \begin{bmatrix} tI & x \\ x^T & t \end{bmatrix} \underset{\mathbb{S}_+^{n+1}}{\succeq} 0 \end{aligned} \quad (459)$$

where $\|x\|_2 = \|x\| \triangleq \sqrt{x^T x} = t^*$.

$$\begin{aligned} \|x\|_\infty = \underset{t \in \mathbb{R}}{\text{minimize}} \quad & t \\ \text{subject to} \quad & -t\mathbf{1} \preceq x \preceq t\mathbf{1} \end{aligned} \quad (460)$$

where $\max\{|x_i|, i=1 \dots n\} = t^*$.

$$\begin{aligned} \|x\|_1 = \underset{\alpha \in \mathbb{R}^n, \beta \in \mathbb{R}^n}{\text{minimize}} \quad & \mathbf{1}^T(\alpha + \beta) \\ \text{subject to} \quad & \alpha, \beta \succeq 0 \\ & x = \alpha - \beta \end{aligned} \quad (461)$$

where $|x| = \alpha^* + \beta^*$ because of complementarity $\alpha^{*T}\beta^* = 0$ at optimality. (458) (460) (461) represent linear programs, (459) is a semidefinite program.

^{3.6}Vector $\mathbf{1}$ may be replaced with any positive [*sic*] vector to get absolute value, theoretically, although $\mathbf{1}$ provides the 1-norm.

Over some convex set \mathcal{C} given vector constant y or matrix constant Y

$$\arg \inf_{x \in \mathcal{C}} \|x - y\|_2 = \arg \inf_{x \in \mathcal{C}} \|x - y\|_2^2 \quad (462)$$

$$\arg \inf_{X \in \mathcal{C}} \|X - Y\|_F = \arg \inf_{X \in \mathcal{C}} \|X - Y\|_F^2 \quad (463)$$

are unconstrained quadratic problems. Optimal solution is norm dependent. [53, p.297]

$$\begin{aligned} \underset{x \in \mathbb{R}^n}{\text{minimize}} \quad & \|x\|_1 \\ \text{subject to} \quad & x \in \mathcal{C} \end{aligned} \quad \equiv \quad \begin{aligned} \underset{x \in \mathbb{R}^n, t \in \mathbb{R}}{\text{minimize}} \quad & \mathbf{1}^\top t \\ \text{subject to} \quad & -t \preceq x \preceq t \\ & x \in \mathcal{C} \end{aligned} \quad (464)$$

$$\begin{aligned} \underset{x \in \mathbb{R}^n}{\text{minimize}} \quad & \|x\|_2 \\ \text{subject to} \quad & x \in \mathcal{C} \end{aligned} \quad \equiv \quad \begin{aligned} \underset{x \in \mathbb{R}^n, t \in \mathbb{R}}{\text{minimize}} \quad & t \\ \text{subject to} \quad & \begin{bmatrix} tI & x \\ x^\top & t \end{bmatrix} \underset{\mathbb{S}_+^{n+1}}{\succeq} 0 \\ & x \in \mathcal{C} \end{aligned} \quad (465)$$

$$\begin{aligned} \underset{x \in \mathbb{R}^n}{\text{minimize}} \quad & \|x\|_\infty \\ \text{subject to} \quad & x \in \mathcal{C} \end{aligned} \quad \equiv \quad \begin{aligned} \underset{x \in \mathbb{R}^n, t \in \mathbb{R}}{\text{minimize}} \quad & t \\ \text{subject to} \quad & -t\mathbf{1} \preceq x \preceq t\mathbf{1} \\ & x \in \mathcal{C} \end{aligned} \quad (466)$$

In \mathbb{R}^n these norms represent: $\|x\|_1$ is length measured along a grid (*taxicab distance*), $\|x\|_2$ is Euclidean length, $\|x\|_\infty$ is maximum |coordinate|.

$$\begin{aligned} \underset{x \in \mathbb{R}^n}{\text{minimize}} \quad & \|x\|_1 \\ \text{subject to} \quad & x \in \mathcal{C} \end{aligned} \quad \equiv \quad \begin{aligned} \underset{\alpha \in \mathbb{R}^n, \beta \in \mathbb{R}^n}{\text{minimize}} \quad & \mathbf{1}^\top(\alpha + \beta) \\ \text{subject to} \quad & \alpha, \beta \succeq 0 \\ & x = \alpha - \beta \\ & x \in \mathcal{C} \end{aligned} \quad (467)$$

These foregoing problems (458)-(467) are convex whenever set \mathcal{C} is.




Figure 64: 1-norm ball \mathcal{B}_1 is convex hull of all cardinality 1 vectors of unit norm (its vertices). Ball boundary contains all points equidistant from origin in 1-norm. Cartesian axes drawn for reference. Plane \mathcal{A} is overhead (drawn truncated). If 1-norm ball is expanded until it *kisses* \mathcal{A} (intersects ball only at boundary), then distance (in 1-norm) from origin to \mathcal{A} is achieved. Euclidean ball would be spherical in this dimension. Only were \mathcal{A} parallel to two axes could there be a minimum cardinality least Euclidean norm solution. Yet 1-norm ball offers infinitely many (but not all) \mathcal{A} -orientations resulting in a minimum cardinality solution. (∞ -norm ball is a cube in this dimension.)

3.1.3.0.1 Example. *Projecting the origin on an affine subset, in 1-norm.* In (1771) we interpret *least norm* solution to linear system $Ax = b$ as orthogonal projection of the origin $\mathbf{0}$ on affine subset $\mathcal{A} = \{x \in \mathbb{R}^n \mid Ax = b\}$ where $A \in \mathbb{R}^{m \times n}$ is fat full-rank. Suppose, instead of the Euclidean metric, we use taxicab distance to do projection. Then the least 1-norm problem is stated, for $b \in \mathcal{R}(A)$

$$\begin{aligned} & \underset{x}{\text{minimize}} \quad \|x\|_1 \\ & \text{subject to} \quad Ax = b \end{aligned} \tag{468}$$

Optimal solution can be interpreted as an *oblique projection* on \mathcal{A} simply because the Euclidean metric is not employed. This problem statement sometimes returns optimal x^* having minimum cardinality; which can be explained intuitively with reference to Figure 64: [18]

Projection of the origin, in 1-norm, on affine subset \mathcal{A} is equivalent to maximization (in this case) of the 1-norm ball until it kisses \mathcal{A} ; rather, a kissing point in \mathcal{A} achieves the distance in 1-norm from the origin to \mathcal{A} . For the example illustrated ($m = 1$, $n = 3$), it appears that a vertex of the ball will be first to touch \mathcal{A} . 1-norm ball vertices in \mathbb{R}^3 represent nontrivial points of minimum cardinality 1, whereas edges represent cardinality 2, while relative interiors of facets represent maximum cardinality 3. By reorienting affine subset \mathcal{A} so it were parallel to an edge or facet, it becomes evident as we expand or contract the ball that a kissing point is not necessarily unique.^{3.7}

The 1-norm ball in \mathbb{R}^n has 2^n facets and $2n$ vertices.^{3.8} For $n > 0$

$$\mathcal{B}_1 = \{x \in \mathbb{R}^n \mid \|x\|_1 \leq 1\} = \text{conv}\{\pm e_i \in \mathbb{R}^n, i = 1 \dots n\} \tag{469}$$

is a vertex-description of the unit 1-norm ball. Maximization of the 1-norm ball until it kisses \mathcal{A} is equivalent to minimization of the 1-norm ball until it no longer intersects \mathcal{A} . Then projection of the origin on affine subset \mathcal{A} is

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} \quad \|x\|_1 \\ & \text{subject to} \quad Ax = b \end{aligned} \equiv \begin{aligned} & \underset{c \in \mathbb{R}, x \in \mathbb{R}^n}{\text{minimize}} \quad c \\ & \text{subject to} \quad x \in c\mathcal{B}_1 \\ & \quad \quad \quad Ax = b \end{aligned} \tag{470}$$

where

$$c\mathcal{B}_1 = \{[I \quad -I]a \mid a^T \mathbf{1} = c, a \succeq 0\} \tag{471}$$

^{3.7}This is unlike the case for the Euclidean ball (1771) where minimum-distance projection on a convex set is unique (§E.9); all kissable faces of the Euclidean ball are single points (vertices).

^{3.8}The ∞ -norm ball in \mathbb{R}^n has $2n$ facets and 2^n vertices.




Figure 65: Exact recovery transition: Respectively signed [100] [102] or positive [107] [105] [106] solutions x to $Ax=b$ with sparsity k below thick curve are recoverable. For Gaussian random matrix $A \in \mathbb{R}^{m \times n}$, thick curve demarcates phase transition in ability to find sparsest solution x by linear programming. These results were empirically reproduced in [34].

Then (470) is equivalent to

$$\begin{aligned}
 & \underset{c \in \mathbb{R}, x \in \mathbb{R}^n, a \in \mathbb{R}^{2n}}{\text{minimize}} && c \\
 & \text{subject to} && x = [I \ -I]a \\
 & && a^T \mathbf{1} = c \\
 & && a \succeq 0 \\
 & && Ax = b
 \end{aligned}
 \quad \equiv \quad
 \begin{aligned}
 & \underset{a \in \mathbb{R}^{2n}}{\text{minimize}} && \|a\|_1 \\
 & \text{subject to} && [A \ -A]a = b \\
 & && a \succeq 0
 \end{aligned}
 \quad (472)$$

where $x^* = [I \ -I]a^*$. (confer(467)) Significance of this result:

- Any vector 1-norm minimization problem may have its variable replaced with a nonnegative variable of twice the length.

All other things being equal, nonnegative variables are easier to solve for sparse solutions. (Figure 65, Figure 83) The compressed sensing problem becomes easier to interpret; *e.g.*,

$$\begin{aligned}
 & \underset{x}{\text{minimize}} && \|x\|_1 \\
 & \text{subject to} && Ax = b \\
 & && x \succeq 0
 \end{aligned}
 \quad \equiv \quad
 \begin{aligned}
 & \underset{x}{\text{minimize}} && \mathbf{1}^T x \\
 & \text{subject to} && Ax = b \\
 & && x \succeq 0
 \end{aligned}
 \quad (473)$$

movement of a hyperplane (Figure 24, Figure 28) over a bounded polyhedron always has a *vertex solution* [82, p.22]. \square

3.1.3.0.2 Exercise. *Combinatorial optimization.*

A device commonly employed to relax combinatorial problems is to arrange desirable solutions at vertices of bounded polyhedra; *e.g.*, the permutation matrices of dimension n , which are factorial in number, are the extreme points of a polyhedron in the nonnegative orthant described by an intersection of $2n$ hyperplanes (§2.3.2.0.4). Minimizing a linear objective function over a bounded polyhedron is a convex problem (a linear program) that always has an optimal solution residing at a vertex. [82]

What about minimizing other functions? Given some nonsingular matrix A , geometrically describe three circumstances under which there are likely to exist vertex solutions to

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && \|Ax\|_1 \\ & \text{subject to} && x \in \mathcal{P} \end{aligned} \tag{474}$$

optimized over some bounded polyhedron \mathcal{P} . ^{3.9} ▼

3.1.3.1 k smallest/largest entries

Sum of the k smallest entries of $x \in \mathbb{R}^n$ is the optimal objective value from:
for $1 \leq k \leq n$

$$\begin{aligned} \sum_{i=n-k+1}^n \pi(x)_i = & \underset{y \in \mathbb{R}^n}{\text{minimize}} && x^T y \\ & \text{subject to} && 0 \preceq y \preceq \mathbf{1} \\ & && \mathbf{1}^T y = k \end{aligned} \quad \text{or} \quad \begin{aligned} \sum_{i=n-k+1}^n \pi(x)_i = & \underset{z \in \mathbb{R}^n, t \in \mathbb{R}}{\text{maximize}} && kt + \mathbf{1}^T z \\ & \text{subject to} && x \succeq t \mathbf{1} + z \\ & && z \preceq 0 \end{aligned} \tag{475}$$

which are dual linear programs, where $\pi(x)_1 = \max\{x_i, i=1 \dots n\}$ where π is a nonlinear permutation-operator sorting its vector argument into nonincreasing order. Finding k smallest entries of an n -length vector x is expressible as an infimum of $n!/(k!(n-k)!)$ linear functions of x . The sum $\sum \pi(x)_i$ is therefore a concave function of x ; in fact, monotonic (§3.1.8.1.1) in this instance.

^{3.9}Hint: Suppose, for example, \mathcal{P} belongs to an orthant and A were orthogonal. Begin with $A=I$ and apply level sets of the objective, as in Figure 62 and Figure 64. Or rewrite the problem as a linear program like (464) and (466) but in a composite variable $\begin{bmatrix} x \\ t \end{bmatrix} \leftarrow y$.

Sum of the k largest entries of $x \in \mathbb{R}^n$ is the optimal objective value from: [53, exer.5.19]

$$\sum_{i=1}^k \pi(x)_i = \begin{array}{ll} \text{maximize} & x^T y \\ \text{subject to} & 0 \preceq y \preceq \mathbf{1} \\ & \mathbf{1}^T y = k \end{array} \quad \text{or} \quad \sum_{i=1}^k \pi(x)_i = \begin{array}{ll} \text{minimize} & k t + \mathbf{1}^T z \\ \text{subject to} & x \preceq t \mathbf{1} + z \\ & z \succeq 0 \end{array} \quad (476)$$

which are dual linear programs. Finding k largest entries of an n -length vector x is expressible as a supremum of $n!/(k!(n-k)!)$ linear functions of x . The summation is therefore a convex function (and monotonic in this instance, §3.1.8.1.1).

Let Πx be a permutation of entries x_i such that their absolute value becomes arranged in nonincreasing order: $|\Pi x|_1 \geq |\Pi x|_2 \geq \dots \geq |\Pi x|_n$. Sum of the k largest entries of $|x| \in \mathbb{R}^n$ is a norm, by properties of vector norm (§3.1.3), and is the optimal objective value of a linear program:

$$\|x\|_k^n \triangleq \sum_{i=1}^k |\Pi x|_i = \begin{array}{ll} \text{minimize} & k t + \mathbf{1}^T z \\ \text{subject to} & -t \mathbf{1} - z \preceq x \preceq t \mathbf{1} + z \\ & z \succeq 0 \end{array} \quad (477)$$

where the norm subscript derives from a binomial coefficient $\binom{n}{k}$, and

$$\begin{aligned} \|x\|_n^n &\triangleq \|x\|_1 \\ \|x\|_1^n &\triangleq \|x\|_\infty \end{aligned} \quad (478)$$

Finding k largest absolute entries of an n -length vector x is expressible as a supremum of $2^k n!/(k!(n-k)!)$ linear functions of x ; hence, this norm is convex (nonmonotonic) in x . [53, exer.6.3(e)]

$$\begin{array}{ll} \text{minimize} & \|x\|_k^n \\ \text{subject to} & x \in \mathcal{C} \end{array} \quad \equiv \quad \begin{array}{ll} \text{minimize} & k t + \mathbf{1}^T z \\ \text{subject to} & -t \mathbf{1} - z \preceq x \preceq t \mathbf{1} + z \\ & z \succeq 0 \\ & x \in \mathcal{C} \end{array} \quad (479)$$

3.1.3.2 clipping

Clipping negative vector entries is accomplished:

$$\begin{aligned} \|x_+\|_1 &= \underset{t \in \mathbb{R}^n}{\text{minimize}} \quad \mathbf{1}^T t \\ &\text{subject to} \quad x \preceq t \\ &\quad \quad \quad 0 \preceq t \end{aligned} \quad (480)$$

where, for $x = [x_i] \in \mathbb{R}^n$

$$x_+ \triangleq t^* = \left[\begin{array}{cc} x_i, & x_i \geq 0 \\ 0, & x_i < 0 \end{array} \right], \quad i=1 \dots n \quad = \frac{1}{2}(x + |x|) \quad (481)$$

(clipping)

$$\begin{aligned} \underset{x \in \mathbb{R}^n}{\text{minimize}} \quad \|x_+\|_1 &\equiv \underset{x \in \mathbb{R}^n, t \in \mathbb{R}^n}{\text{minimize}} \quad \mathbf{1}^T t \\ \text{subject to} \quad x \in \mathcal{C} &\quad \text{subject to} \quad x \preceq t \\ &\quad \quad \quad 0 \preceq t \\ &\quad \quad \quad x \in \mathcal{C} \end{aligned} \quad (482)$$

3.1.4 inverted

We wish to implement objectives of the form x^{-1} . Suppose we have a 2×2 matrix

$$T \triangleq \begin{bmatrix} x & z \\ z & y \end{bmatrix} \in \mathbb{R}^2 \quad (483)$$

which is positive semidefinite by (1423) when

$$T \succeq 0 \quad \Leftrightarrow \quad x > 0 \quad \mathbf{and} \quad xy \geq z^2 \quad (484)$$

A polynomial constraint such as this is therefore called a *conic constraint*.^{3.10}

This means we may formulate convex problems, having inverted variables, as semidefinite programs in Schur-form; *e.g.*,

$$\begin{aligned} \underset{x \in \mathbb{R}}{\text{minimize}} \quad x^{-1} &\equiv \underset{x, y \in \mathbb{R}}{\text{minimize}} \quad y \\ \text{subject to} \quad x > 0 &\quad \text{subject to} \quad \begin{bmatrix} x & 1 \\ 1 & y \end{bmatrix} \succeq 0 \\ &\quad \quad \quad x \in \mathcal{C} \end{aligned} \quad (485)$$

^{3.10}In this dimension, the convex cone formed from the set of all values $\{x, y, z\}$ satisfying constraint (484) is called a *rotated quadratic* or circular cone or positive semidefinite cone.

rather

$$x > 0, \quad y \geq \frac{1}{x} \quad \Leftrightarrow \quad \begin{bmatrix} x & 1 \\ 1 & y \end{bmatrix} \succeq 0 \quad (486)$$

(inverted) For vector $x = [x_i, i = 1 \dots n] \in \mathbb{R}^n$

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n x_i^{-1} \\ \text{subject to} & x \succ 0 \\ & x \in \mathcal{C} \end{array} \quad \equiv \quad \begin{array}{ll} \text{minimize} & y \\ \text{subject to} & \begin{bmatrix} x_i & \sqrt{n} \\ \sqrt{n} & y \end{bmatrix} \succeq 0, \quad i = 1 \dots n \\ & x \in \mathcal{C} \end{array} \quad (487)$$

rather

$$x \succ 0, \quad y \geq \text{tr}(\delta(x)^{-1}) \quad \Leftrightarrow \quad \begin{bmatrix} x_i & \sqrt{n} \\ \sqrt{n} & y \end{bmatrix} \succeq 0, \quad i = 1 \dots n \quad (488)$$

3.1.5 fractional power

[129] To implement an objective of the form x^α for positive α , we quantize α and work instead with that approximation. Choose nonnegative integer q for adequate quantization of α like so:

$$\alpha \triangleq \frac{k}{2^q} \quad (489)$$

where $k \in \{0, 1, 2, \dots, 2^q - 1\}$. Any k from that set may be written $k = \sum_{i=1}^q b_i 2^{i-1}$ where $b_i \in \{0, 1\}$. Define vector $y = [y_i, i = 0 \dots q]$ with $y_0 = 1$.

3.1.5.1 positive

Then we have the equivalent semidefinite program for maximizing a concave function x^α , for quantized $0 \leq \alpha < 1$

$$\begin{array}{ll} \text{maximize} & x^\alpha \\ \text{subject to} & x > 0 \\ & x \in \mathcal{C} \end{array} \quad \equiv \quad \begin{array}{ll} \text{maximize} & y_q \\ \text{subject to} & \begin{bmatrix} y_{i-1} & y_i \\ y_i & x^{b_i} \end{bmatrix} \succeq 0, \quad i = 1 \dots q \\ & x \in \mathcal{C} \end{array} \quad (490)$$

where nonnegativity of y_q is enforced by maximization; *id est*,

$$x > 0, \quad y_q \leq x^\alpha \quad \Leftrightarrow \quad \begin{bmatrix} y_{i-1} & y_i \\ y_i & x^{b_i} \end{bmatrix} \succeq 0, \quad i=1 \dots q \quad (491)$$

3.1.5.2 negative

Is it also desirable implement an objective of the form $x^{-\alpha}$ for positive α . The technique is nearly the same as before: for quantized $0 \leq \alpha < 1$

$$\begin{array}{ll} \text{minimize}_{x \in \mathbb{R}} & x^{-\alpha} \\ \text{subject to} & x > 0 \\ & x \in \mathcal{C} \end{array} \quad \equiv \quad \begin{array}{ll} \text{minimize}_{x, z \in \mathbb{R}, y \in \mathbb{R}^{q+1}} & z \\ \text{subject to} & \begin{bmatrix} y_{i-1} & y_i \\ y_i & x^{b_i} \end{bmatrix} \succeq 0, \quad i=1 \dots q \\ & \begin{bmatrix} z & 1 \\ 1 & y_q \end{bmatrix} \succeq 0 \\ & x \in \mathcal{C} \end{array} \quad (492)$$

rather

$$x > 0, \quad z \geq x^{-\alpha} \quad \Leftrightarrow \quad \begin{array}{l} \begin{bmatrix} y_{i-1} & y_i \\ y_i & x^{b_i} \end{bmatrix} \succeq 0, \quad i=1 \dots q \\ \begin{bmatrix} z & 1 \\ 1 & y_q \end{bmatrix} \succeq 0 \end{array} \quad (493)$$

3.1.5.3 positive inverted

Now define vector $t = [t_i, i=0 \dots q]$ with $t_0 = 1$. To implement an objective $x^{1/\alpha}$ for quantized $0 \leq \alpha < 1$ as in (489)

$$\begin{array}{ll} \text{minimize}_{x \in \mathbb{R}} & x^{1/\alpha} \\ \text{subject to} & x \geq 0 \\ & x \in \mathcal{C} \end{array} \quad \equiv \quad \begin{array}{ll} \text{minimize}_{y \in \mathbb{R}, t \in \mathbb{R}^{q+1}} & y \\ \text{subject to} & \begin{bmatrix} t_{i-1} & t_i \\ t_i & y^{b_i} \end{bmatrix} \succeq 0, \quad i=1 \dots q \\ & x = t_q \geq 0 \\ & x \in \mathcal{C} \end{array} \quad (494)$$

rather

$$x \geq 0, \quad y \geq x^{1/\alpha} \quad \Leftrightarrow \quad \begin{bmatrix} t_{i-1} & t_i \\ t_i & y^{b_i} \end{bmatrix} \succeq 0, \quad i=1 \dots q \quad (495)$$

$$x = t_q \geq 0$$

3.1.6 affine function

A function $f(X)$ is *affine* when it is continuous and has the dimensionally extensible form (confer §2.9.1.0.2)

$$f(X) = AX + B \quad (496)$$

All affine functions are simultaneously convex and concave. This means: both $-AX + B$ and $AX + B$, for example, are convex functions of X . When $B = \mathbf{0}$, $f(X)$ is a *linear function*.

Affine multidimensional functions are recognized by existence of no *multivariate* terms (multiplicative in argument entries) and no polynomial terms of degree higher than 1; *id est*, entries of the function are characterized only by linear combinations of argument entries plus constants.

The real affine function in Figure 66 illustrates hyperplanes, in its domain, constituting contours of equal function-value (*level sets* (500)).

3.1.6.0.1 Example. Engineering control. [347, §2.2]^{3.11}

For $X \in \mathbb{S}^M$ and matrices A, B, Q, R of any compatible dimensions, for example, the expression XAX is not affine in X whereas

$$g(X) = \begin{bmatrix} R & B^T X \\ X B & Q + A^T X + X A \end{bmatrix} \quad (497)$$

is an affine multidimensional function. Such a function is typical in engineering control. [51] [131] □

Any single- or many-valued inverse of an affine function is affine.

^{3.11}The interpretation from this citation of $\{X \in \mathbb{S}^M \mid g(X) \succeq 0\}$ as “an intersection between a linear subspace and the cone of positive semidefinite matrices” is incorrect. (See §2.9.1.0.2 for a similar example.) The conditions they state under which strong duality holds for semidefinite programming are conservative. (confer §4.2.3.0.1)




Figure 66: Cartesian axes in \mathbb{R}^3 and three hyperplanes intersecting convex set $C \subset \mathbb{R}^2$ reproduced from Figure 26. Plotted with third dimension is affine set $\mathcal{A} = f(\mathbb{R}^2)$ a plane. Sequence of hyperplanes, w.r.t domain \mathbb{R}^2 of an affine function $f(z) = a^T z + b : \mathbb{R}^2 \rightarrow \mathbb{R}$, is increasing in direction of gradient a (§3.1.8.0.3) because affine function increases in normal direction (Figure 24).

3.1.6.0.2 Example. *Linear objective.*

Consider minimization of a real affine function $f(z) = a^T z + b$ over the convex feasible set \mathcal{C} in its domain \mathbb{R}^2 illustrated in Figure 66. Since vector b is fixed, the problem posed is the same as the convex optimization

$$\begin{aligned} & \underset{z}{\text{minimize}} && a^T z \\ & \text{subject to} && z \in \mathcal{C} \end{aligned} \tag{498}$$

whose objective of minimization is a real linear function. Were convex set \mathcal{C} polyhedral (§2.12), then this problem would be called a linear program. Were convex set \mathcal{C} an intersection with a positive semidefinite cone, then this problem would be called a semidefinite program.

There are two distinct ways to visualize this problem: one in the objective function's domain \mathbb{R}^2 , the other including the ambient space of the objective function's range as in $\begin{bmatrix} \mathbb{R}^2 \\ \mathbb{R} \end{bmatrix}$. Both visualizations are illustrated in Figure 66. Visualization in the function domain is easier because of lower dimension and because

- level sets (500) of any affine function are affine. (§2.1.9)

In this circumstance, the level sets are parallel hyperplanes with respect to \mathbb{R}^2 . One solves optimization problem (498) graphically by finding that hyperplane intersecting feasible set \mathcal{C} furthest right (in the direction of negative gradient $-a$ (§3.1.8)). \square

When a differentiable convex objective function f is nonlinear, the negative gradient $-\nabla f$ is a viable search direction (replacing $-a$ in (498)). (§2.13.10.1, Figure 62) [133] Then the nonlinear objective function can be replaced with a dynamic linear objective; linear as in (498).

3.1.6.0.3 Example. *Support function.* [177, §C.2.1]

For arbitrary set $\mathcal{Y} \subseteq \mathbb{R}^n$, its *support function* $\sigma_{\mathcal{Y}}(a) : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined

$$\sigma_{\mathcal{Y}}(a) \triangleq \sup_{z \in \mathcal{Y}} a^T z \tag{499}$$

whose range contains $\pm\infty$ [220, p.135] [177, §C.2.3.1]. For each $z \in \mathcal{Y}$, $a^T z$ is a linear function of vector a . Because $\sigma_{\mathcal{Y}}(a)$ is the pointwise supremum of linear functions, it is convex in a . (Figure 67) Application of the support function is illustrated in Figure 27(a) for one particular normal a . \square




Figure 67: Pointwise supremum of convex functions remains a convex function. Illustrated is a supremum of affine functions in variable a evaluated for a particular argument a_p . Topmost affine function is supremum for each value of a .




Figure 68: Quasiconvex function q epigraph is not necessarily convex, but convex function f epigraph is convex in any dimension. Sublevel sets are necessarily convex for either.

3.1.6.0.4 Exercise. *Level sets.*

Given a function f and constant κ , its level sets are defined

$$\mathcal{L}_\kappa^\kappa f \triangleq \{z \mid f(z) = \kappa\} \tag{500}$$

Give two distinct examples of convex function, that are not affine, having convex level sets. ▼

3.1.7 epigraph, sublevel set

It is well established that a continuous real function is convex if and only if its *epigraph* makes a convex set. [177] [271] [318] [331] [220] Thereby, piecewise-continuous convex functions are admitted. Epigraph is the connection between convex sets and convex functions. Its generalization to a vector-valued function $f(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}^M$ is straightforward: [260]

$$\text{epi } f \triangleq \{(X, t) \in \mathbb{R}^{p \times k} \times \mathbb{R}^M \mid X \in \text{dom } f, f(X) \preceq_{\mathbb{R}_+^M} t\} \tag{501}$$

id est,

$$f \text{ convex} \Leftrightarrow \text{epi } f \text{ convex} \tag{502}$$

Necessity is proven: [53, exer.3.60] Given any $(X, u), (Y, v) \in \text{epi } f$, we must show for all $\mu \in [0, 1]$ that $\mu(X, u) + (1-\mu)(Y, v) \in \text{epi } f$; *id est*, we must show

$$f(\mu X + (1-\mu)Y) \underset{\mathbb{R}_+^M}{\preceq} \mu u + (1-\mu)v \quad (503)$$

Yet this holds by definition because $f(\mu X + (1-\mu)Y) \underset{\mathbb{R}_+^M}{\preceq} \mu f(X) + (1-\mu)f(Y)$. The converse also holds. \blacklozenge

3.1.7.0.1 Exercise. Epigraph sufficiency.

Prove that converse: Given any $(X, u), (Y, v) \in \text{epi } f$, if for all $\mu \in [0, 1]$ $\mu(X, u) + (1-\mu)(Y, v) \in \text{epi } f$ holds, then f must be convex. \blacktriangledown

Sublevel sets of a real convex function are convex. Likewise, corresponding to each and every $\nu \in \mathbb{R}^M$

$$\mathcal{L}_\nu f \triangleq \{X \in \text{dom } f \mid f(X) \underset{\mathbb{R}_+^M}{\preceq} \nu\} \subseteq \mathbb{R}^{p \times k} \quad (504)$$

sublevel sets of a vector-valued convex function are convex. As for real functions, the converse does not hold. (Figure 68)

To prove necessity of convex sublevel sets: For any $X, Y \in \mathcal{L}_\nu f$ we must show for each and every $\mu \in [0, 1]$ that $\mu X + (1-\mu)Y \in \mathcal{L}_\nu f$. By definition,

$$f(\mu X + (1-\mu)Y) \underset{\mathbb{R}_+^M}{\preceq} \mu f(X) + (1-\mu)f(Y) \underset{\mathbb{R}_+^M}{\preceq} \nu \quad (505)$$

\blacklozenge

When an epigraph (501) is artificially bounded above, $t \preceq \nu$, then the corresponding sublevel set can be regarded as an orthogonal projection of the epigraph on the function domain.

Sense of the inequality is reversed in (501), for concave functions, and we use instead the nomenclature *hypograph*. Sense of the inequality in (504) is reversed, similarly, with each convex set then called *superlevel set*.

3.1.7.0.2 Example. *Matrix pseudofractional function.*

Consider a real function of two variables

$$f(A, x) : \mathbb{S}^n \times \mathbb{R}^n \rightarrow \mathbb{R} = x^T A^\dagger x \quad (506)$$

on $\text{dom } f = \mathbb{S}_+^n \times \mathcal{R}(A)$. This function is convex simultaneously in both variables when variable matrix A belongs to the entire positive semidefinite cone \mathbb{S}_+^n and variable vector x is confined to range $\mathcal{R}(A)$ of matrix A .

To explain this, we need only demonstrate that the function epigraph is convex. Consider Schur-form (1420) from §A.4: for $t \in \mathbb{R}$

$$\begin{aligned} G(A, z, t) &= \begin{bmatrix} A & z \\ z^T & t \end{bmatrix} \succeq 0 \\ &\Leftrightarrow \\ & z^T(I - AA^\dagger) = \mathbf{0} \\ & t - z^T A^\dagger z \geq 0 \\ & A \succeq 0 \end{aligned} \quad (507)$$

Inverse image of the positive semidefinite cone \mathbb{S}_+^{n+1} under affine mapping $G(A, z, t)$ is convex by Theorem 2.1.9.0.1. Of the equivalent conditions for positive semidefiniteness of G , the first is an equality demanding vector z belong to $\mathcal{R}(A)$. Function $f(A, z) = z^T A^\dagger z$ is convex on convex domain $\mathbb{S}_+^n \times \mathcal{R}(A)$ because the Cartesian product constituting its epigraph

$$\text{epi } f(A, z) = \{(A, z, t) \mid A \succeq 0, z \in \mathcal{R}(A), z^T A^\dagger z \leq t\} = G^{-1}(\mathbb{S}_+^{n+1}) \quad (508)$$

is convex. \square

3.1.7.0.3 Exercise. *Matrix product function.*

Continue Example 3.1.7.0.2 by introducing vector variable x and making the substitution $z \leftarrow Ax$. Because of matrix symmetry (§E), for all $x \in \mathbb{R}^n$

$$f(A, z(x)) = z^T A^\dagger z = x^T A^T A^\dagger A x = x^T A x = f(A, x) \quad (509)$$

whose epigraph is

$$\text{epi } f(A, x) = \{(A, x, t) \mid A \succeq 0, x^T A x \leq t\} \quad (510)$$

Provide two simple explanations why $f(A, x) = x^T A x$ is not a function convex simultaneously in positive semidefinite matrix A and vector x on $\text{dom } f = \mathbb{S}_+^n \times \mathbb{R}^n$. \blacktriangledown

3.1.7.0.4 Example. *Matrix fractional function.* (confer §3.2.2.0.1)

Continuing Example 3.1.7.0.2, now consider a real function of two variables on $\text{dom } f = \mathbb{S}_+^n \times \mathbb{R}^n$ for small positive constant ϵ (confer (1757))

$$f(A, x) = \epsilon x^T(A + \epsilon I)^{-1}x \quad (511)$$

where the inverse always exists by (1364). This function is convex simultaneously in both variables over the entire positive semidefinite cone \mathbb{S}_+^n and all $x \in \mathbb{R}^n$: Consider Schur-form (1423) from §A.4: for $t \in \mathbb{R}$

$$\begin{aligned} G(A, z, t) &= \begin{bmatrix} A + \epsilon I & z \\ z^T & \epsilon^{-1}t \end{bmatrix} \succeq 0 \\ &\Leftrightarrow \\ &t - \epsilon z^T(A + \epsilon I)^{-1}z \geq 0 \\ &A + \epsilon I \succ 0 \end{aligned} \quad (512)$$

Inverse image of the positive semidefinite cone \mathbb{S}_+^{n+1} under affine mapping $G(A, z, t)$ is convex by Theorem 2.1.9.0.1. Function $f(A, z)$ is convex on $\mathbb{S}_+^n \times \mathbb{R}^n$ because its epigraph is that inverse image:

$$\text{epi } f(A, z) = \{(A, z, t) \mid A + \epsilon I \succ 0, \epsilon z^T(A + \epsilon I)^{-1}z \leq t\} = G^{-1}(\mathbb{S}_+^{n+1}) \quad (513)$$

□

3.1.7.1 matrix fractional projector function

Consider nonlinear function f having orthogonal projector W as argument:

$$f(W, x) = \epsilon x^T(W + \epsilon I)^{-1}x \quad (514)$$

Projection matrix W has property $W^\dagger = W^T = W \succeq 0$ (1802). Any orthogonal projector can be decomposed into an outer product of orthonormal matrices $W = UU^T$ where $U^T U = I$ as explained in §E.3.2. From (1757) for any $\epsilon > 0$ and idempotent symmetric W , $\epsilon(W + \epsilon I)^{-1} = I - (1 + \epsilon)^{-1}W$ from which

$$f(W, x) = \epsilon x^T(W + \epsilon I)^{-1}x = x^T(I - (1 + \epsilon)^{-1}W)x \quad (515)$$

Therefore

$$\lim_{\epsilon \rightarrow 0^+} f(W, x) = \lim_{\epsilon \rightarrow 0^+} \epsilon x^T(W + \epsilon I)^{-1}x = x^T(I - W)x \quad (516)$$

where $I - W$ is also an orthogonal projector (§E.2).

We learned from Example 3.1.7.0.4 that $f(W, x) = \epsilon x^T(W + \epsilon I)^{-1}x$ is convex simultaneously in both variables over all $x \in \mathbb{R}^n$ when $W \in \mathbb{S}_+^n$ is confined to the entire positive semidefinite cone (including its boundary). It is now our goal to incorporate f into an optimization problem such that an optimal solution returned always comprises a projection matrix W . The set of orthogonal projection matrices is a nonconvex subset of the positive semidefinite cone. So f cannot be convex on the projection matrices, and its equivalent (for idempotent W)

$$f(W, x) = x^T(I - (1 + \epsilon)^{-1}W)x \quad (517)$$

cannot be convex simultaneously in both variables on either the positive semidefinite or symmetric projection matrices.

Suppose we allow $\text{dom } f$ to constitute the entire positive semidefinite cone but constrain W to a Fantope (83); *e.g.*, for convex set \mathcal{C} and $0 < k < n$ as in

$$\begin{aligned} & \underset{x \in \mathbb{R}^n, W \in \mathbb{S}^n}{\text{minimize}} && \epsilon x^T(W + \epsilon I)^{-1}x \\ & \text{subject to} && 0 \preceq W \preceq I \\ & && \text{tr } W = k \\ & && x \in \mathcal{C} \end{aligned} \quad (518)$$

Although this is a convex problem, there is no guarantee that optimal W is a projection matrix because only extreme points of a Fantope are orthogonal projection matrices UU^T .

Let's try partitioning the problem into two convex parts (one for x and one for W), substitute equivalence (515), and then iterate solution of convex problem

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && x^T(I - (1 + \epsilon)^{-1}W)x \\ & \text{subject to} && x \in \mathcal{C} \end{aligned} \quad (519)$$

with convex problem

$$\begin{aligned} & \underset{W \in \mathbb{S}^n}{\text{minimize}} && x^*{}^T(I - (1 + \epsilon)^{-1}W)x^* \\ & \text{subject to} && 0 \preceq W \preceq I \\ & && \text{tr } W = k \end{aligned} \quad \equiv \quad \begin{aligned} & \underset{W \in \mathbb{S}^n}{\text{maximize}} && x^*{}^T W x^* \\ & \text{subject to} && 0 \preceq W \preceq I \\ & && \text{tr } W = k \end{aligned} \quad (520)$$

until convergence, where x^* represents an optimal solution of (519) from any particular iteration. The idea is to optimally solve for the partitioned

variables which are later combined to solve the original problem (518). What makes this approach sound is that the constraints are separable, the partitioned feasible sets are not interdependent, and the fact that the original problem (though nonlinear) is convex simultaneously in both variables.^{3.12}

But partitioning alone does not guarantee a projector. To make orthogonal projector W a certainty, we must invoke a known analytical optimal solution to problem (520): Diagonalize optimal solution from problem (519) $x^*x^{*\text{T}} \triangleq Q\Lambda Q^{\text{T}}$ (§A.5.2) and set $U^* = Q(:, 1:k) \in \mathbb{R}^{n \times k}$ per (1592c);

$$W = U^*U^{*\text{T}} = \frac{x^*x^{*\text{T}}}{\|x^*\|^2} + Q(:, 2:k)Q(:, 2:k)^{\text{T}} \quad (521)$$

Then optimal solution (x^*, U^*) to problem (518) is found, for small ϵ , by iterating solution to problem (519) with optimal (projector) solution (521) to convex problem (520).

Proof. Optimal vector x^* is orthogonal to the last $n-1$ columns of orthogonal matrix Q , so

$$f_{(519)}^* = \|x^*\|^2(1 - (1 + \epsilon)^{-1}) \quad (522)$$

after each iteration. Convergence of $f_{(519)}^*$ is proven with the observation that iteration (519) (520a) is a nonincreasing sequence that is bounded below by 0. Any bounded monotonic sequence in \mathbb{R} is convergent. [228, §1.2] [37, §1.1] Expression (521) for optimal projector W holds at each iteration, therefore $\|x^*\|^2(1 - (1 + \epsilon)^{-1})$ must also represent the optimal objective value $f_{(519)}^*$ at convergence.

Because the objective $f_{(518)}$ from problem (518) is also bounded below by 0 on the same domain, this convergent optimal objective value $f_{(519)}^*$ (for positive ϵ arbitrarily close to 0) is necessarily optimal for (518); *id est*,

$$f_{(519)}^* \geq f_{(518)}^* \geq 0 \quad (523)$$

by (1575), and

$$\lim_{\epsilon \rightarrow 0^+} f_{(519)}^* = 0 \quad (524)$$

^{3.12}A convex problem has convex feasible set, and the objective *surface* has one and only one global minimum.

Since optimal (x^*, U^*) from problem (519) is feasible to problem (518), and because their objectives are equivalent for projectors by (515), then converged (x^*, U^*) must also be optimal to (518) in the limit. Because problem (518) is convex, this represents a globally optimal solution. \blacklozenge

3.1.7.2 Semidefinite program via Schur

Schur complement (1420) can be used to convert a projection problem to an optimization problem in *epigraph form*. Suppose, for example, we are presented with the constrained projection problem studied by Hayden & Wells in [163] (who provide analytical solution): Given $A \in \mathbb{R}^{M \times M}$ and some full-rank matrix $S \in \mathbb{R}^{M \times L}$ with $L < M$

$$\begin{aligned} & \underset{X \in \mathbb{S}^M}{\text{minimize}} && \|A - X\|_{\text{F}}^2 \\ & \text{subject to} && S^T X S \succeq 0 \end{aligned} \quad (525)$$

Variable X is constrained to be positive semidefinite, but only on a subspace determined by S . First we write the epigraph form:

$$\begin{aligned} & \underset{X \in \mathbb{S}^M, t \in \mathbb{R}}{\text{minimize}} && t \\ & \text{subject to} && \|A - X\|_{\text{F}}^2 \leq t \\ & && S^T X S \succeq 0 \end{aligned} \quad (526)$$

Next we use Schur complement [244, §6.4.3] [219] and matrix vectorization (§2.2):

$$\begin{aligned} & \underset{X \in \mathbb{S}^M, t \in \mathbb{R}}{\text{minimize}} && t \\ & \text{subject to} && \begin{bmatrix} tI & \text{vec}(A - X) \\ \text{vec}(A - X)^T & 1 \end{bmatrix} \succeq 0 \\ & && S^T X S \succeq 0 \end{aligned} \quad (527)$$

This semidefinite program (§4) is an epigraph form in disguise, equivalent to (525); it demonstrates how a quadratic objective or constraint can be converted to a semidefinite constraint.

Were problem (525) instead equivalently expressed without the square

$$\begin{aligned} & \underset{X \in \mathbb{S}^M}{\text{minimize}} && \|A - X\|_{\text{F}} \\ & \text{subject to} && S^T X S \succeq 0 \end{aligned} \quad (528)$$

then we get a subtle variation:

$$\begin{aligned} & \underset{X \in \mathbb{S}^M, t \in \mathbb{R}}{\text{minimize}} && t \\ & \text{subject to} && \|A - X\|_F \leq t \\ & && S^T X S \succeq 0 \end{aligned} \quad (529)$$

that leads to an equivalent for (528) (and for (525) by (463))

$$\begin{aligned} & \underset{X \in \mathbb{S}^M, t \in \mathbb{R}}{\text{minimize}} && t \\ & \text{subject to} && \begin{bmatrix} tI & \text{vec}(A - X) \\ \text{vec}(A - X)^T & t \end{bmatrix} \succeq 0 \\ & && S^T X S \succeq 0 \end{aligned} \quad (530)$$

3.1.7.2.1 Example. Schur anomaly.

Consider a problem abstract in the convex constraint, given symmetric matrix A

$$\begin{aligned} & \underset{X \in \mathbb{S}^M}{\text{minimize}} && \|X\|_F^2 - \|A - X\|_F^2 \\ & \text{subject to} && X \in \mathcal{C} \end{aligned} \quad (531)$$

the minimization of a difference of two quadratic functions each convex in matrix X . Observe equality

$$\|X\|_F^2 - \|A - X\|_F^2 = 2 \text{tr}(XA) - \|A\|_F^2 \quad (532)$$

So problem (531) is equivalent to the convex optimization

$$\begin{aligned} & \underset{X \in \mathbb{S}^M}{\text{minimize}} && \text{tr}(XA) \\ & \text{subject to} && X \in \mathcal{C} \end{aligned} \quad (533)$$

But this problem (531) does not have Schur-form

$$\begin{aligned} & \underset{X \in \mathbb{S}^M, \alpha, t}{\text{minimize}} && t - \alpha \\ & \text{subject to} && X \in \mathcal{C} \\ & && \|X\|_F^2 \leq t \\ & && \|A - X\|_F^2 \geq \alpha \end{aligned} \quad (534)$$

because the constraint in α is nonconvex. (§2.9.1.0.1) \square




Figure 69: Gradient in \mathbb{R}^2 evaluated on grid over some open disc in domain of convex quadratic bowl $f(Y) = Y^T Y : \mathbb{R}^2 \rightarrow \mathbb{R}$ illustrated in Figure 70. Circular contours are level sets; each defined by a constant function-value.

3.1.8 gradient

Gradient ∇f of any differentiable multidimensional function f (formally defined in §D.1) maps each entry f_i to a space having the same dimension as the ambient space of its domain. Notation ∇f is shorthand for gradient $\nabla_x f(x)$ of f with respect to x . $\nabla f(y)$ can mean $\nabla_y f(y)$ or gradient $\nabla_x f(y)$ of $f(x)$ with respect to x evaluated at y ; a distinction that should become clear from context.

Gradient of a differentiable real function $f(x) : \mathbb{R}^K \rightarrow \mathbb{R}$ with respect to its vector argument is defined

$$\nabla f(x) \triangleq \begin{bmatrix} \frac{\partial f(x)}{\partial x_1} \\ \frac{\partial f(x)}{\partial x_2} \\ \vdots \\ \frac{\partial f(x)}{\partial x_K} \end{bmatrix} \in \mathbb{R}^K \quad (1648)$$

while the second-order gradient of the twice differentiable real function with

respect to its vector argument is traditionally called the *Hessian*,^{3.13}

$$\nabla^2 f(x) \triangleq \begin{bmatrix} \frac{\partial^2 f(x)}{\partial^2 x_1} & \frac{\partial^2 f(x)}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f(x)}{\partial x_1 \partial x_K} \\ \frac{\partial^2 f(x)}{\partial x_2 \partial x_1} & \frac{\partial^2 f(x)}{\partial^2 x_2} & \cdots & \frac{\partial^2 f(x)}{\partial x_2 \partial x_K} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f(x)}{\partial x_K \partial x_1} & \frac{\partial^2 f(x)}{\partial x_K \partial x_2} & \cdots & \frac{\partial^2 f(x)}{\partial^2 x_K} \end{bmatrix} \in \mathbb{S}^K \quad (1649)$$

The gradient can be interpreted as a vector pointing in the direction of greatest change. [194, §15.6] The gradient can also be interpreted as that vector normal to a level set; *e.g.*, Figure 71, Figure 62.

For the quadratic bowl in Figure 70, the gradient maps to \mathbb{R}^2 ; illustrated in Figure 69. For a one-dimensional function of real variable, for example, the gradient evaluated at any point in the function domain is just the *slope* (or *derivative*) of that function there. (*confer* §D.1.4.1)

- For any differentiable multidimensional function, zero gradient $\nabla f = \mathbf{0}$ is a necessary condition for its unconstrained minimization [133, §3.2]:

3.1.8.0.1 Example. Projection on a rank-1 subset.

For $A \in \mathbb{S}^N$ having eigenvalues $\lambda(A) = [\lambda_i] \in \mathbb{R}^N$, consider the unconstrained nonconvex optimization that is a projection on the rank-1 subset (§2.9.2.1) of positive semidefinite cone \mathbb{S}_+^N : Defining $\lambda_1 \triangleq \max_i \{\lambda(A)_i\}$ and corresponding eigenvector v_1

$$\begin{aligned} \underset{x}{\text{minimize}} \quad \|xx^T - A\|_F^2 &= \underset{x}{\text{minimize}} \quad \text{tr}(xx^T(x^T x) - 2Axx^T + A^T A) \\ &= \begin{cases} \|\lambda(A)\|^2, & \lambda_1 \leq 0 \\ \|\lambda(A)\|^2 - \lambda_1^2, & \lambda_1 > 0 \end{cases} \end{aligned} \quad (1587)$$

$$\arg \underset{x}{\text{minimize}} \quad \|xx^T - A\|_F^2 = \begin{cases} \mathbf{0}, & \lambda_1 \leq 0 \\ v_1 \sqrt{\lambda_1}, & \lambda_1 > 0 \end{cases} \quad (1588)$$

From (1677) and §D.2.1, the gradient of $\|xx^T - A\|_F^2$ is

$$\nabla_x ((x^T x)^2 - 2x^T A x) = 4(x^T x)x - 4Ax \quad (535)$$

^{3.13} *Jacobian* is the Hessian transpose, so commonly confused in matrix calculus.

Setting the gradient to $\mathbf{0}$

$$Ax = x(x^T x) \quad (536)$$

is necessary for optimal solution. Replace vector x with a normalized eigenvector v_i of $A \in \mathbb{S}^N$, corresponding to a positive eigenvalue λ_i , scaled by square root of that eigenvalue. Then (536) is satisfied

$$x \leftarrow v_i \sqrt{\lambda_i} \Rightarrow Av_i = v_i \lambda_i \quad (537)$$

$xx^T = \lambda_i v_i v_i^T$ is a rank-1 matrix on the positive semidefinite cone boundary, and the minimum is achieved (§7.1.2) when $\lambda_i = \lambda_1$ is the largest positive eigenvalue of A . If A has no positive eigenvalue, then $x = \mathbf{0}$ yields the minimum. \square

Differentiability is a prerequisite neither to convexity or to numerical solution of a convex optimization problem. The gradient provides a necessary and sufficient condition (321) (411) for optimality in the constrained case, nevertheless, as it does in the unconstrained case:

- For any differentiable multidimensional convex function, zero gradient $\nabla f = \mathbf{0}$ is a necessary and sufficient condition for its unconstrained minimization [53, §5.5.3]:

3.1.8.0.2 Example. Pseudoinverse.

The pseudoinverse matrix is the unique solution to an unconstrained convex optimization problem [139, §5.5.4]: given $A \in \mathbb{R}^{m \times n}$

$$\underset{X \in \mathbb{R}^{n \times m}}{\text{minimize}} \|XA - I\|_F^2 \quad (538)$$

where

$$\|XA - I\|_F^2 = \text{tr}(A^T X^T X A - X A - A^T X^T + I) \quad (539)$$

whose gradient (§D.2.3)

$$\nabla_X \|XA - I\|_F^2 = 2(XAA^T - A^T) = \mathbf{0} \quad (540)$$

vanishes when

$$XAA^T = A^T \quad (541)$$

When A is fat full-rank, then AA^T is invertible, $X^* = A^T(AA^T)^{-1}$ is the pseudoinverse A^\dagger , and $AA^\dagger = I$. Otherwise, we can make AA^T invertible by adding a positively scaled identity, for any $A \in \mathbb{R}^{m \times n}$

$$X = A^T(AA^T + tI)^{-1} \quad (542)$$

Invertibility is guaranteed for any finite positive value of t by (1364). Then matrix X becomes the pseudoinverse $X \rightarrow A^\dagger \triangleq X^*$ in the limit $t \rightarrow 0^+$. Minimizing instead $\|AX - I\|_F^2$ yields the second flavor in (1756). \square

3.1.8.0.3 Example. *Hyperplane, line, described by affine function.* Consider the real affine function of vector variable, (confer Figure 66)

$$f(x) : \mathbb{R}^p \rightarrow \mathbb{R} = a^T x + b \quad (543)$$

whose domain is \mathbb{R}^p and whose gradient $\nabla f(x) = a$ is a constant vector (independent of x). This function describes the real line \mathbb{R} (its range), and it describes a *nonvertical* [177, §B.1.2] hyperplane $\partial\mathcal{H}$ in the space $\mathbb{R}^p \times \mathbb{R}$ for any particular vector a (confer §2.4.2);

$$\partial\mathcal{H} = \left\{ \begin{bmatrix} x \\ a^T x + b \end{bmatrix} \mid x \in \mathbb{R}^p \right\} \subset \mathbb{R}^p \times \mathbb{R} \quad (544)$$

having nonzero normal

$$\eta = \begin{bmatrix} a \\ -1 \end{bmatrix} \in \mathbb{R}^p \times \mathbb{R} \quad (545)$$

This equivalence to a hyperplane holds only for real functions.^{3.14} The epigraph of real affine function $f(x)$ is therefore a halfspace in $\begin{bmatrix} \mathbb{R}^p \\ \mathbb{R} \end{bmatrix}$, so we

^{3.14}To prove that, consider a vector-valued affine function

$$f(x) : \mathbb{R}^p \rightarrow \mathbb{R}^M = Ax + b$$

having gradient $\nabla f(x) = A^T \in \mathbb{R}^{p \times M}$: The affine set

$$\left\{ \begin{bmatrix} x \\ Ax + b \end{bmatrix} \mid x \in \mathbb{R}^p \right\} \subset \mathbb{R}^p \times \mathbb{R}^M$$

is perpendicular to

$$\eta \triangleq \begin{bmatrix} \nabla f(x) \\ -I \end{bmatrix} \in \mathbb{R}^{p \times M} \times \mathbb{R}^{M \times M}$$

because

$$\eta^T \left(\begin{bmatrix} x \\ Ax + b \end{bmatrix} - \begin{bmatrix} 0 \\ b \end{bmatrix} \right) = 0 \quad \forall x \in \mathbb{R}^p$$

Yet η is a vector (in $\mathbb{R}^p \times \mathbb{R}^M$) only when $M = 1$. \blacklozenge

have:

The real affine function is to convex functions
as
the hyperplane is to convex sets.

Similarly, the matrix-valued affine function of real variable x , for any particular matrix $A \in \mathbb{R}^{M \times N}$,

$$h(x) : \mathbb{R} \rightarrow \mathbb{R}^{M \times N} = Ax + B \quad (546)$$

describes a line in $\mathbb{R}^{M \times N}$ in direction A

$$\{Ax + B \mid x \in \mathbb{R}\} \subseteq \mathbb{R}^{M \times N} \quad (547)$$

and describes a line in $\mathbb{R} \times \mathbb{R}^{M \times N}$

$$\left\{ \begin{bmatrix} x \\ Ax + B \end{bmatrix} \mid x \in \mathbb{R} \right\} \subset \mathbb{R} \times \mathbb{R}^{M \times N} \quad (548)$$

whose slope with respect to x is A . □

3.1.8.1 monotonic function

A real function of real argument is called *monotonic* when it is exclusively nonincreasing or nondecreasing over the whole of its domain. A real differentiable function of real argument is monotonic when its first derivative (which is not necessarily continuous) maintains sign over the function domain.

3.1.8.1.1 Definition. *Monotonicity.*

Multidimensional function $f(X)$ is

$$\begin{array}{ll} \text{nondecreasing monotonic when} & Y \succeq X \Rightarrow f(Y) \succeq f(X) \\ \text{nonincreasing monotonic when} & Y \succeq X \Rightarrow f(Y) \preceq f(X) \end{array} \quad (549)$$

$\forall X, Y \in \text{dom } f$. △

For vector-valued functions compared with respect to the nonnegative orthant, it is necessary and sufficient for each entry f_i to be monotonic in the same sense.

Any affine function is monotonic. $\text{tr}(Y^T X)$ is a nondecreasing monotonic function of matrix $X \in \mathbb{S}^M$ when constant matrix Y is positive semidefinite, for example; which follows from a result (342) of Fejér.

A convex function can be characterized by another kind of nondecreasing monotonicity of its gradient:

3.1.8.1.2 Theorem. *Gradient monotonicity.* [177, §B.4.1.4] [48, §3.1, exer.20] Given $f(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}$ a real differentiable function with matrix argument on open convex domain, the condition

$$\langle \nabla f(Y) - \nabla f(X), Y - X \rangle \geq 0 \text{ for each and every } X, Y \in \text{dom } f \quad (550)$$

is necessary and sufficient for convexity of f . Strict inequality and *caveat* distinct Y, X provide necessary and sufficient conditions for strict convexity.

◇

3.1.8.1.3 Example. *Composition of functions.* [53, §3.2.4] [177, §B.2.1] Monotonic functions play a vital role determining convexity of functions constructed by transformation. Given functions $g : \mathbb{R}^k \rightarrow \mathbb{R}$ and $h : \mathbb{R}^n \rightarrow \mathbb{R}^k$, their composition $f = g(h) : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$f(x) = g(h(x)), \quad \text{dom } f = \{x \in \text{dom } h \mid h(x) \in \text{dom } g\} \quad (551)$$

is convex when

- g is convex nondecreasing monotonic **and** h is convex
- g is convex nonincreasing monotonic **and** h is concave

and composite function f is concave when

- g is concave nondecreasing monotonic **and** h is concave
- g is concave nonincreasing monotonic **and** h is convex

where ∞ ($-\infty$) is assigned to convex (concave) g when evaluated outside its domain. When functions are differentiable, these rules are consequent to (1678). Convexity (concavity) of any g is preserved when h is affine. □

3.1.8.1.4 Exercise. *Product and ratio of convex functions.* [53, exer.3.32] In general the product or ratio of two convex functions is not convex. [202] However, there are some results that apply to functions on \mathbb{R} [real domain]. Prove the following.^{3.15}

- (a) If f and g are convex, both nondecreasing (or nonincreasing), and positive functions on an interval, then fg is convex.
- (b) If f, g are concave, positive, with one nondecreasing and the other nonincreasing, then fg is concave.
- (c) If f is convex, nondecreasing, and positive, and g is concave, nonincreasing, and positive, then f/g is convex. ▼

3.1.9 first-order convexity condition, real function

Discretization of $w \succeq 0$ in (452) invites refocus to the real-valued function:

3.1.9.0.1 Theorem. *Necessary and sufficient convexity condition.*

[53, §3.1.3] [115, §I.5.2] [350, §1.2.3] [37, §1.2] [293, §4.2] [270, §3] For real differentiable function $f(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}$ with matrix argument on open convex domain, the condition (confer §D.1.7)

$$f(Y) \geq f(X) + \langle \nabla f(X), Y - X \rangle \quad \text{for each and every } X, Y \in \text{dom } f \quad (552)$$

is necessary and sufficient for convexity of f . ◊

When $f(X) : \mathbb{R}^p \rightarrow \mathbb{R}$ is a real differentiable convex function with vector argument on open convex domain, there is simplification of the first-order condition (552); for each and every $X, Y \in \text{dom } f$

$$f(Y) \geq f(X) + \nabla f(X)^T (Y - X) \quad (553)$$

From this we can find $\underline{\partial \mathcal{H}}$ a unique [331, §5.5.4] nonvertical [177, §B.1.2] hyperplane (§2.4), expressed in terms of function gradient, supporting epi f at $\begin{bmatrix} X \\ f(X) \end{bmatrix}$: *videlicet*, defining $f(Y \notin \text{dom } f) \triangleq \infty$ [53, §3.1.7]

$$\begin{bmatrix} Y \\ t \end{bmatrix} \in \text{epi } f \Leftrightarrow t \geq f(Y) \Rightarrow [\nabla f(X)^T \quad -1] \left(\begin{bmatrix} Y \\ t \end{bmatrix} - \begin{bmatrix} X \\ f(X) \end{bmatrix} \right) \leq 0 \quad (554)$$

^{3.15}Hint: Prove §3.1.8.1.4(a) by verifying Jensen's inequality ((451) at $\mu = \frac{1}{2}$).




Figure 70: When a real function f is differentiable at each point in its open domain, there is an intuitive geometric interpretation of function convexity in terms of its gradient ∇f and its epigraph: Drawn is a convex quadratic bowl in $\mathbb{R}^2 \times \mathbb{R}$ (confer Figure 139, p.624); $f(Y) = Y^T Y : \mathbb{R}^2 \rightarrow \mathbb{R}$ versus Y on some open disc in \mathbb{R}^2 . Unique strictly supporting hyperplane $\partial\mathcal{H}_- \in \mathbb{R}^2 \times \mathbb{R}$ (only partially drawn) and its normal vector $[\nabla f(X)^T \ -1]^T$ at the particular point of support $[X^T \ f(X)]^T$ are illustrated. The interpretation: At each and every coordinate Y , there is such a hyperplane containing $[Y^T \ f(Y)]^T$ and supporting the epigraph.

This means, for each and every point X in the domain of a real convex function $f(X)$, there exists a hyperplane $\underline{\partial\mathcal{H}}_-$ in $\mathbb{R}^p \times \mathbb{R}$ having normal $\begin{bmatrix} \nabla f(X) \\ -1 \end{bmatrix}$ supporting the function epigraph at $\begin{bmatrix} X \\ f(X) \end{bmatrix} \in \underline{\partial\mathcal{H}}_-$

$$\underline{\partial\mathcal{H}}_- = \left\{ \begin{bmatrix} Y \\ t \end{bmatrix} \in \begin{bmatrix} \mathbb{R}^p \\ \mathbb{R} \end{bmatrix} \mid [\nabla f(X)^T \quad -1] \left(\begin{bmatrix} Y \\ t \end{bmatrix} - \begin{bmatrix} X \\ f(X) \end{bmatrix} \right) = 0 \right\} \quad (555)$$

One such supporting hyperplane (*confer* Figure 27(a)) is illustrated in Figure 70 for a convex quadratic.

From (553) we deduce, for each and every $X, Y \in \text{dom } f$

$$\nabla f(X)^T(Y - X) \geq 0 \Rightarrow f(Y) \geq f(X) \quad (556)$$

meaning, the gradient at X identifies a supporting hyperplane there in \mathbb{R}^p

$$\{Y \in \mathbb{R}^p \mid \nabla f(X)^T(Y - X) = 0\} \quad (557)$$

to the convex sublevel sets of convex function f (*confer* (504))

$$\mathcal{L}_{f(X)}f \triangleq \{Y \in \text{dom } f \mid f(Y) \leq f(X)\} \subseteq \mathbb{R}^p \quad (558)$$

illustrated for an arbitrary real convex function in Figure 71 and Figure 62.

3.1.10 first-order convexity condition, vector function

Now consider the first-order necessary and sufficient condition for convexity of a vector-valued function: Differentiable function $f(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}^M$ is convex if and only if $\text{dom } f$ is open, convex, and for each and every $X, Y \in \text{dom } f$

$$f(Y) \succeq_{\mathbb{R}_+^M} f(X) + \overset{\rightarrow Y-X}{df(X)} = f(X) + \left. \frac{d}{dt} \right|_{t=0} f(X + t(Y - X)) \quad (559)$$

where $\overset{\rightarrow Y-X}{df(X)}$ is the *directional derivative*^{3.16} [194] [295] of f at X in direction $Y - X$. This, of course, follows from the real-valued function case: by dual generalized inequalities (§2.13.2.0.1),

^{3.16}We extend the traditional definition of directional derivative in §D.1.4 so that direction may be indicated by a vector or a matrix, thereby broadening the scope of the Taylor series (§D.1.7). The right-hand side of the inequality (559) is the first-order Taylor series expansion of f about X .




Figure 71: Shown is a plausible contour plot in \mathbb{R}^2 of some arbitrary real differentiable convex function $f(Z)$ at selected levels α , β , and γ ; contours of equal level f (level sets) drawn in the function's domain. A convex function has convex sublevel sets $\mathcal{L}_{f(X)}f$ (558). [271, §4.6] The sublevel set whose boundary is the level set at α , for instance, comprises all the shaded regions. For any particular convex function, the family comprising all its sublevel sets is nested. [177, p.75] Were the sublevel sets not convex, we may certainly conclude the corresponding function is neither convex. Contour plots of real affine functions are illustrated in Figure 24 and Figure 66.

$$f(Y) - f(X) - \underset{\mathbb{R}_+^M}{df(X)} \succeq 0 \Leftrightarrow \left\langle f(Y) - f(X) - \underset{\mathbb{R}_+^M}{df(X)}, w \right\rangle \geq 0 \quad \forall w \succeq 0 \quad (560)$$

where

$$\underset{\mathbb{R}_+^M}{df(X)} = \begin{bmatrix} \text{tr}(\nabla f_1(X)^\top(Y - X)) \\ \text{tr}(\nabla f_2(X)^\top(Y - X)) \\ \vdots \\ \text{tr}(\nabla f_M(X)^\top(Y - X)) \end{bmatrix} \in \mathbb{R}^M \quad (561)$$

Necessary and sufficient discretization (452) allows relaxation of the semi-infinite number of conditions $w \succeq 0$ instead to $w \in \{e_i, i=1 \dots M\}$ the extreme directions of the nonnegative orthant. Each extreme direction picks out a real entry f_i and $\underset{\mathbb{R}_+^M}{df(X)}_i$ from vector-valued function f and its directional derivative $\underset{\mathbb{R}_+^M}{df(X)}$, then Theorem 3.1.9.0.1 applies.

The vector-valued function case (559) is therefore a straightforward application of the first-order convexity condition for real functions to each entry of the vector-valued function.

3.1.11 second-order convexity condition

Again, by discretization (452), we are obliged only to consider each individual entry f_i of a vector-valued function f ; *id est*, the real functions $\{f_i\}$.

For $f(X) : \mathbb{R}^p \rightarrow \mathbb{R}^M$, a twice differentiable vector-valued function with vector argument on open convex domain,

$$\underset{\mathbb{S}_+^p}{\nabla^2 f_i(X)} \succeq 0 \quad \forall X \in \text{dom } f, \quad i=1 \dots M \quad (562)$$

is a necessary and sufficient condition for convexity of f . Obviously, when $M=1$, this convexity condition also serves for a real function. Condition (562) demands nonnegative *curvature*, intuitively, hence precluding points of inflection as in Figure 72 (p.245).

Strict inequality in (562) provides only a sufficient condition for strict convexity, but that is nothing new; *videlicet*, the strictly convex real function $f_i(x) = x^4$ does not have positive second derivative at each and every $x \in \mathbb{R}$. Quadratic forms constitute a notable exception where the strict-case converse holds reliably.

3.1.11.0.1 Exercise. *Real fractional function.* (confer §3.1.4, §3.1.7.0.4)
 Prove that real function $f(x, y) = x/y$ is not convex on the first quadrant. Also exhibit this in a plot of the function. (f is *quasilinear* (p.247) on $\{y > 0\}$, in fact, and nonmonotonic even on the first quadrant.) ▼

3.1.11.0.2 Exercise. *Stress function.* (confer (1223))

Define $|x - y| \triangleq \sqrt{(x - y)^2}$ and

$$X = [x_1 \cdots x_N] \in \mathbb{R}^{1 \times N} \quad (69)$$

Given symmetric nonnegative data $[h_{ij}] \in \mathbb{S}^N \cap \mathbb{R}_+^{N \times N}$, consider function

$$f(\text{vec } X) = \sum_{i=1}^{N-1} \sum_{j=i+1}^N (|x_i - x_j| - h_{ij})^2 \in \mathbb{R} \quad (563)$$

Take the gradient and Hessian of f . Then explain why f is not a convex function; *id est*, why doesn't second-order condition (562) apply to the constant positive semidefinite Hessian matrix you found. For $N=6$ and h_{ij} data from (1294), apply the *line theorem* from §3.2.3.0.1 to plot f along some arbitrary lines through its domain. ▼

3.1.11.1 second-order \Rightarrow first-order condition

For a twice-differentiable real function $f_i(X) : \mathbb{R}^p \rightarrow \mathbb{R}$ having open domain, a consequence of the *mean value theorem* from calculus allows compression of its complete Taylor series expansion about $X \in \text{dom } f_i$ (§D.1.7) to three terms: On some open interval of $\|Y\|_2$, so that each and every line segment $[X, Y]$ belongs to $\text{dom } f_i$, there exists an $\alpha \in [0, 1]$ such that [350, §1.2.3] [37, §1.1.4]

$$f_i(Y) = f_i(X) + \nabla f_i(X)^\top (Y - X) + \frac{1}{2} (Y - X)^\top \nabla^2 f_i(\alpha X + (1 - \alpha)Y) (Y - X) \quad (564)$$

The first-order condition for convexity (553) follows directly from this and the second-order condition (562).

3.2 Matrix-valued convex function

We need different tools for matrix argument: We are primarily interested in continuous matrix-valued functions $g(X)$. We choose symmetric $g(X) \in \mathbb{S}^M$ because matrix-valued functions are most often compared (565) with respect to the positive semidefinite cone \mathbb{S}_+^M in the ambient space of symmetric matrices.^{3.17}

3.2.0.0.1 Definition. *Convex matrix-valued function:*

1) *Matrix-definition.*

A function $g(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{S}^M$ is convex in X iff $\text{dom } g$ is a convex set and, for each and every $Y, Z \in \text{dom } g$ and all $0 \leq \mu \leq 1$ [190, §2.3.7]

$$g(\mu Y + (1 - \mu)Z) \preceq_{\mathbb{S}_+^M} \mu g(Y) + (1 - \mu)g(Z) \quad (565)$$

Reversing sense of the inequality flips this definition to concavity. Strict convexity is defined less a stroke of the pen in (565) similarly to (454).

2) *Scalar-definition.*

It follows that $g(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{S}^M$ is convex in X iff $w^T g(X) w : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}$ is convex in X for each and every $\|w\| = 1$; shown by substituting the defining inequality (565). By dual generalized inequalities we have the equivalent but more broad criterion, (§2.13.5)

$$g \text{ convex} \Leftrightarrow \langle W, g \rangle \text{ convex} \quad \forall W \succeq_{\mathbb{S}_+^M} 0 \quad (566)$$

Strict convexity on both sides requires *caveat* $W \neq \mathbf{0}$. Because the set of all extreme directions for the positive semidefinite cone (§2.9.2.4) comprises a minimal set of generators for that cone, discretization (§2.13.4.2.1) allows replacement of matrix W with symmetric dyad ww^T as proposed. \triangle

^{3.17}Function symmetry is not a necessary requirement for convexity; indeed, for $A \in \mathbb{R}^{m \times p}$ and $B \in \mathbb{R}^{m \times k}$, $g(X) = AX + B$ is a convex (affine) function in X on domain $\mathbb{R}^{p \times k}$ with respect to the nonnegative orthant $\mathbb{R}_+^{m \times k}$. Symmetric convex functions share the same benefits as symmetric matrices. Horn & Johnson [180, §7.7] liken symmetric matrices to real numbers, and (symmetric) positive definite matrices to positive real numbers.

3.2.0.0.2 Example. *Taxicab distance matrix.*

Consider an n -dimensional vector space \mathbf{R}^n with metric induced by the 1-norm. Then distance between points x_1 and x_2 is the norm of their difference: $\|x_1 - x_2\|_1$. Given a list of points arranged columnar in a matrix

$$X = [x_1 \cdots x_N] \in \mathbf{R}^{n \times N} \quad (69)$$

then we could define a taxicab distance matrix

$$\begin{aligned} \mathbf{D}_1(X) &\triangleq (I \otimes \mathbf{1}_n^T) |\operatorname{vec}(X)\mathbf{1}^T - \mathbf{1} \otimes X| \in \mathbf{S}_h^N \cap \mathbf{R}_+^{N \times N} \\ &= \begin{bmatrix} 0 & \|x_1 - x_2\|_1 & \|x_1 - x_3\|_1 & \cdots & \|x_1 - x_N\|_1 \\ \|x_1 - x_2\|_1 & 0 & \|x_2 - x_3\|_1 & \cdots & \|x_2 - x_N\|_1 \\ \|x_1 - x_3\|_1 & \|x_2 - x_3\|_1 & 0 & & \|x_3 - x_N\|_1 \\ \vdots & \vdots & & \ddots & \vdots \\ \|x_1 - x_N\|_1 & \|x_2 - x_N\|_1 & \|x_3 - x_N\|_1 & \cdots & 0 \end{bmatrix} \end{aligned} \quad (567)$$

where $\mathbf{1}_n$ is a vector of ones having $\dim \mathbf{1}_n = n$ and where \otimes represents Kronecker product. This matrix-valued function is convex with respect to the nonnegative orthant since, for each and every $Y, Z \in \mathbf{R}^{n \times N}$ and all $0 \leq \mu \leq 1$

$$\mathbf{D}_1(\mu Y + (1 - \mu)Z) \underset{\mathbf{R}_+^{N \times N}}{\preceq} \mu \mathbf{D}_1(Y) + (1 - \mu) \mathbf{D}_1(Z) \quad (568)$$

□

3.2.0.0.3 Exercise. *1-norm distance matrix.*

The 1-norm is called *taxicab distance* because to go from one point to another in a city by car, road distance is a sum of grid lengths. Prove (568). ▼

3.2.1 first-order convexity condition, matrix function

From the *scalar-definition* (§3.2.0.0.1) of a convex matrix-valued function, for differentiable function g and for each and every real vector w of unit norm $\|w\| = 1$, we have

$$w^T g(Y) w \geq w^T g(X) w + w^T \overset{\rightarrow Y-X}{dg}(X) w \quad (569)$$

that follows immediately from the first-order condition (552) for convexity of a real function because

$$w^T \overset{\rightarrow Y-X}{dg}(X) w = \langle \nabla_X w^T g(X) w, Y - X \rangle \quad (570)$$

where $\overset{\rightarrow Y-X}{dg}(X)$ is the directional derivative (§D.1.4) of function g at X in direction $Y - X$. By discretized dual generalized inequalities, (§2.13.5)

$$g(Y) - g(X) - \underset{\mathbb{S}_+^M}{\overset{\rightarrow Y-X}{dg}(X)} \succeq 0 \Leftrightarrow \left\langle g(Y) - g(X) - \underset{\mathbb{S}_+^M}{\overset{\rightarrow Y-X}{dg}(X)}, ww^T \right\rangle \geq 0 \quad \forall ww^T (\succeq 0)_{\mathbb{S}_+^M} \quad (571)$$

For each and every $X, Y \in \text{dom } g$ (confer (559))

$$g(Y) \underset{\mathbb{S}_+^M}{\succeq} g(X) + \underset{\mathbb{S}_+^M}{\overset{\rightarrow Y-X}{dg}(X)} \quad (572)$$

must therefore be necessary and sufficient for convexity of a matrix-valued function of matrix variable on open convex domain.

3.2.2 epigraph of matrix-valued function, sublevel sets

We generalize the epigraph to a continuous matrix-valued function $g(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{S}^M$:

$$\text{epi } g \triangleq \{(X, T) \in \mathbb{R}^{p \times k} \times \mathbb{S}^M \mid X \in \text{dom } g, g(X) \preceq_{\mathbb{S}_+^M} T\} \quad (573)$$

from which it follows

$$g \text{ convex} \Leftrightarrow \text{epi } g \text{ convex} \quad (574)$$

Proof of necessity is similar to that in §3.1.7 on page 220.

Sublevel sets of a matrix-valued convex function corresponding to each and every $S \in \mathbb{S}^M$ (confer (504))

$$\mathcal{L}_S g \triangleq \{X \in \text{dom } g \mid g(X) \preceq_{\mathbb{S}_+^M} S\} \subseteq \mathbb{R}^{p \times k} \quad (575)$$

are convex. There is no converse.

3.2.2.0.1 Example. *Matrix fractional function* redux.

Generalizing Example 3.1.7.0.4 consider a matrix-valued function of two variables on $\text{dom } g = \mathbb{S}_+^N \times \mathbb{R}^{n \times N}$ for small positive constant ϵ (*confer* (1757))

$$g(A, X) = \epsilon X(A + \epsilon I)^{-1} X^T \quad (576)$$

where the inverse always exists by (1364). This function is convex simultaneously in both variables over the entire positive semidefinite cone \mathbb{S}_+^N and all $X \in \mathbb{R}^{n \times N}$: Consider Schur-form (1423) from §A.4: for $T \in \mathbb{S}^n$

$$\begin{aligned} G(A, X, T) &= \begin{bmatrix} A + \epsilon I & X^T \\ X & \epsilon^{-1} T \end{bmatrix} \succeq 0 \\ &\Leftrightarrow \\ &T - \epsilon X(A + \epsilon I)^{-1} X^T \succeq 0 \\ &A + \epsilon I \succ 0 \end{aligned} \quad (577)$$

By Theorem 2.1.9.0.1, inverse image of the positive semidefinite cone \mathbb{S}_+^{N+n} under affine mapping $G(A, X, T)$ is convex. Function $g(A, X)$ is convex on $\mathbb{S}_+^N \times \mathbb{R}^{n \times N}$ because its epigraph is that inverse image:

$$\text{epi } g(A, X) = \{(A, X, T) \mid A + \epsilon I \succ 0, \epsilon X(A + \epsilon I)^{-1} X^T \preceq T\} = G^{-1}(\mathbb{S}_+^{N+n}) \quad (578)$$

□

3.2.3 second-order condition, matrix function

The following *line theorem* is a potent tool for establishing convexity of a multidimensional function. To understand it, what is meant by *line* must first be solidified. Given a function $g(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{S}^M$ and particular $X, Y \in \mathbb{R}^{p \times k}$ not necessarily in that function's domain, then we say a line $\{X + tY \mid t \in \mathbb{R}\}$ passes through $\text{dom } g$ when $X + tY \in \text{dom } g$ over some interval of $t \in \mathbb{R}$.

3.2.3.0.1 Theorem. *Line theorem.* [53, §3.1.1]

Matrix-valued function $g(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{S}^M$ is convex in X if and only if it remains convex on the intersection of any line with its domain. \diamond

Now we assume a twice differentiable function.

3.2.3.0.2 Definition. *Differentiable convex matrix-valued function.*

Matrix-valued function $g(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{S}^M$ is convex in X iff $\text{dom } g$ is an open convex set, and its second derivative $g''(X + tY) : \mathbb{R} \rightarrow \mathbb{S}^M$ is positive semidefinite on each point of intersection along every line $\{X + tY \mid t \in \mathbb{R}\}$ that intersects $\text{dom } g$; *id est*, iff for each and every $X, Y \in \mathbb{R}^{p \times k}$ such that $X + tY \in \text{dom } g$ over some open interval of $t \in \mathbb{R}$

$$\frac{d^2}{dt^2} g(X + tY) \succeq_0 \quad (579)$$

Similarly, if

$$\frac{d^2}{dt^2} g(X + tY) \succ_0 \quad (580)$$

then g is strictly convex; the converse is generally false. [53, §3.1.4]^{3.18} \triangle

3.2.3.0.3 Example. *Matrix inverse.* (confer §3.1.5)

The matrix-valued function X^μ is convex on $\text{int } \mathbb{S}_+^M$ for $-1 \leq \mu \leq 0$ or $1 \leq \mu \leq 2$ and concave for $0 \leq \mu \leq 1$. [53, §3.6.2] In particular, the function $g(X) = X^{-1}$ is convex on $\text{int } \mathbb{S}_+^M$. For each and every $Y \in \mathbb{S}^M$ (§D.2.1, §A.3.1.0.5)

$$\frac{d^2}{dt^2} g(X + tY) = 2(X + tY)^{-1} Y (X + tY)^{-1} Y (X + tY)^{-1} \succeq_0 \quad (581)$$

on some open interval of $t \in \mathbb{R}$ such that $X + tY \succ_0$. Hence, $g(X)$ is convex in X . This result is extensible;^{3.19} $\text{tr } X^{-1}$ is convex on that same domain. [180, §7.6, prob.2] [48, §3.1, exer.25] \square

3.2.3.0.4 Example. *Matrix squared.*

Iconic real function $f(x) = x^2$ is strictly convex on \mathbb{R} . The matrix-valued function $g(X) = X^2$ is convex on the domain of symmetric matrices; for $X, Y \in \mathbb{S}^M$ and any open interval of $t \in \mathbb{R}$ (§D.2.1)

$$\frac{d^2}{dt^2} g(X + tY) = \frac{d^2}{dt^2} (X + tY)^2 = 2Y^2 \quad (582)$$

^{3.18}The strict-case converse is reliably true for quadratic forms.

^{3.19} $d/dt \text{tr } g(X + tY) = \text{tr } d/dt g(X + tY)$. [181, p.491]

which is positive semidefinite when Y is symmetric because then $Y^2 = Y^T Y$ (1370).^{3.20}

A more appropriate matrix-valued counterpart for f is $g(X) = X^T X$ which is a convex function on domain $\{X \in \mathbb{R}^{m \times n}\}$, and strictly convex whenever X is skinny-or-square full-rank. This matrix-valued function can be generalized to $g(X) = X^T A X$ which is convex whenever matrix A is positive semidefinite (p.636), and strictly convex when A is positive definite and X is skinny-or-square full-rank (Corollary A.3.1.0.5). \square

3.2.3.0.5 Exercise. Squared maps.

Give seven examples of distinct polyhedra \mathcal{P} for which the set

$$\{X^T X \mid X \in \mathcal{P}\} \subseteq \mathbb{S}_+^n \quad (583)$$

were convex. Is this set convex, in general, for any polyhedron \mathcal{P} ? (confer (1151) (1158)) Is the epigraph of function $g(X) = X^T X$ convex for any polyhedral domain? \blacktriangledown

3.2.3.0.6 Example. Matrix exponential.

The matrix-valued function $g(X) = e^X : \mathbb{S}^M \rightarrow \mathbb{S}^M$ is convex on the subspace of *circulant* [147] symmetric matrices. Applying the *line theorem*, for all $t \in \mathbb{R}$ and circulant $X, Y \in \mathbb{S}^M$, from Table D.2.7 we have

$$\frac{d^2}{dt^2} e^{X+tY} = Y e^{X+tY} Y \underset{\mathbb{S}_+^M}{\succeq} 0, \quad (XY)^T = XY \quad (584)$$

because all circulant matrices are *commutative* and, for symmetric matrices, $XY = YX \Leftrightarrow (XY)^T = XY$ (1387). Given symmetric argument, the matrix exponential always resides interior to the cone of positive semidefinite matrices in the symmetric matrix subspace; $e^A \succ 0 \quad \forall A \in \mathbb{S}^M$ (1754). Then for any matrix Y of compatible dimension, $Y^T e^A Y$ is positive semidefinite. (§A.3.1.0.5)

The subspace of circulant symmetric matrices contains all diagonal matrices. The matrix exponential of any diagonal matrix e^A exponentiates each individual entry on the main diagonal. [221, §5.3] So, changing the function domain to the subspace of real diagonal matrices reduces the

^{3.20}By (1388) in §A.3.1, changing the domain instead to all symmetric and nonsymmetric positive semidefinite matrices, for example, will not produce a convex function.




Figure 72: Iconic unimodal differentiable quasiconvex function of two variables graphed in $\mathbb{R}^2 \times \mathbb{R}$ on some open disc in \mathbb{R}^2 . Note reversal of curvature in direction of gradient.

matrix exponential to a vector-valued function in an isometrically isomorphic subspace \mathbb{R}^M ; known convex (§3.1.1) from the real-valued function case [53, §3.1.5]. \square

There are, of course, multifarious methods to determine function convexity, [53] [37] [115] each of them efficient when appropriate.

3.2.3.0.7 Exercise. *log det function.*

Show by two different methods: $\log \det X$ is concave on the interior of the positive semidefinite cone. \blacktriangledown

3.3 Quasiconvex

Quasiconvex functions [53, §3.4] [177] [293] [331] [215, §2] are useful in practical problem solving because they are *unimodal* (by definition when nonmonotonic); a global minimum is guaranteed to exist over any convex set in the function domain; *e.g.*, Figure 72.

3.3.0.0.1 Definition. *Quasiconvex function.*

$f(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{R}$ is a quasiconvex function of matrix X iff $\text{dom } f$ is a convex set and for each and every $Y, Z \in \text{dom } f$, $0 \leq \mu \leq 1$

$$f(\mu Y + (1 - \mu)Z) \leq \max\{f(Y), f(Z)\} \quad (585)$$

A quasiconcave function is determined:

$$f(\mu Y + (1 - \mu)Z) \geq \min\{f(Y), f(Z)\} \quad (586)$$

△

Unlike convex functions, quasiconvex functions are not necessarily continuous; *e.g.*, quasiconcave $\text{rank}(X)$ on \mathbb{S}_+^M (§2.9.2.6.2) and $\text{card}(x)$ on \mathbb{R}_+^M . Although insufficient for convex functions, convexity of each and every sublevel set serves as a definition of quasiconvexity:

3.3.0.0.2 Definition. *Quasiconvex multidimensional function.*

Scalar-, vector-, or matrix-valued function $g(X) : \mathbb{R}^{p \times k} \rightarrow \mathbb{S}^M$ is a quasiconvex function of matrix X iff $\text{dom } g$ is a convex set and the sublevel set corresponding to each and every $S \in \mathbb{S}^M$

$$\mathcal{L}_S g = \{X \in \text{dom } g \mid g(X) \preceq S\} \subseteq \mathbb{R}^{p \times k} \quad (575)$$

is convex. Vectors are compared with respect to the nonnegative orthant \mathbb{R}_+^M while matrices are with respect to the positive semidefinite cone \mathbb{S}_+^M .

Convexity of the *superlevel set* corresponding to each and every $S \in \mathbb{S}^M$, likewise

$$\mathcal{L}^S g = \{X \in \text{dom } g \mid g(X) \succeq S\} \subseteq \mathbb{R}^{p \times k} \quad (587)$$

is necessary and sufficient for quasiconcavity.

△

3.3.0.0.3 Exercise. *Nonconvexity of matrix product.*

Consider real function f on a positive definite domain

$$f(X) = \text{tr}(X_1 X_2), \quad X \triangleq \begin{bmatrix} X_1 \\ X_2 \end{bmatrix} \in \text{dom } f \triangleq \begin{bmatrix} \text{rel int } \mathbb{S}_+^N \\ \text{rel int } \mathbb{S}_+^N \end{bmatrix} \quad (588)$$

with superlevel sets

$$\mathcal{L}^s f = \{X \in \text{dom } f \mid \langle X_1, X_2 \rangle \geq s\} \quad (589)$$

Prove: $f(X)$ is not quasiconcave except when $N = 1$, nor is it quasiconvex unless $X_1 = X_2$. ▼

3.3.1 bilinear

Bilinear function^{3.21} $x^T y$ of vectors x and y is quasiconcave (monotonic) on the entirety of the nonnegative orthants only when vectors are of dimension 1.

3.3.2 quasilinear

When a function is simultaneously quasiconvex and quasiconcave, it is called *quasilinear*. Quasilinear functions are completely determined by convex level sets. One-dimensional function $f(x) = x^3$ and vector-valued signum function $\text{sgn}(x)$, for example, are quasilinear. Any monotonic function is quasilinear^{3.22} (but not *vice versa*, Exercise 3.1.11.0.1).

3.4 Salient properties of convex and quasiconvex functions

1.
 - A convex (or concave) function is assumed continuous but not necessarily differentiable on the relative interior of its domain. [271, §10]
 - A quasiconvex (or quasiconcave) function is not necessarily a continuous function.
2. convex epigraph \Leftrightarrow convexity \Rightarrow quasiconvexity \Leftrightarrow convex sublevel sets.
 convex hypograph \Leftrightarrow concavity \Rightarrow quasiconcavity \Leftrightarrow convex superlevel.
 monotonicity \Rightarrow quasilinearity \Leftrightarrow convex level sets.
3. log-convex \Rightarrow convex \Rightarrow quasiconvex. 3.23
 concave \Rightarrow quasiconcave \Leftarrow log-concave \Leftarrow positive concave.
4.
 - (*homogeneity*) Convexity, concavity, quasiconvexity, and quasiconcavity are invariant to nonnegative scaling of function.
 - g convex $\Leftrightarrow -g$ concave
 g quasiconvex $\Leftrightarrow -g$ quasiconcave
 g log-convex $\Leftrightarrow 1/g$ log-concave

^{3.21}Convex envelope of bilinear functions is well known. [2]

^{3.22}*e.g.*, a monotonic concave function is quasiconvex, but dare not confuse these terms.

^{3.23}*Log-convex* means: logarithm of function f is convex on $\text{dom } f$.

5. The *line theorem* (§3.2.3.0.1) translates identically to quasiconvexity (quasiconcavity). [53, §3.4.2]
6. (*affine transformation of argument*) Composition $g(h(X))$ of convex (concave) function g with any affine function $h: \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^{p \times k}$ remains convex (concave) in $X \in \mathbb{R}^{m \times n}$, where $h(\mathbb{R}^{m \times n}) \cap \text{dom } g \neq \emptyset$. [177, §B.2.1] Likewise for the quasiconvex (quasiconcave) functions g .
7.
 - – Nonnegatively weighted sum of convex (concave) functions remains convex (concave). (§3.1.1.0.1)
 - Nonnegatively weighted nonzero sum of strictly convex (concave) functions remains strictly convex (concave).
 - Nonnegatively weighted maximum (minimum) of convex (concave) functions remains convex (concave).
 - Pointwise supremum (infimum) of convex (concave) functions remains convex (concave). (Figure 67) [271, §5]
 - – Sum of quasiconvex functions not necessarily quasiconvex.
 - Nonnegatively weighted maximum (minimum) of quasiconvex (quasiconcave) functions remains quasiconvex (quasiconcave).
 - Pointwise supremum (infimum) of quasiconvex (quasiconcave) functions remains quasiconvex (quasiconcave).

Chapter 4

Semidefinite programming

Prior to 1984, linear and nonlinear programming,^{4.1} one a subset of the other, had evolved for the most part along unconnected paths, without even a common terminology. (The use of “programming” to mean “optimization” serves as a persistent reminder of these differences.)

–Forsgren, Gill, & Wright (2002) [126]

Given some application of convex analysis, it may at first seem puzzling why the search for its solution ends abruptly with a formalized statement of the problem itself as a constrained optimization. The explanation is: typically we do not seek analytical solution because there are relatively few. (§C, §3.1.7.2) If a problem can be expressed in convex form, rather, then there exist computer programs providing efficient numerical global solution. [146] [343] [344] [342] [350] The goal, then, becomes conversion of a given problem (perhaps a nonconvex or combinatorial problem statement) to an equivalent convex form or to an *alternation* of convex subproblems convergent to a solution of the original problem:

By the *fundamental theorem of Convex Optimization*, any locally optimal point of a convex problem is globally optimal. [53, §4.2.2] [270, §1] Given convex real objective function g and convex feasible set $\mathcal{C} \subseteq \text{dom } g$, which is the set of all variable values satisfying the problem constraints, we pose a

^{4.1} nascence of *interior-point methods* of solution [323] [340].
Linear programming \subset (convex \cap nonlinear) programming.

generic convex optimization problem

$$\begin{aligned} & \underset{X}{\text{minimize}} && g(X) \\ & \text{subject to} && X \in \mathcal{C} \end{aligned} \tag{590}$$

where constraints are abstract here in the membership of variable X to feasible set \mathcal{C} . Inequality constraint functions of a convex optimization problem are convex while equality constraint functions are conventionally affine, but not necessarily so. Affine equality constraint functions (convex), as opposed to the larger set of all convex equality constraint functions having convex level sets (§3.1.6.0.4), make convex optimization tractable.

Similarly, the problem

$$\begin{aligned} & \underset{X}{\text{maximize}} && g(X) \\ & \text{subject to} && X \in \mathcal{C} \end{aligned} \tag{591}$$

is convex were g a real concave function. As conversion to convex form is not always possible, there is much ongoing research to determine which problem classes have convex expression or relaxation. [31] [51] [131] [244] [307] [129]

4.1 Conic problem

Still, we are surprised to see the relatively small number of submissions to semidefinite programming (SDP) solvers, as this is an area of significant current interest to the optimization community. We speculate that semidefinite programming is simply experiencing the fate of most new areas: Users have yet to understand how to pose their problems as semidefinite programs, and the lack of support for SDP solvers in popular modelling languages likely discourages submissions.

—SIAM News, 2002. [98, p.9]

Consider a *conic problem* (p) and its dual (d): [259, §3.3.1] [211, §2.1] [212] (*confer* p.150)

$$\begin{array}{ll} \text{(p)} & \underset{x}{\text{minimize}} \quad c^T x \\ & \text{subject to} \quad x \in \mathcal{K} \\ & \quad \quad \quad Ax = b \end{array} \qquad \begin{array}{ll} & \underset{y,s}{\text{maximize}} \quad b^T y \\ & \text{subject to} \quad s \in \mathcal{K}^* \\ & \quad \quad \quad A^T y + s = c \end{array} \tag{275}$$

where \mathcal{K} is a closed convex cone, \mathcal{K}^* is its dual, matrix A is fixed, and the remaining quantities are vectors.

When \mathcal{K} is a polyhedral cone (§2.12.1), then each conic problem becomes a *linear program* [82]; the self-dual nonnegative orthant providing the prototypical primal linear program and its dual. More generally, each optimization problem is convex when \mathcal{K} is a closed convex cone. Unlike the optimal objective value, a solution to each problem is not necessarily unique; in other words, the optimal solution set $\{x^*\}$ or $\{y^*, s^*\}$ is convex and may comprise more than a single point although the corresponding optimal objective value is unique when the feasible set is nonempty.

4.1.1 a Semidefinite program

When \mathcal{K} is the self-dual cone of positive semidefinite matrices in the subspace of symmetric matrices, then each conic problem is called a *semidefinite program* (SDP); [244, §6.4] primal problem (P) having matrix variable $X \in \mathbb{S}^n$ while corresponding dual (D) has matrix *slack variable* $S \in \mathbb{S}^n$ and vector variable $y = [y_i] \in \mathbb{R}^m$: [9] [10, §2] [350, §1.3.8]

$$\begin{array}{ll}
 \text{minimize}_{X \in \mathbb{S}^n} & \langle C, X \rangle \\
 \text{(P) subject to} & X \succeq 0 \\
 & A \text{ svec } X = b
 \end{array}
 \qquad
 \begin{array}{ll}
 \text{maximize}_{y \in \mathbb{R}^m, S \in \mathbb{S}^n} & \langle b, y \rangle \\
 \text{subject to} & S \succeq 0 \\
 & \text{svec}^{-1}(A^T y) + S = C
 \end{array}
 \quad \text{(D)}$$

$$A \text{ svec } X = b \qquad \text{svec}^{-1}(A^T y) + S = C \quad (592)$$

This is the prototypical semidefinite program and its dual, where matrix $C \in \mathbb{S}^n$ and vector $b \in \mathbb{R}^m$ are fixed, as is

$$A \triangleq \begin{bmatrix} \text{svec}(A_1)^T \\ \vdots \\ \text{svec}(A_m)^T \end{bmatrix} \in \mathbb{R}^{m \times n(n+1)/2} \quad (593)$$

where $A_i \in \mathbb{S}^n$, $i=1 \dots m$, are given. Thus

$$\begin{aligned}
 A \text{ svec } X &= \begin{bmatrix} \langle A_1, X \rangle \\ \vdots \\ \langle A_m, X \rangle \end{bmatrix} \\
 \text{svec}^{-1}(A^T y) &= \sum_{i=1}^m y_i A_i
 \end{aligned}
 \quad (594)$$

The vector inner-product for matrices is defined in the Euclidean/Frobenius sense in the isomorphic vector space $\mathbb{R}^{n(n+1)/2}$; *id est*,

$$\langle C, X \rangle \triangleq \text{tr}(C^T X) = \text{svec}(C)^T \text{svec} X \quad (32)$$

where $\text{svec} X$ defined by (50) denotes symmetric vectorization.

Linear programming, invented by Dantzig in 1947, is now integral to modern technology. But the same cannot yet be said of semidefinite programming introduced by Bellman & Fan in 1963. [86] Numerical performance of contemporary general-purpose semidefinite program solvers is limited: Computational intensity varies as $O(n^6)$ based on interior-point methods that produce solutions no more relatively accurate than 1E-8. There are no solvers available capable of handling in excess of $n=100$ variables without significant, sometimes crippling, loss of precision or time.^{4.2}

Nevertheless, semidefinite programming has emerged recently to prominence primarily because it admits a new class of problem previously unsolvable by convex optimization techniques, [51] secondarily because it theoretically subsumes other convex techniques such as linear, quadratic, and *second-order cone programming*. Determination of the Riemann mapping function from complex analysis [253] [28, §8, §13], for example, can be posed as a semidefinite program.

4.1.1.1 Maximal complementarity

It has been shown that contemporary *interior-point methods* (developed circa 1990 [131]) [53, §11] [341] [256] [244] [10] [126] for numerical solution of semidefinite programs can converge to a solution of *maximal complementarity*; [155, §5] [349] [223] [138] not a vertex-solution but a solution of highest cardinality or rank among all optimal solutions.^{4.3} [350, §2.5.3]

^{4.2}*Heuristics are not ruled out by SIOPT; indeed I would suspect that most successful methods have (appropriately described) heuristics under the hood - my codes certainly do. ... Of course, there are still questions relating to high-accuracy and speed, but for many applications a few digits of accuracy suffices and overnight runs for non-real-time delivery is acceptable.*

–Nicholas I. M. Gould, Editor-in-Chief, SIOPT

^{4.3}This characteristic might be regarded as a disadvantage to this method of numerical solution, but this behavior is not certain and depends on solver implementation.

4.1.1.2 Reduced-rank solution

A simple rank reduction algorithm, for construction of a primal optimal solution X^* to (592P) satisfying an upper bound on rank governed by Proposition 2.9.3.0.1, is presented in §4.3. That proposition asserts existence of feasible solutions with an upper bound on their rank; [25, §II.13.1] specifically, it asserts an extreme point (§2.6.0.0.1) of the primal feasible set $\mathcal{A} \cap \mathbb{S}_+^n$ satisfies upper bound

$$\text{rank } X \leq \left\lfloor \frac{\sqrt{8m+1} - 1}{2} \right\rfloor \quad (246)$$

where, given $A \in \mathbb{R}^{m \times n(n+1)/2}$ and $b \in \mathbb{R}^m$

$$\mathcal{A} \triangleq \{X \in \mathbb{S}^n \mid A \text{ svec } X = b\} \quad (595)$$

is the affine subset from primal problem (592P).

4.1.1.3 Coexistence of low- and high-rank solutions; analogy

That low-rank and high-rank optimal solutions $\{X^*\}$ of (592P) coexist may be grasped with the following analogy: We compare a proper polyhedral cone \mathcal{S}_+^3 in \mathbb{R}^3 (illustrated in Figure 73) to the positive semidefinite cone \mathbb{S}_+^3 in isometrically isomorphic \mathbb{R}^6 , difficult to visualize. The analogy is good:

- $\text{int } \mathbb{S}_+^3$ is constituted by rank-3 matrices
 $\text{int } \mathcal{S}_+^3$ has three dimensions
- boundary $\partial \mathbb{S}_+^3$ contains rank-0, rank-1, and rank-2 matrices
 boundary $\partial \mathcal{S}_+^3$ contains 0-, 1-, and 2-dimensional faces
- the only rank-0 matrix resides in the vertex at the origin
- Rank-1 matrices are in one-to-one correspondence with extreme directions of \mathbb{S}_+^3 and \mathcal{S}_+^3 . The set of all rank-1 symmetric matrices in this dimension

$$\{G \in \mathbb{S}_+^3 \mid \text{rank } G = 1\} \quad (596)$$

is not a connected set.




Figure 73: Visualizing positive semidefinite cone in high dimension: Proper polyhedral cone $\mathcal{S}_+^3 \subset \mathbb{R}^3$ representing positive semidefinite cone $\mathbb{S}_+^3 \subset \mathbb{S}^3$; analogizing its intersection with hyperplane $\mathbb{S}_+^3 \cap \partial\mathcal{H}$. Number of facets is arbitrary (analogy is not inspired by eigen decomposition). The rank-0 positive semidefinite matrix corresponds to the origin in \mathbb{R}^3 , rank-1 positive semidefinite matrices correspond to the edges of the polyhedral cone, rank-2 to the facet relative interiors, and rank-3 to the polyhedral cone interior. Vertices Γ_1 and Γ_2 are extreme points of polyhedron $\mathcal{P} = \partial\mathcal{H} \cap \mathcal{S}_+^3$, and extreme directions of \mathcal{S}_+^3 . A given vector C is normal to another hyperplane (not illustrated but independent w.r.t $\partial\mathcal{H}$) containing line segment $\overline{\Gamma_1\Gamma_2}$ minimizing real linear function $\langle C, X \rangle$ on \mathcal{P} . (confer Figure 24, Figure 28)

- In any SDP feasibility problem, an SDP feasible solution with the lowest rank must be an extreme point of the feasible set. Thus, there must exist an SDP objective function such that this lowest-rank feasible solution uniquely optimizes it. –Ye, 2006 [211, §2.4] [212]
- Rank of a sum of members $F + G$ in Lemma 2.9.2.6.1 and location of a difference $F - G$ in §2.9.2.9.1 similarly hold for \mathbb{S}_+^3 and \mathcal{S}_+^3 .
- Euclidean distance from any particular rank-3 positive semidefinite matrix (in the cone interior) to the closest rank-2 positive semidefinite matrix (on the boundary) is generally less than the distance to the closest rank-1 positive semidefinite matrix. (§7.1.2)
- distance from any point in $\partial\mathbb{S}_+^3$ to $\text{int}\mathbb{S}_+^3$ is infinitesimal (§2.1.7.1.1)
distance from any point in $\partial\mathcal{S}_+^3$ to $\text{int}\mathcal{S}_+^3$ is infinitesimal
- faces of \mathbb{S}_+^3 correspond to faces of \mathcal{S}_+^3 (confer Table 2.9.2.3.1)

	k	$\dim \mathcal{F}(\mathbb{S}_+^3)$	$\dim \mathcal{F}(\mathcal{S}_+^3)$	$\dim \mathcal{F}(\mathbb{S}_+^3 \ni \text{rank-}k \text{ matrix})$
boundary	0	0	0	0
	1	1	1	1
	2	2	3	3
interior	3	3	6	6

Integer k indexes k -dimensional faces \mathcal{F} of \mathcal{S}_+^3 . Positive semidefinite cone \mathbb{S}_+^3 has four kinds of faces, including cone itself ($k = 3$, boundary + interior), whose dimensions in isometrically isomorphic \mathbb{R}^6 are listed under $\dim \mathcal{F}(\mathbb{S}_+^3)$. Smallest face $\mathcal{F}(\mathbb{S}_+^3 \ni \text{rank-}k \text{ matrix})$ that contains a rank- k positive semidefinite matrix has dimension $k(k + 1)/2$ by (205).

- For \mathcal{A} equal to intersection of m hyperplanes having independent normals, and for $X \in \mathcal{S}_+^3 \cap \mathcal{A}$, we have $\text{rank } X \leq m$; the analogue to (246).

Proof. With reference to Figure 73: Assume one ($m = 1$) hyperplane $\mathcal{A} = \partial\mathcal{H}$ intersects the polyhedral cone. Every intersecting plane contains at least one matrix having rank less than or equal to 1; *id est*, from all $X \in \partial\mathcal{H} \cap \mathcal{S}_+^3$ there exists an X such that $\text{rank } X \leq 1$. Rank 1 is therefore an upper bound in this case.

Now visualize intersection of the polyhedral cone with two ($m = 2$) hyperplanes having linearly independent normals. The hyperplane intersection \mathcal{A} makes a line. Every intersecting line contains at least one matrix having rank less than or equal to 2, providing an upper bound. In other words, there exists a positive semidefinite matrix X belonging to any line intersecting the polyhedral cone such that $\text{rank } X \leq 2$.

In the case of three independent intersecting hyperplanes ($m = 3$), the hyperplane intersection \mathcal{A} makes a point that can reside anywhere in the polyhedral cone. The upper bound on a point in \mathcal{S}_+^3 is also the greatest upper bound: $\text{rank } X \leq 3$. \blacklozenge

4.1.1.3.1 Example. *Optimization on $\mathcal{A} \cap \mathcal{S}_+^3$.*

Consider minimization of the real linear function $\langle C, X \rangle$ on

$$\mathcal{P} \triangleq \mathcal{A} \cap \mathcal{S}_+^3 \quad (597)$$

a polyhedral feasible set;

$$\begin{aligned} f_0^* &\triangleq \underset{X}{\text{minimize}} \quad \langle C, X \rangle \\ &\text{subject to} \quad X \in \mathcal{A} \cap \mathcal{S}_+^3 \end{aligned} \quad (598)$$

As illustrated for particular vector C and hyperplane $\mathcal{A} = \partial\mathcal{H}$ in Figure 73, this linear function is minimized on any X belonging to the face of \mathcal{P} containing extreme points $\{\Gamma_1, \Gamma_2\}$ and all the rank-2 matrices in between; *id est*, on any X belonging to the face of \mathcal{P}

$$\mathcal{F}(\mathcal{P}) = \{X \mid \langle C, X \rangle = f_0^*\} \cap \mathcal{A} \cap \mathcal{S}_+^3 \quad (599)$$

exposed by the hyperplane $\{X \mid \langle C, X \rangle = f_0^*\}$. In other words, the set of all optimal points X^* is a face of \mathcal{P}

$$\{X^*\} = \mathcal{F}(\mathcal{P}) = \overline{\Gamma_1\Gamma_2} \quad (600)$$

comprising rank-1 and rank-2 positive semidefinite matrices. Rank 1 is the upper bound on existence in the feasible set \mathcal{P} for this case $m = 1$ hyperplane constituting \mathcal{A} . The rank-1 matrices Γ_1 and Γ_2 in face $\mathcal{F}(\mathcal{P})$ are extreme points of that face and (by transitivity (§2.6.1.2)) extreme points of the intersection \mathcal{P} as well. As predicted by analogy to Barvinok's

Proposition 2.9.3.0.1, the upper bound on rank of X existent in the feasible set \mathcal{P} is satisfied by an extreme point. The upper bound on rank of an optimal solution X^* existent in $\mathcal{F}(\mathcal{P})$ is thereby also satisfied by an extreme point of \mathcal{P} precisely because $\{X^*\}$ constitutes $\mathcal{F}(\mathcal{P})$;^{4.4} in particular,

$$\{X^* \in \mathcal{P} \mid \text{rank } X^* \leq 1\} = \{\Gamma_1, \Gamma_2\} \subseteq \mathcal{F}(\mathcal{P}) \quad (601)$$

As all linear functions on a polyhedron are minimized on a face, [82] [222] [243] [246] by analogy we so demonstrate coexistence of optimal solutions X^* of (592P) having assorted rank. \square

4.1.1.4 Previous work

Barvinok showed [23, §2.2] when given a positive definite matrix C and an arbitrarily small neighborhood of C comprising positive definite matrices, there exists a matrix \tilde{C} from that neighborhood such that optimal solution X^* to (592P) (substituting \tilde{C}) is an extreme point of $\mathcal{A} \cap \mathbb{S}_+^n$ and satisfies upper bound (246).^{4.5} Given arbitrary positive definite C , this means nothing inherently guarantees that an optimal solution X^* to problem (592P) satisfies (246); certainly nothing given any symmetric matrix C , as the problem is posed. This can be proved by example:

4.1.1.4.1 Example. (Ye) *Maximal Complementarity.*

Assume dimension n to be an even positive number. Then the particular instance of problem (592P),

$$\begin{aligned} & \underset{X \in \mathbb{S}^n}{\text{minimize}} && \left\langle \begin{bmatrix} I & \mathbf{0} \\ \mathbf{0} & 2I \end{bmatrix}, X \right\rangle \\ & \text{subject to} && X \succeq 0 \\ & && \langle I, X \rangle = n \end{aligned} \quad (602)$$

has optimal solution

$$X^* = \begin{bmatrix} 2I & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \in \mathbb{S}^n \quad (603)$$

^{4.4}and every face contains a subset of the extreme points of \mathcal{P} by the *extreme existence theorem* (§2.6.0.0.2). This means: because the affine subset \mathcal{A} and hyperplane $\{X \mid \langle C, X \rangle = f_0^*\}$ must intersect a whole face of \mathcal{P} , calculation of an upper bound on rank of X^* ignores counting the hyperplane when determining m in (246).

^{4.5}Further, the set of all such \tilde{C} in that neighborhood is open and dense.

with an equal number of twos and zeros along the main diagonal. Indeed, optimal solution (603) is a terminal solution along the *central path* taken by the interior-point method as implemented in [350, §2.5.3]; it is also a solution of highest rank among all optimal solutions to (602). Clearly, rank of this primal optimal solution exceeds by far a rank-1 solution predicted by upper bound (246). \square

4.1.1.5 Later developments

This rational example (602) indicates the need for a more generally applicable and simple algorithm to identify an optimal solution X^* satisfying Barvinok's Proposition 2.9.3.0.1. We will review such an algorithm in §4.3, but first we provide more background.

4.2 Framework

4.2.1 Feasible sets

Denote by \mathcal{C} and \mathcal{C}^* the convex sets of primal and dual points respectively satisfying the primal and dual constraints in (592), each assumed nonempty;

$$\mathcal{C} = \left\{ X \in \mathbb{S}_+^n \mid \begin{bmatrix} \langle A_1, X \rangle \\ \vdots \\ \langle A_m, X \rangle \end{bmatrix} = b \right\} = \mathcal{A} \cap \mathbb{S}_+^n \quad (604)$$

$$\mathcal{C}^* = \left\{ S \in \mathbb{S}_+^n, y = [y_i] \in \mathbb{R}^m \mid \sum_{i=1}^m y_i A_i + S = C \right\}$$

These are the *primal feasible set* and *dual feasible set*. Geometrically, primal feasible $\mathcal{A} \cap \mathbb{S}_+^n$ represents an intersection of the positive semidefinite cone \mathbb{S}_+^n with an affine subset \mathcal{A} of the subspace of symmetric matrices \mathbb{S}^n in isometrically isomorphic $\mathbb{R}^{n(n+1)/2}$. The affine subset has dimension $n(n+1)/2 - m$ when the A_i are linearly independent. Dual feasible set \mathcal{C}^* is the Cartesian product of the positive semidefinite cone with its inverse image (§2.1.9.0.1) under affine transformation $C - \sum y_i A_i$.^{4.6} Both feasible

^{4.6}The inequality $C - \sum y_i A_i \succeq 0$ follows directly from (592D) (§2.9.0.1.1) and is known as a *linear matrix inequality*. (§2.13.5.1.1) Because $\sum y_i A_i \preceq C$, matrix S is known as a slack variable (a term borrowed from linear programming [82]) since its inclusion raises this inequality to equality.

sets are closed and convex, and the objective functions on a Euclidean vector space are linear. Hence, convex optimization problems (592P) and (592D).

4.2.1.1 $\mathcal{A} \cap \mathbb{S}_+^n$ emptiness determination via Farkas' lemma

4.2.1.1.1 Lemma. *Semidefinite Farkas' lemma.*

Given an arbitrary set $\{A_i \in \mathbb{S}^n, i=1 \dots m\}$ and a vector $b = [b_i] \in \mathbb{R}^m$, define the affine subset

$$\mathcal{A} = \{X \in \mathbb{S}^n \mid \langle A_i, X \rangle = b_i, i=1 \dots m\} \quad (595)$$

Primal feasible set $\mathcal{A} \cap \mathbb{S}_+^n$ is nonempty if and only if $y^T b \geq 0$ holds for each and every vector $y = [y_i] \in \mathbb{R}^m$ such that $\sum_{i=1}^m y_i A_i \succeq 0$.

Equivalently, primal feasible set $\mathcal{A} \cap \mathbb{S}_+^n$ is nonempty if and only if $y^T b \geq 0$ holds for each and every norm-1 vector $\|y\|=1$ such that $\sum_{i=1}^m y_i A_i \succeq 0$. \diamond

Semidefinite Farkas' lemma follows directly from a membership relation (§2.13.2.0.1) and the closed convex cones from *linear matrix inequality example 2.13.5.1.1*; given convex cone \mathcal{K} and its dual

$$\mathcal{K} = \{A \text{ svec } X \mid X \succeq 0\} \quad (344)$$

$$\mathcal{K}^* = \{y \mid \sum_{j=1}^m y_j A_j \succeq 0\} \quad (349)$$

where

$$A = \begin{bmatrix} \text{svec}(A_1)^T \\ \vdots \\ \text{svec}(A_m)^T \end{bmatrix} \in \mathbb{R}^{m \times n(n+1)/2} \quad (593)$$

then we have membership relation

$$b \in \mathcal{K} \Leftrightarrow \langle y, b \rangle \geq 0 \quad \forall y \in \mathcal{K}^* \quad (290)$$

and equivalents

$$b \in \mathcal{K} \Leftrightarrow \exists X \succeq 0 \ni A \text{ svec } X = b \Leftrightarrow \mathcal{A} \cap \mathbb{S}_+^n \neq \emptyset \quad (605)$$

$$b \in \mathcal{K} \Leftrightarrow \langle y, b \rangle \geq 0 \quad \forall y \in \mathcal{K}^* \Leftrightarrow \mathcal{A} \cap \mathbb{S}_+^n \neq \emptyset \quad (606)$$

Semidefinite Farkas' lemma provides the conditions required for a set of hyperplanes to have a nonempty intersection $\mathcal{A} \cap \mathbb{S}_+^n$ with the positive semidefinite cone. While the lemma as stated is correct, Ye points out [350, §1.3.8] that a positive definite version of this lemma is required for semidefinite programming because any feasible point in the relative interior $\mathcal{A} \cap \text{int } \mathbb{S}_+^n$ is required by *Slater's condition*^{4.7} to achieve 0 duality gap (primal–dual objective difference §4.2.3, Figure 54). In our circumstance, assuming a nonempty intersection, a positive definite lemma is required to insure a point of intersection closest to the origin is not at infinity; *e.g.*, Figure 41. Then given $A \in \mathbb{R}^{m \times n(n+1)/2}$ having rank m , we wish to detect existence of a nonempty relative interior of the primal feasible set;^{4.8}

$$b \in \text{int } \mathcal{K} \Leftrightarrow \langle y, b \rangle > 0 \quad \forall y \in \mathcal{K}^*, \quad y \neq \mathbf{0} \Leftrightarrow \mathcal{A} \cap \text{int } \mathbb{S}_+^n \neq \emptyset \quad (607)$$

A positive definite Farkas' lemma can easily be constructed from this membership relation (296) and these proper cones \mathcal{K} (344) and \mathcal{K}^* (349):

4.2.1.1.2 Lemma. *Positive definite Farkas' lemma.*

Given a linearly independent set $\{A_i \in \mathbb{S}^n, i=1 \dots m\}$ and a vector $b = [b_i] \in \mathbb{R}^m$, define the affine subset

$$\mathcal{A} = \{X \in \mathbb{S}^n \mid \langle A_i, X \rangle = b_i, i=1 \dots m\} \quad (595)$$

Primal feasible set relative interior $\mathcal{A} \cap \text{int } \mathbb{S}_+^n$ is nonempty if and only if $y^T b > 0$ holds for each and every vector $y = [y_i] \neq \mathbf{0}$ such that $\sum_{i=1}^m y_i A_i \succeq 0$.

Equivalently, primal feasible set relative interior $\mathcal{A} \cap \text{int } \mathbb{S}_+^n$ is nonempty if and only if $y^T b > 0$ holds for each and every norm-1 vector $\|y\| = 1$ such that $\sum_{i=1}^m y_i A_i \succeq 0$. \diamond

^{4.7}Slater's sufficient condition is satisfied whenever any primal *strictly feasible* point exists; *id est*, any point feasible with the affine equality (or affine inequality) constraint functions and relatively interior to convex cone \mathcal{K} . [293, §6.6] [36, p.325] If cone \mathcal{K} were polyhedral, then Slater's condition is satisfied when any feasible point exists relatively interior to \mathcal{K} or on its relative boundary. [53, §5.2.3]

^{4.8}Detection of $\mathcal{A} \cap \text{int } \mathbb{S}_+^n$ by examining \mathcal{K} interior is a trick need not be lost.

4.2.1.1.3 Example. “New” Farkas’ lemma.

In 1995, Lasserre [204, §III] presented an example originally offered by Ben-Israel in 1969 [30, p.378] as evidence of failure in *semidefinite Farkas’ Lemma* 4.2.1.1.1:

$$A \triangleq \begin{bmatrix} \text{svec}(A_1)^T \\ \text{svec}(A_2)^T \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad (608)$$

The intersection $\mathcal{A} \cap \mathbb{S}_+^n$ is practically empty because the solution set

$$\{X \succeq 0 \mid A \text{ svec } X = b\} = \left\{ \begin{bmatrix} \alpha & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & 0 \end{bmatrix} \succeq 0 \mid \alpha \in \mathbb{R} \right\} \quad (609)$$

is positive semidefinite only asymptotically ($\alpha \rightarrow \infty$). Yet the dual system $\sum_{i=1}^m y_i A_i \succeq 0 \Rightarrow y^T b \geq 0$ indicates nonempty intersection; *videlicet*, for $\|y\| = 1$

$$y_1 \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & 0 \end{bmatrix} + y_2 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \succeq 0 \Leftrightarrow y = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \Rightarrow y^T b = 0 \quad (610)$$

On the other hand, *positive definite Farkas’ Lemma* 4.2.1.1.2 shows $\mathcal{A} \cap \text{int } \mathbb{S}_+^n$ is empty; what we need to know for semidefinite programming.

Based on Ben-Israel’s example, Lasserre suggested addition of another condition to *semidefinite Farkas’ Lemma* 4.2.1.1.1 to make a “new” lemma. Ye recommends *positive definite Farkas’ Lemma* 4.2.1.1.2 instead; which is simpler and obviates Lasserre’s proposed additional condition. \square

4.2.1.2 Theorem of the alternative for semidefinite programming

Because these Farkas’ lemmas follow from membership relations, we may construct alternative systems from them. Applying the method of §2.13.2.1.1, then from *positive definite Farkas’ lemma*, for example, we get

$$\begin{aligned} & \mathcal{A} \cap \text{int } \mathbb{S}_+^n \neq \emptyset \\ & \text{or in the alternative} \\ & y^T b \leq 0, \quad \sum_{i=1}^m y_i A_i \succeq 0, \quad y \neq \mathbf{0} \end{aligned} \quad (611)$$

Any single vector y satisfying the alternative certifies $\mathcal{A} \cap \text{int } \mathbb{S}_+^n$ is empty. Such a vector can be found as a solution to another semidefinite program: for linearly independent set $\{A_i \in \mathbb{S}^n, i=1 \dots m\}$

$$\begin{aligned} & \underset{y}{\text{minimize}} && y^T b \\ & \text{subject to} && \sum_{i=1}^m y_i A_i \succeq 0 \\ & && \|y\|^2 \leq 1 \end{aligned} \tag{612}$$

If an optimal vector $y^* \neq \mathbf{0}$ can be found such that $y^{*T} b \leq 0$, then relative interior of the primal feasible set $\mathcal{A} \cap \text{int } \mathbb{S}_+^n$ from (604) is empty.

4.2.1.3 Boundary-membership criterion

(confer (606) (607)) From boundary-membership relation (300) for proper cones and from *linear matrix inequality* cones \mathcal{K} (344) and \mathcal{K}^* (349)

$$b \in \partial\mathcal{K} \Leftrightarrow \exists y \neq \mathbf{0} \ni \langle y, b \rangle = 0, y \in \mathcal{K}^*, b \in \mathcal{K} \Leftrightarrow \partial\mathbb{S}_+^n \supset \mathcal{A} \cap \mathbb{S}_+^n \neq \emptyset \tag{613}$$

Whether vector $b \in \partial\mathcal{K}$ belongs to cone \mathcal{K} boundary, that is a determination we can indeed make; one that is certainly expressible as a feasibility problem: assuming $b \in \mathcal{K}$ (605) given linearly independent set $\{A_i \in \mathbb{S}^n, i=1 \dots m\}$ ^{4.9}

$$\begin{aligned} & \text{find} && y \neq \mathbf{0} \\ & \text{subject to} && y^T b = 0 \\ & && \sum_{i=1}^m y_i A_i \succeq 0 \end{aligned} \tag{614}$$

Any such feasible nonzero vector y certifies that affine subset \mathcal{A} (595) intersects the positive semidefinite cone \mathbb{S}_+^n only on its boundary; in other words, nonempty feasible set $\mathcal{A} \cap \mathbb{S}_+^n$ belongs to the positive semidefinite cone boundary $\partial\mathbb{S}_+^n$.

^{4.9}From the results of Example 2.13.5.1.1, vector b on the boundary of \mathcal{K} cannot be detected simply by looking for 0 eigenvalues in matrix X .

4.2.2 Duals

The dual objective function from (592D) evaluated at any feasible point represents a lower bound on the primal optimal objective value from (592P). We can see this by direct substitution: Assume the feasible sets $\mathcal{A} \cap \mathbb{S}_+^n$ and \mathcal{C}^* are nonempty. Then it is always true:

$$\begin{aligned} \langle C, X \rangle &\geq \langle b, y \rangle \\ \left\langle \sum_i y_i A_i + S, X \right\rangle &\geq [\langle A_1, X \rangle \cdots \langle A_m, X \rangle] y \\ \langle S, X \rangle &\geq 0 \end{aligned} \quad (615)$$

The converse also follows because

$$X \succeq 0, S \succeq 0 \Rightarrow \langle S, X \rangle \geq 0 \quad (1395)$$

Optimal value of the dual objective thus represents the greatest lower bound on the primal. This fact is known as the *weak duality theorem* for semidefinite programming, [350, §1.3.8] and can be used to detect convergence in any primal/dual numerical method of solution.

4.2.3 Optimality conditions

When any primal feasible point exists relatively interior to $\mathcal{A} \cap \mathbb{S}_+^n$ in \mathbb{S}^n , or when any dual feasible point exists relatively interior to \mathcal{C}^* in $\mathbb{S}^n \times \mathbb{R}^m$, then by Slater's sufficient condition (p.260) these two problems (592P) (592D) become *strong duals*. In other words, the primal optimal objective value becomes equivalent to the dual optimal objective value: there is no duality gap (Figure 54); *id est*, if $\exists X \in \mathcal{A} \cap \text{int } \mathbb{S}_+^n$ or $\exists S, y \in \text{rel int } \mathcal{C}^*$ then

$$\begin{aligned} \langle C, X^* \rangle &= \langle b, y^* \rangle \\ \left\langle \sum_i y_i^* A_i + S^*, X^* \right\rangle &= [\langle A_1, X^* \rangle \cdots \langle A_m, X^* \rangle] y^* \\ \langle S^*, X^* \rangle &= 0 \end{aligned} \quad (616)$$

where S^*, y^* denote a dual optimal solution.^{4.10} We summarize this:

^{4.10}Optimality condition $\langle S^*, X^* \rangle = 0$ is called a *complementary slackness condition*, in keeping with linear programming tradition [82], that forbids dual inequalities in (592) to simultaneously hold strictly. [270, §4]

4.2.3.0.1 Corollary. *Optimality and strong duality.* [319, §3.1] [350, §1.3.8] For semidefinite programs (592P) and (592D), assume primal and dual feasible sets $\mathcal{A} \cap \mathbb{S}_+^n \subset \mathbb{S}^n$ and $\mathcal{C}^* \subset \mathbb{S}^n \times \mathbb{R}^m$ (604) are nonempty. Then

- X^* is optimal for (P)
- S^*, y^* are optimal for (D)
- the duality gap $\langle C, X^* \rangle - \langle b, y^* \rangle$ is 0

if and only if

$$\text{i) } \exists X \in \mathcal{A} \cap \text{int } \mathbb{S}_+^n \quad \text{or} \quad \exists S, y \in \text{rel int } \mathcal{C}^*$$

and

$$\text{ii) } \langle S^*, X^* \rangle = 0 \quad \diamond$$

For symmetric positive semidefinite matrices, requirement **ii** is equivalent to the *complementarity* (§A.7.4)

$$\langle S^*, X^* \rangle = 0 \quad \Leftrightarrow \quad S^* X^* = X^* S^* = \mathbf{0} \quad (617)$$

Commutativity of diagonalizable matrices is a necessary and sufficient condition [180, §1.3.12] for these two optimal symmetric matrices to be simultaneously diagonalizable. Therefore

$$\text{rank } X^* + \text{rank } S^* \leq n \quad (618)$$

Proof. To see that, the product of symmetric optimal matrices $X^*, S^* \in \mathbb{S}^n$ must itself be symmetric because of commutativity. (1387) The symmetric product has diagonalization [10, cor.2.11]

$$S^* X^* = X^* S^* = Q \Lambda_{S^*} \Lambda_{X^*} Q^T = \mathbf{0} \quad \Leftrightarrow \quad \Lambda_{X^*} \Lambda_{S^*} = \mathbf{0} \quad (619)$$

where Q is an orthogonal matrix. The product of the nonnegative diagonal Λ matrices can be $\mathbf{0}$ if their main diagonal zeros are complementary or coincide. Due only to symmetry, $\text{rank } X^* = \text{rank } \Lambda_{X^*}$ and $\text{rank } S^* = \text{rank } \Lambda_{S^*}$ for these optimal primal and dual solutions. (1373) So, because of the complementarity, the total number of nonzero diagonal entries from both Λ cannot exceed n . \blacklozenge

When equality is attained in (618)

$$\text{rank } X^* + \text{rank } S^* = n \quad (620)$$

there are no coinciding main diagonal zeros in $\Lambda_{X^*}\Lambda_{S^*}$, and so we have what is called *strict complementarity*.^{4.11} Logically it follows that a necessary and sufficient condition for strict complementarity of an optimal primal and dual solution is

$$X^* + S^* \succ 0 \quad (621)$$

4.2.3.1 solving primal problem via dual

The beauty of Corollary 4.2.3.0.1 is its conjugacy; *id est*, one can solve either the primal or dual problem in (592) and then find a solution to the other via the optimality conditions. When a dual optimal solution is known, for example, a primal optimal solution belongs to the hyperplane $\{X \mid \langle S^*, X \rangle = 0\}$.

4.2.3.1.1 Example. *Minimum cardinality Boolean.* [81] [31, §4.3.4] [307] (confer Example 4.5.1.2.1) Consider finding a *minimum cardinality* Boolean solution x to the classic linear algebra problem $Ax = b$ given noiseless data $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$;

$$\begin{aligned} & \underset{x}{\text{minimize}} && \|x\|_0 \\ & \text{subject to} && Ax = b \\ & && x_i \in \{0, 1\}, \quad i = 1 \dots n \end{aligned} \quad (622)$$

where $\|x\|_0$ denotes cardinality of vector x (a.k.a, *0-norm*; not a convex function).

A minimum cardinality solution answers the question: “Which fewest linear combination of columns in A constructs vector b ?” *Cardinality problems* have extraordinarily wide appeal, arising in many fields of science and across many disciplines. [282] [189] [149] [150] Yet designing an efficient algorithm to optimize cardinality has proved difficult. In this example, we also constrain the variable to be Boolean. The Boolean constraint forces

^{4.11} distinct from maximal complementarity (§4.1.1.1).

an identical solution were the norm in problem (622) instead the 1-norm or 2-norm; *id est*, the two problems

$$(622) \quad \begin{array}{ll} \underset{x}{\text{minimize}} & \|x\|_0 \\ \text{subject to} & Ax = b \\ & x_i \in \{0, 1\}, \quad i=1 \dots n \end{array} = \begin{array}{ll} \underset{x}{\text{minimize}} & \|x\|_1 \\ \text{subject to} & Ax = b \\ & x_i \in \{0, 1\}, \quad i=1 \dots n \end{array} \quad (623)$$

are the same. The Boolean constraint makes the 1-norm problem nonconvex.

Given data^{4.12}

$$A = \begin{bmatrix} -1 & 1 & 8 & 1 & 1 & 0 \\ -3 & 2 & 8 & \frac{1}{2} & \frac{1}{3} & \frac{1}{2} - \frac{1}{3} \\ -9 & 4 & 8 & \frac{1}{4} & \frac{1}{9} & \frac{1}{4} - \frac{1}{9} \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ \frac{1}{2} \\ \frac{1}{4} \end{bmatrix} \quad (624)$$

the obvious and desired solution to the problem posed,

$$x^* = e_4 \in \mathbb{R}^6 \quad (625)$$

has norm $\|x^*\|_2 = 1$ and minimum cardinality; the minimum number of nonzero entries in vector x . The MATLAB [backslash command](#) `x=A\b`, for example, finds

$$x_M = \begin{bmatrix} \frac{2}{128} \\ 0 \\ \frac{5}{128} \\ 0 \\ \frac{90}{128} \\ 0 \end{bmatrix} \quad (626)$$

having norm $\|x_M\|_2 = 0.7044$. Coincidentally, x_M is a 1-norm solution; *id est*, an optimal solution to

$$\begin{array}{ll} \underset{x}{\text{minimize}} & \|x\|_1 \\ \text{subject to} & Ax = b \end{array} \quad (468)$$

^{4.12}This particular matrix A is full-rank having three-dimensional nullspace (but the columns are not conically independent (§2.10)).

The pseudoinverse solution (rounded)

$$x_{\mathbf{P}} = A^{\dagger}b = \begin{bmatrix} -0.0456 \\ -0.1881 \\ 0.0623 \\ 0.2668 \\ 0.3770 \\ -0.1102 \end{bmatrix} \quad (627)$$

has least norm $\|x_{\mathbf{P}}\|_2 = 0.5165$; *id est*, the optimal solution to (§E.0.1.0.1)

$$\begin{aligned} & \underset{x}{\text{minimize}} && \|x\|_2 \\ & \text{subject to} && Ax = b \end{aligned} \quad (628)$$

Certainly none of the traditional methods provide $x^* = e_4$ (625) because, and in general, for $Ax = b$

$$\|\arg \inf \|x\|_2\|_2 \leq \|\arg \inf \|x\|_1\|_2 \leq \|\arg \inf \|x\|_0\|_2 \quad (629)$$

We can reformulate this minimum cardinality Boolean problem (622) as a semidefinite program: First transform the variable

$$x \triangleq (\hat{x} + \mathbf{1})\frac{1}{2} \quad (630)$$

so $\hat{x}_i \in \{-1, 1\}$; equivalently,

$$\begin{aligned} & \underset{\hat{x}}{\text{minimize}} && \|(\hat{x} + \mathbf{1})\frac{1}{2}\|_0 \\ & \text{subject to} && A(\hat{x} + \mathbf{1})\frac{1}{2} = b \\ & && \delta(\hat{x}\hat{x}^T) = \mathbf{1} \end{aligned} \quad (631)$$

where δ is the main-diagonal linear operator (§A.1). By assigning (§B.1)

$$G = \begin{bmatrix} \hat{x} \\ 1 \end{bmatrix} [\hat{x}^T \ 1] = \begin{bmatrix} X & \hat{x} \\ \hat{x}^T & 1 \end{bmatrix} \triangleq \begin{bmatrix} \hat{x}\hat{x}^T & \hat{x} \\ \hat{x}^T & 1 \end{bmatrix} \in \mathbb{S}^{n+1} \quad (632)$$

problem (631) becomes equivalent to: (Theorem A.3.1.0.7)

$$\begin{aligned}
 & \underset{X \in \mathbb{S}^n, \hat{x} \in \mathbb{R}^n}{\text{minimize}} && \mathbf{1}^T \hat{x} \\
 & \text{subject to} && A(\hat{x} + \mathbf{1})_{\frac{1}{2}} = b \\
 & && G = \begin{bmatrix} X & \hat{x} \\ \hat{x}^T & 1 \end{bmatrix} \\
 & && \delta(X) = \mathbf{1} \\
 & && (G \succeq 0) \\
 & && \text{rank } G = 1
 \end{aligned} \tag{633}$$

where solution is confined to rank-1 vertices of the *elliptope* in \mathbb{S}^{n+1} (§5.9.1.0.1) by the rank constraint, the positive semidefiniteness, and the equality constraints $\delta(X) = \mathbf{1}$. The rank constraint makes this problem nonconvex; by removing it^{4.13} we get the semidefinite program

$$\begin{aligned}
 & \underset{X \in \mathbb{S}^n, \hat{x} \in \mathbb{R}^n}{\text{minimize}} && \mathbf{1}^T \hat{x} \\
 & \text{subject to} && A(\hat{x} + \mathbf{1})_{\frac{1}{2}} = b \\
 & && G = \begin{bmatrix} X & \hat{x} \\ \hat{x}^T & 1 \end{bmatrix} \succeq 0 \\
 & && \delta(X) = \mathbf{1}
 \end{aligned} \tag{634}$$

whose optimal solution x^* (630) is identical to that of minimum cardinality Boolean problem (622) if and only if $\text{rank } G^* = 1$. Hope^{4.14} of acquiring a rank-1 solution is not ill-founded because 2^n elliptope vertices have rank 1, and we are minimizing an affine function on a subset of the elliptope (Figure 109) containing rank-1 vertices; *id est*, by assumption that the feasible set of minimum cardinality Boolean problem (622) is nonempty, a desired solution resides on the elliptope relative boundary at a rank-1 vertex.^{4.15}

^{4.13}Relaxed problem (634) can also be derived via Lagrange duality; it is a dual of a dual program [*sic*] to (633). [268] [53, §5, exer.5.39] [336, §IV] [130, §11.3.4] The relaxed problem must therefore be convex having a larger feasible set; its optimal objective value represents a generally *loose* lower bound (1575) on the optimal objective of problem (633).

^{4.14}A more deterministic approach to constraining rank and cardinality is developed in §4.6.0.0.9.

^{4.15}Confinement to the elliptope can be regarded as a kind of normalization akin to matrix A column normalization suggested in [103] and explored in Example 4.2.3.1.2.

For the data given in (624), our semidefinite program solver [343] [344] (accurate in solution to approximately 1E-8)^{4.16} finds optimal solution to (634)

$$\text{round}(G^*) = \begin{bmatrix} 1 & 1 & 1 & -1 & 1 & 1 & -1 \\ 1 & 1 & 1 & -1 & 1 & 1 & -1 \\ 1 & 1 & 1 & -1 & 1 & 1 & -1 \\ -1 & -1 & -1 & 1 & -1 & -1 & 1 \\ 1 & 1 & 1 & -1 & 1 & 1 & -1 \\ 1 & 1 & 1 & -1 & 1 & 1 & -1 \\ -1 & -1 & -1 & 1 & -1 & -1 & 1 \end{bmatrix} \quad (635)$$

near a rank-1 vertex of the ellipsope in \mathbb{S}^{n+1} (§5.9.1.0.2); its sorted eigenvalues,

$$\lambda(G^*) = \begin{bmatrix} 6.99999977799099 \\ 0.00000022687241 \\ 0.00000002250296 \\ 0.00000000262974 \\ -0.00000000999738 \\ -0.00000000999875 \\ -0.00000001000000 \end{bmatrix} \quad (636)$$

The negative eigenvalues are undoubtedly finite-precision effects. Because the largest eigenvalue predominates by many orders of magnitude, we can expect to find a good approximation to a minimum cardinality Boolean solution by truncating all smaller eigenvalues. By so doing we find, indeed,

$$x^* = \text{round} \left(\begin{bmatrix} 0.00000000127947 \\ 0.00000000527369 \\ 0.00000000181001 \\ 0.99999997469044 \\ 0.00000001408950 \\ 0.00000000482903 \end{bmatrix} \right) = e_4 \quad (637)$$

the desired result (625). □

^{4.16}A typically ignored limitation of interior-point methods of solution is their relative accuracy of only about 1E-8 on a machine using 64-bit (*double precision*) floating-point arithmetic; *id est*, optimal solution x^* cannot be more accurate than square root of machine epsilon ($\epsilon = 2.2204\text{E-}16$). Duality gap is not a good measure of solution accuracy.

4.2.3.1.2 Example. *Optimization on ellipsope versus 1-norm polyhedron for minimum cardinality Boolean Example 4.2.3.1.1.*

A minimum cardinality problem is typically formulated via, what is by now, the standard practice [103] [59, §3.2, §3.4] of column normalization applied to a 1-norm problem surrogate like (468). Suppose we define a diagonal matrix

$$\Lambda \triangleq \begin{bmatrix} \|A(:, 1)\|_2 & & & & & \mathbf{0} \\ & \|A(:, 2)\|_2 & & & & \\ & & \ddots & & & \\ & & & \ddots & & \\ \mathbf{0} & & & & & \|A(:, 6)\|_2 \end{bmatrix} \in \mathbb{S}^6 \quad (638)$$

used to normalize the columns (assumed nonzero) of given noiseless data matrix A . Then approximate the minimum cardinality Boolean problem

$$\begin{aligned} & \underset{x}{\text{minimize}} && \|x\|_0 \\ & \text{subject to} && Ax = b \\ & && x_i \in \{0, 1\}, \quad i = 1 \dots n \end{aligned} \quad (622)$$

as

$$\begin{aligned} & \underset{\tilde{y}}{\text{minimize}} && \|\tilde{y}\|_1 \\ & \text{subject to} && A\Lambda^{-1}\tilde{y} = b \\ & && \mathbf{1} \succeq \Lambda^{-1}\tilde{y} \succeq 0 \end{aligned} \quad (639)$$

where optimal solution

$$y^* = \text{round}(\Lambda^{-1}\tilde{y}^*) \quad (640)$$

The inequality in (639) relaxes Boolean constraint $y_i \in \{0, 1\}$ from (622); bounding any solution y^* to a nonnegative unit hypercube whose vertices are binary numbers. Convex problem (639) is justified by the *convex envelope*

$$\text{cenv } \|x\|_0 \text{ on } \{x \in \mathbb{R}^n \mid \|x\|_\infty \leq \kappa\} = \frac{1}{\kappa} \|x\|_1 \quad (1280)$$

Donoho concurs with this particular formulation, equivalently expressible as a linear program via (464).

Approximation (639) is therefore equivalent to minimization of an affine function (§3.1.3) on a bounded polyhedron, whereas semidefinite program

$$\begin{aligned} & \underset{X \in \mathbb{S}^n, \hat{x} \in \mathbb{R}^n}{\text{minimize}} && \mathbf{1}^T \hat{x} \\ & \text{subject to} && A(\hat{x} + \mathbf{1})^{\frac{1}{2}} = b \\ & && G = \begin{bmatrix} X & \hat{x} \\ \hat{x}^T & 1 \end{bmatrix} \succeq 0 \\ & && \delta(X) = \mathbf{1} \end{aligned} \quad (634)$$

minimizes an affine function on an intersection of the ellipsope with hyperplanes. Although the same Boolean solution is obtained from this approximation (639) as compared with semidefinite program (634), when given that particular data from Example 4.2.3.1.1, Singer confides a counter-example: Instead, given data

$$A = \begin{bmatrix} 1 & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & \frac{1}{\sqrt{2}} \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad (641)$$

then solving approximation (639) yields

$$y^* = \text{round} \left(\begin{bmatrix} 1 - \frac{1}{\sqrt{2}} \\ 1 - \frac{1}{\sqrt{2}} \\ 1 \end{bmatrix} \right) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \quad (642)$$

(infeasible, with or without rounding, with respect to original problem (622)) whereas solving semidefinite program (634) produces

$$\text{round}(G^*) = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & 1 & -1 & 1 \\ -1 & -1 & 1 & -1 \\ 1 & 1 & -1 & 1 \end{bmatrix} \quad (643)$$

with sorted eigenvalues

$$\lambda(G^*) = \begin{bmatrix} 3.99999965057264 \\ 0.00000035942736 \\ -0.00000000000000 \\ -0.00000001000000 \end{bmatrix} \quad (644)$$

Truncating all but the largest eigenvalue, from (630) we obtain (*confer* y^*)

$$x^* = \text{round} \left(\begin{bmatrix} 0.9999999625299 \\ 0.9999999625299 \\ 0.00000001434518 \end{bmatrix} \right) = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad (645)$$

the desired minimum cardinality Boolean result. \square

4.2.3.1.3 Exercise. *Minimum cardinality Boolean art.*

Assess general performance of standard-practice approximation (639) as compared with the proposed semidefinite program (634). \blacktriangledown

4.3 Rank reduction

...it is not clear generally how to predict $\text{rank } X^*$ or $\text{rank } S^*$ before solving the SDP problem.

–Farid Alizadeh (1995) [10, p.22]

The premise of rank reduction in semidefinite programming is: an optimal solution found does not satisfy Barvinok's upper bound (246) on rank. The particular numerical algorithm solving a semidefinite program may have instead returned a high-rank optimal solution (§4.1.1.1; e.g., (603)) when a lower-rank optimal solution was expected.

4.3.1 Posit a perturbation of X^*

Recall from §4.1.1.2, there is an extreme point of $\mathcal{A} \cap \mathbb{S}_+^n$ (595) satisfying upper bound (246) on rank. [23, §2.2] It is therefore sufficient to locate an extreme point of the intersection whose primal objective value (592P) is optimal.^{4.17} [96, §31.5.3] [211, §2.4] [212] [6, §3] [257]

Consider again the affine subset

$$\mathcal{A} = \{X \in \mathbb{S}^n \mid A \text{ svec } X = b\} \quad (595)$$

where for $A_i \in \mathbb{S}^n$

$$A \triangleq \begin{bmatrix} \text{svec}(A_1)^T \\ \vdots \\ \text{svec}(A_m)^T \end{bmatrix} \in \mathbb{R}^{m \times n(n+1)/2} \quad (593)$$

Given any optimal solution X^* to

$$\begin{aligned} & \underset{X \in \mathbb{S}^n}{\text{minimize}} && \langle C, X \rangle \\ & \text{subject to} && X \in \mathcal{A} \cap \mathbb{S}_+^n \end{aligned} \quad (592)(P)$$

^{4.17} There is no known construction for Barvinok's tighter result (251). –Monique Laurent

whose rank does not satisfy upper bound (246), we posit existence of a set of perturbations

$$\{t_j B_j \mid t_j \in \mathbb{R}, B_j \in \mathbb{S}^n, j=1 \dots n\} \quad (646)$$

such that, for some $0 \leq i \leq n$ and scalars $\{t_j, j=1 \dots i\}$,

$$X^* + \sum_{j=1}^i t_j B_j \quad (647)$$

becomes an extreme point of $\mathcal{A} \cap \mathbb{S}_+^n$ and remains an optimal solution of (592P). Membership of (647) to affine subset \mathcal{A} is secured for the i^{th} perturbation by demanding

$$\langle B_i, A_j \rangle = 0, \quad j=1 \dots m \quad (648)$$

while membership to the positive semidefinite cone \mathbb{S}_+^n is insured by small perturbation (657). In this manner feasibility is insured. Optimality is proved in §4.3.3.

The following simple algorithm has very low computational intensity and locates an optimal extreme point, assuming a nontrivial solution:

4.3.1.0.1 Procedure. *Rank reduction.* ([Wikimization](#))

initialize: $B_i = \mathbf{0} \quad \forall i$

for iteration $i=1 \dots n$

{

1. compute a nonzero perturbation matrix B_i of $X^* + \sum_{j=1}^{i-1} t_j B_j$

2. maximize t_i

subject to $X^* + \sum_{j=1}^i t_j B_j \in \mathbb{S}_+^n$

}

¶

A rank-reduced optimal solution is then

$$X^* \leftarrow X^* + \sum_{j=1}^i t_j B_j \quad (649)$$

4.3.2 Perturbation form

The perturbations are independent of constants $C \in \mathbb{S}^n$ and $b \in \mathbb{R}^m$ in primal and dual programs (592). Numerical accuracy of any rank-reduced result, found by perturbation of an initial optimal solution X^* , is therefore quite dependent upon initial accuracy of X^* .

4.3.2.0.1 Definition. *Matrix step function.* (confer §A.6.5.0.1)
Define the signum-like quasiconcave real function $\psi : \mathbb{S}^n \rightarrow \mathbb{R}$

$$\psi(Z) \triangleq \begin{cases} 1, & Z \succeq 0 \\ -1, & \text{otherwise} \end{cases} \quad (650)$$

The value -1 is taken for indefinite or nonzero negative semidefinite argument. \triangle

Deza & Laurent [96, §31.5.3] prove: every perturbation matrix B_i , $i=1 \dots n$, is of the form

$$B_i = -\psi(Z_i) R_i Z_i R_i^T \in \mathbb{S}^n \quad (651)$$

where

$$X^* \triangleq R_1 R_1^T, \quad X^* + \sum_{j=1}^{i-1} t_j B_j \triangleq R_i R_i^T \in \mathbb{S}^n \quad (652)$$

where the t_j are scalars and $R_i \in \mathbb{R}^{n \times \rho}$ is full-rank and skinny where

$$\rho \triangleq \text{rank} \left(X^* + \sum_{j=1}^{i-1} t_j B_j \right) \quad (653)$$

and where matrix $Z_i \in \mathbb{S}^\rho$ is found at each iteration i by solving a very simple feasibility problem: [4.18](#)

$$\begin{aligned} & \text{find } Z_i \in \mathbb{S}^\rho \\ & \text{subject to } \langle Z_i, R_i^T A_j R_i \rangle = 0, \quad j=1 \dots m \end{aligned} \quad (654)$$

Were there a sparsity pattern common to each member of the set $\{R_i^T A_j R_i \in \mathbb{S}^\rho, j=1 \dots m\}$, then a good choice for Z_i has 1 in each entry corresponding to a 0 in the pattern; *id est*, a sparsity pattern complement. At iteration i

$$X^* + \sum_{j=1}^{i-1} t_j B_j + t_i B_i = R_i (I - t_i \psi(Z_i) Z_i) R_i^T \quad (655)$$

By fact ([1363](#)), therefore

$$X^* + \sum_{j=1}^{i-1} t_j B_j + t_i B_i \succeq 0 \Leftrightarrow \mathbf{1} - t_i \psi(Z_i) \lambda(Z_i) \succeq 0 \quad (656)$$

where $\lambda(Z_i) \in \mathbb{R}^\rho$ denotes the eigenvalues of Z_i .

Maximization of each t_i in step 2 of the Procedure reduces rank of (655) and locates a new point on the boundary $\partial(\mathcal{A} \cap \mathbb{S}_+^n)$. [4.19](#) Maximization of t_i thereby has closed form;

[4.18](#) A simple method of solution is closed-form projection of a random nonzero point on that proper subspace of isometrically isomorphic $\mathbb{R}^{\rho(\rho+1)/2}$ specified by the constraints. (§[E.5.0.0.6](#)) Such a solution is nontrivial assuming the specified intersection of hyperplanes is not the origin; guaranteed by $\rho(\rho+1)/2 > m$. Indeed, this geometric intuition about forming the perturbation is what bounds any solution's rank from below; m is fixed by the number of equality constraints in ([592P](#)) while rank ρ decreases with each iteration i . Otherwise, we might iterate indefinitely.

[4.19](#) This holds because rank of a positive semidefinite matrix in \mathbb{S}^n is diminished below n by the number of its 0 eigenvalues ([1373](#)), and because a positive semidefinite matrix having one or more 0 eigenvalues corresponds to a point on the PSD cone boundary ([176](#)). Necessity and sufficiency are due to the facts: R_i can be completed to a nonsingular matrix (§[A.3.1.0.5](#)), and $I - t_i \psi(Z_i) Z_i$ can be padded with zeros while maintaining equivalence in ([655](#)).

$$(t_i^*)^{-1} = \max \{ \psi(Z_i) \lambda(Z_i)_j, j=1 \dots \rho \} \quad (657)$$

When Z_i is indefinite, the direction of perturbation (determined by $\psi(Z_i)$) is arbitrary. We may take an early exit from the Procedure were Z_i to become $\mathbf{0}$ or were

$$\text{rank} [\text{svec } R_i^T A_1 R_i \quad \text{svec } R_i^T A_2 R_i \quad \cdots \quad \text{svec } R_i^T A_m R_i] = \rho(\rho + 1)/2 \quad (658)$$

which characterizes the rank ρ of any [sic] extreme point in $\mathcal{A} \cap \mathbb{S}_+^n$. [211, §2.4] [212]

Proof. Assuming the form of every perturbation matrix is indeed (651), then by (654)

$$\text{svec } Z_i \perp [\text{svec}(R_i^T A_1 R_i) \quad \text{svec}(R_i^T A_2 R_i) \quad \cdots \quad \text{svec}(R_i^T A_m R_i)] \quad (659)$$

By orthogonal complement we have

$$\begin{aligned} & \text{rank} [\text{svec}(R_i^T A_1 R_i) \quad \cdots \quad \text{svec}(R_i^T A_m R_i)]^\perp \\ & + \text{rank} [\text{svec}(R_i^T A_1 R_i) \quad \cdots \quad \text{svec}(R_i^T A_m R_i)] = \rho(\rho + 1)/2 \end{aligned} \quad (660)$$

When Z_i can only be $\mathbf{0}$, then the perturbation is null because an extreme point has been found; thus

$$[\text{svec}(R_i^T A_1 R_i) \quad \cdots \quad \text{svec}(R_i^T A_m R_i)]^\perp = \mathbf{0} \quad (661)$$

from which the stated result (658) directly follows. \blacklozenge

4.3.3 Optimality of perturbed X^*

We show that the optimal objective value is unaltered by perturbation (651); *id est*,

$$\langle C, X^* + \sum_{j=1}^i t_j B_j \rangle = \langle C, X^* \rangle \quad (662)$$

Proof. From Corollary 4.2.3.0.1 we have the necessary and sufficient relationship between optimal primal and dual solutions under the assumption of existence of a relatively interior feasible point:

$$S^* X^* = S^* R_1 R_1^T = X^* S^* = R_1 R_1^T S^* = \mathbf{0} \quad (663)$$

This means $\mathcal{R}(R_1) \subseteq \mathcal{N}(S^*)$ and $\mathcal{R}(S^*) \subseteq \mathcal{N}(R_1^T)$. From (652) and (655) we get the sequence:

$$\begin{aligned} X^* &= R_1 R_1^T \\ X^* + t_1 B_1 &= R_2 R_2^T = R_1 (I - t_1 \psi(Z_1) Z_1) R_1^T \\ X^* + t_1 B_1 + t_2 B_2 &= R_3 R_3^T = R_2 (I - t_2 \psi(Z_2) Z_2) R_2^T = R_1 (I - t_1 \psi(Z_1) Z_1) (I - t_2 \psi(Z_2) Z_2) R_1^T \\ &\vdots \\ X^* + \sum_{j=1}^i t_j B_j &= R_1 \left(\prod_{j=1}^i (I - t_j \psi(Z_j) Z_j) \right) R_1^T \end{aligned} \quad (664)$$

Substituting $C = \text{svec}^{-1}(A^T y^*) + S^*$ from (592),

$$\begin{aligned} \langle C, X^* + \sum_{j=1}^i t_j B_j \rangle &= \left\langle \text{svec}^{-1}(A^T y^*) + S^*, R_1 \left(\prod_{j=1}^i (I - t_j \psi(Z_j) Z_j) \right) R_1^T \right\rangle \\ &= \left\langle \sum_{k=1}^m y_k^* A_k, X^* + \sum_{j=1}^i t_j B_j \right\rangle \\ &= \left\langle \sum_{k=1}^m y_k^* A_k + S^*, X^* \right\rangle = \langle C, X^* \rangle \end{aligned} \quad (665)$$

because $\langle B_i, A_j \rangle = 0 \quad \forall i, j$ by design (648). \blacklozenge

4.3.3.0.1 Example. $A\delta(X) = b$.

This academic example demonstrates that a solution found by rank reduction can certainly have rank less than Barvinok's upper bound (246): Assume a given vector $b \in \mathbb{R}^m$ belongs to the conic hull of columns of a given matrix $A \in \mathbb{R}^{m \times n}$;

$$A = \begin{bmatrix} -1 & 1 & 8 & 1 & 1 \\ -3 & 2 & 8 & \frac{1}{2} & \frac{1}{3} \\ -9 & 4 & 8 & \frac{1}{4} & \frac{1}{9} \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ \frac{1}{2} \\ \frac{1}{4} \end{bmatrix} \quad (666)$$

Consider the convex optimization problem

$$\begin{aligned} & \underset{X \in \mathbb{S}^5}{\text{minimize}} && \text{tr } X \\ & \text{subject to} && X \succeq 0 \\ & && A\delta(X) = b \end{aligned} \quad (667)$$

that minimizes the 1-norm of the main diagonal; *id est*, problem (667) is the same as

$$\begin{aligned} & \underset{X \in \mathbb{S}^5}{\text{minimize}} && \|\delta(X)\|_1 \\ & \text{subject to} && X \succeq 0 \\ & && A\delta(X) = b \end{aligned} \quad (668)$$

that finds a solution to $A\delta(X) = b$. Rank-3 solution $X^* = \delta(x_{\mathbf{M}})$ is optimal, where (*confer* (626))

$$x_{\mathbf{M}} = \begin{bmatrix} \frac{2}{128} \\ 0 \\ \frac{5}{128} \\ 0 \\ \frac{90}{128} \end{bmatrix} \quad (669)$$

Yet upper bound (246) predicts existence of at most a

$$\text{rank-}\left(\left\lfloor \frac{\sqrt{8m+1}-1}{2} \right\rfloor = 2\right) \quad (670)$$

feasible solution from $m=3$ equality constraints. To find a lower rank ρ optimal solution to (667) (barring combinatorics), we invoke Procedure 4.3.1.0.1:

Initialize:

$$\left\{ \begin{array}{l} C=I, \quad \rho=3, \quad A_j \triangleq \delta(A(j, :)), \quad j=1, 2, 3, \quad X^* = \delta(x_M), \quad m=3, \quad n=5. \end{array} \right.$$

Iteration $i=1$:

$$\text{Step 1: } R_1 = \begin{bmatrix} \sqrt{\frac{2}{128}} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \sqrt{\frac{5}{128}} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \sqrt{\frac{90}{128}} \end{bmatrix}.$$

$$\begin{array}{l} \text{find } Z_1 \in \mathbb{S}^3 \\ \text{subject to } \langle Z_1, R_1^T A_j R_1 \rangle = 0, \quad j=1, 2, 3 \end{array} \quad (671)$$

A nonzero randomly selected matrix Z_1 having $\mathbf{0}$ main diagonal is feasible and yields a nonzero perturbation matrix. Choose, arbitrarily,

$$Z_1 = \mathbf{1}\mathbf{1}^T - I \in \mathbb{S}^3 \quad (672)$$

then (rounding)

$$B_1 = \begin{bmatrix} 0 & 0 & 0.0247 & 0 & 0.1048 \\ 0 & 0 & 0 & 0 & 0 \\ 0.0247 & 0 & 0 & 0 & 0.1657 \\ 0 & 0 & 0 & 0 & 0 \\ 0.1048 & 0 & 0.1657 & 0 & 0 \end{bmatrix} \quad (673)$$

Step 2: $t_1^* = 1$ because $\lambda(Z_1) = [-1 \ -1 \ 2]^T$. So,

$$X^* \leftarrow \delta(x_M) + B_1 = \begin{bmatrix} \frac{2}{128} & 0 & 0.0247 & 0 & 0.1048 \\ 0 & 0 & 0 & 0 & 0 \\ 0.0247 & 0 & \frac{5}{128} & 0 & 0.1657 \\ 0 & 0 & 0 & 0 & 0 \\ 0.1048 & 0 & 0.1657 & 0 & \frac{90}{128} \end{bmatrix} \quad (674)$$

has rank $\rho \leftarrow 1$ and produces the same optimal objective value.

}

□

4.3.3.0.2 Exercise. *Rank reduction of maximal complementarity.*

Apply rank reduction Procedure 4.3.1.0.1 to the *maximal complementarity example* (§4.1.1.4.1). Demonstrate a rank-1 solution; which can certainly be found (by Barvinok’s Proposition 2.9.3.0.1) because there is only one equality constraint. ▼

4.3.4 thoughts regarding rank reduction

Because the *rank reduction procedure* is guaranteed only to produce another optimal solution conforming to Barvinok’s upper bound (246), the Procedure will not necessarily produce solutions of arbitrarily low rank; but if they exist, the Procedure can. Arbitrariness of search direction when matrix Z_i becomes indefinite, mentioned on page 276, and the enormity of choices for Z_i (654) are liabilities for this algorithm.

4.3.4.1 Inequality constraints

The question naturally arises: what to do when a semidefinite program (not in prototypical form (592))^{4.20} has linear inequality constraints of the form

$$\alpha_i^T \text{svec } X \preceq \beta_i, \quad i = 1 \dots k \quad (675)$$

where $\{\beta_i\}$ are given scalars and $\{\alpha_i\}$ are given vectors. One expedient way to handle this circumstance is to convert the inequality constraints to equality constraints by introducing a slack variable γ ; *id est*,

$$\alpha_i^T \text{svec } X + \gamma_i = \beta_i, \quad i = 1 \dots k, \quad \gamma \succeq 0 \quad (676)$$

thereby converting the problem to prototypical form.

^{4.20}Contemporary numerical packages for solving semidefinite programs can solve a range of problems wider than prototype (592). Generally, they do so by transforming a given problem into prototypical form by introducing new constraints and variables. [10] [344] We are momentarily considering a departure from the primal prototype that augments the constraint set with linear inequalities.

Alternatively, we say the i^{th} inequality constraint is *active* when it is met with equality; *id est*, when for particular i in (675), $\alpha_i^T \text{svec } X^* = \beta_i$. An optimal high-rank solution X^* is, of course, feasible satisfying all the constraints. But for the purpose of rank reduction, inactive inequality constraints are ignored while active inequality constraints are interpreted as equality constraints. In other words, we take the union of active inequality constraints (as equalities) with equality constraints $A \text{svec } X = b$ to form a composite affine subset \hat{A} substituting for (595). Then we proceed with rank reduction of X^* as though the semidefinite program were in prototypical form (592P).

4.4 Rank-constrained semidefinite program

A technique for finding low-rank optimal solutions to semidefinite programs of a more general form was discovered in October 2005:

4.4.1 rank constraint by convex iteration

Consider a semidefinite feasibility problem of the form

$$\begin{aligned} & \underset{G \in \mathbb{S}^N}{\text{find}} && G \\ & \text{subject to} && G \in \mathcal{C} \\ & && G \succeq 0 \\ & && \text{rank } G \leq n \end{aligned} \tag{677}$$

where \mathcal{C} is a convex set presumed to contain positive semidefinite matrices of rank n or less; *id est*, \mathcal{C} intersects the positive semidefinite cone boundary. We propose that this rank-constrained feasibility problem is equivalently expressed as the problem sequence (678) (1592a) having convex constraints:

$$\begin{aligned} & \underset{G \in \mathbb{S}^N}{\text{minimize}} && \langle G, W \rangle \\ & \text{subject to} && G \in \mathcal{C} \\ & && G \succeq 0 \end{aligned} \tag{678}$$

where *direction matrix*^{4.21} W is an optimal solution to semidefinite program, for $0 \leq n \leq N - 1$

^{4.21}Search *direction* W is a hyperplane-normal pointing opposite to direction of movement describing minimization of a real linear function $\langle G, W \rangle$ (p.71).

$$\begin{aligned} \sum_{i=n+1}^N \lambda(G^*)_i &= \underset{W \in \mathbb{S}^N}{\text{minimize}} && \langle G^*, W \rangle && (1592a) \\ &\text{subject to} && 0 \preceq W \preceq I \\ &&& \text{tr } W = N - n \end{aligned}$$

whose feasible set is a Fantope (§2.3.2.0.1), and where G^* is an optimal solution to problem (678) given some iterate W . The idea is to iterate solution of (678) and (1592a) until convergence as defined in §4.4.1.2. ^{4.22} Optimal matrix W^* is defined as any direction matrix yielding optimal solution G^* of rank n or less to then convex equivalent (678) of feasibility problem (677); *id est*, any direction matrix for which the last $N - n$ nonincreasingly ordered eigenvalues λ of G^* are zero: (p.603)

$$\sum_{i=n+1}^N \lambda(G^*)_i = \langle G^*, W^* \rangle \triangleq 0 \quad (679)$$

We emphasize that convex problem (678) is not a relaxation of the rank-constrained feasibility problem (677); at convergence, *convex iteration* (678) (1592a) makes it instead an *equivalent problem*. ^{4.23}

4.4.1.1 direction interpretation

We make no assumption regarding uniqueness of direction matrix W . The feasible set of direction matrices in (1592a) is the convex hull of outer product of all rank- $(N - n)$ orthonormal matrices; *videlicet*,

$$\text{conv}\{UU^T \mid U \in \mathbb{R}^{N \times N-n}, U^T U = I\} = \{A \in \mathbb{S}^N \mid I \succeq A \succeq 0, \langle I, A \rangle = N - n\} \quad (83)$$

Set $\{UU^T \mid U \in \mathbb{R}^{N \times N-n}, U^T U = I\}$ comprises the extreme points of this Fantope (83). An optimal solution W to (1592a), that is an extreme point, is known in closed form (p.603). By (206), that particular direction ($-W$) can be regarded as pointing toward positive semidefinite rank- n matrices that

^{4.22}The proposed iteration is not an *alternating projection*. (confer Figure 143)

^{4.23}Terminology *equivalent problem* meaning, optimal solution to one problem can be derived from optimal solution to another. Terminology *same problem* means: optimal solution set for one problem is identical to the optimal solution set of another (without transformation).

are simultaneously diagonalizable with G^* . When $W = I$, as in the trace heuristic for example, then $-W$ points directly at the origin (the rank-0 PSD matrix). Inner product of an optimization variable with the direction matrix W is therefore a generalization of the trace heuristic (§7.2.2.1); $-W$ is instead trained toward the boundary of the positive semidefinite cone.

4.4.1.2 convergence

We study convergence to ascertain conditions under which a direction matrix will reveal a feasible G matrix of rank n or less in semidefinite program (678). Denote by W^* a particular optimal direction matrix from semidefinite program (1592a) such that (679) holds. Then we define *global convergence* of the iteration (678) (1592a) to correspond with this vanishing vector inner-product (679) of optimal solutions.

Because this iterative technique for constraining rank is not a projection method, it can find a rank- n solution G^* ((679) will be satisfied) only if at least one exists in the feasible set of program (678).

4.4.1.2.1 Proof. Suppose $\langle G^*, W \rangle = \tau$ is satisfied for some nonnegative constant τ after any particular iteration (678) (1592a) of the two minimization problems. Once a particular value of τ is achieved, it can never be exceeded by subsequent iterations because existence of feasible G and W having that vector inner-product τ has been established simultaneously in each problem. Because the infimum of vector inner-product of two positive semidefinite matrix variables is zero, the nonincreasing sequence of iterations is thus bounded below hence convergent because any bounded monotonic sequence in \mathbb{R} is convergent. [228, §1.2] [37, §1.1] *Local convergence* to some τ is thereby established. \blacklozenge

When a rank- n feasible solution to (678) exists, it remains to be shown under what conditions $\langle G^*, W^* \rangle = 0$ (679) is achieved by iterative solution of semidefinite programs (678) and (1592a).^{4.24} Then pair (G^*, W^*) becomes a *fixed point* of iteration.

A nonexistent feasible rank- n solution would mean certain failure to converge by definition (679) but, as proved, *convex iteration* always converges locally if not globally. Now, an application:

^{4.24}The case of affine constraints \mathcal{C} is considered in [265].

4.4.1.2.2 Example. *Sensor-Network Localization and Wireless Location.* Heuristic solution to a sensor-network localization problem, proposed by Carter, Jin, Saunders, & Ye in [63],^{4.25} is limited to two Euclidean dimensions and applies semidefinite programming (SDP) to little subproblems. There, a large network is partitioned into smaller subnetworks (as small as one *sensor* – a mobile point, whereabouts unknown) and then semidefinite programming and heuristics called SPASELOC are applied to localize each and every partition by two-dimensional distance geometry. Their partitioning procedure is one-pass, yet termed *iterative*; a term applicable only in so far as adjoining partitions can share localized sensors and *anchors* (absolute sensor positions known *a priori*). But there is no iteration on the entire network, hence the term “iterative” is perhaps inappropriate. As partitions are selected based on “rule sets” (heuristics, not geographics), they also term the partitioning *adaptive*. But no adaptation of a partition actually occurs once it has been determined.

One can reasonably argue that semidefinite programming methods are unnecessary for localization of small partitions of large sensor networks. In the past, these nonlinear localization problems were solved algebraically and computed by least squares solution to hyperbolic equations; called *multilateration*.^{4.26} Indeed, practical contemporary numerical methods for global positioning by satellite (GPS) do not rely on semidefinite programming.

Modern distance geometry is inextricably melded with semidefinite programming. The beauty of semidefinite programming, as relates to localization, lies in convex expression of classical multilateration: So & Ye showed [284] that the problem of finding unique solution, to a noiseless nonlinear system describing the common point of intersection of hyperspheres in real Euclidean vector space, can be expressed as a semidefinite program via distance geometry.

But the need for SDP methods in Carter & Jin *et alii* is enigmatic for two more reasons: 1) guessing solution to a partition whose intersensor measurement data or connectivity is inadequate for localization by distance geometry, 2) reliance on complicated and extensive heuristics for partitioning

^{4.25}The paper constitutes [Jin’s dissertation](#) for University of Toronto [191] although her name appears as second author. Ye’s authorship is honorary.

^{4.26}**Multilateration** – literally, *having many sides*; shape of a geometric figure formed by nearly intersecting lines of position. In navigation systems, therefore: Obtaining a *fix* from multiple lines of position.

a large network that could instead be solved whole by one semidefinite program. While partitions range in size between 2 and 10 sensors, 5 sensors optimally, heuristics provided are only for two spatial dimensions (no higher-dimensional heuristics are proposed). For these small numbers it remains unclarified as to precisely what advantage is gained over traditional least squares.

Partitioning of large sensor networks is a compromise to rapid growth of SDP computational complexity with problem size. But when impact of noise on distance measurement is of most concern, one is averse to a partitioning scheme because noise-effects vary inversely with problem size. [46, §2.2] (§5.13.2) Since an individual partition's solution is not iterated in Carter & Jin and is interdependent with adjoining partitions, we expect errors to propagate from one partition to the next; the ultimate partition solved, expected to suffer most.

Heuristics often fail on real-world data because of unanticipated circumstances. When heuristics fail, generally they are repaired by adding more heuristics. Tenuous is any presumption, for example, that distance measurement errors have distribution characterized by circular contours of equal probability about an unknown sensor-location. That presumption effectively appears within Carter & Jin's optimization problem statement as affine equality constraints relating unknowns to distance measurements that are corrupted by noise. Yet in most all urban environments, this measurement noise is more aptly characterized by ellipsoids of varying orientation and eccentricity as one recedes from a sensor. Each unknown sensor must therefore instead be bound to its own particular range of distance, primarily determined by the terrain.^{4.27} The nonconvex problem we must instead solve is:

$$\begin{aligned} & \underset{i, j \in \mathcal{I}}{\text{find}} && \{x_i, x_j\} \\ & \text{subject to} && \underline{d}_{ij} \leq \|x_i - x_j\|^2 \leq \overline{d}_{ij} \end{aligned} \quad (680)$$

where x_i represents sensor location, and where \underline{d}_{ij} and \overline{d}_{ij} respectively represent lower and upper bounds on measured distance-square from i^{th} to j^{th} sensor (or from sensor to anchor). Figure 78 illustrates contours of equal sensor-location uncertainty. By establishing these individual upper and lower bounds, orientation and eccentricity can effectively be incorporated into the problem statement.

^{4.27}A distinct contour map corresponding to each anchor is required in practice.

Generally speaking, there can be no unique solution to the sensor-network localization problem because there is no unique formulation; that is the art of Optimization. Any optimal solution obtained depends on whether or how the network is partitioned and how the problem is formulated. When a particular formulation is a convex optimization problem, then the set of all optimal solutions forms a convex set containing the actual or true localization. Measurement noise precludes equality constraints representing distance. The optimal solution set is consequently expanded; necessitated by introduction of distance inequalities admitting more and higher-rank solutions. Even were the optimal solution set a single point, it is not necessarily the true localization because there is little hope of exact localization by any algorithm once significant noise is introduced.

Carter & Jin gauge performance of their heuristics to the SDP formulation of author Biswas whom they regard as vanguard to the art. [13, §1] Biswas posed localization as an optimization problem minimizing a distance measure. [42] [40] Intuitively, minimization of any distance measure yields compacted solutions; (*confer* §6.4.0.0.1) precisely the anomaly motivating Carter & Jin. Their two-dimensional heuristics outperformed Biswas' localizations both in execution-time and proximity to the desired result. Perhaps, instead of heuristics, Biswas' approach to localization can be improved: [39] [41].

The sensor-network localization problem is considered difficult. [13, §2] Rank constraints in optimization are considered more difficult. Control of Euclidean dimension in Carter & Jin is suboptimal because of implicit projection on \mathbb{R}^2 . In what follows, we present the localization problem as a semidefinite program (equivalent to (680)) having an explicit rank constraint which controls Euclidean dimension of an optimal solution. We show how to achieve that rank constraint only if the feasible set contains a matrix of desired rank. Our problem formulation is extensible to any spatial dimension.

proposed standardized test

Jin proposes an academic test in two-dimensional real Euclidean space \mathbb{R}^2 that we adopt. In essence, this test is a localization of sensors and anchors arranged in a regular triangular lattice. Lattice connectivity is solely determined by sensor radio range; a connectivity graph is assumed incomplete. In the interest of test standardization, we propose adoption of a few small examples: Figure 74 through Figure 77 and their particular connectivity represented by matrices (681) through (684) respectively.




Figure 74: 2-lattice in \mathbb{R}^2 , hand-drawn. Nodes 3 and 4 are anchors; remaining nodes are sensors. Radio range of sensor 1 indicated by arc.

$$\begin{array}{cccc}
 0 & \bullet & ? & \bullet \\
 \bullet & 0 & \bullet & \bullet \\
 ? & \bullet & 0 & \circ \\
 \bullet & \bullet & \circ & 0
 \end{array} \tag{681}$$

Matrix entries *dot* \bullet indicate measurable distance between nodes while unknown distance is denoted by $?$ (*question mark*). Matrix entries *hollow dot* \circ represent known distance between anchors (to very high accuracy) while zero distance is denoted 0. Because measured distances are quite unreliable in practice, our solution to the localization problem substitutes a distinct range of possible distance for each measurable distance; equality constraints exist only for anchors.

Anchors are chosen so as to increase difficulty for algorithms dependent on existence of sensors in their convex hull. The challenge is to find a solution in two dimensions close to the true sensor positions given incomplete noisy intersensor distance information.




Figure 75: 3-lattice in \mathbb{R}^2 , hand-drawn. Nodes 7, 8, and 9 are anchors; remaining nodes are sensors. Radio range of sensor 1 indicated by arc.

$$\begin{array}{cccccccc}
 0 & \bullet & \bullet & ? & \bullet & ? & ? & \bullet & \bullet \\
 \bullet & 0 & \bullet & \bullet & ? & \bullet & ? & \bullet & \bullet \\
 \bullet & \bullet & 0 & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\
 ? & \bullet & \bullet & 0 & ? & \bullet & \bullet & \bullet & \bullet \\
 \bullet & ? & \bullet & ? & 0 & \bullet & \bullet & \bullet & \bullet \\
 ? & \bullet & \bullet & \bullet & \bullet & 0 & \bullet & \bullet & \bullet \\
 ? & ? & \bullet & \bullet & \bullet & \bullet & 0 & \circ & \circ \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \circ & 0 & \circ \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \circ & \circ & 0
 \end{array}$$

(682)




Figure 76: 4-lattice in \mathbb{R}^2 , hand-drawn. Nodes 13, 14, 15, and 16 are anchors; remaining nodes are sensors. Radio range of sensor 1 indicated by arc.

$$\begin{array}{cccccccccccccccc}
 0 & ? & ? & \bullet & ? & ? & \bullet & ? & ? & ? & ? & ? & ? & ? & \bullet & \bullet \\
 ? & 0 & \bullet & \bullet & \bullet & \bullet & ? & \bullet & ? & ? & ? & ? & ? & \bullet & \bullet & \bullet \\
 ? & \bullet & 0 & ? & \bullet & \bullet & ? & ? & \bullet & ? & ? & ? & ? & ? & \bullet & \bullet \\
 \bullet & \bullet & ? & 0 & \bullet & ? & \bullet & \bullet & ? & \bullet & ? & ? & \bullet & \bullet & \bullet & \bullet \\
 ? & \bullet & \bullet & \bullet & 0 & \bullet & ? & \bullet & \bullet & ? & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\
 ? & \bullet & \bullet & ? & \bullet & 0 & ? & \bullet & \bullet & ? & \bullet & \bullet & ? & ? & ? & ? \\
 \bullet & ? & ? & \bullet & ? & ? & 0 & ? & ? & \bullet & ? & ? & \bullet & \bullet & \bullet & \bullet \\
 ? & \bullet & ? & \bullet & \bullet & \bullet & ? & 0 & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\
 ? & ? & \bullet & ? & \bullet & \bullet & ? & \bullet & 0 & ? & \bullet & \bullet & ? & \bullet & ? \\
 ? & ? & ? & \bullet & ? & ? & \bullet & \bullet & ? & 0 & \bullet & ? & \bullet & \bullet & \bullet & ? \\
 ? & ? & ? & ? & \bullet & \bullet & ? & \bullet & \bullet & \bullet & 0 & \bullet & \bullet & \bullet & \bullet & ? \\
 ? & ? & ? & ? & \bullet & \bullet & ? & \bullet & \bullet & ? & \bullet & 0 & ? & ? & ? & ? \\
 ? & ? & ? & \bullet & \bullet & ? & \bullet & \bullet & \bullet & \bullet & ? & 0 & \circ & \circ & \circ & \circ \\
 ? & \bullet & ? & \bullet & \bullet & ? & \bullet & \bullet & ? & \bullet & \bullet & ? & \circ & 0 & \circ & \circ \\
 \bullet & \bullet & \bullet & \bullet & \bullet & ? & \bullet & \bullet & \bullet & \bullet & \bullet & ? & \circ & \circ & 0 & \circ \\
 \bullet & \bullet & \bullet & \bullet & \bullet & ? & \bullet & \bullet & ? & ? & ? & ? & \circ & \circ & \circ & 0
 \end{array} \tag{683}$$




Figure 77: 5-lattice in \mathbb{R}^2 . Nodes 21 through 25 are anchors.

0 ● ? ? ● ● ? ? ● ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ?
 ● 0 ? ? ● ● ? ? ? ● ? ? ? ? ? ? ? ? ? ? ? ? ? ● ●
 ? ? 0 ● ? ● ● ● ? ? ● ● ? ? ? ? ? ? ? ? ? ? ● ● ●
 ? ? ● 0 ? ? ● ● ? ? ? ● ? ? ? ? ? ? ? ? ? ? ? ● ?
 ● ● ? ? 0 ● ? ? ● ● ? ? ● ● ? ? ● ? ? ? ? ? ● ? ●
 ● ● ● ? ● 0 ● ? ● ● ● ? ? ● ? ? ? ? ? ? ? ? ● ● ●
 ? ? ● ● ? ● 0 ● ? ? ● ● ? ? ● ● ? ? ? ? ? ? ? ● ● ●
 ? ? ● ● ? ? ● 0 ? ? ● ● ? ? ● ● ? ? ? ? ? ? ? ? ● ?
 ● ? ? ? ● ● ? ? 0 ● ? ? ● ● ? ? ● ● ? ? ? ? ? ? ? ?
 ? ● ? ? ● ● ? ? ● 0 ● ? ● ● ? ? ? ● ? ? ● ● ● ● ●
 ? ? ● ? ? ● ● ● ? ● 0 ● ? ● ● ● ? ? ● ? ? ● ● ● ●
 ? ? ● ● ? ? ● ● ? ? ● 0 ? ? ● ● ? ? ● ● ? ● ● ● ?
 ? ? ? ? ● ? ? ? ● ● ? ? 0 ● ? ? ● ● ? ? ● ● ? ? ?
 ? ? ? ? ● ● ? ? ● ● ● ? ● 0 ● ? ● ● ● ? ● ● ● ● ?
 ? ? ? ? ? ? ● ● ? ? ● ● ? ? ● 0 ● ? ? ● ● ● ● ● ?
 ? ? ? ? ? ? ● ● ? ? ● ● ? ? ● 0 ? ? ● ● ? ● ? ? ? ?
 ? ? ? ? ● ? ? ? ● ? ? ? ● ● ? ? 0 ● ? ? ● ? ? ? ? ?
 ? ? ? ? ? ? ? ? ? ● ● ? ? ● ● ? ? ● 0 ● ● ? ? ? ?
 ? ? ? ? ? ? ? ? ? ? ● ? ? ● ● ● ? ● ● ● ● 0 ○ ○ ○ ○
 ? ? ? ? ? ? ? ? ? ? ● ● ● ● ● ● ? ● ● ● ● 0 ○ ○ ○
 ? ? ● ? ● ● ● ? ? ● ● ● ? ● ● ? ? ● ● ? ○ ○ 0 ○ ○
 ? ● ● ● ? ● ● ● ? ● ● ● ? ● ● ? ? ? ? ? ? ○ ○ 0 ○ ○
 ? ● ● ? ● ● ● ? ? ● ● ? ? ? ? ? ? ? ? ? ? ○ ○ ○ ○ 0




Figure 78: Location uncertainty ellipsoid in \mathbb{R}^2 for each of 15 sensors \bullet within three city blocks in downtown San Francisco. Data by Polaris Wireless. [285]

problem statement

Ascribe points in a list $\{x_\ell \in \mathbb{R}^n, \ell = 1 \dots N\}$ to the columns of a matrix X ;

$$X = [x_1 \cdots x_N] \in \mathbb{R}^{n \times N} \quad (69)$$

where N is regarded as cardinality of list X . Positive semidefinite matrix $X^T X$, formed from inner product of the list, is a *Gram matrix*; [220, §3.6]

$$G \triangleq X^T X = \begin{bmatrix} \|x_1\|^2 & x_1^T x_2 & x_1^T x_3 & \cdots & x_1^T x_N \\ x_2^T x_1 & \|x_2\|^2 & x_2^T x_3 & \cdots & x_2^T x_N \\ x_3^T x_1 & x_3^T x_2 & \|x_3\|^2 & \ddots & x_3^T x_N \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ x_N^T x_1 & x_N^T x_2 & x_N^T x_3 & \cdots & \|x_N\|^2 \end{bmatrix} \in \mathbb{S}_+^N \quad (817)$$

where \mathbb{S}_+^N is the convex cone of $N \times N$ positive semidefinite matrices in the symmetric matrix subspace \mathbb{S}^N .

Existence of noise precludes measured distance from the input data. We instead assign measured distance to a range estimate specified by individual upper and lower bounds: \bar{d}_{ij} is an upper bound on distance-square from i^{th}

to j^{th} sensor, while \underline{d}_{ij} is a lower bound. These bounds become the input data. Each measurement range is presumed different from the others because of measurement uncertainty; *e.g.*, Figure 78.

Our mathematical treatment of anchors and sensors is not dichotomized.^{4.28} A known sensor position to high accuracy \tilde{x}_i is an anchor. Then the sensor-network localization problem (680) can be expressed equivalently: Given a number of anchors m , and \mathcal{I} a set of indices (corresponding to all measurable distances \bullet), for $0 < m < N$

$$\begin{aligned}
& \underset{G \in \mathbb{S}^N, X \in \mathbb{R}^{n \times N}}{\text{minimize}} && \text{tr } Z \\
& \text{subject to} && \underline{d}_{ij} \leq \langle G, (e_i - e_j)(e_i - e_j)^{\text{T}} \rangle \leq \overline{d}_{ij} \quad \forall (i, j) \in \mathcal{I} \\
& && \langle G, e_i e_i^{\text{T}} \rangle = \|\tilde{x}_i\|^2, \quad i = N - m + 1 \dots N \\
& && \langle G, (e_i e_j^{\text{T}} + e_j e_i^{\text{T}})/2 \rangle = \tilde{x}_i^{\text{T}} \tilde{x}_j, \quad i < j, \quad \forall i, j \in \{N - m + 1 \dots N\} \\
& && X(:, N - m + 1 : N) = [\tilde{x}_{N-m+1} \ \cdots \ \tilde{x}_N] \\
& && Z = \begin{bmatrix} I & X \\ X^{\text{T}} & G \end{bmatrix} \succeq 0 \\
& && \text{rank } Z = n
\end{aligned} \tag{685}$$

where e_i is the i^{th} member of the standard basis for \mathbb{R}^N . Distance-square

$$d_{ij} = \|x_i - x_j\|_2^2 \triangleq \langle x_i - x_j, x_i - x_j \rangle \tag{804}$$

is related to Gram matrix entries $G \triangleq [g_{ij}]$ by vector inner-product

$$\begin{aligned}
d_{ij} &= g_{ii} + g_{jj} - 2g_{ij} \\
&= \langle G, (e_i - e_j)(e_i - e_j)^{\text{T}} \rangle \triangleq \text{tr}(G^{\text{T}}(e_i - e_j)(e_i - e_j)^{\text{T}})
\end{aligned} \tag{819}$$

hence the scalar inequalities. The objective function $\text{tr } Z$ is a heuristic whose sole purpose is to represent the convex envelope of $\text{rank } Z$. (§7.2.2.1.1) By Schur complement (§A.4), any feasible G and X provide a comparison with respect to the positive semidefinite cone

$$G \succeq X^{\text{T}}X \tag{852}$$

^{4.28} *Wireless location* problem thus stated identically; difference being: fewer sensors.

which is a convex relaxation of the desired equality constraint

$$\begin{bmatrix} I & X \\ X^T & G \end{bmatrix} = \begin{bmatrix} I \\ X^T \end{bmatrix} [I \ X] \quad (853)$$

The rank constraint insures this equality holds thus restricting solution to \mathbb{R}^n .

convex equivalent problem statement

Problem statement (685) is nonconvex because of the rank constraint. We do not eliminate or ignore the rank constraint; rather, we find a convex way to enforce it: for $0 < n < N$

$$\begin{aligned} & \underset{G \in \mathbb{S}^N, X \in \mathbb{R}^{n \times N}}{\text{minimize}} && \langle Z, W \rangle \\ & \text{subject to} && \overline{d_{ij}} \leq \langle G, (e_i - e_j)(e_i - e_j)^T \rangle \leq \overline{d_{ij}} \quad \forall (i, j) \in \mathcal{I} \\ & && \langle G, e_i e_i^T \rangle = \|\check{x}_i\|^2, \quad i = N - m + 1 \dots N \\ & && \langle G, (e_i e_j^T + e_j e_i^T)/2 \rangle = \check{x}_i^T \check{x}_j, \quad i < j, \quad \forall i, j \in \{N - m + 1 \dots N\} \\ & && X(:, N - m + 1 : N) = [\check{x}_{N-m+1} \ \cdots \ \check{x}_N] \\ & && Z = \begin{bmatrix} I & X \\ X^T & G \end{bmatrix} \succeq 0 \end{aligned} \quad (686)$$

Each linear equality constraint in $G \in \mathbb{S}^N$ represents a hyperplane in isometrically isomorphic Euclidean vector space $\mathbb{R}^{N(N+1)/2}$, while each linear inequality pair represents a convex Euclidean body known as *slab* (an intersection of two parallel but opposing halfspaces, Figure 11). In this convex optimization problem (686), a semidefinite program, we substitute a vector inner-product objective function for trace from nonconvex problem (685);

$$\langle Z, I \rangle = \text{tr } Z \leftarrow \langle Z, W \rangle \quad (687)$$

a generalization of the known trace heuristic [232] [118] for minimizing convex envelope of rank, where $W \in \mathbb{S}_+^{N+n}$ is constant with respect to (686). Matrix W is normal to a hyperplane in \mathbb{S}^{N+n} minimized over a convex feasible set specified by the constraints in (686). Matrix W is chosen so $-W$ points in the direction of a feasible rank- n Gram matrix. Thus the purpose of vector inner-product objective (687) is to locate a feasible rank- n Gram matrix that is presumed existent on the boundary of positive semidefinite cone \mathbb{S}_+^N .

direction matrix W

Denote by Z^* an optimal composite matrix from semidefinite program (686). Then for $Z^* \in \mathbb{S}^{N+n}$ whose eigenvalues $\lambda(Z^*) \in \mathbb{R}^{N+n}$ are arranged in nonincreasing order, (Ky Fan)

$$\begin{aligned} \sum_{i=n+1}^{N+n} \lambda(Z^*)_i &= \underset{W \in \mathbb{S}^{N+n}}{\text{minimize}} \quad \langle Z^*, W \rangle & (1592a) \\ &\text{subject to} \quad 0 \preceq W \preceq I \\ &\quad \text{tr } W = N \end{aligned}$$

which has an optimal solution that is known in closed form. This eigenvalue sum is zero when Z^* has rank n or less.

Foreknowledge of optimal Z^* , to make possible this search for W , implies iteration; *id est*, semidefinite program (686) is solved for Z^* initializing $W=I$ or $W=0$. Once found, Z^* becomes constant in semidefinite program (1592a) where a new normal direction W is found as its optimal solution. Then the cycle (686) (1592a) iterates until convergence. When rank $Z^* = n$, solution via this *convex iteration* is optimal for sensor-network localization problem (680) and its equivalent (685).

numerical solution

In all examples to follow, number of anchors

$$m = \sqrt{N} \tag{688}$$

equals square root of cardinality N of list X . Indices set \mathcal{I} identifying all measurable distances \bullet is ascertained from connectivity matrix (681), (682), (683), or (684). We solve iteration (686) (1592a) in dimension $n=2$ for each respective example illustrated in Figure 74 through Figure 77.

In presence of negligible noise, true position is reliably localized for every standardized example; noteworthy in so far as each example represents an incomplete graph. This means the set of all solutions having lowest rank is a single point, to within a rigid transformation.

To make the examples interesting and consistent with previous work, we randomize each range of distance-square that bounds $\langle G, (e_i - e_j)(e_i - e_j)^T \rangle$ in (686); *id est*, for each and every $(i, j) \in \mathcal{I}$

$$\begin{aligned} \overline{d_{ij}} &= d_{ij}(1 + \sqrt{3}\eta\chi_l)^2 \\ \underline{d_{ij}} &= d_{ij}(1 - \sqrt{3}\eta\chi_{l+1})^2 \end{aligned} \tag{689}$$

where $\eta = 0.1$ is a constant noise factor, χ_l is the l^{th} sample of a noise process realization uniformly distributed in the interval $[0, 1]$ like `rand(1)` from MATLAB, and d_{ij} is actual distance-square from i^{th} to j^{th} sensor. Because of distinct function calls to `rand()`, each range of distance-square $[\underline{d}_{ij}, \overline{d}_{ij}]$ is not necessarily centered on actual distance-square d_{ij} . Unit stochastic variance is provided by factor $\sqrt{3}$.

Figure 79 through Figure 82 each illustrate one realization of numerical solution to the standardized lattice problems posed by Figure 74 through Figure 77 respectively. Exact localization is impossible because of measurement noise. Certainly, by inspection of their published graphical data, our new results are competitive with those of Carter & Jin. Obviously our solutions do not suffer from those compaction-type errors (clustering of localized sensors) exhibited by Biswas' graphical results for the same noise factor η ; which is all we intended to demonstrate.

localization example conclusion

Solution to this sensor-network localization problem became apparent by understanding geometry of optimization. Trace of a matrix, to a student of linear algebra, is perhaps a sum of eigenvalues. But to us, trace represents the normal I to some hyperplane in Euclidean vector space.

Numerical solution to noisy problems containing sensor variables in excess of 100 becomes difficult via the holistic semidefinite program we propose. When problem size is within reach of contemporary general-purpose semidefinite program solvers, then the equivalent program presented inherently overcomes limitations of Carter & Jin with respect to both noise performance and ability to constrain localization to any desired affine dimension.

The legacy of Carter, Jin, Saunders, & Ye [63] is a sobering demonstration of the need for more efficient methods for solution of semidefinite programs, while that of So & Ye [284] is the bonding of distance geometry to semidefinite programming. Elegance of our semidefinite problem statement (686), for constraining affine dimension of sensor-network localization, should provide some *impetus* to focus more research on computational intensity of general-purpose semidefinite program solvers. An approach different from interior-point methods is required; higher speed and greater accuracy from a simplex-like solver is what is needed. \square




Figure 79: Typical solution for 2-lattice in Figure 74 with noise factor $\eta = 0.1$. Two red rightmost nodes are anchors; two remaining nodes are sensors. Radio range of sensor 1 indicated by arc; radius = 1.14. Actual sensor indicated by target \circ while its localization is indicated by bullet \bullet . Rank-2 solution found in 1 iteration (686) (1592a) subject to reflection error.




Figure 80: Typical solution for 3-lattice in Figure 75 with noise factor $\eta = 0.1$. Three red vertical middle nodes are anchors; remaining nodes are sensors. Radio range of sensor 1 indicated by arc; radius = 1.12. Actual sensor indicated by target \circ while its localization is indicated by bullet \bullet . Rank-2 solution found in 2 iterations (686) (1592a).




Figure 81: Typical solution for 4-lattice in Figure 76 with noise factor $\eta = 0.1$. Four red vertical middle-left nodes are anchors; remaining nodes are sensors. Radio range of sensor 1 indicated by arc; radius = 0.75. Actual sensor indicated by target \odot while its localization is indicated by bullet \bullet . Rank-2 solution found in 7 iterations (686) (1592a).




Figure 82: Typical solution for 5-lattice in Figure 77 with noise factor $\eta = 0.1$. Five red vertical middle nodes are anchors; remaining nodes are sensors. Radio range of sensor 1 indicated by arc; radius = 0.56. Actual sensor indicated by target \odot while its localization is indicated by bullet \bullet . Rank-2 solution found in 3 iterations (686) (1592a).

We numerically tested the foregoing technique for constraining rank on a wide range of problems including localization of randomized positions, *stress* (§7.2.2.7.1), *ball packing* (§5.4.2.2.3), and cardinality problems. We have had some success introducing the direction matrix inner-product (687) as a *regularization* term (a *multiobjective* optimization) whose purpose is to constrain rank, affine dimension, or cardinality:

4.5 Constraining cardinality

The *convex iteration* technique for constraining rank, discovered in 2005, was soon applied to cardinality problems:

4.5.1 nonnegative variable

Our goal is to reliably constrain rank in a semidefinite program. There is a direct analogy to linear programming that is simpler to present but, perhaps, more difficult to solve. In Optimization, that analogy is known as the *cardinality problem*.

Consider a feasibility problem $Ax = b$, but with an upper bound k on cardinality $\|x\|_0$ of a nonnegative solution x : for $A \in \mathbb{R}^{m \times n}$ and vector $b \in \mathcal{R}(A)$

$$\begin{aligned} & \text{find } x \in \mathbb{R}^n \\ & \text{subject to } Ax = b \\ & \quad x \succeq 0 \\ & \quad \|x\|_0 \leq k \end{aligned} \tag{690}$$

where $\|x\|_0 \leq k$ means^{4.29} vector x has at most k nonzero entries; such a vector is presumed existent in the feasible set. Nonnegativity constraint $x \succeq 0$ is analogous to positive semidefiniteness; the notation means vector x belongs to the nonnegative orthant \mathbb{R}_+^n . Cardinality is quasiconcave on \mathbb{R}_+^n just as rank is quasiconcave on \mathbb{S}_+^n . [53, §3.4.2]

We propose that cardinality-constrained feasibility problem (690) is equivalently expressed as a sequence of convex problems: for $0 \leq k \leq n-1$

^{4.29} Although it is a metric (§5.2), cardinality $\|x\|_0$ cannot be a norm (§3.1.3) because it is not positively homogeneous.

$$\begin{aligned}
& \underset{x \in \mathbb{R}^n}{\text{minimize}} && \langle x, y \rangle \\
& \text{subject to} && Ax = b \\
& && x \succeq 0
\end{aligned} \tag{144}$$

$$\begin{aligned}
\sum_{i=k+1}^n \pi(x^*)_i &= \underset{y \in \mathbb{R}^n}{\text{minimize}} && \langle x^*, y \rangle \\
& \text{subject to} && 0 \preceq y \preceq \mathbf{1} \\
& && y^T \mathbf{1} = n - k
\end{aligned} \tag{475}$$

where π is the (nonincreasing) presorting function. This sequence is iterated until $x^{*\top} y^*$ vanishes; *id est*, until desired cardinality is achieved.

Problem (475) is analogous to the rank constraint problem; (Ky Fan, *confer* p.282)

$$\begin{aligned}
\sum_{i=k+1}^N \lambda(G^*)_i &= \underset{W \in \mathbb{S}^N}{\text{minimize}} && \langle G^*, W \rangle \\
& \text{subject to} && 0 \preceq W \preceq I \\
& && \text{tr } W = N - k
\end{aligned} \tag{1592a}$$

Linear program (475) sums the $n-k$ smallest entries from vector x . In context of problem (690), we want $n-k$ entries of x to sum to zero; *id est*, we want a globally optimal objective $x^{*\top} y^*$ to vanish: more generally,

$$\sum_{i=k+1}^n \pi(|x^*|)_i = \langle |x^*|, y^* \rangle \stackrel{\Delta}{=} 0 \tag{691}$$

Then $n-k$ entries of x^* are themselves zero whenever their absolute sum is, and cardinality of $x^* \in \mathbb{R}^n$ is at most k .

4.5.1.1 direction vector interpretation

Vector y may be interpreted as a negative search direction; it points opposite to direction of movement of hyperplane $\{x \mid \langle x, y \rangle = \tau\}$ in a minimization of real linear function $\langle x, y \rangle$ over the feasible set in linear program (144). (p.71)

Direction vector y is not unique. The feasible set of direction vectors in (475) is the convex hull of all cardinality- $(n-k)$ one-vectors; *videlicet*,

$$\text{conv}\{u \in \mathbb{R}^n \mid \text{card } u = n-k, u_i \in \{0, 1\}\} = \{a \in \mathbb{R}^n \mid \mathbf{1} \succeq a \succeq 0, \langle \mathbf{1}, a \rangle = n-k\} \tag{692}$$

The set $\{u \in \mathbb{R}^n \mid \text{card } u = n-k, u_i \in \{0, 1\}\}$ comprises the extreme points of set (692) which is a nonnegative *hypercube slice*. An optimal solution y to (475), that is an extreme point of its feasible set, is known in closed form: it has 1 in each entry corresponding to the $n-k$ smallest entries of x^* and has 0 elsewhere. By Proposition 7.1.3.0.3, the polar of that particular direction $-y$ can be interpreted^{4.30} as pointing toward the set of all vectors having the same ordering as x^* and cardinality- k . When $y = \mathbf{1}$, as in 1-norm minimization for example, then polar direction $-y$ points directly at the origin (the cardinality-0 vector).

4.5.1.2 convergence

Convex iteration (144) (475) always converges to a locally optimal solution by virtue of a monotonically nonincreasing real objective sequence. [228, §1.2] [37, §1.1] There can be no proof of global optimality, defined by (691). Constraining cardinality, solution to problem (690), can often be achieved but simple examples can be contrived that cause an iteration to *stall* at a solution of undesirable cardinality. The direction vector is then manipulated to steer out of local minima:

4.5.1.2.1 Example. Sparsest solution to $Ax = b$. [61] [101]

Problem (624) has sparsest solution not easily recoverable by least 1-norm; *id est*, not by *compressed sensing* because of proximity to a theoretical lower bound on number of *measurements* m depicted in Figure 83: for $A \in \mathbb{R}^{m \times n}$

- Given data from Example 4.2.3.1.1, for $m=3$, $n=6$, $k=1$

$$A = \begin{bmatrix} -1 & 1 & 8 & 1 & 1 & 0 \\ -3 & 2 & 8 & \frac{1}{2} & \frac{1}{3} & \frac{1}{2} - \frac{1}{3} \\ -9 & 4 & 8 & \frac{1}{4} & \frac{1}{9} & \frac{1}{4} - \frac{1}{9} \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ \frac{1}{2} \\ \frac{1}{4} \end{bmatrix} \quad (624)$$

the sparsest solution to classical linear equation $Ax = b$ is $x = e_4 \in \mathbb{R}^6$ (*confer* (637)).

Although the sparsest solution is recoverable by inspection, we discern it instead by *convex iteration*; namely, by iterating problem sequence

^{4.30} *Convex iteration* (144) (475) is not a projection method because there is no thresholding or discard of variable-vector x entries.




Figure 83: For Gaussian random matrix $A \in \mathbb{R}^{m \times n}$, graph illustrates Donoho/Tanner least lower bound on number of measurements m below which recovery of k -sparse n -length signal x by linear programming fails with overwhelming probability. Hard problems are below curve, but not the reverse; *id est*, failure above depends on proximity. Inequality demarcates *approximation* (dashed) empirically observed in [22]. Problems having nonnegativity constraint (dotted) are easier to solve. [105] [106]

(144) (475) on page 299. From the numerical data given, cardinality $\|x\|_0 = 1$ is expected. Iteration continues until $x^T y$ vanishes (to within some numerical precision); *id est*, until desired cardinality is achieved. But this comes not without a stall.

Stalling, whose occurrence is sensitive to initial conditions of *convex iteration*, is a consequence of finding a local minimum of a multimodal objective $\langle x, y \rangle$ when regarded as simultaneously variable in x and y . Stalls are simply detected as solutions x of undesirable cardinality, sometimes remedied by reinitializing direction vector y to a random positive state.

Bolstered by success in breaking out of a stall, we then apply *convex iteration* to 22,000 randomized problems:

- Given random data for $m=3$, $n=6$, $k=1$, in MATLAB notation

$$A = \text{randn}(3, 6), \quad \text{index} = \text{round}(5 * \text{rand}(1)) + 1, \quad b = \text{rand}(1) * A(:, \text{index}) \quad (693)$$

the sparsest solution $x \propto e_{\text{index}}$ is a scaled standard basis vector.

Without *convex iteration* or a nonnegativity constraint $x \succeq 0$, rate of failure for this minimum cardinality problem $Ax = b$ by 1-norm minimization of x is 22%. That failure rate drops to 6% with a nonnegativity constraint. If we then engage *convex iteration*, detect stalls, and randomly reinitialize the direction vector, failure rate drops to 0% but the amount of computation is approximately doubled. \square

Stalling is not an inevitable behavior. For some problem types (beyond mere $Ax = b$), *convex iteration* succeeds nearly all the time. Here is a cardinality problem, with noise, whose statement is just a bit more intricate but easy to solve in a few convex iterations:

4.5.1.2.2 Example. *Signal dropout.* [104, §6.2]

Signal dropout is an old problem; well studied from both an industrial and academic perspective. Essentially *dropout* means momentary loss or gap in a signal, while passing through some channel, caused by some man-made or natural phenomenon. The signal lost is assumed completely destroyed somehow. What remains within the time-gap is system or idle channel noise. The signal could be voice over Internet protocol (VoIP), for example, audio data from a *compact disc* (CD) or video data from a digital video disc (DVD),

a television transmission over cable or the airwaves, or a typically ravaged cell phone communication, *etcetera*.

Here we consider signal dropout in a discrete-time signal corrupted by additive white noise assumed uncorrelated to the signal. The linear channel is assumed to introduce no filtering. We create a discretized windowed signal for this example by positively combining k randomly chosen vectors from a *discrete cosine transform* (DCT) basis denoted $\Psi \in \mathbb{R}^{n \times n}$. Frequency increases, in the Fourier sense, from DC toward Nyquist as column index of basis Ψ increases. Otherwise, details of the basis are unimportant except for its orthogonality $\Psi^T = \Psi^{-1}$. Transmitted signal is denoted

$$s = \Psi z \in \mathbb{R}^n \quad (694)$$

whose upper bound on DCT basis coefficient cardinality $\text{card } z \leq k$ is assumed known;^{4.31} hence a critical assumption: transmitted signal s is sparsely supported ($k < n$) on the DCT basis. It is further assumed that nonzero signal coefficients in vector z place each chosen basis vector above the noise floor.

We also assume that the gap's beginning and ending in time are precisely localized to within a sample; *id est*, index ℓ locates the last sample prior to the gap's onset, while index $n - \ell + 1$ locates the first sample subsequent to the gap: for rectangularly windowed received signal g possessing a time-gap loss and additive noise $\eta \in \mathbb{R}^n$

$$g = \begin{bmatrix} s_{1:\ell} & + & \eta_{1:\ell} \\ & & \eta_{\ell+1:n-\ell} \\ s_{n-\ell+1:n} & + & \eta_{n-\ell+1:n} \end{bmatrix} \in \mathbb{R}^n \quad (695)$$

The window is thereby centered on the gap and short enough so that the DCT spectrum of signal s can be assumed invariant over the window's duration n . Signal to noise ratio within this window is defined

$$\text{SNR} \triangleq 20 \log \frac{\left\| \begin{bmatrix} s_{1:\ell} \\ s_{n-\ell+1:n} \end{bmatrix} \right\|}{\|\eta\|} \quad (696)$$

In absence of noise, knowing the signal DCT basis and having a good estimate of basis coefficient cardinality makes perfectly reconstructing

^{4.31}This simplifies exposition, although it may be an unrealistic assumption in many applications.

gap-loss easy: it amounts to solving a linear system of equations and requires little or no optimization; with caveat, number of equations exceeds cardinality of signal representation (roughly $\ell \geq k$) with respect to DCT basis.

But addition of a significant amount of noise η increases level of difficulty dramatically; a 1-norm based method of reducing cardinality, for example, almost always returns DCT basis coefficients numbering in excess of minimum cardinality. We speculate that is because signal cardinality 2ℓ becomes the predominant cardinality. DCT basis coefficient cardinality is an explicit constraint to the optimization problem we shall pose: In presence of noise, constraints equating reconstructed signal f to received signal g are not possible. We can instead formulate the dropout recovery problem as a best approximation:

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && \left\| \begin{bmatrix} f_{1:\ell} - g_{1:\ell} \\ f_{n-\ell+1:n} - g_{n-\ell+1:n} \end{bmatrix} \right\| \\ & \text{subject to} && f = \Psi x \\ & && x \succeq 0 \\ & && \text{card } x \leq k \end{aligned} \tag{697}$$

We propose solving this nonconvex problem (697) by moving the cardinality constraint to the objective as a regularization term as explained in §4.5; *id est*, by iteration of two convex problems until convergence:

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && \langle x, y \rangle + \left\| \begin{bmatrix} f_{1:\ell} - g_{1:\ell} \\ f_{n-\ell+1:n} - g_{n-\ell+1:n} \end{bmatrix} \right\| \\ & \text{subject to} && f = \Psi x \\ & && x \succeq 0 \end{aligned} \tag{698}$$

and

$$\begin{aligned} & \underset{y \in \mathbb{R}^n}{\text{minimize}} && \langle x^*, y \rangle \\ & \text{subject to} && 0 \preceq y \preceq \mathbf{1} \\ & && y^T \mathbf{1} = n - k \end{aligned} \tag{475}$$

Signal cardinality 2ℓ is implicit to the problem statement. When number of samples in the dropout region exceeds half the window size, then that deficient cardinality of signal remaining becomes a source of degradation to reconstruction in presence of noise. Thus, by observation, we divine a reconstruction rule for this signal dropout problem to attain good





Figure 84: **(a)** Signal dropout in signal s corrupted by noise η (SNR = 10dB, $g = s + \eta$). Flatline indicates duration of signal dropout. **(b)** Reconstructed signal f (red) overlaid with corrupted signal g .




Figure 85: (a) Error signal power (reconstruction f less original noiseless signal s) is 36dB below s . (b) Original signal s overlaid with reconstruction f (red) from signal g having dropout plus noise.

noise suppression: ℓ must exceed a maximum of cardinality bounds; $2\ell \geq \max\{2k, n/2\}$.

Figure 84 and Figure 85 show one realization of this dropout problem. Original signal s is created by adding four ($k=4$) randomly selected DCT basis vectors, from Ψ ($n=500$ in this example), whose amplitudes are randomly selected from a uniform distribution above the noise floor; in the interval $[10^{-10/20}, 1]$. Then a 240-sample dropout is realized ($\ell=130$) and Gaussian noise η added to make corrupted signal g (from which a best approximation f will be made) having 10dB signal to noise ratio (696). The time gap contains much noise, as apparent from Figure 84a. But in only a few iterations (698) (475), original signal s is recovered with relative error power 36dB down; illustrated in Figure 85. Correct cardinality is also recovered ($\text{card } x = \text{card } z$) along with the basis vector indices used to make original signal s . Approximation error is due to DCT basis coefficient estimate error. When this experiment is repeated 1000 times on noisy signals averaging 10dB SNR, the correct cardinality and indices are recovered 99% of the time with average relative error power 30dB down. Without noise, we get perfect reconstruction in one iteration. (MATLAB code: [Wikimization](#))

□

4.5.1.3 Compressed sensing geometry with a nonnegative variable

It is well known that cardinality problem (690), on page 298, is easier to solve by linear programming when variable x is nonnegatively constrained than when not; *e.g.*, Figure 65, Figure 83. We postulate a simple geometrical explanation: Figure 64 illustrates 1-norm ball \mathcal{B}_1 in \mathbb{R}^3 and affine subset $\mathcal{A} = \{x \mid Ax = b\}$. Prototypical compressed sensing problem, for $A \in \mathbb{R}^{m \times n}$

$$\begin{aligned} & \underset{x}{\text{minimize}} && \|x\|_1 \\ & \text{subject to} && Ax = b \end{aligned} \quad (468)$$

is solved when the 1-norm ball \mathcal{B}_1 kisses affine subset \mathcal{A} .

- If variable x is constrained to the nonnegative orthant,

$$\begin{aligned} & \underset{x}{\text{minimize}} && \|x\|_1 \\ & \text{subject to} && Ax = b \\ & && x \succeq 0 \end{aligned} \quad (699)$$

then 1-norm ball \mathcal{B}_1 (Figure 64) becomes simplex \mathcal{S} in Figure 86.




Figure 86: Line \mathcal{A} intersecting two-dimensional (cardinality-2) face \mathcal{F} of nonnegative simplex \mathcal{S} from Figure 49 emerges at a (cardinality-1) vertex. \mathcal{S} is first quadrant of 1-norm ball \mathcal{B}_1 from Figure 64. Kissing point (\bullet on edge) is achieved as ball contracts or expands under optimization.

Whereas 1-norm ball \mathcal{B}_1 has only six vertices in \mathbb{R}^3 corresponding to cardinality-1 solutions, simplex \mathcal{S} has three edges (along the Cartesian axes) containing an infinity of cardinality-1 solutions. And whereas \mathcal{B}_1 has twelve edges containing cardinality-2 solutions, \mathcal{S} has three (out of total four) two-dimensional faces constituting cardinality-2 solutions. In other words, likelihood of a low-cardinality solution is higher by kissing nonnegative simplex \mathcal{S} (699) than by kissing 1-norm ball \mathcal{B}_1 (468) because facial dimension (corresponding to given cardinality) is higher in \mathcal{S} .

Empirically, this conclusion also holds in other Euclidean dimensions (Figure 65, Figure 83).

4.5.1.4 cardinality-1 compressed sensing problem always solvable

In the special case of cardinality-1 solution to prototypical compressed sensing problem (699), there is a geometrical interpretation that leads to an algorithm more efficient than *convex iteration*. Figure 86 shows a vertex-solution to problem (699) when desired cardinality is 1. But first-quadrant \mathcal{S} of 1-norm ball \mathcal{B}_1 does not kiss line \mathcal{A} ; which requires explanation: Under the assumption that nonnegative cardinality-1 solutions exist in feasible set \mathcal{A} , it so happens,

- columns of *measurement matrix* A , corresponding to any particular solution (to (699)) of cardinality greater than 1, may be deprecated and the problem solved again with those columns missing. Such columns are recursively removed from A until a cardinality-1 solution is found.

Either a solution to problem (699) is cardinality-1 or column indices of A , corresponding to a higher cardinality solution, do not intersect that index corresponding to the cardinality-1 solution. When problem (699) is first solved, in the example of Figure 86, solution is cardinality-2 at the kissing point \bullet on the indicated edge of simplex \mathcal{S} . Imagining that the corresponding cardinality-2 face \mathcal{F} has been removed, then the simplex collapses to a line segment along one of the Cartesian axes. When that line segment kisses line \mathcal{A} , then the cardinality-1 vertex-solution illustrated has been found. A similar argument holds for any orientation of line \mathcal{A} and point of entry to simplex \mathcal{S} .

Because signed compressed sensing problem (468) can be equivalently expressed in a nonnegative variable, as we learned in Example 3.1.3.0.1 (p.209), and because a cardinality-1 constraint in (468) transforms to

a cardinality-1 constraint in its nonnegative equivalent (472), then this cardinality-1 recursive reconstruction algorithm continues to hold for a signed variable as in (468). This cardinality-1 reconstruction algorithm also holds more generally when affine subset \mathcal{A} has any higher dimension $n-m$. But we leave proof of this curious result pending because it is practically useless; by that we mean, it is more efficient to search over the columns of matrix A for a cardinality-1 solution known *a priori* to exist.

4.5.2 constraining cardinality of signed variable

Now consider a feasibility problem equivalent to the classical problem from linear algebra $Ax = b$, but with an upper bound k on cardinality $\|x\|_0$: for vector $b \in \mathcal{R}(A)$

$$\begin{aligned} & \text{find } x \in \mathbb{R}^n \\ & \text{subject to } Ax = b \\ & \|x\|_0 \leq k \end{aligned} \tag{700}$$

where $\|x\|_0 \leq k$ means vector x has at most k nonzero entries; such a vector is presumed existent in the feasible set. *Convex iteration* (§4.5.1) works with a nonnegative variable; absolute value $|x|$ is therefore needed here. We propose that nonconvex problem (700) can be equivalently written as a sequence of convex problems that move the cardinality constraint to the objective:

$$\begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} \quad \langle |x|, y \rangle \\ & \text{subject to } Ax = b \end{aligned} \quad \equiv \quad \begin{aligned} & \underset{x \in \mathbb{R}^n, t \in \mathbb{R}^n}{\text{minimize}} \quad \langle t, y + \varepsilon \mathbf{1} \rangle \\ & \text{subject to } Ax = b \\ & -t \preceq x \preceq t \end{aligned} \tag{701}$$

$$\begin{aligned} & \underset{y \in \mathbb{R}^n}{\text{minimize}} \quad \langle t^*, y + \varepsilon \mathbf{1} \rangle \\ & \text{subject to } 0 \preceq y \preceq \mathbf{1} \\ & y^T \mathbf{1} = n - k \end{aligned} \tag{475}$$

where ε is a relatively small positive constant. This sequence is iterated until a direction vector y is found that makes $|x^*|^T y^*$ vanish. The term $\langle t, \varepsilon \mathbf{1} \rangle$ in (701) is necessary to determine absolute value $|x^*| = t^*$ (§3.1.3) because vector y can have zero-valued entries. By initializing y to $(1-\varepsilon)\mathbf{1}$,

the first iteration of problem (701) is a 1-norm problem (464); *id est*,

$$\begin{aligned} & \underset{x \in \mathbb{R}^n, t \in \mathbb{R}^n}{\text{minimize}} && \langle t, \mathbf{1} \rangle \\ & \text{subject to} && Ax = b \\ & && -t \preceq x \preceq t \end{aligned} \quad \equiv \quad \begin{aligned} & \underset{x \in \mathbb{R}^n}{\text{minimize}} && \|x\|_1 \\ & \text{subject to} && Ax = b \end{aligned} \quad (468)$$

Subsequent iterations of problem (701) engaging cardinality term $\langle t, y \rangle$ can be interpreted as corrections to this 1-norm problem leading to a 0-norm solution; vector y can be interpreted as a direction of search.

4.5.2.1 local convergence

As before (§4.5.1.2), *convex iteration* (701) (475) always converges to a locally optimal solution.

4.5.2.2 simple variations on a signed variable

Several useful equivalents to linear programs (701) (475) are easily devised, but their geometrical interpretation is not as apparent: *e.g.*, equivalent in the limit $\varepsilon \rightarrow 0^+$

$$\begin{aligned} & \underset{x \in \mathbb{R}^n, t \in \mathbb{R}^n}{\text{minimize}} && \langle t, y \rangle \\ & \text{subject to} && Ax = b \\ & && -t \preceq x \preceq t \end{aligned} \quad (702)$$

$$\begin{aligned} & \underset{y \in \mathbb{R}^n}{\text{minimize}} && \langle |x^*|, y \rangle \\ & \text{subject to} && 0 \preceq y \preceq \mathbf{1} \\ & && y^T \mathbf{1} = n - k \end{aligned} \quad (475)$$

We get another equivalent to linear programs (701) (475), in the limit, by interpreting problem (468) as infimum to a vertex-description of the 1-norm ball (Figure 64, Example 3.1.3.0.1, *confer*(467)):

$$\begin{aligned} & \underset{a \in \mathbb{R}^{2n}}{\text{minimize}} && \langle a, y \rangle \\ & \text{subject to} && [A \ -A]a = b \\ & && a \succeq 0 \end{aligned} \quad (703)$$

$$\begin{aligned} & \underset{y \in \mathbb{R}^{2n}}{\text{minimize}} && \langle a^*, y \rangle \\ & \text{subject to} && 0 \preceq y \preceq \mathbf{1} \\ & && y^T \mathbf{1} = 2n - k \end{aligned} \quad (475)$$

where $x^* = [I \ -I]a^*$; from which it may be construed that any vector 1-norm minimization problem has equivalent expression in a nonnegative variable.

4.6 Cardinality and rank constraint examples

4.6.0.0.1 Example. *Projection on an ellipsoid boundary.* [45] [128, §5.1] [218, §2] Consider classical linear equation $Ax = b$ but with a constraint on norm of solution x , given matrices C , fat A , and vector $b \in \mathcal{R}(A)$

$$\begin{aligned} \text{find } & x \in \mathbb{R}^N \\ \text{subject to } & Ax = b \\ & \|Cx\| = 1 \end{aligned} \tag{704}$$

The set $\{x \mid \|Cx\| = 1\}$ (2) describes an ellipsoid boundary (Figure 13). This is a nonconvex problem because solution is constrained to that boundary. Assign

$$G = \begin{bmatrix} Cx \\ 1 \end{bmatrix} [x^T C^T \ 1] = \begin{bmatrix} X & Cx \\ x^T C^T & 1 \end{bmatrix} \triangleq \begin{bmatrix} Cxx^T C^T & Cx \\ x^T C^T & 1 \end{bmatrix} \in \mathbb{S}^{N+1} \tag{705}$$

Any rank-1 solution must have this form. (§B.1.0.2) Ellipsoidally constrained feasibility problem (704) is equivalent to:

$$\begin{aligned} \text{find}_{X \in \mathbb{S}^N} & x \in \mathbb{R}^N \\ \text{subject to } & Ax = b \\ & G = \begin{bmatrix} X & Cx \\ x^T C^T & 1 \end{bmatrix} \\ & (G \succeq 0) \\ & \text{rank } G = 1 \\ & \text{tr } X = 1 \end{aligned} \tag{706}$$

This is transformed to an equivalent convex problem by moving the rank constraint to the objective: We iterate solution of

$$\begin{aligned}
& \underset{X \in \mathbb{S}^N, x \in \mathbb{R}^N}{\text{minimize}} && \langle G, Y \rangle \\
& \text{subject to} && Ax = b \\
& && G = \begin{bmatrix} X & Cx \\ x^T C^T & 1 \end{bmatrix} \succeq 0 \\
& && \text{tr } X = 1
\end{aligned} \tag{707}$$

with

$$\begin{aligned}
& \underset{Y \in \mathbb{S}^{N+1}}{\text{minimize}} && \langle G^*, Y \rangle \\
& \text{subject to} && 0 \preceq Y \preceq I \\
& && \text{tr } Y = N
\end{aligned} \tag{708}$$

until convergence. Direction matrix $Y \in \mathbb{S}^{N+1}$, initially $\mathbf{0}$, controls rank. (1592a) Singular value decomposition $G^* = U\Sigma Q^T \in \mathbb{S}_+^{N+1}$ (§A.6) provides a new direction matrix $Y = U(:, 2:N+1)U(:, 2:N+1)^T$ that optimally solves (708) at each iteration. An optimal solution to (704) is thereby found in a few iterations, making convex problem (707) its equivalent.

It remains possible for the iteration to stall; were a rank-1 G matrix not found. In that case, the current search direction is momentarily reversed with an added random element:

$$Y = -U(:, 2:N+1) * (U(:, 2:N+1)' + \text{randn}(N, 1) * U(:, 1)') \tag{709}$$

in MATLAB notation. This heuristic is quite effective for problem (704) which is exceptionally easy to solve by *convex iteration*.

When $b \notin \mathcal{R}(A)$ then problem (704) must be restated as a projection:

$$\begin{aligned}
& \underset{x \in \mathbb{R}^N}{\text{minimize}} && \|Ax - b\| \\
& \text{subject to} && \|Cx\| = 1
\end{aligned} \tag{710}$$

This is a projection of point b on an ellipsoid boundary because any affine transformation of an ellipsoid remains an ellipsoid. Problem (707) in turn becomes

$$\begin{aligned}
& \underset{X \in \mathbb{S}^N, x \in \mathbb{R}^N}{\text{minimize}} && \langle G, Y \rangle + \|Ax - b\| \\
& \text{subject to} && G = \begin{bmatrix} X & Cx \\ x^T C^T & 1 \end{bmatrix} \succeq 0 \\
& && \text{tr } X = 1
\end{aligned} \tag{711}$$

We iterate this with calculation (708) of direction matrix Y as before until a rank-1 G matrix is found. \square

4.6.0.0.2 Example. *Procrustes problem.* [45]

Example 4.6.0.0.1 is extensible. An orthonormal matrix $Q \in \mathbb{R}^{n \times p}$ is completely characterized by $Q^T Q = I$. Consider the particular case $Q = [x \ y] \in \mathbb{R}^{n \times 2}$ as variable to a Procrustes problem (§C.3): given $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{m \times 2}$

$$\begin{aligned} & \underset{Q \in \mathbb{R}^{n \times 2}}{\text{minimize}} && \|AQ - B\|_F \\ & \text{subject to} && Q^T Q = I \end{aligned} \quad (712)$$

which is nonconvex. By vectorizing matrix Q we can make the assignment:

$$G = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \begin{bmatrix} x^T & y^T & 1 \end{bmatrix} = \begin{bmatrix} X & Z & x \\ Z^T & Y & y \\ x^T & y^T & 1 \end{bmatrix} \triangleq \begin{bmatrix} xx^T & xy^T & x \\ yx^T & yy^T & y \\ x^T & y^T & 1 \end{bmatrix} \in \mathbb{S}^{2n+1} \quad (713)$$

Now orthonormal Procrustes problem (712) can be equivalently restated:

$$\begin{aligned} & \underset{X, Y, Z, x, y}{\text{minimize}} && \|A[x \ y] - B\|_F \\ & \text{subject to} && G = \begin{bmatrix} X & Z & x \\ Z^T & Y & y \\ x^T & y^T & 1 \end{bmatrix} \\ & && (G \succeq 0) \\ & && \text{rank } G = 1 \\ & && \text{tr } X = 1 \\ & && \text{tr } Y = 1 \\ & && \text{tr } Z = 0 \end{aligned} \quad (714)$$

To solve this, we form the convex problem sequence:

$$\begin{aligned} & \underset{X, Y, Z, x, y}{\text{minimize}} && \|A[x \ y] - B\|_F + \langle G, W \rangle \\ & \text{subject to} && G = \begin{bmatrix} X & Z & x \\ Z^T & Y & y \\ x^T & y^T & 1 \end{bmatrix} \succeq 0 \\ & && \text{tr } X = 1 \\ & && \text{tr } Y = 1 \\ & && \text{tr } Z = 0 \end{aligned} \quad (715)$$

and

$$\begin{aligned} & \underset{W \in \mathbb{S}^{2n+1}}{\text{minimize}} && \langle G^*, W \rangle \\ & \text{subject to} && 0 \preceq W \preceq I \\ & && \text{tr } W = 2n \end{aligned} \tag{716}$$

which has an optimal solution W that is known in closed form (p.603). These two problems are iterated until convergence and a rank-1 G matrix is found. A good initial value for direction matrix W is $\mathbf{0}$. Optimal Q^* equals $[x^* \ y^*]$.

Numerically, this Procrustes problem is easy to solve; a solution seems always to be found in one or few iterations. This problem formulation is extensible, of course, to orthogonal (square) matrices Q . \square

4.6.0.0.3 Example. Combinatorial Procrustes problem.

In case $A, B \in \mathbb{R}^n$, when vector $A = \Xi B$ is known to be a permutation of vector B , solution to orthogonal Procrustes problem

$$\begin{aligned} & \underset{X \in \mathbb{R}^{n \times n}}{\text{minimize}} && \|A - XB\|_F \\ & \text{subject to} && X^T = X^{-1} \end{aligned} \tag{1604}$$

is not necessarily a permutation matrix Ξ even though an optimal objective value of 0 is found by the known analytical solution (§C.3). The simplest method of solution finds permutation matrix $X^* = \Xi$ simply by sorting vector B with respect to A .

Instead of sorting, we design two different convex problems each of whose optimal solution is a permutation matrix: one design is based on rank constraint, the other on cardinality. Because permutation matrices are sparse by definition, we depart from a traditional Procrustes problem by instead demanding a vector 1-norm which is known to produce solutions more sparse than Frobenius norm.

There are two principal facts exploited by the first *convex iteration* design (§4.4.1) we propose. Permutation matrices Ξ constitute:

- 1) the set of all nonnegative orthogonal matrices,
- 2) all points extreme to the polyhedron (92) of doubly stochastic matrices.

That means:




Figure 87: Permutation matrix i^{th} column-sum and column-norm constraint in two dimensions when rank constraint is satisfied. Optimal solutions reside at intersection of hyperplane with unit circle.

- 1) norm of each row and column is 1, [4.32](#)

$$\|\Xi(:, i)\| = 1, \quad \|\Xi(i, :)\| = 1, \quad i = 1 \dots n \quad (717)$$

- 2) sum of each nonnegative row and column is 1, ([§2.3.2.0.4](#))

$$\Xi^T \mathbf{1} = \mathbf{1}, \quad \Xi \mathbf{1} = \mathbf{1}, \quad \Xi \geq \mathbf{0} \quad (718)$$

solution via rank constraint

The idea is to individually constrain each column of variable matrix X to have unity norm. Matrix X must also belong to that polyhedron, (92) in the nonnegative orthant, implied by constraints (718); so each row-sum and

^{4.32}This fact would be superfluous were the objective of minimization linear, because the permutation matrices reside at the extreme points of a polyhedron (92) implied by (718). But as posed, only either rows or columns need be constrained to unit norm because matrix orthogonality implies transpose orthogonality. (§B.5.1) Absence of vanishing inner product constraints that help define orthogonality, like $\text{tr } Z = 0$ from Example 4.6.0.0.2, is a consequence of nonnegativity; *id est*, the only orthogonal matrices having exclusively nonnegative entries are permutations of the identity.

column-sum of X must also be unity. It is this combination of nonnegativity, sum, and sum square constraints that extracts the permutation matrices: (Figure 87) given nonzero vectors A, B

$$\begin{aligned}
 & \underset{X \in \mathbb{R}^{n \times n}, G_i \in \mathbb{S}^{n+1}}{\text{minimize}} && \|A - XB\|_1 + w \sum_{i=1}^n \langle G_i, W_i \rangle \\
 & \text{subject to} && G_i = \left[\begin{array}{cc} G_i(1:n, 1:n) & X(:, i) \\ X(:, i)^T & 1 \end{array} \right] \succeq 0 \\
 & && \text{tr } G_i = 2 \\
 & && X^T \mathbf{1} = \mathbf{1} \\
 & && X \mathbf{1} = \mathbf{1} \\
 & && X \geq \mathbf{0}
 \end{aligned} \tag{719}$$

where $w \approx 10$ positively weights the rank regularization term. Optimal solutions G_i^* are key to finding direction matrices W_i for the next iteration of semidefinite programs (719) (720):

$$\begin{aligned}
 & \underset{W_i \in \mathbb{S}^{n+1}}{\text{minimize}} && \langle G_i^*, W_i \rangle \\
 & \text{subject to} && 0 \preceq W_i \preceq I \\
 & && \text{tr } W_i = n
 \end{aligned} \tag{720}$$

Direction matrices thus found lead toward rank-1 matrices G_i^* on subsequent iterations. Constraint on trace of G_i^* normalizes the i^{th} column of X^* to unity because (confer p.371)

$$G_i^* = \begin{bmatrix} X^*(:, i) \\ 1 \end{bmatrix} [X^*(:, i)^T \ 1] \tag{721}$$

at convergence. Binary-valued X^* column entries result from the further sum constraint $X \mathbf{1} = \mathbf{1}$. Columnar orthogonality is a consequence of the further transpose-sum constraint $X^T \mathbf{1} = \mathbf{1}$ in conjunction with nonnegativity constraint $X \geq \mathbf{0}$; but we leave proof of orthogonality an exercise. The optimal objective value is 0 for both semidefinite programs when vectors A and B are related by permutation. In any case, optimal solution X^* becomes a permutation matrix Ξ .

Because there are n direction matrices W_i to find, it can be advantageous to invoke a known closed-form solution for each from page 603. What makes this combinatorial problem more tractable are relatively small semidefinite

constraints in (719). (*confer* (715)) When a permutation A of vector B exists, number of iterations can be as small as 1. But this combinatorial Procrustes problem can be made even more challenging when vector A has repeated entries.

solution via cardinality constraint

Now the idea is to force solution at a vertex of permutation polyhedron (92) by finding a solution of desired sparsity. Because permutation matrix X is n -sparse by assumption, this combinatorial Procrustes problem may instead be formulated as a compressed sensing problem with *convex iteration* on cardinality of vectorized X (§4.5.1): given nonzero vectors A, B

$$\begin{aligned} & \underset{X \in \mathbb{R}^{n \times n}}{\text{minimize}} && \|A - XB\|_1 + w\langle X, Y \rangle \\ & \text{subject to} && X^T \mathbf{1} = \mathbf{1} \\ & && X \mathbf{1} = \mathbf{1} \\ & && X \geq \mathbf{0} \end{aligned} \tag{722}$$

where direction vector Y is an optimal solution to

$$\begin{aligned} & \underset{Y \in \mathbb{R}^{n \times n}}{\text{minimize}} && \langle X^*, Y \rangle \\ & \text{subject to} && \mathbf{0} \leq Y \leq \mathbf{1} \\ & && \mathbf{1}^T Y \mathbf{1} = n^2 - n \end{aligned} \tag{475}$$

each a linear program. In this circumstance, use of closed-form solution for direction vector Y is discouraged. When vector A is a permutation of B , both linear programs have objectives that converge to 0. When vectors A and B are permutations and no entries of A are repeated, optimal solution X^* can be found as soon as the first iteration.

In any case, $X^* = \Xi$ is a permutation matrix. □

4.6.0.0.4 Exercise. *Combinatorial Procrustes constraints.*

Assume that the objective of semidefinite program (719) is 0 at optimality. Prove that the constraints in program (719) are necessary and sufficient to produce a permutation matrix as optimal solution. Alternatively and equivalently, prove those constraints necessary and sufficient to optimally produce a nonnegative orthogonal matrix. ▼

4.6.0.0.5 Example. *Tractable polynomial constraint.*

The ability to handle rank constraints makes polynomial constraints (generally nonconvex) transformable to convex constraints. *All optimization problems having polynomial objective and polynomial constraints can be reformulated as a semidefinite program with a rank-1 constraint.* [252] Suppose we require

$$3 + 2x - xy \leq 0 \tag{723}$$

Assign

$$G = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \begin{bmatrix} x & y & 1 \end{bmatrix} = \begin{bmatrix} X & z \\ z^T & 1 \end{bmatrix} \triangleq \begin{bmatrix} x^2 & xy & x \\ xy & y^2 & y \\ x & y & 1 \end{bmatrix} \in \mathbb{S}^3 \tag{724}$$

The polynomial constraint (723) is equivalent to the constraint set (§B.1.0.2)

$$\begin{aligned} \text{tr}(GA) &\leq 0 \\ G &= \begin{bmatrix} X & z \\ z^T & 1 \end{bmatrix} (\succeq 0) \\ \text{rank } G &= 1 \end{aligned} \tag{725}$$

in symmetric variable matrix $X \in \mathbb{S}^2$ and variable vector $z \in \mathbb{R}^2$ where

$$A = \begin{bmatrix} 0 & -\frac{1}{2} & 1 \\ -\frac{1}{2} & 0 & 0 \\ 1 & 0 & 3 \end{bmatrix} \tag{726}$$

Then the method of *convex iteration* from §4.4.1 is applied to implement the rank constraint. □

4.6.0.0.6 Example. *Boolean vector feasible to $Ax \preceq b$.* (confer §4.2.3.1.1)

Now we consider solution to a discrete problem whose only known analytical method of solution is combinatorial in complexity: given $A \in \mathbb{R}^{M \times N}$ and $b \in \mathbb{R}^M$

$$\begin{aligned} \text{find} \quad & x \in \mathbb{R}^N \\ \text{subject to} \quad & Ax \preceq b \\ & \delta(xx^T) = \mathbf{1} \end{aligned} \tag{727}$$

This nonconvex problem demands a Boolean solution $[x_i = \pm 1, i = 1 \dots N]$.

Assign a rank-1 matrix of variables; symmetric variable matrix X and solution vector x :

$$G = \begin{bmatrix} x \\ 1 \end{bmatrix} [x^T \ 1] = \begin{bmatrix} X & x \\ x^T & 1 \end{bmatrix} \triangleq \begin{bmatrix} xx^T & x \\ x^T & 1 \end{bmatrix} \in \mathbb{S}^{N+1} \quad (728)$$

Then design an equivalent semidefinite feasibility problem to find a Boolean solution to $Ax \preceq b$:

$$\begin{aligned} & \underset{X \in \mathbb{S}^N}{\text{find}} && x \in \mathbb{R}^N \\ & \text{subject to} && Ax \preceq b \\ & && G = \begin{bmatrix} X & x \\ x^T & 1 \end{bmatrix} \\ & && \text{rank } G = 1 \\ & && (G \succeq 0) \\ & && \delta(X) = \mathbf{1} \end{aligned} \quad (729)$$

where $x_i^* \in \{-1, 1\}$, $i=1 \dots N$. The two variables X and x are made dependent via their assignment to rank-1 matrix G . By (1517), an optimal rank-1 matrix G^* must take the form (728).

As before, we regularize the rank constraint by introducing a direction matrix Y into the objective:

$$\begin{aligned} & \underset{X \in \mathbb{S}^N, x \in \mathbb{R}^N}{\text{minimize}} && \langle G, Y \rangle \\ & \text{subject to} && Ax \preceq b \\ & && G = \begin{bmatrix} X & x \\ x^T & 1 \end{bmatrix} \succeq 0 \\ & && \delta(X) = \mathbf{1} \end{aligned} \quad (730)$$

Solution of this semidefinite program is iterated with calculation of the direction matrix Y from semidefinite program (708). At convergence, in the sense (679), convex problem (730) becomes equivalent to nonconvex Boolean problem (727).

Direction matrix Y can be an orthogonal projector having closed-form expression, by (1592a), although *convex iteration* is not a projection method. Given randomized data A and b for a large problem, we find that stalling becomes likely (convergence of the iteration to a positive fixed point

$\langle G^*, Y \rangle$). To overcome this behavior, we introduce a heuristic into the implementation on *Wikimization* that momentarily reverses direction of search (like (709)) upon stall detection. We find that rate of convergence can be sped significantly by detecting stalls early. \square

4.6.0.0.7 Example. *Variable-vector normalization.*

Suppose, within some convex optimization problem, we want vector variables $x, y \in \mathbb{R}^N$ constrained by a nonconvex equality:

$$x\|y\| = y \tag{731}$$

id est, $\|x\| = 1$ and x points in the same direction as $y \neq \mathbf{0}$; e.g.,

$$\begin{aligned} & \underset{x, y}{\text{minimize}} && f(x, y) \\ & \text{subject to} && (x, y) \in \mathcal{C} \\ & && x\|y\| = y \end{aligned} \tag{732}$$

where f is some convex function and \mathcal{C} is some convex set. We can realize the nonconvex equality by constraining rank and adding a regularization term to the objective. Make the assignment:

$$G = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \begin{bmatrix} x^T & y^T & 1 \end{bmatrix} = \begin{bmatrix} X & Z & x \\ Z & Y & y \\ x^T & y^T & 1 \end{bmatrix} \triangleq \begin{bmatrix} xx^T & xy^T & x \\ yx^T & yy^T & y \\ x^T & y^T & 1 \end{bmatrix} \in \mathbb{S}^{2N+1} \tag{733}$$

where $X, Y \in \mathbb{S}^N$, also $Z \in \mathbb{S}^N$ [sic]. Any rank-1 solution must take the form of (733). (§B.1) The problem statement equivalent to (732) is then written

$$\begin{aligned} & \underset{X, Y, Z, x, y}{\text{minimize}} && f(x, y) + \|X - Y\|_F \\ & \text{subject to} && (x, y) \in \mathcal{C} \\ & && G = \begin{bmatrix} X & Z & x \\ Z & Y & y \\ x^T & y^T & 1 \end{bmatrix} \\ & && \text{rank } G = 1 \\ & && (G \succeq 0) \\ & && \text{tr}(X) = 1 \\ & && \delta(Z) \succeq 0 \end{aligned} \tag{734}$$

The trace constraint on X normalizes vector x while the diagonal constraint on Z maintains sign between respective entries of x and y . Regularization term $\|X - Y\|_F$ then makes x equal to y to within a real scalar; (§C.2.0.0.2) in this case, a positive scalar. To make this program solvable by *convex iteration*, as explained in Example 4.4.1.2.2 and other previous examples, we move the rank constraint to the objective

$$\begin{aligned}
 & \underset{X, Y, Z, x, y}{\text{minimize}} && f(x, y) + \|X - Y\|_F + \langle G, W \rangle \\
 & \text{subject to} && (x, y) \in \mathcal{C} \\
 & && G = \begin{bmatrix} X & Z & x \\ Z & Y & y \\ x^T & y^T & 1 \end{bmatrix} \succeq 0 \\
 & && \text{tr}(X) = 1 \\
 & && \delta(Z) \succeq 0
 \end{aligned} \tag{735}$$

by introducing a direction matrix W found from (1592a):

$$\begin{aligned}
 & \underset{W \in \mathbb{S}^{2N+1}}{\text{minimize}} && \langle G^*, W \rangle \\
 & \text{subject to} && 0 \preceq W \preceq I \\
 & && \text{tr } W = 2N
 \end{aligned} \tag{736}$$

This semidefinite program has an optimal solution that is known in closed form. Iteration (735) (736) terminates when $\text{rank } G = 1$ and linear regularization $\langle G, W \rangle$ vanishes to within some numerical tolerance in (735); typically, in two iterations. If function f competes too much with the regularization, positively weighting each regularization term will become required. At convergence, problem (735) becomes a convex equivalent to the original nonconvex problem (732). \square

4.6.0.0.8 Example. FAST MAX CUT. [96]

Let Γ be an n -node graph, and let the arcs (i, j) of the graph be associated with $[\]$ weights a_{ij} . The problem is to find a cut of the largest possible weight, i.e., to partition the set of nodes into two parts $\mathcal{S}, \mathcal{S}'$ in such a way that the total weight of all arcs linking \mathcal{S} and \mathcal{S}' (i.e., with one incident node in \mathcal{S} and the other one in \mathcal{S}' [Figure 88]) is as large as possible. [31, §4.3.3]




Figure 88: A CUT partitions nodes $\{i=1 \dots 16\}$ of this graph into \mathcal{S} and \mathcal{S}' . Linear arcs have circled weights. The problem is to find a cut maximizing total weight of all arcs linking partitions made by the cut.

Literature on the MAX CUT problem is vast because this problem has elegant primal and dual formulation, its solution is very difficult, and there exist many commercial applications; *e.g.*, semiconductor design [108], quantum computing [346].

Our purpose here is to demonstrate how iteration of two simple convex problems can quickly converge to an optimal solution of the MAX CUT problem with a 98% success rate, on average.^{4.33} MAX CUT is stated:

$$\begin{aligned} & \underset{x}{\text{maximize}} \quad \sum_{1 \leq i < j \leq n} a_{ij} (1 - x_i x_j) \frac{1}{2} \\ & \text{subject to} \quad \delta(xx^T) = \mathbf{1} \end{aligned} \tag{737}$$

where $[a_{ij}]$ are real arc weights, and binary vector $x = [x_i] \in \mathbb{R}^n$ corresponds to the n nodes; specifically,

$$\begin{aligned} \text{node } i \in \mathcal{S} & \Leftrightarrow x_i = 1 \\ \text{node } i \in \mathcal{S}' & \Leftrightarrow x_i = -1 \end{aligned} \tag{738}$$

If nodes i and j have the same binary value x_i and x_j , then they belong to the same partition and contribute nothing to the cut. Arc (i, j) traverses the cut, otherwise, adding its weight a_{ij} to the cut.

MAX CUT statement (737) is the same as, for $A = [a_{ij}] \in \mathbb{S}^n$

$$\begin{aligned} & \underset{x}{\text{maximize}} \quad \frac{1}{4} \langle \mathbf{1}\mathbf{1}^T - xx^T, A \rangle \\ & \text{subject to} \quad \delta(xx^T) = \mathbf{1} \end{aligned} \tag{739}$$

Because of Boolean assumption $\delta(xx^T) = \mathbf{1}$

$$\langle \mathbf{1}\mathbf{1}^T - xx^T, A \rangle = \langle xx^T, \delta(A\mathbf{1}) - A \rangle \tag{740}$$

so problem (739) is the same as

$$\begin{aligned} & \underset{x}{\text{maximize}} \quad \frac{1}{4} \langle xx^T, \delta(A\mathbf{1}) - A \rangle \\ & \text{subject to} \quad \delta(xx^T) = \mathbf{1} \end{aligned} \tag{741}$$

This MAX CUT problem is combinatorial (nonconvex).

^{4.33}We term our solution to MAX CUT *fast* because we sacrifice a little accuracy to achieve speed; *id est*, only about two or three convex iterations, achieved by heavily weighting a rank regularization term.

Because an estimate of upper bound to MAX CUT is needed to ascertain convergence when vector x has large dimension, we digress to derive the dual problem: Directly from (741), the *Lagrangian* is [53, §5.1.5] (1331)

$$\begin{aligned} \mathfrak{L}(x, \nu) &= \frac{1}{4}\langle xx^T, \delta(A\mathbf{1}) - A \rangle + \langle \nu, \delta(xx^T) - \mathbf{1} \rangle \\ &= \frac{1}{4}\langle xx^T, \delta(A\mathbf{1}) - A \rangle + \langle \delta(\nu), xx^T \rangle - \langle \nu, \mathbf{1} \rangle \\ &= \frac{1}{4}\langle xx^T, \delta(A\mathbf{1} + 4\nu) - A \rangle - \langle \nu, \mathbf{1} \rangle \end{aligned} \tag{742}$$

where quadratic $x^T(\delta(A\mathbf{1} + 4\nu) - A)x$ has supremum 0 if $\delta(A\mathbf{1} + 4\nu) - A$ is negative semidefinite, and has supremum ∞ otherwise. The finite supremum of dual function

$$g(\nu) = \sup_x \mathfrak{L}(x, \nu) = \begin{cases} -\langle \nu, \mathbf{1} \rangle, & A - \delta(A\mathbf{1} + 4\nu) \succeq 0 \\ \infty & \text{otherwise} \end{cases} \tag{743}$$

is chosen to be the objective of minimization to dual (convex) problem

$$\begin{aligned} &\underset{\nu}{\text{minimize}} && -\nu^T \mathbf{1} \\ &\text{subject to} && A - \delta(A\mathbf{1} + 4\nu) \succeq 0 \end{aligned} \tag{744}$$

whose optimal value provides a least upper bound to MAX CUT, but is not tight ($\frac{1}{4}\langle xx^T, \delta(A\mathbf{1}) - A \rangle < g(\nu)$, duality gap is nonzero). [137] In fact, we find that the bound’s variance with problem instance is too large to be useful for this problem; thus ending our digression. ^{4.34}

To transform MAX CUT to its convex equivalent, first define

$$X = xx^T \in \mathbb{S}^n \tag{749}$$

then MAX CUT (741) becomes

$$\begin{aligned} &\underset{X \in \mathbb{S}^n}{\text{maximize}} && \frac{1}{4}\langle X, \delta(A\mathbf{1}) - A \rangle \\ &\text{subject to} && \delta(X) = \mathbf{1} \\ &&& (X \succeq 0) \\ &&& \text{rank } X = 1 \end{aligned} \tag{745}$$

^{4.34}Taking the dual of dual problem (744) would provide (745) but without the rank constraint. [130]

whose rank constraint can be regularized as in

$$\begin{aligned} & \underset{X \in \mathbb{S}^n}{\text{maximize}} && \frac{1}{4} \langle X, \delta(A\mathbf{1}) - A \rangle - w \langle X, W \rangle \\ & \text{subject to} && \delta(X) = \mathbf{1} \\ & && X \succeq 0 \end{aligned} \tag{746}$$

where $w \approx 1000$ is a nonnegative fixed weight, and W is a direction matrix determined from

$$\begin{aligned} \sum_{i=2}^n \lambda(X^*)_i &= \underset{W \in \mathbb{S}^n}{\text{minimize}} && \langle X^*, W \rangle \\ & \text{subject to} && 0 \preceq W \preceq I \\ & && \text{tr } W = n - 1 \end{aligned} \tag{1592a}$$

which has an optimal solution that is known in closed form. These two problems (746) and (1592a) are iterated until convergence as defined on page 282.

Because convex problem statement (746) is so elegant, it is numerically solvable for large binary vectors within reasonable time.^{4.35} To test our convex iterative method, we compare an optimal convex result to an actual solution of the MAX CUT problem found by performing a brute force combinatorial search of (741)^{4.36} for a tight upper bound. Search-time limits binary vector lengths to 24 bits (about five days CPU time). Accuracy obtained, 98%, is independent of binary vector length (12, 13, 20, 24) when averaged over more than 231 problem instances including planar, randomized, and toroidal graphs.^{4.37} When failure occurred, large and small errors were manifest. That same 98% average accuracy is presumed maintained when binary vector length is further increased. A MATLAB program is provided on [Wikimization](#). \square

^{4.35}We solved for a length-250 binary vector in only a few minutes and convex iterations on a Dell Precision computer, model M90.

^{4.36}more computationally intensive than the proposed *convex iteration* by many orders of magnitude. Solving MAX CUT by searching over all binary vectors of length 100, for example, would occupy a contemporary supercomputer for a million years.

^{4.37}Existence of a polynomial-time approximation to MAX CUT with accuracy better than 94.11% would refute proof of NP-hardness, which Håstad believes to be highly unlikely. [160, thm.8.2] [161]

4.6.0.0.9 Example. *Cardinality/rank problem.*

d'Aspremont *et alii* [83] propose approximating a positive semidefinite matrix $A \in \mathbb{S}_+^N$ by a rank-one matrix having a constraint on cardinality c : for $0 < c < N$

$$\begin{aligned} & \underset{z}{\text{minimize}} && \|A - zz^T\|_F \\ & \text{subject to} && \text{card } z \leq c \end{aligned} \tag{747}$$

which, they explain, is a hard problem equivalent to

$$\begin{aligned} & \underset{x}{\text{maximize}} && x^T A x \\ & \text{subject to} && \|x\| = 1 \\ & && \text{card } x \leq c \end{aligned} \tag{748}$$

where $z \triangleq \sqrt{\lambda} x$ and where optimal solution x^* is a *principal eigenvector* (1586) (§A.5) of A and $\lambda = x^{*T} A x^*$ is the *principal eigenvalue* [139, p.331] when c is true cardinality of that eigenvector. This is *principal component analysis* with a cardinality constraint which controls solution sparsity. Define the matrix variable

$$X \triangleq xx^T \in \mathbb{S}^N \tag{749}$$

whose desired rank is 1, and whose desired diagonal cardinality

$$\text{card } \delta(X) \equiv \text{card } x \tag{750}$$

is equivalent to cardinality c of vector x . Then we can transform cardinality problem (748) to an equivalent problem in new variable X :^{4.38}

$$\begin{aligned} & \underset{X \in \mathbb{S}^N}{\text{maximize}} && \langle X, A \rangle \\ & \text{subject to} && \langle X, I \rangle = 1 \\ & && (X \succeq 0) \\ & && \text{rank } X = 1 \\ & && \text{card } \delta(X) \leq c \end{aligned} \tag{751}$$

We transform problem (751) to an equivalent convex problem by introducing two direction matrices into regularization terms: W to achieve

^{4.38}A semidefiniteness constraint $X \succeq 0$ is not required, theoretically, because positive semidefiniteness of a rank-1 matrix is enforced by symmetry. (Theorem A.3.1.0.7)

desired cardinality $\text{card } \delta(X)$, and Y to find an approximating rank-one matrix X :

$$\begin{aligned} & \underset{X \in \mathbb{S}^N}{\text{maximize}} && \langle X, A - w_1 Y \rangle - w_2 \langle \delta(X), \delta(W) \rangle \\ & \text{subject to} && \langle X, I \rangle = 1 \\ & && X \succeq 0 \end{aligned} \tag{752}$$

where w_1 and w_2 are positive scalars respectively weighting $\text{tr}(XY)$ and $\delta(X)^T \delta(W)$ just enough to insure that they vanish to within some numerical precision, where direction matrix Y is an optimal solution to semidefinite program

$$\begin{aligned} & \underset{Y \in \mathbb{S}^N}{\text{minimize}} && \langle X^*, Y \rangle \\ & \text{subject to} && 0 \preceq Y \preceq I \\ & && \text{tr } Y = N - 1 \end{aligned} \tag{753}$$

and where diagonal direction matrix $W \in \mathbb{S}^N$ optimally solves linear program

$$\begin{aligned} & \underset{W = \delta^2(W)}{\text{minimize}} && \langle \delta(X^*), \delta(W) \rangle \\ & \text{subject to} && 0 \preceq \delta(W) \preceq \mathbf{1} \\ & && \text{tr } W = N - c \end{aligned} \tag{754}$$

Both direction matrix programs are derived from (1592a) whose analytical solution is known but is not necessarily unique. We emphasize (*confer* p.282): because this iteration (752) (753) (754) (initial $Y, W = \mathbf{0}$) is not a projection method, success relies on existence of matrices in the feasible set of (752) having desired rank and diagonal cardinality. In particular, the feasible set of convex problem (752) is a Fantope (84) whose extreme points constitute the set of all normalized rank-one matrices; among those are found rank-one matrices of any desired diagonal cardinality.

Convex problem (752) is neither a relaxation of cardinality problem (748); instead, problem (752) is a convex equivalent to (748) at convergence of iteration (752) (753) (754). Because the feasible set of convex problem (752) contains all normalized (§B.1) symmetric rank-one matrices of every nonzero diagonal cardinality, a constraint too low or high in cardinality c will not prevent solution. An optimal rank-one solution X^* , whose diagonal cardinality is equal to cardinality of a principal eigenvector of matrix A , will produce the lowest residual Frobenius norm (to within machine noise processes) in the original problem statement (747). \square

phantom(256)

Figure 89: Shepp-Logan phantom from MATLAB *image processing toolbox*.**4.6.0.0.10 Example.** *Compressive sampling of a phantom.*

In summer of 2004, Candès, Romberg, & Tao [61] and Donoho [101] released papers on perfect signal reconstruction from samples that stand in violation of Shannon's classical sampling theorem. These defiant signals are assumed sparse inherently or under some sparsifying affine transformation. Essentially, they proposed *sparse sampling theorems* asserting average sample rate independent of signal bandwidth and less than Shannon's rate.

MINIMUM SAMPLING RATE:

- OF Ω -BANDLIMITED SIGNAL: 2Ω (Shannon)
- OF k -SPARSE LENGTH- n SIGNAL: $k \log_2(1+n/k)$ (Candès/Donoho)

(Figure 83). Certainly, much was already known about nonuniform or random sampling [32] [184] and about subsampling or *multirate systems* [80] [317]. Vetterli, Marziliano, & Blu [326] had congealed a theory of noiseless signal reconstruction, in May 2001, from samples that violate the Shannon rate. They anticipated the sparsifying transform by recognizing: it is the *innovation* (onset) of functions constituting a (not necessarily bandlimited) signal that determines minimum sampling rate for perfect reconstruction. Average onset (*sparsity*) Vetterli *et alii* call *rate of innovation*. The vector inner-products that Candès/Donoho call *samples* or *measurements*, Vetterli calls *projections*. From those projections Vetterli demonstrates reconstruction (by digital signal processing

and “root finding”) of a Dirac comb, the very same prototypical signal from which Candès probabilistically derives minimum sampling rate (*Compressive Sampling and Frontiers in Signal Processing*, University of Minnesota, June 6, 2007). Combining their terminology, we paraphrase a sparse sampling theorem:

- Minimum sampling rate, asserted by Candès/Donoho, is proportional to Vetterli’s rate of innovation (a.k.a, *information rate, degrees of freedom, ibidem* June 5, 2007).

What distinguishes these researchers are their methods of reconstruction.

Properties of the 1-norm were also well understood by June 2004 finding applications in *deconvolution* of linear systems [74], *penalized linear regression (Lasso)* [282], and *basis pursuit* [188]. But never before had there been a formalized and rigorous sense that perfect reconstruction were possible by convex optimization of 1-norm when information lost in a subsampling process became nonrecoverable by classical methods. Donoho named this discovery *compressed sensing* to describe a nonadaptive perfect reconstruction method by means of linear programming. By the time Candès’ and Donoho’s landmark papers were finally published by IEEE in 2006, compressed sensing was old news that had spawned intense research which still persists; notably, from prominent members of the *wavelet* community.

Reconstruction of the *Shepp-Logan phantom* (Figure 89), from a severely aliased image (Figure 91) obtained by Magnetic Resonance Imaging (MRI), was the *impetus* driving Candès’ quest for a sparse sampling theorem. He realized that line segments appearing in the aliased image were regions of high *total variation*. There is great motivation, in the medical community, to apply compressed sensing to MRI because it translates to reduced scan-time which brings great technological and physiological benefits. MRI is now about 35 years old, beginning in 1973 with Nobel laureate Paul Lauterbur from Stony Brook. There has been much progress in MRI and compressed sensing since 2004, but there have also been two beacons pointing toward abandonment of the 1-norm (indigenous to reconstruction by compressed sensing) in favor of criteria closer to 0-norm because of a correspondingly smaller number of measurements required to accurately reconstruct a sparse signal:^{4.39}

^{4.39}Efficient techniques continually emerge that urge abandonment of the 1-norm criterion; [71] *e.g.*, five techniques for compressed sensing are compared in [33] demonstrating that 1-norm performance limits for cardinality minimization can be reliably exceeded.

- June 2004: Candès, Romberg, & Tao publish in cyberspace: 5481 complex samples (22 radial lines, ≈ 256 complex samples per) to reconstruct noiseless 256×256 -pixel Shepp-Logan phantom by 1-norm minimization of an image-gradient integral estimate called *total variation*; *id est*, 8.4% subsampling of 65536 data. [61, §1.1] [60, §3.2]
- March 2006: Chartrand discovers^{4.40} that reconstruction of the Shepp-Logan phantom is possible with only 2521 complex samples (10 radial lines, Figure 90); 3.8% subsampled data input to a (nonconvex) $\frac{1}{2}$ -norm total-variation minimization. [67, §IIIA]
- April 2007: Trzasko & Manduca recover the phantom using same subsampling as Chartrand, but realize an accurate 0-norm total-variation minimization algorithm that is faster^{4.41} than prior methods. [315, §IID] [316]

Passage of these few years witnessed an algorithmic speedup and dramatic reduction in minimum number of samples required for perfect reconstruction of the noiseless Shepp-Logan phantom. But minimization of total variation is ideally suited to recovery of any piecewise-constant image, like a phantom, because the gradient of such images is highly sparse by design.

There is no inherent characteristic of real-life MRI images that would make reasonable an expectation of sparse gradient. Sparsification of an image-gradient tends to preserve edges. Then minimization of total variation seeks an image having fewest edges. There is no deeper theoretical foundation than that. When applied to human brain scan or angiogram, with as much as 20% of 256×256 Fourier samples, we have observed^{4.42} a 30dB image/(reconstruction)error ratio^{4.43} barrier that seems impenetrable by

^{4.40}*I try using $p = \frac{1}{2}$ in place of $p = 1$ [-norm] to reconstruct the Shepp-Logan phantom, and am very surprised to see how well it works. This is by alternating gradient descent with projection onto the constraint. In hindsight, one aspect appears to have been crucial to its success: regularization of the cusp with a relatively large epsilon, then decreasing epsilon after convergence. At the time I did this solely to try to speed up convergence, based on experience with total-variation minimization algorithms (like lagged diffusivity). But it now seems to be what allows the algorithm to avoid local minima, as these are filled in by the larger epsilon.*
 –Rick Chartrand

^{4.41}Their MATLAB code runs in a minute, on an Intel-based dual-processor 2006 vintage laptop computer, capable of recovery at 50dB image/error from 3.2% subsampled data.

^{4.42}Work on real-life images was performed in conjunction with Christine Law at Lucas Center for Imaging, Stanford University.

^{4.43}Noise considered here is due only to the reconstruction process itself; *id est*, noise in excess of that produced by the best reconstruction of an image from a complete set

the total-variation objective. Total-variation minimization has met with moderate success, in retrospect, only because some medical images are moderately piecewise-constant signals. One simply hopes a reconstruction, that is in some sense equal to a known subset of samples and whose gradient is most sparse, is that unique image we seek.^{4.44}

The total-variation objective, operating on an image, is expressible as norm of a linear transformation (773). It is natural to ask whether there exist other sparsifying transforms that might break the real-life 30dB barrier (@20% 256×256 data) in MRI. There has been much research into application of wavelets, discrete cosine transform (DCT), randomized orthogonal bases, splines, *etcetera*, but with suspiciously little focus on objective measures like image/error or illustration of difference images; the predominant basis of comparison instead being subjectively visual (Duensing & Huang, ISMRM Toronto 2008).^{4.45} Despite choice of transform, there seems yet to have been a breakthrough of the 30dB barrier. Application of compressed sensing to MRI, therefore, remains fertile in 2008 for continued research.

Lagrangian form of compressed sensing in imaging

We now repeat Candès' image reconstruction experiment from 2004 [61] which led to discovery of sparse sampling theorems. But we achieve perfect reconstruction with an algorithm based on vanishing gradient of the compressed sensing problem's Lagrangian, which is computationally efficient. Our *contraction method* (p.339) is faster than contemporary methods also

of samples in the sense of Shannon. At less than 30dB image/error, artifacts generally remain visible to the naked eye. We estimate about 50dB is required to eliminate noticeable distortion in a visual A/B comparison.

^{4.44}In vascular radiology, diagnoses are almost exclusively based on morphology of vessels and, in particular, presence of stenoses. There is a compelling argument for total-variation reconstruction of magnetic resonance angiogram because it helps isolate structures of particular interest.

^{4.45}*I have never calculated the PSNR of these reconstructed images* [of Barbara].

–Jean-Luc Starck

The sparsity of the image is the percentage of transform coefficients sufficient for diagnostic-quality reconstruction. Of course the term "diagnostic quality" is subjective. ... I have yet to see an "objective" measure of image quality. Difference images, in my experience, definitely do not tell the whole story. Often I would show people some of my results and get mixed responses, but when I add artificial Gaussian noise to an image, often people say that it looks better.

–Michael Lustig

because matrix multiplications are replaced by fast Fourier transforms and number of constraints is cut in half by sampling symmetrically. *Convex iteration* for cardinality minimization (§4.5) is incorporated which allows perfect reconstruction of a phantom at 4.1% subsampling rate; 50% Candès' rate. By making neighboring-pixel selection adaptive, *convex iteration* reduces image-gradient sparsity of the Shepp-Logan phantom to 1.9%; 33% lower than previously reported.

We demonstrate application of image-gradient sparsification to the $n \times n = 256 \times 256$ Shepp-Logan phantom, simulating idealized acquisition of MRI data by radial sampling in the Fourier domain (Figure 90).^{4,46} Define a Nyquist-centric *discrete Fourier transform* (DFT) matrix

$$F \triangleq \begin{bmatrix} 1 & 1 & 1 & 1 & \cdots & 1 \\ 1 & e^{-j2\pi/n} & e^{-j4\pi/n} & e^{-j6\pi/n} & \cdots & e^{-j(n-1)2\pi/n} \\ 1 & e^{-j4\pi/n} & e^{-j8\pi/n} & e^{-j12\pi/n} & \cdots & e^{-j(n-1)4\pi/n} \\ 1 & e^{-j6\pi/n} & e^{-j12\pi/n} & e^{-j18\pi/n} & \cdots & e^{-j(n-1)6\pi/n} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & e^{-j(n-1)2\pi/n} & e^{-j(n-1)4\pi/n} & e^{-j(n-1)6\pi/n} & \cdots & e^{-j(n-1)^2 2\pi/n} \end{bmatrix} \frac{1}{\sqrt{n}} \in \mathbb{C}^{n \times n} \quad (755)$$

a symmetric (nonHermitian) unitary matrix characterized

$$\begin{aligned} F &= F^T \\ F^{-1} &= F^H \end{aligned} \quad (756)$$

Denoting an unknown image $\mathcal{U} \in \mathbb{R}^{n \times n}$, its two-dimensional discrete Fourier transform \mathfrak{F} is

$$\mathfrak{F}(\mathcal{U}) \triangleq F\mathcal{U}F \quad (757)$$

hence the inverse discrete transform

$$\mathcal{U} = F^H \mathfrak{F}(\mathcal{U}) F^H \quad (758)$$

From §A.1.1 no.31 we have a vectorized two-dimensional DFT via Kronecker product \otimes

$$\text{vec } \mathfrak{F}(\mathcal{U}) \triangleq (F \otimes F) \text{vec } \mathcal{U} \quad (759)$$

and from (758) its inverse [145, p.24]

$$\text{vec } \mathcal{U} = (F^H \otimes F^H)(F \otimes F) \text{vec } \mathcal{U} = (F^H F \otimes F^H F) \text{vec } \mathcal{U} \quad (760)$$




Figure 90: MRI radial sampling pattern, in DC-centric Fourier domain, representing 4.1% (10 lines) subsampled data. Only half of these complex samples, in theory, need be acquired for a real image because of conjugate symmetry. Due to MRI machine imperfections, samples are generally taken over full extent of each radial line segment. MRI acquisition time is proportional to number of lines.

Idealized radial sampling in the Fourier domain can be simulated by Hadamard product \circ with a binary mask $\Phi \in \mathbb{R}^{n \times n}$ whose nonzero entries could, for example, correspond with the radial line segments in Figure 90. To make the mask Nyquist-centric, like DFT matrix F , define a circulant [147] symmetric permutation matrix^{4.47}

$$\Theta \triangleq \begin{bmatrix} \mathbf{0} & I \\ I & \mathbf{0} \end{bmatrix} \in \mathbb{S}^n \quad (761)$$

Then given subsampled Fourier domain (MRI \mathbf{k} -space) measurements in incomplete $K \in \mathbb{C}^{n \times n}$, we might constrain $\mathfrak{F}(\mathcal{U})$ thus:

$$\Theta \Phi \Theta \circ F \mathcal{U} F = K \quad (762)$$

and in vector form, (36) (1675)

$$\delta(\text{vec } \Theta \Phi \Theta)(F \otimes F) \text{vec } \mathcal{U} = \text{vec } K \quad (763)$$

^{4.46} \mathbf{k} -space is conventional acquisition terminology indicating domain of the continuous raw data provided by an MRI machine. An image is reconstructed by inverse discrete Fourier transform of that data sampled on a Cartesian grid in two dimensions.

^{4.47} MATLAB `fftshift()`

Because measurements K are complex, there are actually twice the number of equality constraints as there are measurements.

We can cut that number of constraints in half via vertical and horizontal mask Φ symmetry which forces the imaginary inverse transform to $\mathbf{0}$: The inverse subsampled transform in matrix form is

$$F^H(\Theta\Phi\Theta \circ F\mathcal{U}F)F^H = F^HKF^H \tag{764}$$

and in vector form

$$(F^H \otimes F^H)\delta(\text{vec } \Theta\Phi\Theta)(F \otimes F) \text{vec } \mathcal{U} = (F^H \otimes F^H) \text{vec } K \tag{765}$$

later abbreviated

$$P \text{vec } \mathcal{U} = f \tag{766}$$

where

$$P \triangleq (F^H \otimes F^H)\delta(\text{vec } \Theta\Phi\Theta)(F \otimes F) \in \mathbb{C}^{n^2 \times n^2} \tag{767}$$

Because of idempotence $P = P^2$, P is a projection matrix. Because of its Hermitian symmetry [145, p.24]

$$P = (F^H \otimes F^H)\delta(\text{vec } \Theta\Phi\Theta)(F \otimes F) = (F \otimes F)^H \delta(\text{vec } \Theta\Phi\Theta)(F^H \otimes F^H)^H = P^H \tag{768}$$

P is an orthogonal projector.^{4.48} $P \text{vec } \mathcal{U}$ is real when P is real; *id est*, when for positive even integer n

$$\Phi = \begin{bmatrix} \Phi_{11} & \Phi(1, 2:n)\Xi \\ \Xi\Phi(2:n, 1) & \Xi\Phi(2:n, 2:n)\Xi \end{bmatrix} \in \mathbb{R}^{n \times n} \tag{769}$$

where $\Xi \in \mathbb{S}^{n-1}$ is the order-reversing permutation matrix (1619). In words, this necessary and sufficient condition on Φ (for a real inverse subsampled transform [251, p.53]) demands vertical symmetry about row $\frac{n}{2}+1$ and horizontal symmetry^{4.49} about column $\frac{n}{2}+1$.

^{4.48} (767) is a diagonalization of matrix P whose binary eigenvalues are $\delta(\text{vec } \Theta\Phi\Theta)$ while the corresponding eigenvectors constitute the columns of unitary matrix $F^H \otimes F^H$.

^{4.49}This condition on Φ applies to both DC- and Nyquist-centric DFT matrices.




Figure 91: Aliasing of Shepp-Logan phantom in Figure 89 resulting from \mathbf{k} -space subsampling pattern in Figure 90. This image is real because binary mask Φ is vertically and horizontally symmetric. It is remarkable that the phantom can be reconstructed, by *convex iteration*, given only $\mathcal{U}_0 = \text{vec}^{-1} f$.

Define

$$\Delta \triangleq \begin{bmatrix} 1 & 0 & & & \mathbf{0} \\ -1 & 1 & 0 & & \\ & -1 & 1 & \ddots & \\ & & \ddots & \ddots & \ddots \\ & & & \ddots & 1 & 0 \\ \mathbf{0}^T & & & & -1 & 1 \end{bmatrix} \in \mathbb{R}^{n \times n} \quad (770)$$

Express an image-gradient estimate

$$\nabla \mathcal{U} \triangleq \begin{bmatrix} \mathcal{U} \Delta \\ \mathcal{U} \Delta^T \\ \Delta \mathcal{U} \\ \Delta^T \mathcal{U} \end{bmatrix} \in \mathbb{R}^{4n \times n} \quad (771)$$

that is a simple first-order difference of neighboring pixels (Figure 92) to the right, left, above, and below.^{4.50} By §A.1.1 no.31, its vectorization:

^{4.50}There is significant improvement in reconstruction quality by augmentation of a normally two-dimensional image-gradient to a four-dimensional estimate per pixel by

for $\Psi_i \in \mathbb{R}^{n^2 \times n^2}$

$$\text{vec } \nabla \mathcal{U} = \begin{bmatrix} \Delta^T \otimes I \\ \Delta \otimes I \\ I \otimes \Delta \\ I \otimes \Delta^T \end{bmatrix} \text{vec } \mathcal{U} \triangleq \begin{bmatrix} \Psi_1 \\ \Psi_1^T \\ \Psi_2 \\ \Psi_2^T \end{bmatrix} \text{vec } \mathcal{U} \triangleq \Psi \text{vec } \mathcal{U} \in \mathbb{R}^{4n^2} \quad (772)$$

where $\Psi \in \mathbb{R}^{4n^2 \times n^2}$. A total-variation minimization for reconstructing MRI image \mathcal{U} , that is known suboptimal [62], may be concisely posed

$$\begin{aligned} & \underset{\mathcal{U}}{\text{minimize}} && \|\Psi \text{vec } \mathcal{U}\|_1 \\ & \text{subject to} && P \text{vec } \mathcal{U} = f \end{aligned} \quad (773)$$

where

$$f = (F^H \otimes F^H) \text{vec } K \in \mathbb{C}^{n^2} \quad (774)$$

is the known inverse subsampled Fourier data (a vectorized aliased image, Figure 91), and where a norm of image-gradient $\nabla \mathcal{U}$ is equivalently expressed as norm of a linear transformation $\Psi \text{vec } \mathcal{U}$.

Although this simple problem statement (773) is equivalent to a linear program (§3.1.3), its numerical solution is beyond the capability of even the most highly regarded of contemporary commercial solvers.^{4.51} Our only recourse is to recast the problem in Lagrangian form and write customized code to solve it: [315, §IIIA] [222, p.398] [53, p.244]

$$\begin{aligned} & \underset{\mathcal{U}}{\text{minimize}} && \langle |\Psi \text{vec } \mathcal{U}|, y \rangle && \text{(a)} \\ & \text{subject to} && P \text{vec } \mathcal{U} = f && \\ & && \equiv && \\ & \underset{\mathcal{U}}{\text{minimize}} && \langle |\Psi \text{vec } \mathcal{U}|, y \rangle + \frac{1}{2} \lambda \|P \text{vec } \mathcal{U} - f\|_2^2 && \text{(b)} \end{aligned} \quad (775)$$

where multiobjective parameter $\lambda \in \mathbb{R}_+$ is quite large ($\lambda \approx 1\text{E}8$) so as to enforce the equality constraint: $P \text{vec } \mathcal{U} - f = \mathbf{0} \Leftrightarrow \|P \text{vec } \mathcal{U} - f\|_2^2 = 0$ (§A.7.1). We

inclusion of two polar directions. We find small improvement on real-life images, $\approx 1\text{dB}$ empirically, by further augmentation with diagonally adjacent pixel differences.

^{4.51} for images as small as 128×128 pixels. Obstacle to numerical solution is not a computer resource: *e.g.*, execution time, memory. The obstacle is, in fact, inadequate numerical precision. Even when all dependent equality constraints are manually removed, the best commercial solvers fail simply because computer numerics become nonsense; *id est*, numerical errors enter significant digits and the algorithm exits prematurely, loops indefinitely, or produces an infeasible solution.




Figure 92: Neighboring-pixel candidates from [316] for estimating image-gradient. Our implementation selects adaptively from darkest four • about central.

introduce a direction vector $y \in \mathbb{R}_+^{4n^2}$ as part of a *convex iteration* (§4.5.2) to overcome that known suboptimal minimization of image-gradient cardinality: *id est*, there exists a vector y^* having entries $y_i^* \in \{0, 1\}$ such that

$$\begin{aligned} \underset{\mathcal{U}}{\text{minimize}} \quad & \|\Psi \text{vec} \mathcal{U}\|_0 \\ \text{subject to} \quad & P \text{vec} \mathcal{U} = f \end{aligned} \quad \equiv \quad \underset{\mathcal{U}}{\text{minimize}} \quad \langle |\Psi \text{vec} \mathcal{U}|, y^* \rangle + \frac{1}{2} \lambda \|P \text{vec} \mathcal{U} - f\|_2^2 \quad (776)$$

Existence of such a y^* , complementary to an optimal vector $\Psi \text{vec} \mathcal{U}^*$, is obvious by definition of global optimality $\langle |\Psi \text{vec} \mathcal{U}^*|, y^* \rangle = 0$ (691) under which a cardinality- c optimal objective is assumed to exist.

Because (775b) is an unconstrained convex problem, a zero in the objective function's gradient is necessary and sufficient for optimality (§2.13.3); *id est*, (§D.2.1)

$$\Psi^T \delta(y) \text{sgn}(\Psi \text{vec} \mathcal{U}) + \lambda P^H (P \text{vec} \mathcal{U} - f) = \mathbf{0} \quad (777)$$

Because of P idempotence and Hermitian symmetry and $\text{sgn}(\kappa) = \kappa/|\kappa|$, this is equivalent to

$$\lim_{\epsilon \rightarrow 0} (\Psi^T \delta(y) \delta(|\Psi \text{vec} \mathcal{U}| + \epsilon \mathbf{1})^{-1} \Psi + \lambda P) \text{vec} \mathcal{U} = \lambda P f \quad (778)$$

where small positive constant $\epsilon \in \mathbb{R}_+$ has been introduced for invertibility. When small enough for practical purposes^{4.52} ($\epsilon \approx 1\text{E-}3$), we may ignore the limiting operation. Then by Weyl's theorem (§A.3.1.0.2) the mapping, for $\lambda > 1$ and $0 \preceq y \preceq \mathbf{1}$

$$\text{vec} \mathcal{U}_{t+1} = (\Psi^T \delta(y) \delta(|\Psi \text{vec} \mathcal{U}_t| + \epsilon \mathbf{1})^{-1} \Psi + \lambda P)^{-1} \lambda P f \quad (779)$$

is a *contraction* in \mathcal{U}_t that can be solved recursively in t for its unique *fixed point*; *id est*, until $\mathcal{U}_{t+1} \rightarrow \mathcal{U}_t$. [201, p.300] [181, p.155] [315, §IIIA] Calculating this inversion directly is not possible for large matrices on contemporary computers because of numerical precision, so instead we apply the *conjugate gradient* method of solution to

$$(\Psi^T \delta(y) \delta(|\Psi \text{vec} \mathcal{U}_t| + \epsilon \mathbf{1})^{-1} \Psi + \lambda P) \text{vec} \mathcal{U}_{t+1} = \lambda P f \quad (780)$$

which is linear in \mathcal{U}_{t+1} at each recursion in the [MATLAB program](#).^{4.53}

Observe that P (767), in the equality constraint from problem (775a), is not a fat matrix.^{4.54} Although number of Fourier samples taken is equal to the number of nonzero entries in binary mask Φ , matrix P is square but never actually formed during computation. Rather, a two-dimensional fast Fourier transform of \mathcal{U} is computed followed by masking with $\Theta\Phi\Theta$ and then an inverse fast Fourier transform. This technique significantly reduces memory requirements and, together with contraction method of solution, is the principal reason for relatively fast computation when compared to prior methods.

convex iteration

By *convex iteration* we mean alternation of solution to (775a) and (781) until convergence. Direction vector y is initialized to $\mathbf{1}$ until the first fixed point is found; which means, the contraction recursion begins calculating a (1-norm) solution \mathcal{U}^* to (773) via problem (775b). Once \mathcal{U}^* is found, vector y is updated according to an estimate of image-gradient cardinality c : Sum

^{4.52}We are looking for at least 50dB image/error ratio from only 4.1% subsampled data (10 radial lines in \mathbf{k} -space). With this setting of ϵ , we actually attain in excess of 100dB from a very simple MATLAB program in about a minute on a 2006 vintage laptop. By trading execution time and treating image-gradient cardinality as a known quantity for this phantom, over 160dB is achievable.

^{4.53}Conjugate gradient method requires positive definiteness. [133, §4.8.3.2]

^{4.54}Fat is typical of compressed sensing problems; *e.g.*, [60] [67].

of the $4n^2 - c$ smallest entries of $|\Psi \text{vec } \mathcal{U}^*| \in \mathbb{R}^{4n^2}$ is the optimal objective value from a linear program, for $0 \leq c \leq 4n^2 - 1$ (475)

$$\begin{aligned} \sum_{i=c+1}^{4n^2} \pi(|\Psi \text{vec } \mathcal{U}^*|)_i &= \underset{y \in \mathbb{R}^{4n^2}}{\text{minimize}} && \langle |\Psi \text{vec } \mathcal{U}^*|, y \rangle \\ &\text{subject to} && 0 \preceq y \preceq \mathbf{1} \\ &&& y^T \mathbf{1} = 4n^2 - c \end{aligned} \quad (781)$$

where π is the nonlinear permutation-operator sorting its vector argument into nonincreasing order. An optimal solution y to (781), that is an extreme point of its feasible set, is known in closed form: it has 1 in each entry corresponding to the $4n^2 - c$ smallest entries of $|\Psi \text{vec } \mathcal{U}^*|$ and has 0 elsewhere (page 300). The updated image \mathcal{U}^* is assigned to \mathcal{U}_t , the contraction is recomputed solving (775b), direction vector y is updated again, and so on until convergence which is guaranteed by virtue of a monotonically nonincreasing real sequence of objective values in (775a) and (781).

There are two features that distinguish problem formulation (775b) and our particular implementation of it (MATLAB on [Wikimization](#)):

- 1) An image-gradient estimate may engage any combination of four adjacent pixels. In other words, the algorithm is not locked into a four-point gradient estimate (Figure 92); number of points constituting an estimate is determined by direction vector y . Indeed, we find only $c = 5092$ zero entries in y^* for the Shepp-Logan phantom; meaning, image-gradient sparsity is actually closer to 1.9% than the 3% reported elsewhere; *e.g.*, [315, §IIB].
- 2) Numerical precision ($\approx 1\text{E-}2$) of the fixed point of contraction (779) is a parameter to the implementation; meaning, direction vector y is typically updated after fixed-point recursion begins but prior to its culmination. Impact of this idiosyncrasy tends toward simultaneous optimization in variables \mathcal{U} and y while insuring that vector y settles on a boundary point of the feasible set (nonnegative hypercube slice) in (781) at every iteration; for only a boundary point^{4.55} can yield the sum of smallest entries in $|\Psi \text{vec } \mathcal{U}^*|$.

Reconstruction of the Shepp-Logan phantom at 103dB image/error is achieved in a MATLAB minute with 4.1% subsampled data; well below an

^{4.55}Simultaneous optimization of these two variables \mathcal{U} and y should never be a pinnacle of aspiration; for then, optimal y might not attain a boundary point.



Figure 93: Massachusetts Institute of Technology (MIT) logo, including its white boundary, may be interpreted as a rank-5 matrix. (Stanford University logo rank is much higher;) This constitutes *Scene Y* observed by the one-pixel camera in Figure 94 for Example 4.6.0.0.11.

11% least lower bound predicted by the sparse sampling theorem. Because reconstruction approaches optimal solution to a 0-norm problem, minimum number of Fourier-domain samples is bounded below by cardinality of the image-gradient at 1.9%. \square

4.6.0.0.11 Example. *Compressed sensing, compressive sampling.* [265] *As our modern technology-driven civilization acquires and exploits ever-increasing amounts of data, everyone now knows that most of the data we acquire can be thrown away with almost no perceptual loss – witness the broad success of lossy compression formats for sounds, images, and specialized technical data. The phenomenon of ubiquitous compressibility raises very natural questions: Why go to so much effort to acquire all the data when most of what we get will be thrown away? Can't we just directly measure the part that won't end up being thrown away?* –David Donoho [101]

Lossy data compression techniques are popular, but it is also well known that compression artifacts become quite perceptible with signal processing that goes beyond mere playback of a compressed signal. [195] [216] Spatial or audio frequencies presumed masked by a simultaneity, for example, become perceptible with significant post-filtering of the compressed signal. Further, there can be no universally acceptable and unique metric of perception for gauging exactly how much data can be tossed. For these reasons, there will always be need for raw (uncompressed) data.




Figure 94: One-pixel camera. [306] [330] *Compressive imaging camera block diagram. Incident lightfield (corresponding to the desired image Y) is reflected off a digital micromirror device (DMD) array whose mirror orientations are modulated in the pseudorandom pattern supplied by the random number generators (RNG). Each different mirror pattern produces a voltage at the single photodiode that corresponds to one measurement y_i .*

In this example we throw out only so much information as to leave perfect reconstruction within reach. Specifically, the MIT logo in Figure 93 is perfectly reconstructed from 700 time-sequential samples $\{y_i\}$ acquired by the one-pixel camera illustrated in Figure 94. The MIT-logo image in this example effectively impinges a 46×81 array micromirror device. This mirror array is modulated by a pseudonoise source that independently positions all the individual mirrors. A single photodiode (one pixel) integrates incident light from all mirrors. After stabilizing the mirrors to a fixed but pseudorandom pattern, light so collected is then digitized into one sample y_i by analog-to-digital (A/D) conversion. This sampling process is repeated with the micromirror array modulated to a new pseudorandom pattern.

The most important questions are: How many samples do we need for perfect reconstruction? Does that number of samples represent compression of the original data?

We claim that perfect reconstruction of the MIT logo can be reliably achieved with as few as 700 samples $y = [y_i] \in \mathbb{R}^{700}$ from this one-pixel camera. That number represents only 19% of information obtainable from

3726 micromirrors.^{4.56}

Our approach to reconstruction is to look for low-rank solution to an *underdetermined* system:

$$\begin{aligned} & \text{find} && X \\ & \text{subject to} && A \text{vec } X = y \\ & && \text{rank } X \leq 5 \end{aligned} \tag{782}$$

where $\text{vec } X$ is vectorized matrix $X \in \mathbb{R}^{46 \times 81}$ (stacked columns). Each row of fat matrix A is one realization of a pseudorandom pattern applied to the micromirrors. Since these patterns are deterministic (known), then the i^{th} sample y_i equals $A(i, :) \text{vec } Y$; *id est*, $y = A \text{vec } Y$. Perfect reconstruction means optimal solution X^* equals scene $Y \in \mathbb{R}^{46 \times 81}$ to within machine precision.

Because variable matrix X is generally not square or positive semidefinite, we constrain its rank by rewriting the problem equivalently

$$\begin{aligned} & \text{find} && X \\ & \text{subject to} && A \text{vec } X = y \\ & && \text{rank} \begin{bmatrix} W_1 & X \\ X^T & W_2 \end{bmatrix} \leq 5 \end{aligned} \tag{783}$$

This rank constraint on the composite matrix insures $\text{rank } X \leq 5$ for any choice of dimensionally compatible matrices W_1 and W_2 . But to solve this problem by *convex iteration*, we alternate solution of semidefinite program

$$\begin{aligned} & \underset{W_1, W_2, X}{\text{minimize}} && \text{tr} \left(\begin{bmatrix} W_1 & X \\ X^T & W_2 \end{bmatrix} Z \right) \\ & \text{subject to} && A \text{vec } X = y \\ & && \begin{bmatrix} W_1 & X \\ X^T & W_2 \end{bmatrix} \succeq 0 \end{aligned} \tag{784}$$

with semidefinite program

$$\begin{aligned} & \underset{Z}{\text{minimize}} && \text{tr} \left(\begin{bmatrix} W_1 & X \\ X^T & W_2 \end{bmatrix}^* Z \right) \\ & \text{subject to} && 0 \preceq Z \preceq I \\ & && \text{tr } Z = 46 + 81 - 5 \end{aligned} \tag{785}$$

^{4.56}That number (700 samples) is difficult to achieve, as reported in [265, §6]. If a minimal basis for the MIT logo were instead constructed, only five rows or columns worth of data (from a 46×81 matrix) are independent. This means a lower bound on achievable compression is about 230 samples; which corresponds to 6% of the original information.

(which has an optimal solution that is known in closed form, p.603) until a rank-5 composite matrix is found. The semidefiniteness constraints imply that matrices W_1 and W_2 and Z are symmetric.

With 1000 samples $\{y_i\}$, convergence occurs in two iterations; 700 samples require more than ten, but reconstruction remains perfect. Reconstruction is independent of pseudorandom sequence parameters; *e.g.*, binary sequences succeed with the same efficiency as Gaussian or uniformly distributed sequences. \square

4.6.1 rank-constraint midsummary

We find that this *direction matrix* idea works well and quite independently of desired rank or affine dimension. This idea of direction matrix is good principally because of its simplicity: When confronted with a problem otherwise convex if not for a rank or cardinality constraint, then that constraint becomes a linear regularization term in the objective.

There exists a common thread through all these Examples; that being, convex iteration with a direction matrix as normal to a linear regularization (a generalization of the well-known trace heuristic). But each problem type (per Example) possesses its own idiosyncrasies that slightly modify how a rank-constrained optimal solution is actually obtained: The *ball packing* problem in Chapter 5.4.2.2.3, for example, requires a problem sequence in a progressively larger number of balls to find a good initial value for the direction matrix, whereas many of the examples in the present chapter require an initial value of $\mathbf{0}$. Finding a feasible Boolean vector in Example 4.6.0.0.6 requires a procedure to detect stalls, when other problems have no such requirement. The combinatorial Procrustes problem in Example 4.6.0.0.3 allows use of a known closed-form solution for direction vector when solved via rank constraint, but not when solved via cardinality constraint. Some problems require a careful weighting of the regularization term, whereas other problems do not, and so on. It would be nice if there were a universally applicable method for constraining rank; one that is less susceptible to quirks of a particular problem type.

Poor initialization of the direction matrix from the regularization can lead to an erroneous result. We speculate one reason to be a simple dearth of optimal solutions of desired rank or cardinality;^{4.57} an unfortunate choice

^{4.57}Recall that, in Convex Optimization, an optimal solution generally comes from a

of initial search direction leading astray. Ease of solution by *convex iteration* occurs when optimal solutions abound. With this speculation in mind, we now propose a further generalization of *convex iteration* for constraining rank that attempts to ameliorate quirks and unify problem types:

4.7 Convex Iteration rank-1

We now develop a general method for constraining rank that first decomposes a given problem via standard diagonalization of matrices (§A.5). This method is motivated by observation (§4.4.1.1) that an optimal direction matrix can be diagonalizable simultaneously with an optimal variable matrix. This suggests minimization of an objective function directly in terms of eigenvalues. A second motivating observation is that variable orthogonal matrices seem easily found by *convex iteration*; e.g., Procrustes Example 4.6.0.0.2.

It turns out that this general method always requires solution to a rank-1 constrained problem regardless of desired rank from the original problem. To demonstrate, we pose a semidefinite feasibility problem

$$\begin{aligned} \text{find} \quad & X \in \mathbb{S}^n \\ \text{subject to} \quad & A \text{ svec } X = b \\ & X \succeq 0 \\ & \text{rank } X \leq \rho \end{aligned} \tag{786}$$

given an upper bound $0 < \rho < n$ on rank, vector $b \in \mathbb{R}^m$, and typically fat full-rank

$$A \triangleq \begin{bmatrix} \text{svec}(A_1)^T \\ \vdots \\ \text{svec}(A_m)^T \end{bmatrix} \in \mathbb{R}^{m \times n(n+1)/2} \tag{593}$$

where $A_i \in \mathbb{S}^n$, $i = 1 \dots m$ are also given. Symmetric matrix vectorization *svec* is defined in (50). Thus

$$A \text{ svec } X = \begin{bmatrix} \text{tr}(A_1 X) \\ \vdots \\ \text{tr}(A_m X) \end{bmatrix} \tag{594}$$

convex set of optimal solutions that can be large.

This program (786) states the classical problem of finding a matrix of specified rank in the intersection of the positive semidefinite cone with a number m of hyperplanes in the subspace of symmetric matrices \mathbb{S}^n . [25, §II.13] [23, §2.2]

Express the nonincreasingly ordered diagonalization of variable matrix

$$X \triangleq Q\Lambda Q^T = \sum_{i=1}^n \lambda_i Q_{ii} \in \mathbb{S}^n \quad (787)$$

which is a sum of rank-1 orthogonal projection matrices Q_{ii} weighted by eigenvalues λ_i where $Q_{ij} \triangleq q_i q_j^T \in \mathbb{R}^{n \times n}$, $Q = [q_1 \cdots q_n] \in \mathbb{R}^{n \times n}$, $Q^T = Q^{-1}$, $\Lambda_{ii} = \lambda_i \in \mathbb{R}$, and

$$\Lambda = \begin{bmatrix} \lambda_1 & & & \mathbf{0} \\ & \lambda_2 & & \\ & & \ddots & \\ \mathbf{0}^T & & & \lambda_n \end{bmatrix} \in \mathbb{S}^n \quad (788)$$

The fact

$$\Lambda \succeq 0 \Leftrightarrow X \succeq 0 \quad (1363)$$

allows splitting semidefinite feasibility problem (786) into two parts:

$$\begin{aligned} & \underset{\Lambda}{\text{minimize}} \quad \left\| A \operatorname{svec} \left(\sum_{i=1}^{\rho} \lambda_i Q_{ii} \right) - b \right\| \\ & \text{subject to} \quad \delta(\Lambda) \in \begin{bmatrix} \mathbb{R}_+^{\rho} \\ \mathbf{0} \end{bmatrix} \end{aligned} \quad (789)$$

$$\begin{aligned} & \underset{Q}{\text{minimize}} \quad \left\| A \operatorname{svec} \left(\sum_{i=1}^{\rho} \lambda_i^* Q_{ii} \right) - b \right\| \\ & \text{subject to} \quad Q^T = Q^{-1} \end{aligned} \quad (790)$$

The linear equality constraint $A \operatorname{svec} X = b$ has been moved to the objective within a norm because these two problems (789) (790) are iterated; equality might only become feasible near convergence. This iteration always converges to a local minimum because the sequence of objective values is monotonic and nonincreasing; any monotonically nonincreasing real sequence converges. [228, §1.2] [37, §1.1] A rank- ρ matrix X feasible to the original problem (786) is found when the objective converges to 0. Positive semidefiniteness of

matrix X with an upper bound ρ on rank is assured by the constraint on eigenvalue matrix Λ in convex problem (789); it means, the main diagonal of Λ must belong to the nonnegative orthant in a ρ -dimensional subspace of \mathbb{R}^n .

The second problem (790) in the iteration is not convex. We propose solving it by *convex iteration*: Make the assignment

$$\begin{aligned} G &= \begin{bmatrix} q_1 \\ \vdots \\ q_\rho \end{bmatrix} [q_1^T \cdots q_\rho^T] \in \mathbb{S}^{n\rho} \\ &= \begin{bmatrix} Q_{11} & \cdots & Q_{1\rho} \\ \vdots & \ddots & \vdots \\ Q_{1\rho}^T & \cdots & Q_{\rho\rho} \end{bmatrix} \triangleq \begin{bmatrix} q_1 q_1^T & \cdots & q_1 q_\rho^T \\ \vdots & \ddots & \vdots \\ q_\rho q_1^T & \cdots & q_\rho q_\rho^T \end{bmatrix} \end{aligned} \quad (791)$$

Given ρ eigenvalues λ_i^* , then a problem equivalent to (790) is

$$\begin{aligned} &\underset{Q_{ij} \in \mathbb{R}^{n \times n}}{\text{minimize}} \quad \left\| A \operatorname{svec} \left(\sum_{i=1}^{\rho} \lambda_i^* Q_{ii} \right) - b \right\| \\ &\text{subject to} \quad \operatorname{tr} Q_{ii} = 1, & i = 1 \dots \rho \\ &\quad \operatorname{tr} Q_{ij} = 0, & i < j = 2 \dots \rho \\ &\quad G = \begin{bmatrix} Q_{11} & \cdots & Q_{1\rho} \\ \vdots & \ddots & \vdots \\ Q_{1\rho}^T & \cdots & Q_{\rho\rho} \end{bmatrix} (\succeq 0) \\ &\quad \operatorname{rank} G = 1 \end{aligned} \quad (792)$$

This new problem always enforces a rank-1 constraint on matrix G ; *id est*, regardless of upper bound on rank ρ of variable matrix X , this equivalent problem (792) always poses a rank-1 constraint. Given some fixed weighting factor w , we propose solving (792) by iteration of the convex sequence

$$\begin{aligned} &\underset{Q_{ij} \in \mathbb{R}^{n \times n}}{\text{minimize}} \quad \left\| A \operatorname{svec} \left(\sum_{i=1}^{\rho} \lambda_i^* Q_{ii} \right) - b \right\| + w \operatorname{tr}(G W) \\ &\text{subject to} \quad \operatorname{tr} Q_{ii} = 1, & i = 1 \dots \rho \\ &\quad \operatorname{tr} Q_{ij} = 0, & i < j = 2 \dots \rho \\ &\quad G = \begin{bmatrix} Q_{11} & \cdots & Q_{1\rho} \\ \vdots & \ddots & \vdots \\ Q_{1\rho}^T & \cdots & Q_{\rho\rho} \end{bmatrix} \succeq 0 \end{aligned} \quad (793)$$

with

$$\begin{aligned}
 & \underset{W \in \mathbb{S}^{n\rho}}{\text{minimize}} && \text{tr}(G^* W) \\
 & \text{subject to} && 0 \preceq W \preceq I \\
 & && \text{tr } W = n\rho - 1
 \end{aligned} \tag{794}$$

the latter providing direction of search W for a rank-1 matrix G in (793). An optimal solution to (794) has closed form (p.603). These convex problems (793) (794) are iterated with (789) until a rank-1 G matrix is found and the objective of (793) vanishes.

For a fixed number of equality constraints m and upper bound ρ , the feasible solution set in rank-constrained semidefinite feasibility problem (786) grows with matrix dimension n . Empirically, this semidefinite feasibility problem becomes easier to solve by method of *convex iteration* as n increases. We find a good value of weight $w \approx 5$ for randomized problems. Initial value of direction matrix W is not critical. With dimension $n = 26$ and number of equality constraints $m = 10$, convergence to a rank- $\rho = 2$ solution to semidefinite feasibility problem (786) is achieved for nearly all realizations in less than 10 iterations. (This finding is significant in so far as Barvinok's Proposition 2.9.3.0.1 predicts existence of matrices having rank 4 or less in this intersection, but his upper bound can be no tighter than 3.) For small n , stall detection is required; one numerical implementation is disclosed on [Wikimization](#).

This diagonalization decomposition technique is extensible to other problem types, of course, including optimization problems having nontrivial objectives. Because of a greatly expanded feasible set, for example,

$$\begin{aligned}
 & \text{find} && X \in \mathbb{S}^n \\
 & \text{subject to} && A \text{ svec } X \succeq b \\
 & && X \succeq 0 \\
 & && \text{rank } X \leq \rho
 \end{aligned} \tag{795}$$

this relaxation of (786) [199, §III] is more easily solved by *convex iteration* rank-1. (The inequality in A remains in the constraints after diagonalization.) Because of the nonconvex nature of a rank constrained problem, more generally, there can be no proof of global convergence of *convex iteration* from an arbitrary initialization; although, local convergence is guaranteed by virtue of monotonically nonincreasing real objective sequences. [228, §1.2] [37, §1.1]

Chapter 5

Euclidean Distance Matrix

These results [(827)] were obtained by Schoenberg (1935), a surprisingly late date for such a fundamental property of Euclidean geometry.

– John Clifford Gower [141, §3]

By itself, distance information between many points in Euclidean space is lacking. We might want to know more; such as, relative or absolute position or dimension of some hull. A question naturally arising in some fields (*e.g.*, geodesy, economics, genetics, psychology, biochemistry, engineering) [88] asks what facts can be deduced given only distance information. What can we know about the underlying points that the distance information purports to describe? We also ask what it means when given distance information is incomplete; or suppose the distance information is not reliable, available, or specified only by certain tolerances (affine inequalities). These questions motivate a study of interpoint distance, well represented in any spatial dimension by a simple matrix from linear algebra.^{5.1} In what follows, we will answer some of these questions via Euclidean distance matrices.

^{5.1} *e.g.*, $\sqrt{D} \in \mathbb{R}^{N \times N}$, a classical two-dimensional matrix representation of absolute interpoint distance because its entries (in ordered rows and columns) can be written neatly on a piece of paper. Matrix D will be reserved throughout to hold distance-square.




Figure 95: Convex hull of three points ($N = 3$) is shaded in \mathbb{R}^3 ($n = 3$). Dotted lines are imagined vectors to points whose affine dimension is 2.

5.1 EDM

Euclidean space \mathbb{R}^n is a finite-dimensional real vector space having an inner product defined on it, inducing a *metric*. [201, §3.1] A Euclidean distance matrix, an EDM in $\mathbb{R}_+^{N \times N}$, is an exhaustive table of distance-square d_{ij} between points taken by pair from a list of N points $\{x_\ell, \ell = 1 \dots N\}$ in \mathbb{R}^n ; the squared metric, the measure of distance-square:

$$d_{ij} = \|x_i - x_j\|_2^2 \triangleq \langle x_i - x_j, x_i - x_j \rangle \quad (796)$$

Each point is labelled ordinally, hence the row or column index of an EDM, i or $j = 1 \dots N$, individually addresses all the points in the list.

Consider the following example of an EDM for the case $N = 3$:

$$D = [d_{ij}] = \begin{bmatrix} d_{11} & d_{12} & d_{13} \\ d_{21} & d_{22} & d_{23} \\ d_{31} & d_{32} & d_{33} \end{bmatrix} = \begin{bmatrix} 0 & d_{12} & d_{13} \\ d_{12} & 0 & d_{23} \\ d_{13} & d_{23} & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 5 \\ 1 & 0 & 4 \\ 5 & 4 & 0 \end{bmatrix} \quad (797)$$

Matrix D has N^2 entries but only $N(N-1)/2$ pieces of information. In Figure 95 are shown three points in \mathbb{R}^3 that can be arranged in a list to

correspond to D in (797). But such a list is not unique because any rotation, reflection, or translation (§5.5) of those points would produce the same EDM D .

5.2 First metric properties

For $i, j = 1 \dots N$, distance between points x_i and x_j must satisfy the defining requirements imposed upon any *metric space*: [201, §1.1] [228, §1.7] namely, for Euclidean metric $\sqrt{d_{ij}}$ (§5.4) in \mathbb{R}^n

1. $\sqrt{d_{ij}} \geq 0, \quad i \neq j$ nonnegativity
2. $\sqrt{d_{ij}} = 0 \Leftrightarrow x_i = x_j$ self-distance
3. $\sqrt{d_{ij}} = \sqrt{d_{ji}}$ symmetry
4. $\sqrt{d_{ij}} \leq \sqrt{d_{ik}} + \sqrt{d_{kj}}, \quad i \neq j \neq k$ triangle inequality

Then all entries of an EDM must be in concord with these Euclidean metric properties: specifically, each entry must be nonnegative,^{5.2} the main diagonal must be $\mathbf{0}$,^{5.3} and an EDM must be symmetric. The fourth property provides upper and lower bounds for each entry. Property 4 is true more generally when there are no restrictions on indices i, j, k , but furnishes no new information.

5.3 \exists fifth Euclidean metric property

The four properties of the Euclidean metric provide information insufficient to certify that a bounded convex polyhedron more complicated than a triangle has a Euclidean realization. [141, §2] Yet any list of points or the vertices of any bounded convex polyhedron must conform to the properties.

^{5.2}Implicit from the terminology, $\sqrt{d_{ij}} \geq 0 \Leftrightarrow d_{ij} \geq 0$ is always assumed.

^{5.3}What we call self-distance, Marsden calls *nondegeneracy*. [228, §1.6] Kreyszig calls these first metric properties *axioms of the metric*; [201, p.4] Blumenthal refers to them as *postulates*. [44, p.15]

5.3.0.0.1 Example. *Triangle.*

Consider the EDM in (797), but missing one of its entries:

$$D = \begin{bmatrix} 0 & 1 & d_{13} \\ 1 & 0 & 4 \\ d_{31} & 4 & 0 \end{bmatrix} \quad (798)$$

Can we determine unknown entries of D by applying the metric properties? Property 1 demands $\sqrt{d_{13}}, \sqrt{d_{31}} \geq 0$, property 2 requires the main diagonal be $\mathbf{0}$, while property 3 makes $\sqrt{d_{31}} = \sqrt{d_{13}}$. The fourth property tells us

$$1 \leq \sqrt{d_{13}} \leq 3 \quad (799)$$

Indeed, described over that closed interval $[1, 3]$ is a family of triangular polyhedra whose angle at vertex x_2 varies from 0 to π radians. So, yes we can determine the unknown entries of D , but they are not unique; nor should they be from the information given for this example. \square

5.3.0.0.2 Example. *Small completion problem, I.*

Now consider the polyhedron in Figure 96(b) formed from an unknown list $\{x_1, x_2, x_3, x_4\}$. The corresponding EDM less one critical piece of information, d_{14} , is given by

$$D = \begin{bmatrix} 0 & 1 & 5 & d_{14} \\ 1 & 0 & 4 & 1 \\ 5 & 4 & 0 & 1 \\ d_{14} & 1 & 1 & 0 \end{bmatrix} \quad (800)$$

From metric property 4 we may write a few inequalities for the two triangles common to d_{14} ; we find

$$\sqrt{5}-1 \leq \sqrt{d_{14}} \leq 2 \quad (801)$$

We cannot further narrow those bounds on $\sqrt{d_{14}}$ using only the four metric properties (§5.8.3.1.1). Yet there is only one possible choice for $\sqrt{d_{14}}$ because points x_2, x_3, x_4 must be collinear. All other values of $\sqrt{d_{14}}$ in the interval $[\sqrt{5}-1, 2]$ specify impossible distances in any dimension; *id est*, in this particular example the triangle inequality does not yield an interval for $\sqrt{d_{14}}$ over which a family of convex polyhedra can be reconstructed. \square




Figure 96: (a) Complete *dimensionless EDM graph*. (b) Emphasizing obscured segments $\overline{x_2x_4}$, $\overline{x_4x_3}$, and $\overline{x_2x_3}$, now only five $(2N - 3)$ absolute distances are specified. EDM so represented is incomplete, missing d_{14} as in (800), yet the isometric reconstruction (§5.4.2.2.6) is unique as proved in §5.9.3.0.1 and §5.14.4.1.1. First four properties of Euclidean metric are not a recipe for reconstruction of this polyhedron.

We will return to this simple Example 5.3.0.0.2 to illustrate more elegant methods of solution in §5.8.3.1.1, §5.9.3.0.1, and §5.14.4.1.1. Until then, we can deduce some general principles from the foregoing examples:

- Unknown d_{ij} of an EDM are not necessarily uniquely determinable.
- The triangle inequality does not produce necessarily *tight* bounds.^{5.4}
- Four Euclidean metric properties are insufficient for reconstruction.

^{5.4}The term *tight* with reference to an inequality means equality is achievable.

5.3.1 Lookahead

There must exist at least one requirement more than the four properties of the Euclidean metric that makes them altogether necessary and sufficient to certify realizability of bounded convex polyhedra. Indeed, there are infinitely many more; there are precisely $N+1$ necessary and sufficient Euclidean metric requirements for N points constituting a generating list (§2.3.2). Here is the fifth requirement:

5.3.1.0.1 Fifth Euclidean metric property. *Relative-angle inequality.* (confer §5.14.2.1.1) Augmenting the four fundamental properties of the Euclidean metric in \mathbb{R}^n , for all $i, j, \ell \neq k \in \{1 \dots N\}$, $i < j < \ell$, and for $N \geq 4$ distinct points $\{x_k\}$, the inequalities

$$\begin{aligned} \cos(\theta_{ik\ell} + \theta_{\ell kj}) &\leq \cos \theta_{ikj} \leq \cos(\theta_{ik\ell} - \theta_{\ell kj}) \\ 0 &\leq \theta_{ik\ell}, \theta_{\ell kj}, \theta_{ikj} \leq \pi \end{aligned} \tag{802}$$

where $\theta_{ikj} = \theta_{jki}$ is the angle between vectors at vertex x_k (869) (Figure 97), must be satisfied at each point x_k regardless of affine dimension. \diamond

We will explore this in §5.14. One of our early goals is to determine matrix criteria that subsume all the Euclidean metric properties and any further requirements. Looking ahead, we will find (1150) (827) (832)

$$\begin{aligned} -z^T D z &\geq 0 \\ \mathbf{1}^T z &= 0 \\ (\forall \|z\| = 1) &\Leftrightarrow D \in \text{EDM}^N \\ D &\in \mathbb{S}_h^N \end{aligned} \tag{803}$$

where the convex cone of Euclidean distance matrices $\text{EDM}^N \subseteq \mathbb{S}_h^N$ belongs to the subspace of symmetric hollow^{5.5} matrices (§2.2.3.0.1. Numerical test `isedm(D)` is provided on [Wikimization](#).) Having found equivalent matrix criteria, we will see there is a bridge from bounded convex polyhedra to EDMs in §5.9.^{5.6}

^{5.5} $\mathbf{0}$ main diagonal.

^{5.6} From an EDM, a generating list (§2.3.2, §2.12.2) for a polyhedron can be found (§5.12) correct to within a rotation, reflection, and translation (§5.5).




Figure 97: Fifth Euclidean metric property nomenclature. Each angle θ is made by a vector pair at vertex k while i, j, k, l index four points at the vertices of a generally irregular tetrahedron. The fifth property is necessary for realization of four or more points; a reckoning by three angles in any dimension. Together with the first four Euclidean metric properties, this fifth property is necessary and sufficient for realization of four points.

Now we develop some invaluable concepts, moving toward a link of the Euclidean metric properties to matrix criteria.

5.4 EDM definition

Ascribe points in a list $\{x_\ell \in \mathbb{R}^n, \ell=1 \dots N\}$ to the columns of a matrix

$$X = [x_1 \ \cdots \ x_N] \in \mathbb{R}^{n \times N} \quad (69)$$

where N is regarded as *cardinality* of list X . When matrix $D = [d_{ij}]$ is an EDM, its entries must be related to those points constituting the list by the Euclidean distance-square: for $i, j=1 \dots N$ (§A.1.1 no.33)

$$\begin{aligned} d_{ij} &= \|x_i - x_j\|^2 = (x_i - x_j)^T(x_i - x_j) = \|x_i\|^2 + \|x_j\|^2 - 2x_i^T x_j \\ &= \begin{bmatrix} x_i^T & x_j^T \end{bmatrix} \begin{bmatrix} I & -I \\ -I & I \end{bmatrix} \begin{bmatrix} x_i \\ x_j \end{bmatrix} \\ &= \text{vec}(X)^T (\Phi_{ij} \otimes I) \text{vec } X = \langle \Phi_{ij}, X^T X \rangle \end{aligned} \quad (804)$$

where

$$\text{vec } X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_N \end{bmatrix} \in \mathbb{R}^{nN} \quad (805)$$

and where \otimes signifies Kronecker product (§D.1.2.1). $\Phi_{ij} \otimes I$ is positive semidefinite (1393) having $I \in \mathbb{S}^n$ in its ii^{th} and jj^{th} block of entries while $-I \in \mathbb{S}^n$ fills its ij^{th} and ji^{th} block; *id est*,

$$\begin{aligned} \Phi_{ij} &\triangleq \delta((e_i e_j^T + e_j e_i^T) \mathbf{1}) - (e_i e_j^T + e_j e_i^T) \in \mathbb{S}_+^N \\ &= e_i e_i^T + e_j e_j^T - e_i e_j^T - e_j e_i^T \\ &= (e_i - e_j)(e_i - e_j)^T \end{aligned} \quad (806)$$

where $\{e_i \in \mathbb{R}^N, i=1 \dots N\}$ is the set of standard basis vectors. Thus each entry d_{ij} is a convex quadratic function (§A.4.0.0.1) of $\text{vec } X$ (31). [271, §6]

The collection of all Euclidean distance matrices \mathbb{EDM}^N is a convex subset of $\mathbb{R}_+^{N \times N}$ called the *EDM cone* (§6, Figure 131, p.502);

$$\mathbf{0} \in \mathbb{EDM}^N \subseteq \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N} \subset \mathbb{S}^N \quad (807)$$

An EDM D must be expressible as a function of some list X ; *id est*, it must have the form

$$\mathbf{D}(X) \triangleq \delta(X^T X) \mathbf{1}^T + \mathbf{1} \delta(X^T X)^T - 2X^T X \in \mathbb{EDM}^N \quad (808)$$

$$= [\text{vec}(X)^T (\Phi_{ij} \otimes I) \text{vec } X, \quad i, j = 1 \dots N] \quad (809)$$

Function $\mathbf{D}(X)$ will make an EDM given any $X \in \mathbb{R}^{n \times N}$, conversely, but $\mathbf{D}(X)$ is not a convex function of X (§5.4.1). Now the EDM cone may be described:

$$\mathbb{EDM}^N = \{\mathbf{D}(X) \mid X \in \mathbb{R}^{N-1 \times N}\} \quad (810)$$

Expression $\mathbf{D}(X)$ is a matrix definition of EDM and so conforms to the Euclidean metric properties:

Nonnegativity of EDM entries (property 1, §5.2) is obvious from the distance-square definition (804), so holds for any D expressible in the form $\mathbf{D}(X)$ in (808).

When we say D is an EDM, reading from (808), it implicitly means the main diagonal must be $\mathbf{0}$ (property 2, self-distance) and D must be symmetric (property 3); $\delta(D) = \mathbf{0}$ and $D^T = D$ or, equivalently, $D \in \mathbb{S}_h^N$ are necessary matrix criteria.

5.4.0.1 homogeneity

Function $\mathbf{D}(X)$ is homogeneous in the sense, for $\zeta \in \mathbb{R}$

$$\sqrt{\circ} \mathbf{D}(\zeta X) = |\zeta| \sqrt{\circ} \mathbf{D}(X) \quad (811)$$

where the positive square root is entrywise (\circ).

Any nonnegatively scaled EDM remains an EDM; *id est*, the matrix class EDM is invariant to nonnegative scaling ($\alpha \mathbf{D}(X)$ for $\alpha \geq 0$) because all EDMs of dimension N constitute a convex cone \mathbb{EDM}^N (§6, Figure 118).

5.4.1 $-V_{\mathcal{N}}^T \mathbf{D}(X) V_{\mathcal{N}}$ convexity

We saw that EDM entries $d_{ij} \left(\begin{bmatrix} x_i \\ x_j \end{bmatrix} \right)$ are convex quadratic functions. Yet $-\mathbf{D}(X)$ (808) is not a quasiconvex function of matrix $X \in \mathbb{R}^{n \times N}$ because the second directional derivative (§3.3)

$$-\frac{d^2}{dt^2} \Big|_{t=0} \mathbf{D}(X + tY) = 2(-\delta(Y^T Y) \mathbf{1}^T - \mathbf{1} \delta(Y^T Y)^T + 2Y^T Y) \quad (812)$$

is indefinite for any $Y \in \mathbb{R}^{n \times N}$ since its main diagonal is $\mathbf{0}$. [139, §4.2.8] [180, §7.1, prob.2] Hence $-\mathbf{D}(X)$ can neither be convex in X .

The outcome is different when instead we consider

$$-V_{\mathcal{N}}^T \mathbf{D}(X) V_{\mathcal{N}} = 2V_{\mathcal{N}}^T X^T X V_{\mathcal{N}} \quad (813)$$

where we introduce the full-rank skinny *Schoenberg auxiliary matrix* (§B.4.2)

$$V_{\mathcal{N}} \triangleq \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & -1 & \cdots & -1 \\ 1 & & & \mathbf{0} \\ & 1 & & \\ & & \ddots & \\ \mathbf{0} & & & 1 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} -\mathbf{1}^T \\ I \end{bmatrix} \in \mathbb{R}^{N \times N-1} \quad (814)$$

($\mathcal{N}(V_{\mathcal{N}}) = \mathbf{0}$) having range

$$\mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^T), \quad V_{\mathcal{N}}^T \mathbf{1} = \mathbf{0} \quad (815)$$

Matrix-valued function (813) meets the criterion for convexity in §3.2.3.0.2 over its domain that is all of $\mathbb{R}^{n \times N}$; *videlicet*, for any $Y \in \mathbb{R}^{n \times N}$

$$-\frac{d^2}{dt^2} V_{\mathcal{N}}^T \mathbf{D}(X + tY) V_{\mathcal{N}} = 4V_{\mathcal{N}}^T Y^T Y V_{\mathcal{N}} \succeq 0 \quad (816)$$

Quadratic matrix-valued function $-V_{\mathcal{N}}^T \mathbf{D}(X) V_{\mathcal{N}}$ is therefore convex in X achieving its minimum, with respect to a positive semidefinite cone (§2.7.2.2), at $X = \mathbf{0}$. When the penultimate number of points exceeds the dimension of the space $n < N - 1$, strict convexity of the quadratic (813) becomes impossible because (816) could not then be positive definite.

5.4.2 Gram-form EDM definition

Positive semidefinite matrix $X^T X$ in (808), formed from inner product of list X , is known as a *Gram matrix*; [220, §3.6]

$$\begin{aligned}
 G \triangleq X^T X &= \begin{bmatrix} x_1^T \\ \vdots \\ x_N^T \end{bmatrix} [x_1 \cdots x_N] = \begin{bmatrix} \|x_1\|^2 & x_1^T x_2 & x_1^T x_3 & \cdots & x_1^T x_N \\ x_2^T x_1 & \|x_2\|^2 & x_2^T x_3 & \cdots & x_2^T x_N \\ x_3^T x_1 & x_3^T x_2 & \|x_3\|^2 & \ddots & x_3^T x_N \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ x_N^T x_1 & x_N^T x_2 & x_N^T x_3 & \cdots & \|x_N\|^2 \end{bmatrix} \in \mathbb{S}_+^N \\
 &= \delta \left(\begin{bmatrix} \|x_1\| \\ \|x_2\| \\ \vdots \\ \|x_N\| \end{bmatrix} \right) \begin{bmatrix} 1 & \cos \psi_{12} & \cos \psi_{13} & \cdots & \cos \psi_{1N} \\ \cos \psi_{12} & 1 & \cos \psi_{23} & \cdots & \cos \psi_{2N} \\ \cos \psi_{13} & \cos \psi_{23} & 1 & \ddots & \cos \psi_{3N} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ \cos \psi_{1N} & \cos \psi_{2N} & \cos \psi_{3N} & \cdots & 1 \end{bmatrix} \delta \left(\begin{bmatrix} \|x_1\| \\ \|x_2\| \\ \vdots \\ \|x_N\| \end{bmatrix} \right) \\
 &\triangleq \sqrt{\delta^2(G)} \Psi \sqrt{\delta^2(G)} \tag{817}
 \end{aligned}$$

where ψ_{ij} (837) is angle between vectors x_i and x_j , and where δ^2 denotes a diagonal matrix in this case. Positive semidefiniteness of *interpoint angle matrix* Ψ implies positive semidefiniteness of Gram matrix G ; [53, §8.3]

$$G \succeq 0 \Leftrightarrow \Psi \succeq 0 \tag{818}$$

When $\delta^2(G)$ is nonsingular, then $G \succeq 0 \Leftrightarrow \Psi \succeq 0$. (§A.3.1.0.5)

Distance-square d_{ij} (804) is related to Gram matrix entries $G^T = G \triangleq [g_{ij}]$

$$\begin{aligned}
 d_{ij} &= g_{ii} + g_{jj} - 2g_{ij} \\
 &= \langle \Phi_{ij}, G \rangle \tag{819}
 \end{aligned}$$

where Φ_{ij} is defined in (806). Hence the linear EDM definition

$$\left. \begin{aligned}
 \mathbf{D}(G) &\triangleq \delta(G)\mathbf{1}^T + \mathbf{1}\delta(G)^T - 2G \in \mathbb{EDM}^N \\
 &= [\langle \Phi_{ij}, G \rangle, i, j = 1 \dots N]
 \end{aligned} \right\} \Leftrightarrow G \succeq 0 \tag{820}$$

The EDM cone may be described, (*confer* (904)(910))

$$\mathbb{EDM}^N = \{ \mathbf{D}(G) \mid G \in \mathbb{S}_+^N \} \tag{821}$$

5.4.2.1 First point at origin

Assume the first point x_1 in an unknown list X resides at the origin;

$$Xe_1 = \mathbf{0} \Leftrightarrow Ge_1 = \mathbf{0} \quad (822)$$

Consider the symmetric translation $(I - \mathbf{1}e_1^T)\mathbf{D}(G)(I - e_1\mathbf{1}^T)$ that shifts the first row and column of $\mathbf{D}(G)$ to the origin; setting Gram-form EDM operator $\mathbf{D}(G) = D$ for convenience,

$$-(D - (De_1\mathbf{1}^T + \mathbf{1}e_1^TD) + \mathbf{1}e_1^TD e_1\mathbf{1}^T)^{\frac{1}{2}} = G - (Ge_1\mathbf{1}^T + \mathbf{1}e_1^TG) + \mathbf{1}e_1^TG e_1\mathbf{1}^T \quad (823)$$

where

$$e_1 \triangleq \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \quad (824)$$

is the first vector from the standard basis. Then it follows for $D \in \mathbb{S}_h^N$

$$\begin{aligned} G &= -(D - (De_1\mathbf{1}^T + \mathbf{1}e_1^TD))^{\frac{1}{2}}, \quad x_1 = \mathbf{0} \\ &= -[\mathbf{0} \quad \sqrt{2}V_{\mathcal{N}}]^T D [\mathbf{0} \quad \sqrt{2}V_{\mathcal{N}}]^{\frac{1}{2}} \\ &= \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & -V_{\mathcal{N}}^T D V_{\mathcal{N}} \end{bmatrix} \\ V_{\mathcal{N}}^T G V_{\mathcal{N}} &= -V_{\mathcal{N}}^T D V_{\mathcal{N}}^{\frac{1}{2}} \quad \forall X \end{aligned} \quad (825)$$

where

$$I - e_1\mathbf{1}^T = \begin{bmatrix} \mathbf{0} & \sqrt{2}V_{\mathcal{N}} \end{bmatrix} \quad (826)$$

is a projector nonorthogonally projecting (§E.1) on

$$\begin{aligned} \mathbb{S}_1^N &= \{G \in \mathbb{S}^N \mid Ge_1 = \mathbf{0}\} \\ &= \left\{ \begin{bmatrix} \mathbf{0} & \sqrt{2}V_{\mathcal{N}} \end{bmatrix}^T Y \begin{bmatrix} \mathbf{0} & \sqrt{2}V_{\mathcal{N}} \end{bmatrix} \mid Y \in \mathbb{S}^N \right\} \end{aligned} \quad (1889)$$

in the Euclidean sense. From (825) we get sufficiency of the first matrix criterion for an EDM proved by Schoenberg in 1935; [276]^{5.7}

^{5.7}From (815) we know $\mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^T)$, so (827) is the same as (803). In fact, any matrix V in place of $V_{\mathcal{N}}$ will satisfy (827) whenever $\mathcal{R}(V) = \mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^T)$. But $V_{\mathcal{N}}$ is the matrix implicit in Schoenberg's seminal exposition.

$$D \in \text{EDM}^N \Leftrightarrow \begin{cases} -V_N^T D V_N \in \mathbb{S}_+^{N-1} \\ D \in \mathbb{S}_h^N \end{cases} \quad (827)$$

We provide a rigorous complete more geometric proof of this fundamental *Schoenberg criterion* in §5.9.1.0.4. (`isedm(D)` on [Wikimization](#).)

By substituting $G = \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & -V_N^T D V_N \end{bmatrix}$ (825) into $\mathbf{D}(G)$ (820), assuming $x_1 = \mathbf{0}$

$$D = \begin{bmatrix} 0 \\ \delta(-V_N^T D V_N) \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & \delta(-V_N^T D V_N)^T \end{bmatrix} - 2 \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & -V_N^T D V_N \end{bmatrix} \quad (828)$$

We provide details of this bijection in §5.6.2.

5.4.2.2 0 geometric center

Assume the *geometric center* (§5.5.1.0.1) of an unknown list X is the origin;

$$X\mathbf{1} = \mathbf{0} \Leftrightarrow G\mathbf{1} = \mathbf{0} \quad (829)$$

Now consider the calculation $(I - \frac{1}{N}\mathbf{1}\mathbf{1}^T)\mathbf{D}(G)(I - \frac{1}{N}\mathbf{1}\mathbf{1}^T)$, a geometric centering or projection operation. (§E.7.2.0.2) Setting $\mathbf{D}(G) = D$ for convenience as in §5.4.2.1,

$$\begin{aligned} G &= -\left(D - \frac{1}{N}(D\mathbf{1}\mathbf{1}^T + \mathbf{1}\mathbf{1}^T D) + \frac{1}{N^2}\mathbf{1}\mathbf{1}^T D \mathbf{1}\mathbf{1}^T\right) \frac{1}{2}, \quad X\mathbf{1} = \mathbf{0} \\ &= -VDV \frac{1}{2} \\ VGV &= -VDV \frac{1}{2} \quad \forall X \end{aligned} \quad (830)$$

where more properties of the auxiliary (*geometric centering*, projection) matrix

$$V \triangleq I - \frac{1}{N}\mathbf{1}\mathbf{1}^T \in \mathbb{S}^N \quad (831)$$

are found in §B.4. VGV may be regarded as a *covariance matrix* of means 0. From (830) and the assumption $D \in \mathbb{S}_h^N$ we get sufficiency of the more popular form of Schoenberg's criterion:

$$D \in \text{EDM}^N \Leftrightarrow \begin{cases} -VDV \in \mathbb{S}_+^N \\ D \in \mathbb{S}_h^N \end{cases} \quad (832)$$

Of particular utility when $D \in \text{EDM}^N$ is the fact, (§B.4.2 no.20) (804)

$$\begin{aligned} \text{tr}(-VDV\frac{1}{2}) &= \frac{1}{2N} \sum_{i,j} d_{ij} &= \frac{1}{2N} \text{vec}(X)^T \left(\sum_{i,j} \Phi_{ij} \otimes I \right) \text{vec} X \\ &= \text{tr}(VGV), \quad G \succeq 0 & (833) \\ &= \text{tr} G &= \sum_{\ell=1}^N \|x_\ell\|^2 = \|X\|_F^2, \quad X\mathbf{1} = \mathbf{0} \end{aligned}$$

where $\sum \Phi_{ij} \in \mathbb{S}_+^N$ (806), therefore convex in $\text{vec} X$. We will find this trace useful as a heuristic to minimize affine dimension of an unknown list arranged columnar in X (§7.2.2), but it tends to facilitate reconstruction of a list configuration having least energy; *id est*, it compacts a reconstructed list by minimizing total norm-square of the vertices.

By substituting $G = -VDV\frac{1}{2}$ (830) into $\mathbf{D}(G)$ (820), assuming $X\mathbf{1} = \mathbf{0}$ (confer §5.6.1)

$$D = \delta(-VDV\frac{1}{2})\mathbf{1}^T + \mathbf{1}\delta(-VDV\frac{1}{2})^T - 2(-VDV\frac{1}{2}) \quad (834)$$

These relationships will allow combination of distance and Gram constraints in any optimization problem we may pose:

- Constraining all main diagonal entries of a Gram matrix to 1, for example, is equivalent to the constraint that all points lie on a hypersphere (§5.9.1.0.3) of radius 1 centered at the origin. This is equivalent to the EDM constraint: $D\mathbf{1} = 2N\mathbf{1}$. [78, p.116] Any further constraint on that Gram matrix then applies only to interpoint angle Ψ .
- More generally, interpoint angle Ψ can be constrained by fixing all the individual point lengths $\delta(G)^{1/2}$; then

$$\Psi = -\frac{1}{2}\delta^2(G)^{-1/2}VDV\delta^2(G)^{-1/2} \quad (835)$$

5.4.2.2.1 Example. *List-member constraints via Gram matrix.*

Capitalizing on identity (830) relating Gram and EDM D matrices, a constraint set such as

$$\left. \begin{aligned} \operatorname{tr}\left(-\frac{1}{2}VDVe_i e_i^T\right) &= \|x_i\|^2 \\ \operatorname{tr}\left(-\frac{1}{2}VDV(e_i e_j^T + e_j e_i^T)\frac{1}{2}\right) &= x_i^T x_j \\ \operatorname{tr}\left(-\frac{1}{2}VDVe_j e_j^T\right) &= \|x_j\|^2 \end{aligned} \right\} \quad (836)$$

relates list member x_i to x_j to within an isometry through inner-product identity [339, §1-7]

$$\cos \psi_{ij} = \frac{x_i^T x_j}{\|x_i\| \|x_j\|} \quad (837)$$

where ψ_{ij} is angle between the two vectors as in (817). For M list members, there total $M(M+1)/2$ such constraints. Angle constraints are incorporated in Example 5.4.2.2.2 and Example 5.4.2.2.9. \square

Consider the academic problem of finding a Gram matrix subject to constraints on each and every entry of the corresponding EDM:

$$\begin{aligned} &\underset{D \in \mathbb{S}_h^N}{\text{find}} \quad -VDV\frac{1}{2} \in \mathbb{S}^N \\ &\text{subject to} \quad \langle D, (e_i e_j^T + e_j e_i^T)\frac{1}{2} \rangle = \check{d}_{ij}, \quad i, j = 1 \dots N, \quad i < j \\ &\quad \quad \quad -VDV \succeq 0 \end{aligned} \quad (838)$$

where the \check{d}_{ij} are given nonnegative constants. EDM D can, of course, be replaced with the equivalent Gram-form (820). Requiring only the self-adjointness property (1331) of the main-diagonal linear operator δ we get, for $A \in \mathbb{S}^N$

$$\langle D, A \rangle = \langle \delta(G)\mathbf{1}^T + \mathbf{1}\delta(G)^T - 2G, A \rangle = \langle G, \delta(A\mathbf{1}) - A \rangle 2 \quad (839)$$

Then the problem equivalent to (838) becomes, for $G \in \mathbb{S}_c^N \Leftrightarrow G\mathbf{1} = \mathbf{0}$

$$\begin{aligned} &\underset{G \in \mathbb{S}_c^N}{\text{find}} \quad G \in \mathbb{S}^N \\ &\text{subject to} \quad \left\langle G, \delta((e_i e_j^T + e_j e_i^T)\mathbf{1}) - (e_i e_j^T + e_j e_i^T) \right\rangle = \check{d}_{ij}, \quad i, j = 1 \dots N, \quad i < j \\ &\quad \quad \quad G \succeq 0 \end{aligned} \quad (840)$$




Figure 98: Arbitrary hexagon in \mathbb{R}^3 whose vertices are labelled clockwise.

Barvinok's Proposition 2.9.3.0.1 predicts existence for either formulation (838) or (840) such that implicit equality constraints induced by subspace membership are ignored

$$\text{rank } G, \text{ rank } VDV \leq \left\lfloor \frac{\sqrt{8(N(N-1)/2) + 1} - 1}{2} \right\rfloor = N - 1 \quad (841)$$

because, in each case, the Gram matrix is confined to a face of positive semidefinite cone \mathbb{S}_+^N isomorphic with \mathbb{S}_+^{N-1} (§6.7.1). (§E.7.2.0.2) This bound is tight (§5.7.1.1) and is the greatest upper bound.^{5.8}

5.4.2.2.2 Example. Hexagon.

Barvinok [24, §2.6] poses a problem in *geometric realizability* of an arbitrary hexagon (Figure 98) having:

1. prescribed (one-dimensional) face-lengths l
2. prescribed angles φ between the three pairs of opposing faces
3. a constraint on the sum of norm-square of each and every vertex x

^{5.8} $-VDV|_{N \leftarrow 1} = 0$ (§B.4.1)

ten affine equality constraints in all on a Gram matrix $G \in \mathbb{S}^6$ (830). Let's realize this as a convex feasibility problem (with constraints written in the same order) also assuming $\mathbf{0}$ geometric center (829):

$$\begin{aligned}
 & \underset{D \in \mathbb{S}_h^6}{\text{find}} && -VDV \frac{1}{2} \in \mathbb{S}^6 \\
 \text{subject to} &&& \text{tr}(D(e_i e_j^T + e_j e_i^T) \frac{1}{2}) = l_{ij}^2, && j-1 = (i=1 \dots 6) \bmod 6 \\
 &&& \text{tr}(-\frac{1}{2}VDV(A_i + A_i^T) \frac{1}{2}) = \cos \varphi_i, && i = 1, 2, 3 \\
 &&& \text{tr}(-\frac{1}{2}VDV) = 1 \\
 &&& -VDV \succeq 0
 \end{aligned} \tag{842}$$

where, for $A_i \in \mathbb{R}^{6 \times 6}$ (837)

$$\begin{aligned}
 A_1 &= (e_1 - e_6)(e_3 - e_4)^T / (l_{61} l_{34}) \\
 A_2 &= (e_2 - e_1)(e_4 - e_5)^T / (l_{12} l_{45}) \\
 A_3 &= (e_3 - e_2)(e_5 - e_6)^T / (l_{23} l_{56})
 \end{aligned} \tag{843}$$

and where the first constraint on length-square l_{ij}^2 can be equivalently written as a constraint on the Gram matrix $-VDV \frac{1}{2}$ via (839). We show how to numerically solve such a problem by *alternating projection* in §E.10.2.1.1. Barvinok's Proposition 2.9.3.0.1 asserts existence of a list, corresponding to Gram matrix G solving this feasibility problem, whose affine dimension (§5.7.1.1) does not exceed 3 because the convex feasible set is bounded by the third constraint $\text{tr}(-\frac{1}{2}VDV) = 1$ (833). \square

5.4.2.2.3 Example. *Kissing-number of sphere packing.*

Two nonoverlapping Euclidean balls are said to *kiss* if they touch. An elementary geometrical problem can be posed: Given hyperspheres, each having the same diameter 1, how many hyperspheres can simultaneously kiss one central hypersphere? [353] The noncentral hyperspheres are allowed, but not required, to kiss.

As posed, the problem seeks the maximal number of spheres K kissing a central sphere in a particular dimension. The total number of spheres is $N = K + 1$. In one dimension the answer to this kissing problem is 2. In two dimensions, 6. (Figure 7)

The question was presented in three dimensions to [Isaac Newton](#) in the context of celestial mechanics, and became controversy with David Gregory on the campus of Cambridge University in 1694. Newton correctly identified




Figure 99: Sphere-packing illustration from [333, *kissing number*]. Translucent balls illustrated all have the same diameter.

the kissing number as 12 (Figure 99) while Gregory argued for 13. Their dispute was finally resolved in 1953 by Schütte & van der Waerden. [263] In 2003, Oleg Musin tightened the upper bound on kissing number K in four dimensions from 25 to $K = 24$ by refining a method by Philippe Delsarte from 1973 providing an infinite number [15] of linear inequalities necessary for converting a rank-constrained semidefinite program^{5.9} to a linear program.^{5.10} [242]

There are no proofs known for kissing number in higher dimension excepting dimensions eight and twenty four.

Translating this problem to an *EDM graph* realization (Figure 96, Figure 100) is suggested by Pfender & Ziegler. Imagine the centers of each sphere are connected by line segments. Then the distance between centers must obey simple criteria: Each sphere touching the central sphere has a line segment of length exactly 1 joining its center to the central sphere's center. All spheres, excepting the central sphere, must have centers separated by a distance of at least 1.

From this perspective, the kissing problem can be posed as a semidefinite program. Assign index 1 to the central sphere, and assume a total of N

^{5.9} whose feasible set belongs to that subset of an ellipsope (§5.9.1.0.1) bounded above by some desired rank.

^{5.10} Simplex-method solvers for linear programs produce numerically better results than contemporary log-barrier (interior-point method) solvers for semidefinite programs by about 7 orders of magnitude.

spheres:

$$\begin{aligned}
 & \underset{D \in \mathbb{S}^N}{\text{minimize}} && -\text{tr} W V_N^T D V_N \\
 & \text{subject to} && D_{1j} = 1, \quad j = 2 \dots N \\
 & && D_{ij} \geq 1, \quad 2 \leq i < j = 3 \dots N \\
 & && D \in \text{EDM}^N
 \end{aligned} \tag{844}$$

Then the kissing number

$$K = N_{\max} - 1 \tag{845}$$

is found from the maximal number of spheres N that solve this semidefinite program in a given affine dimension r . Matrix W can be interpreted as the direction of search through the positive semidefinite cone \mathbb{S}_+^{N-1} for a rank- r optimal solution $-V_N^T D^* V_N$; it is constant, in this program, determined by a method disclosed in §4.4.1. In two dimensions,

$$W = \begin{bmatrix} 4 & 1 & 2 & -1 & -1 & 1 \\ 1 & 4 & -1 & -1 & 2 & 1 \\ 2 & -1 & 4 & 1 & 1 & -1 \\ -1 & -1 & 1 & 4 & 1 & 2 \\ -1 & 2 & 1 & 1 & 4 & -1 \\ 1 & 1 & -1 & 2 & -1 & 4 \end{bmatrix} \frac{1}{6} \tag{846}$$

In three dimensions,

$$W = \begin{bmatrix} 9 & 1 & -2 & -1 & 3 & -1 & -1 & 1 & 2 & 1 & -2 & 1 \\ 1 & 9 & 3 & -1 & -1 & 1 & 1 & -2 & 1 & 2 & -1 & -1 \\ -2 & 3 & 9 & 1 & 2 & -1 & -1 & 2 & -1 & -1 & 1 & 2 \\ -1 & -1 & 1 & 9 & 1 & -1 & 1 & -1 & 3 & 2 & -1 & 1 \\ 3 & -1 & 2 & 1 & 9 & 1 & 1 & -1 & -1 & -1 & 1 & -1 \\ -1 & 1 & -1 & -1 & 1 & 9 & 2 & -1 & 2 & -1 & 2 & 3 \\ -1 & 1 & -1 & 1 & 1 & 2 & 9 & 3 & -1 & 1 & -2 & -1 \\ 1 & -2 & 2 & -1 & -1 & -1 & 3 & 9 & 2 & -1 & 1 & 1 \\ 2 & 1 & -1 & 3 & -1 & 2 & -1 & 2 & 9 & -1 & 1 & -1 \\ 1 & 2 & -1 & 2 & -1 & -1 & 1 & -1 & -1 & 9 & 3 & 1 \\ -2 & -1 & 1 & -1 & 1 & 2 & -2 & 1 & 1 & 3 & 9 & -1 \\ 1 & -1 & 2 & 1 & -1 & 3 & -1 & 1 & -1 & 1 & -1 & 9 \end{bmatrix} \frac{1}{12} \tag{847}$$

A four-dimensional solution has rational direction matrix as well, but these direction matrices are not unique and their precision not critical. Here is an optimal point list^{5.11} in MATLAB output format:

^{5.11}An optimal five-dimensional point list is known: *The answer was known at least 175*

Columns 1 through 6

X =	0	-0.1983	-0.4584	0.1657	0.9399	0.7416
	0	0.6863	0.2936	0.6239	-0.2936	0.3927
	0	-0.4835	0.8146	-0.6448	0.0611	-0.4224
	0	0.5059	0.2004	-0.4093	-0.1632	0.3427

Columns 7 through 12

-0.4815	-0.9399	-0.7416	0.1983	0.4584	-0.2832
0	0.2936	-0.3927	-0.6863	-0.2936	-0.6863
-0.8756	-0.0611	0.4224	0.4835	-0.8146	-0.3922
-0.0372	0.1632	-0.3427	-0.5059	-0.2004	-0.5431

Columns 13 through 18

0.2832	-0.2926	-0.6473	0.0943	0.3640	-0.3640
0.6863	0.9176	-0.6239	-0.2313	-0.0624	0.0624
0.3922	0.1698	-0.2309	-0.6533	-0.1613	0.1613
0.5431	-0.2088	0.3721	0.7147	-0.9152	0.9152

Columns 19 through 25

-0.0943	0.6473	-0.1657	0.2926	-0.5759	0.5759	0.4815
0.2313	0.6239	-0.6239	-0.9176	0.2313	-0.2313	0
0.6533	0.2309	0.6448	-0.1698	-0.2224	0.2224	0.8756
-0.7147	-0.3721	0.4093	0.2088	-0.7520	0.7520	0.0372

This particular optimal solution was found by solving a problem sequence in increasing number of spheres.

Numerical problems begin to arise with matrices of this size due to

years ago. I believe Gauss knew it. Moreover, Korkine & Zolotarev proved in 1882 that D_5 is the densest lattice in five dimensions. So they proved that if a kissing arrangement in five dimensions can be extended to some lattice, then $k(5)=40$. Of course, the conjecture in the general case also is: $k(5)=40$. You would like to see coordinates? Easily. Let $A=\sqrt{2}$. Then $p(1)=(A, A, 0, 0, 0)$, $p(2)=(-A, A, 0, 0, 0)$, $p(3)=(A, -A, 0, 0, 0)$, ... $p(40)=(0, 0, 0, -A, -A)$; i.e., we are considering points with coordinates that have two A and three 0 with any choice of signs and any ordering of the coordinates; the same coordinates-expression in dimensions 3 and 4.

The first miracle happens in dimension 6. There are better packings than D_6 (Conjecture: $k(6)=72$). It's a real miracle how dense the packing is in eight dimensions (E_8 =Korkine & Zolotarev packing that was discovered in 1880s) and especially in dimension 24, that is the so-called Leech lattice.

Actually, people in coding theory have conjectures on the kissing numbers for dimensions up to 32 (or even greater?). However, sometimes they found better lower bounds. I know that Ericson & Zinoviev a few years ago discovered (by hand, no computer) in dimensions 13 and 14 better kissing arrangements than were known before. —Oleg Musin

interior-point methods of solution. By eliminating some equality constraints for this particular problem, matrix size can be reduced: From §5.8.3 we have

$$-V_{\mathcal{N}}^T D V_{\mathcal{N}} = \mathbf{1}\mathbf{1}^T - [\mathbf{0} \ I] D \begin{bmatrix} \mathbf{0}^T \\ I \end{bmatrix} \frac{1}{2} \quad (848)$$

(which does not hold more generally) where identity matrix $I \in \mathbb{S}^{N-1}$ has one less dimension than EDM D . By defining an EDM principal submatrix

$$\hat{D} \triangleq [\mathbf{0} \ I] D \begin{bmatrix} \mathbf{0}^T \\ I \end{bmatrix} \in \mathbb{S}_h^{N-1} \quad (849)$$

we get a convex problem equivalent to (844)

$$\begin{aligned} & \underset{\hat{D} \in \mathbb{S}^{N-1}}{\text{minimize}} && -\text{tr}(W\hat{D}) \\ & \text{subject to} && \hat{D}_{ij} \geq 1, \quad 1 \leq i < j = 2 \dots N-1 \\ & && \mathbf{1}\mathbf{1}^T - \hat{D} \frac{1}{2} \succeq 0 \\ & && \delta(\hat{D}) = \mathbf{0} \end{aligned} \quad (850)$$

Any feasible matrix $\mathbf{1}\mathbf{1}^T - \hat{D} \frac{1}{2}$ belongs to an elliptope (§5.9.1.0.1). \square

This next example shows how finding the common point of intersection for three circles in a plane, a nonlinear problem, has convex expression.

5.4.2.2.4 Example. *So \mathcal{E} Ye trilateration in wireless sensor network.* Given three known absolute point positions in \mathbb{R}^2 (three *anchors* $\check{x}_2, \check{x}_3, \check{x}_4$) and only one unknown point (one *sensor* $x_1 \in \mathbb{R}^2$), the sensor's absolute position is determined from its noiseless measured distance-square \check{d}_{i1} to each of three anchors (Figure 2, Figure 100(a)). This trilateration can be expressed as a convex optimization problem in terms of list $X \triangleq [x_1 \ \check{x}_2 \ \check{x}_3 \ \check{x}_4] \in \mathbb{R}^{2 \times 4}$ and Gram matrix $G \in \mathbb{S}^4$ (817):

$$\begin{aligned} & \underset{G \in \mathbb{S}^4, X \in \mathbb{R}^{2 \times 4}}{\text{minimize}} && \text{tr } G \\ & \text{subject to} && \text{tr}(G\Phi_{i1}) = \check{d}_{i1}, \quad i = 2, 3, 4 \\ & && \text{tr}(Ge_i e_i^T) = \|\check{x}_i\|^2, \quad i = 2, 3, 4 \\ & && \text{tr}(G(e_i e_j^T + e_j e_i^T)/2) = \check{x}_i^T \check{x}_j, \quad 2 \leq i < j = 3, 4 \\ & && X(:, 2:4) = [\check{x}_2 \ \check{x}_3 \ \check{x}_4] \\ & && \begin{bmatrix} I & X \\ X^T & G \end{bmatrix} \succeq 0 \end{aligned} \quad (851)$$




Figure 100: (a) Given three distances indicated with absolute point positions $\check{x}_2, \check{x}_3, \check{x}_4$ known and noncollinear, absolute position of x_1 in \mathbb{R}^2 can be precisely and uniquely determined by *trilateration*; solution to a system of nonlinear equations. Dimensionless EDM graphs (b) (c) (d) represent EDMs in various states of completion. Line segments represent known absolute distances and may cross without vertex at intersection. (b) Four-point list must always be embeddable in affine subset having dimension $\text{rank } V_N^T D V_N$ not exceeding 3. To determine relative position of x_2, x_3, x_4 , triangle inequality is necessary and sufficient (§5.14.1). Knowing all distance information, then (by injectivity of \mathbf{D} (§5.6)) point position x_1 is uniquely determined to within an isometry in any dimension. (c) When fifth point is introduced, only distances to x_3, x_4, x_5 are required to determine relative position of x_2 in \mathbb{R}^2 . Graph represents first instance of missing distance information; $\sqrt{d_{12}}$. (d) Three distances are absent ($\sqrt{d_{12}}, \sqrt{d_{13}}, \sqrt{d_{23}}$) from complete set of interpoint distances, yet unique isometric reconstruction (§5.4.2.2.6) of six points in \mathbb{R}^2 is certain.

where

$$\Phi_{ij} = (e_i - e_j)(e_i - e_j)^T \in \mathbb{S}_+^N \quad (806)$$

and where the constraint on distance-square \check{d}_{i1} is equivalently written as a constraint on the Gram matrix via (819). There are 9 independent affine equality constraints on that Gram matrix while the sensor is constrained only by dimensioning to lie in \mathbb{R}^2 . Although $\text{tr} G$ the objective of minimization^{5.12} insures a solution on the boundary of positive semidefinite cone \mathbb{S}_+^4 , for this problem, we claim that the set of feasible Gram matrices forms a line (§2.5.1.1) in isomorphic \mathbb{R}^{10} tangent (§2.1.7.2) to the positive semidefinite cone boundary. (§5.4.2.2.5, confer §4.2.1.3)

By Schur complement (§A.4, §2.9.1.0.1), any feasible G and X provide

$$G \succeq X^T X \quad (852)$$

which is a convex *relaxation* of the desired (nonconvex) equality constraint

$$\begin{bmatrix} I & X \\ X^T & G \end{bmatrix} = \begin{bmatrix} I \\ X^T \end{bmatrix} [I \ X] \quad (853)$$

expected positive semidefinite rank-2 under noiseless conditions. But, by (1400), the relaxation admits

$$(3 \geq) \text{rank } G \geq \text{rank } X \quad (854)$$

(a third dimension corresponding to an intersection of three spheres, not circles, were there noise). If rank of an optimal solution equals 2,

$$\text{rank} \begin{bmatrix} I & X^* \\ X^{*T} & G^* \end{bmatrix} = 2 \quad (855)$$

then $G^* = X^{*T} X^*$ by Theorem A.4.0.0.5.

As posed, this *localization* problem does not require affinely independent (Figure 25, three noncollinear) anchors. Assuming the anchors exhibit no rotational or reflective symmetry in their affine hull (§5.5.2) and assuming the sensor x_1 lies in that affine hull, then sensor position solution $x_1^* = X^*(:, 1)$ is unique under noiseless measurement. [284] \square

^{5.12}Trace ($\text{tr} G = \langle I, G \rangle$) minimization is a heuristic for rank minimization. (§7.2.2.1) It may be interpreted as squashing G which is bounded below by $X^T X$ as in (852); *id est*, $G - X^T X \succeq 0 \Rightarrow \text{tr} G \geq \text{tr} X^T X$ (1398). $\delta(G - X^T X) = \mathbf{0} \Leftrightarrow G = X^T X$ (§A.7.2) $\Rightarrow \text{tr} G = \text{tr} X^T X$ which is a condition necessary for equality.

This preceding transformation of trilateration to a semidefinite program works all the time ((855) holds) despite relaxation (852) because the optimal solution set is a unique point.

5.4.2.2.5 Proof (sketch). Only the sensor location x_1 is unknown. The objective function together with the equality constraints make a linear system of equations in Gram matrix variable G

$$\begin{aligned} \operatorname{tr} G &= \|x_1\|^2 + \|\check{x}_2\|^2 + \|\check{x}_3\|^2 + \|\check{x}_4\|^2 \\ \operatorname{tr}(G\Phi_{i1}) &= \check{d}_{i1}, & i = 2, 3, 4 \\ \operatorname{tr}(Ge_i e_i^T) &= \|\check{x}_i\|^2, & i = 2, 3, 4 \\ \operatorname{tr}(G(e_i e_j^T + e_j e_i^T)/2) &= \check{x}_i^T \check{x}_j, & 2 \leq i < j = 3, 4 \end{aligned} \quad (856)$$

which is invertible:

$$\operatorname{svec} G = \begin{bmatrix} \operatorname{svec}(I)^T \\ \operatorname{svec}(\Phi_{21})^T \\ \operatorname{svec}(\Phi_{31})^T \\ \operatorname{svec}(\Phi_{41})^T \\ \operatorname{svec}(e_2 e_2^T)^T \\ \operatorname{svec}(e_3 e_3^T)^T \\ \operatorname{svec}(e_4 e_4^T)^T \\ \operatorname{svec}((e_2 e_3^T + e_3 e_2^T)/2)^T \\ \operatorname{svec}((e_2 e_4^T + e_4 e_2^T)/2)^T \\ \operatorname{svec}((e_3 e_4^T + e_4 e_3^T)/2)^T \end{bmatrix}^{-1} \begin{bmatrix} \|x_1\|^2 + \|\check{x}_2\|^2 + \|\check{x}_3\|^2 + \|\check{x}_4\|^2 \\ \check{d}_{21} \\ \check{d}_{31} \\ \check{d}_{41} \\ \|\check{x}_2\|^2 \\ \|\check{x}_3\|^2 \\ \|\check{x}_4\|^2 \\ \check{x}_2^T \check{x}_3 \\ \check{x}_2^T \check{x}_4 \\ \check{x}_3^T \check{x}_4 \end{bmatrix} \quad (857)$$

That line in the ambient space \mathbb{S}^4 of G , claimed on page 371, is traced by $\|x_1\|^2 \in \mathbb{R}$ on the right-hand side, as it turns out. One must show this line to be tangential (§2.1.7.2) to \mathbb{S}_+^4 in order to prove uniqueness. Tangency is possible for affine dimension 1 or 2 while its occurrence depends completely on the known measurement data. ■

But as soon as significant noise is introduced or whenever distance data is incomplete, such problems can remain convex although the set of all optimal solutions generally becomes a convex set bigger than a single point (and still containing the noiseless solution).

5.4.2.2.6 Definition. *Isometric reconstruction.* (confer §5.5.3)

Isometric reconstruction from an EDM means building a list X correct to within a rotation, reflection, and translation; in other terms, reconstruction of relative position, correct to within an isometry, or to within a rigid transformation. \triangle

How much distance information is needed to uniquely localize a sensor (to recover actual relative position)? The narrative in Figure 100 helps dispel any notion of distance data proliferation in *low affine dimension* ($r < N - 2$).^{5.13} Huang, Liang, and Pardalos [183, §4.2] claim $O(2N)$ distances is a least lower bound (independent of affine dimension r) for unique isometric reconstruction; achievable under certain noiseless conditions on graph connectivity and point position. Alfakih shows how to ascertain uniqueness over all affine dimensions via *Gale matrix*. [8] [3] [4] Figure 96(b) (p.353, from *small completion problem* Example 5.3.0.0.2) is an example in \mathbb{R}^2 requiring only $2N - 3 = 5$ known symmetric entries for unique isometric reconstruction, although the four-point example in Figure 100(b) will not yield a unique reconstruction when any one of the distances is left unspecified.

The list represented by the particular *dimensionless EDM graph* in Figure 101, having only $2N - 3 = 9$ absolute distances specified, has only one realization in \mathbb{R}^2 but has more realizations in higher dimensions. For sake of reference, we provide the complete corresponding EDM:

$$D = \begin{bmatrix} 0 & 50641 & 56129 & 8245 & 18457 & 26645 \\ 50641 & 0 & 49300 & 25994 & 8810 & 20612 \\ 56129 & 49300 & 0 & 24202 & 31330 & 9160 \\ 8245 & 25994 & 24202 & 0 & 4680 & 5290 \\ 18457 & 8810 & 31330 & 4680 & 0 & 6658 \\ 26645 & 20612 & 9160 & 5290 & 6658 & 0 \end{bmatrix} \quad (858)$$

We consider paucity of distance information in this next example which shows it is possible to recover exact relative position given incomplete noiseless distance information. An *ad hoc* method for recovery of the lowest-rank optimal solution under noiseless conditions is introduced:

^{5.13}When affine dimension r reaches $N - 2$, then all distances-square in the EDM must be known for unique isometric reconstruction in \mathbb{R}^r ; going the other way, when $r = 1$ then the condition that the dimensionless EDM graph be connected is necessary and sufficient. [166, §2.2]




Figure 101: Incomplete EDM corresponding to this dimensionless EDM graph provides unique isometric reconstruction in \mathbb{R}^2 . (drawn freehand, no symmetry intended)




Figure 102: Two sensors \bullet and three anchors \circ in \mathbb{R}^2 . (Ye) Connecting line-segments denote known absolute distances. Incomplete EDM corresponding to this dimensionless EDM graph provides unique isometric reconstruction in \mathbb{R}^2 .




Figure 103: Given in red \circ are two discrete linear trajectories of sensors x_1 and x_2 in \mathbb{R}^2 localized by algorithm (859) as indicated by blue bullets \bullet . Anchors \check{x}_3 , \check{x}_4 , \check{x}_5 corresponding to Figure 102 are indicated by \otimes . When targets \circ and bullets \bullet coincide under these noiseless conditions, localization is successful. On this run, two visible localization errors are due to rank-3 Gram optimal solutions. These errors can be corrected by choosing a different normal in objective of minimization.

5.4.2.2.7 Example. *Tandem trilateration in wireless sensor network.* Given three known absolute point-positions in \mathbb{R}^2 (three anchors $\check{x}_3, \check{x}_4, \check{x}_5$) and two unknown sensors $x_1, x_2 \in \mathbb{R}^2$, the sensors' absolute positions are determinable from their noiseless distances-square (as indicated in Figure 102) assuming the anchors exhibit no rotational or reflective symmetry in their affine hull (§5.5.2). This example differs from Example 5.4.2.2.4 in so far as trilateration of each sensor is now in terms of one unknown position, the other sensor. We express this localization as a convex optimization problem (a semidefinite program, §4.1) in terms of list $X \triangleq [x_1 \ x_2 \ \check{x}_3 \ \check{x}_4 \ \check{x}_5] \in \mathbb{R}^{2 \times 5}$ and Gram matrix $G \in \mathbb{S}^5$ (817) via relaxation (852):

$$\begin{aligned}
& \underset{G \in \mathbb{S}^5, X \in \mathbb{R}^{2 \times 5}}{\text{minimize}} && \text{tr } G \\
& \text{subject to} && \text{tr}(G \Phi_{i1}) = \check{d}_{i1}, \quad i = 2, 4, 5 \\
& && \text{tr}(G \Phi_{i2}) = \check{d}_{i2}, \quad i = 3, 5 \\
& && \text{tr}(G e_i e_i^T) = \|\check{x}_i\|^2, \quad i = 3, 4, 5 \\
& && \text{tr}(G(e_i e_j^T + e_j e_i^T)/2) = \check{x}_i^T \check{x}_j, \quad 3 \leq i < j = 4, 5 \\
& && X(:, 3:5) = [\check{x}_3 \ \check{x}_4 \ \check{x}_5] \\
& && \begin{bmatrix} I & X \\ X^T & G \end{bmatrix} \succeq 0
\end{aligned} \tag{859}$$

where

$$\Phi_{ij} = (e_i - e_j)(e_i - e_j)^T \in \mathbb{S}_+^N \tag{806}$$

This problem realization is fragile because of the unknown distances between sensors and anchors. Yet there is no more information we may include beyond the 11 independent equality constraints on the Gram matrix (nonredundant constraints not antithetical) to reduce the feasible set^{5.14}. (By virtue of their dimensioning, the sensors are already constrained to \mathbb{R}^2 the affine hull of the anchors.)

Exhibited in Figure 103 are two mistakes in solution $X^*(:, 1:2)$ due to a rank-3 optimal Gram matrix G^* . The trace objective is a heuristic minimizing convex envelope of quasiconcave function^{5.15} $\text{rank } G$. (§2.9.2.6.2,

^{5.14} the presumably nonempty convex set of all points G and X satisfying the constraints.

^{5.15} Projection on that nonconvex subset of all $N \times N$ -dimensional positive semidefinite matrices, in an affine subset, whose rank does not exceed 2 is a problem considered difficult to solve. [313, §4]

§7.2.2.1) A rank-2 optimal Gram matrix can be found and the errors corrected by choosing a different normal for the linear objective function, now implicitly the identity matrix I ; *id est*,

$$\operatorname{tr} G = \langle G, I \rangle \leftarrow \langle G, \delta(u) \rangle \quad (860)$$

where vector $u \in \mathbb{R}^5$ is randomly selected. A random search for a good normal $\delta(u)$ in only a few iterations is quite easy and effective because the problem is small, an optimal solution is known *a priori* to exist in two dimensions, a good normal direction is not necessarily unique, and (we speculate) because the feasible affine-subset slices the positive semidefinite cone thinly in the Euclidean sense.^{5.16} \square

We explore ramifications of noise and incomplete data throughout; their individual effect being to expand the optimal solution set, introducing more solutions and higher-rank solutions. Hence our focus shifts in §4.4 to discovery of a reliable means for diminishing the optimal solution set by introduction of a rank constraint.

Now we illustrate how a problem in distance geometry can be solved without equality constraints representing measured distance; instead, we have only upper and lower bounds on distances measured:

5.4.2.2.8 Example. *Wireless location in a cellular telephone network.*

Utilizing measurements of distance, time of flight, angle of arrival, or signal power, *multilateration* is the process of localizing (determining absolute position of) a radio signal source \bullet by inferring geometry relative to multiple fixed *base stations* \circ whose locations are known.

We consider localization of a cellular telephone by distance geometry, so we assume distance to any particular base station can be inferred from received signal power. On a large open flat expanse of terrain, signal-power measurement corresponds well with inverse distance. But it is not uncommon for measurement of signal power to suffer 20 decibels in loss caused by factors such as *multipath* interference (signal reflections), mountainous terrain, man-made structures, turning one's head, or rolling the windows up in an automobile. Consequently, contours of equal signal power are no longer

^{5.16}The log det rank-heuristic from §7.2.2.4 does not work here because it chooses the wrong normal. Rank reduction (§4.1.1.2) is unsuccessful here because Barvinok's upper bound (§2.9.3.0.1) on rank of G^* is 4.




Figure 104: Regions of coverage by base stations \circ in a cellular telephone network. The term *cellular* arises from packing of regions best covered by neighboring base stations. Illustrated is a pentagonal *cell* best covered by base station \check{x}_2 . Like a Voronoi diagram, cell geometry depends on base-station arrangement. In some US urban environments, it is not unusual to find base stations spaced approximately 1 mile apart. There can be as many as 20 base-station antennae capable of receiving signal from any given cell phone \bullet ; practically, that number is closer to 6.




Figure 105: Some fitted contours of equal signal power in \mathbb{R}^2 transmitted from a commercial cellular telephone \bullet over about 1 mile suburban terrain outside San Francisco in 2005. Data courtesy Polaris Wireless. [285]

circular; their geometry is irregular and would more aptly be approximated by translated ellipsoids of graduated orientation and eccentricity as in Figure 105.

Depicted in Figure 104 is one cell phone x_1 whose signal power is automatically and repeatedly measured by 6 base stations \circ nearby.^{5.17} Those signal power measurements are transmitted from that cell phone to base station \check{x}_2 who decides whether to transfer (*hand-off* or *hand-over*) responsibility for that call should the user roam outside its cell.^{5.18}

Due to noise, at least one distance measurement more than the minimum number of measurements is required for reliable localization in practice; 3 measurements are minimum in two dimensions, 4 in three.^{5.19} Existence of noise precludes measured distance from the input data. We instead assign measured distance to a range estimate specified by individual upper and lower bounds: \overline{d}_{i1} is the upper bound on distance-square from the cell phone to i^{th} base station, while \underline{d}_{i1} is the lower bound. These bounds become the input data. Each measurement range is presumed different from the others.

Then convex problem (851) takes the form:

$$\begin{aligned}
 & \underset{G \in \mathbb{S}^7, X \in \mathbb{R}^{2 \times 7}}{\text{minimize}} && \text{tr } G \\
 \text{subject to} &&& \underline{d}_{i1} \leq \text{tr}(G \Phi_{i1}) \leq \overline{d}_{i1}, && i = 2 \dots 7 \\
 &&& \text{tr}(G e_i e_i^T) = \|\check{x}_i\|^2, && i = 2 \dots 7 \\
 &&& \text{tr}(G(e_i e_j^T + e_j e_i^T)/2) = \check{x}_i^T \check{x}_j, && 2 \leq i < j = 3 \dots 7 \\
 &&& X(:, 2:7) = [\check{x}_2 \ \check{x}_3 \ \check{x}_4 \ \check{x}_5 \ \check{x}_6 \ \check{x}_7] \\
 &&& \begin{bmatrix} I & X \\ X^T & G \end{bmatrix} \succeq 0 && (861)
 \end{aligned}$$

where

$$\Phi_{ij} = (e_i - e_j)(e_i - e_j)^T \in \mathbb{S}_+^N \quad (806)$$

^{5.17}Cell phone signal power is typically encoded logarithmically with 1-decibel increment and 64-decibel dynamic range.

^{5.18}Because distance to base station is quite difficult to infer from signal power measurements in an urban environment, localization of a particular cell phone • by distance geometry would be far easier were the whole cellular system instead conceived so cell phone x_1 also transmits (to base station \check{x}_2) its signal power as received by all other cell phones within range.

^{5.19}In Example 4.4.1.2.2, we explore how this convex optimization algorithm fares in the face of measurement noise.




Figure 106: A depiction of [molecular conformation](#). [99]

This semidefinite program realizes the wireless location problem illustrated in Figure 104. Location $X^*(:, 1)$ is taken as solution, although measurement noise will often cause $\text{rank } G^*$ to exceed 2. Randomized search for a rank-2 optimal solution is not so easy here as in Example 5.4.2.2.7. We introduce a method in §4.4 for enforcing the stronger rank-constraint (855). To formulate this same problem in three dimensions, point list X is simply redimensioned in the semidefinite program. \square

5.4.2.2.9 Example. (Biswas, Nigam, Ye) *Molecular Conformation.*

The subatomic measurement technique called *nuclear magnetic resonance spectroscopy* (NMR) is employed to ascertain physical conformation of molecules; *e.g.*, Figure 3, Figure 106. From this technique, distance, angle, and dihedral angle measurements can be obtained. Dihedral angles arise consequent to a phenomenon where atom subsets are physically constrained to Euclidean planes.

In the rigid covalent geometry approximation, the bond lengths and angles are treated as completely fixed, so that a given spatial structure can be described very compactly indeed by a list of torsion angles alone. . . . These are the dihedral angles between the planes spanned by the two consecutive triples in a chain of four covalently bonded atoms. [77, §1.1]

Crippen & Havel recommend working exclusively with distance data because they consider angle data to be mathematically cumbersome. The present example shows instead how inclusion of dihedral angle data into a problem statement can be made elegant and convex.

As before, ascribe position information to the matrix

$$X = [x_1 \cdots x_N] \in \mathbb{R}^{3 \times N} \quad (69)$$

and introduce a matrix \aleph holding normals η to planes respecting dihedral angles φ :

$$\aleph \triangleq [\eta_1 \cdots \eta_M] \in \mathbb{R}^{3 \times M} \quad (862)$$

As in the other examples, we preferentially work with Gram matrices G because of the bridge they provide between other variables; we define

$$\begin{bmatrix} G_{\aleph} & Z \\ Z^T & G_X \end{bmatrix} \triangleq \begin{bmatrix} \aleph^T \aleph & \aleph^T X \\ X^T \aleph & X^T X \end{bmatrix} = \begin{bmatrix} \aleph^T \\ X^T \end{bmatrix} [\aleph \ X] \in \mathbb{R}^{N+M \times N+M} \quad (863)$$

whose rank is 3 by assumption. So our problem's variables are the two Gram matrices G_X and G_{\aleph} and matrix $Z = \aleph^T X$ of cross products. Then measurements of distance-square d can be expressed as linear constraints on G_X as in (861), dihedral angle φ measurements can be expressed as linear constraints on G_{\aleph} by (837), and normal-vector η conditions can be expressed by vanishing linear constraints on cross-product matrix Z : Consider three points x labelled 1, 2, 3 assumed to lie in the ℓ^{th} plane whose normal is η_{ℓ} . There might occur, for example, the independent constraints

$$\begin{aligned} \eta_{\ell}^T(x_1 - x_2) &= 0 \\ \eta_{\ell}^T(x_2 - x_3) &= 0 \end{aligned} \quad (864)$$

which are expressible in terms of constant matrices $A \in \mathbb{R}^{M \times N}$;

$$\begin{aligned} \langle Z, A_{\ell 12} \rangle &= 0 \\ \langle Z, A_{\ell 23} \rangle &= 0 \end{aligned} \quad (865)$$

Although normals η can be constrained exactly to unit length,

$$\delta(G_{\aleph}) = \mathbf{1} \quad (866)$$

NMR data is noisy; so measurements are given as upper and lower bounds. Given bounds on dihedral angles respecting $0 \leq \varphi_j \leq \pi$ and bounds on distances d_i and given constant matrices A_k (865) and symmetric matrices Φ_i (806) and B_j per (837), then a molecular conformation problem can be expressed:

$$\begin{aligned}
& \underset{G_{\mathbb{N}} \in \mathbb{S}^M, G_X \in \mathbb{S}^N, Z \in \mathbb{R}^{M \times N}}{\text{find}} && G_X \\
& \text{subject to} && \underline{d}_i \leq \text{tr}(G_X \Phi_i) \leq \overline{d}_i, \quad \forall i \in \mathcal{I}_1 \\
& && \underline{\cos \varphi_j} \leq \text{tr}(G_{\mathbb{N}} B_j) \leq \overline{\cos \varphi_j}, \quad \forall j \in \mathcal{I}_2 \\
& && \langle Z, A_k \rangle = 0, \quad \forall k \in \mathcal{I}_3 \\
& && G_X \mathbf{1} = \mathbf{0} \\
& && \delta(G_{\mathbb{N}}) = \mathbf{1} \\
& && \begin{bmatrix} G_{\mathbb{N}} & Z \\ Z^T & G_X \end{bmatrix} \succeq 0 \\
& && \text{rank} \begin{bmatrix} G_{\mathbb{N}} & Z \\ Z^T & G_X \end{bmatrix} = 3
\end{aligned} \tag{867}$$

where $G_X \mathbf{1} = \mathbf{0}$ provides a geometrically centered list X (§5.4.2.2). Ignoring the rank constraint would tend to force cross-product matrix Z to zero. What binds these three variables is the rank constraint; we show how to satisfy it in §4.4. \square

5.4.3 Inner-product form EDM definition

[p.20] *We might, for example, realize a constellation given only interstellar distance (or, equivalently, distance from Earth and relative angular measurement; the Earth as vertex to two stars).*

Equivalent to (804) is [339, §1-7] [294, §3.2]

$$\begin{aligned}
d_{ij} &= d_{ik} + d_{kj} - 2\sqrt{d_{ik}d_{kj}} \cos \theta_{ikj} \\
&= \begin{bmatrix} \sqrt{d_{ik}} & \sqrt{d_{kj}} \end{bmatrix} \begin{bmatrix} 1 & -e^{i\theta_{ikj}} \\ -e^{-i\theta_{ikj}} & 1 \end{bmatrix} \begin{bmatrix} \sqrt{d_{ik}} \\ \sqrt{d_{kj}} \end{bmatrix}
\end{aligned} \tag{868}$$

called the *law of cosines*, where $i \triangleq \sqrt{-1}$, i, k, j are positive integers, and θ_{ikj} is the angle at vertex x_k formed by vectors $x_i - x_k$ and $x_j - x_k$;

$$\cos \theta_{ikj} = \frac{\frac{1}{2}(d_{ik} + d_{kj} - d_{ij})}{\sqrt{d_{ik}d_{kj}}} = \frac{(x_i - x_k)^T(x_j - x_k)}{\|x_i - x_k\| \|x_j - x_k\|} \tag{869}$$

where the numerator forms an inner product of vectors. Distance-square $d_{ij} \left(\begin{bmatrix} \sqrt{d_{ik}} \\ \sqrt{d_{kj}} \end{bmatrix} \right)$ is a convex quadratic function^{5.20} on \mathbb{R}_+^2 whereas $d_{ij}(\theta_{ikj})$ is quasiconvex (§3.3) minimized over domain $\{-\pi \leq \theta_{ikj} \leq \pi\}$ by $\theta_{ikj}^* = 0$, we get the *Pythagorean theorem* when $\theta_{ikj} = \pm\pi/2$, and $d_{ij}(\theta_{ikj})$ is maximized when $\theta_{ikj}^* = \pm\pi$;

$$\begin{aligned} d_{ij} &= (\sqrt{d_{ik}} + \sqrt{d_{kj}})^2, & \theta_{ikj} &= \pm\pi \\ d_{ij} &= d_{ik} + d_{kj}, & \theta_{ikj} &= \pm\frac{\pi}{2} \\ d_{ij} &= (\sqrt{d_{ik}} - \sqrt{d_{kj}})^2, & \theta_{ikj} &= 0 \end{aligned} \quad (870)$$

so

$$|\sqrt{d_{ik}} - \sqrt{d_{kj}}| \leq \sqrt{d_{ij}} \leq \sqrt{d_{ik}} + \sqrt{d_{kj}} \quad (871)$$

Hence the triangle inequality, Euclidean metric property 4, holds for any EDM D .

We may construct an inner-product form of the EDM definition for matrices by evaluating (868) for $k=1$: By defining

$$\Theta^T \Theta \triangleq \begin{bmatrix} d_{12} & \sqrt{d_{12}d_{13}} \cos \theta_{213} & \sqrt{d_{12}d_{14}} \cos \theta_{214} & \cdots & \sqrt{d_{12}d_{1N}} \cos \theta_{21N} \\ \sqrt{d_{12}d_{13}} \cos \theta_{213} & d_{13} & \sqrt{d_{13}d_{14}} \cos \theta_{314} & \cdots & \sqrt{d_{13}d_{1N}} \cos \theta_{31N} \\ \sqrt{d_{12}d_{14}} \cos \theta_{214} & \sqrt{d_{13}d_{14}} \cos \theta_{314} & d_{14} & \ddots & \sqrt{d_{14}d_{1N}} \cos \theta_{41N} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ \sqrt{d_{12}d_{1N}} \cos \theta_{21N} & \sqrt{d_{13}d_{1N}} \cos \theta_{31N} & \sqrt{d_{14}d_{1N}} \cos \theta_{41N} & \cdots & d_{1N} \end{bmatrix} \in \mathbb{S}^{N-1} \quad (872)$$

then any EDM may be expressed

$$\begin{aligned} \mathbf{D}(\Theta) &\triangleq \begin{bmatrix} 0 \\ \delta(\Theta^T \Theta) \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & \delta(\Theta^T \Theta)^T \end{bmatrix} - 2 \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \Theta^T \Theta \end{bmatrix} \in \text{EDM}^N \\ &= \begin{bmatrix} 0 & \delta(\Theta^T \Theta)^T \\ \delta(\Theta^T \Theta) & \delta(\Theta^T \Theta) \mathbf{1}^T + \mathbf{1} \delta(\Theta^T \Theta)^T - 2\Theta^T \Theta \end{bmatrix} \end{aligned} \quad (873)$$

$$\text{EDM}^N = \{ \mathbf{D}(\Theta) \mid \Theta \in \mathbb{R}^{N-1 \times N-1} \} \quad (874)$$

for which all Euclidean metric properties hold. Entries of $\Theta^T \Theta$ result from vector inner-products as in (869); *id est*,

^{5.20} $\begin{bmatrix} 1 & -e^{\gamma\theta_{ikj}} \\ -e^{-\gamma\theta_{ikj}} & 1 \end{bmatrix} \succeq 0$, having eigenvalues $\{0, 2\}$. Minimum is attained for $\begin{bmatrix} \sqrt{d_{ik}} \\ \sqrt{d_{kj}} \end{bmatrix} = \begin{cases} \mu \mathbf{1}, & \mu \geq 0, \theta_{ikj} = 0 \\ \mathbf{0}, & -\pi \leq \theta_{ikj} \leq \pi, \theta_{ikj} \neq 0 \end{cases}$. (§D.2.1, [53, exmp.4.5])

$$\Theta = [x_2 - x_1 \quad x_3 - x_1 \quad \cdots \quad x_N - x_1] = X\sqrt{2}V_N \in \mathbb{R}^{n \times N-1} \quad (875)$$

Inner product $\Theta^T\Theta$ is obviously related to a Gram matrix (817),

$$G = \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \Theta^T\Theta \end{bmatrix}, \quad x_1 = \mathbf{0} \quad (876)$$

For $D = \mathbf{D}(\Theta)$ and no condition on the list X (confer (825) (830))

$$\Theta^T\Theta = -V_N^T D V_N \in \mathbb{R}^{N-1 \times N-1} \quad (877)$$

5.4.3.1 Relative-angle form

The inner-product form EDM definition is not a unique definition of Euclidean distance matrix; there are approximately five flavors distinguished by their argument to operator \mathbf{D} . Here is another one:

Like $\mathbf{D}(X)$ (808), $\mathbf{D}(\Theta)$ will make an EDM given any $\Theta \in \mathbb{R}^{n \times N-1}$, it is neither a convex function of Θ (§5.4.3.2), and it is homogeneous in the sense (811). Scrutinizing $\Theta^T\Theta$ (872) we find that because of the arbitrary choice $k=1$, distances therein are all with respect to point x_1 . Similarly, relative angles in $\Theta^T\Theta$ are between all vector pairs having vertex x_1 . Yet picking arbitrary θ_{i1j} to fill $\Theta^T\Theta$ will not necessarily make an EDM; inner product (872) must be positive semidefinite.

$$\Theta^T\Theta = \sqrt{\delta(d)} \Omega \sqrt{\delta(d)} \triangleq \begin{bmatrix} \sqrt{d_{12}} & & & \mathbf{0} \\ & \sqrt{d_{13}} & & \\ & & \ddots & \\ \mathbf{0} & & & \sqrt{d_{1N}} \end{bmatrix} \begin{bmatrix} 1 & \cos \theta_{213} & \cdots & \cos \theta_{21N} \\ \cos \theta_{213} & 1 & \ddots & \cos \theta_{31N} \\ \vdots & \ddots & \ddots & \vdots \\ \cos \theta_{21N} & \cos \theta_{31N} & \cdots & 1 \end{bmatrix} \begin{bmatrix} \sqrt{d_{12}} & & & \mathbf{0} \\ & \sqrt{d_{13}} & & \\ & & \ddots & \\ \mathbf{0} & & & \sqrt{d_{1N}} \end{bmatrix} \quad (878)$$

Expression $\mathbf{D}(\Theta)$ defines an EDM for any positive semidefinite *relative-angle matrix*

$$\Omega = [\cos \theta_{i1j}, \quad i, j = 2 \dots N] \in \mathbb{S}^{N-1} \quad (879)$$

and any nonnegative distance vector

$$d = [d_{1j}, \quad j = 2 \dots N] = \delta(\Theta^T\Theta) \in \mathbb{R}^{N-1} \quad (880)$$

because (§A.3.1.0.5)

$$\Omega \succeq 0 \Rightarrow \Theta^T \Theta \succeq 0 \quad (881)$$

Decomposition (878) and the *relative-angle matrix inequality* $\Omega \succeq 0$ lead to a different expression of an inner-product form EDM definition (873)

$$\begin{aligned} \mathbf{D}(\Omega, d) &\triangleq \begin{bmatrix} 0 \\ d \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & d^T \end{bmatrix} - 2\sqrt{\delta\left(\begin{bmatrix} 0 \\ d \end{bmatrix}\right)} \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \Omega \end{bmatrix} \sqrt{\delta\left(\begin{bmatrix} 0 \\ d \end{bmatrix}\right)} \\ &= \begin{bmatrix} 0 & & d^T \\ d & d\mathbf{1}^T + \mathbf{1}d^T - 2\sqrt{\delta(d)}\Omega\sqrt{\delta(d)} & \end{bmatrix} \in \mathbb{EDM}^N \end{aligned} \quad (882)$$

and another expression of the EDM cone:

$$\mathbb{EDM}^N = \left\{ \mathbf{D}(\Omega, d) \mid \Omega \succeq 0, \sqrt{\delta(d)} \succeq 0 \right\} \quad (883)$$

In the particular circumstance $x_1 = \mathbf{0}$, we can relate interpoint angle matrix Ψ from the Gram decomposition in (817) to relative-angle matrix Ω in (878). Thus,

$$\Psi \equiv \begin{bmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & \Omega \end{bmatrix}, \quad x_1 = \mathbf{0} \quad (884)$$

5.4.3.2 Inner-product form $-V_{\mathcal{N}}^T \mathbf{D}(\Theta) V_{\mathcal{N}}$ convexity

On page 383 we saw that each EDM entry d_{ij} is a convex quadratic function of $\left[\begin{smallmatrix} \sqrt{d_{ik}} \\ \sqrt{d_{kj}} \end{smallmatrix} \right]$ and a quasiconvex function of θ_{ikj} . Here the situation for inner-product form EDM operator $\mathbf{D}(\Theta)$ (873) is identical to that in §5.4.1 for list-form $\mathbf{D}(X)$; $-\mathbf{D}(\Theta)$ is not a quasiconvex function of Θ by the same reasoning, and from (877)

$$-V_{\mathcal{N}}^T \mathbf{D}(\Theta) V_{\mathcal{N}} = \Theta^T \Theta \quad (885)$$

is a convex quadratic function of Θ on domain $\mathbb{R}^{n \times N-1}$ achieving its minimum at $\Theta = \mathbf{0}$.

5.4.3.3 Inner-product form, discussion

We deduce that knowledge of interpoint distance is equivalent to knowledge of distance and angle from the perspective of one point, x_1 in our chosen case. The total amount of information $N(N-1)/2$ in $\Theta^T\Theta$ is unchanged^{5.21} with respect to EDM D .

5.5 Invariance

When D is an EDM, there exist an infinite number of corresponding N -point lists X (69) in Euclidean space. All those lists are related by isometric transformation: rotation, reflection, and translation (*offset* or *shift*).

5.5.1 Translation

Any translation common among all the points x_ℓ in a list will be cancelled in the formation of each d_{ij} . Proof follows directly from (804). Knowing that translation α in advance, we may remove it from the list constituting the columns of X by subtracting $\alpha\mathbf{1}^T$. Then it stands to reason by list-form definition (808) of an EDM, for any translation $\alpha \in \mathbb{R}^n$

$$\mathbf{D}(X - \alpha\mathbf{1}^T) = \mathbf{D}(X) \quad (886)$$

In words, interpoint distances are unaffected by offset; EDM D is *translation invariant*. When $\alpha = x_1$ in particular,

$$[x_2 - x_1 \quad x_3 - x_1 \quad \cdots \quad x_N - x_1] = X\sqrt{2}V_N \in \mathbb{R}^{n \times N-1} \quad (875)$$

and so

$$\mathbf{D}(X - x_1\mathbf{1}^T) = \mathbf{D}(X - Xe_1\mathbf{1}^T) = \mathbf{D}\left(X \begin{bmatrix} \mathbf{0} & \sqrt{2}V_N \end{bmatrix}\right) = \mathbf{D}(X) \quad (887)$$

^{5.21}The reason for amount $O(N^2)$ information is because of the relative measurements. Use of a fixed reference in measurement of angles and distances would reduce required information but is antithetical. In the particular case $n=2$, for example, ordering all points x_ℓ (in a length- N list) by increasing angle of vector $x_\ell - x_1$ with respect to $x_2 - x_1$, θ_{i1j} becomes equivalent to $\sum_{k=i}^{j-1} \theta_{k,1,k+1} \leq 2\pi$ and the amount of information is reduced to $2N-3$; rather, $O(N)$.

5.5.1.0.1 Example. *Translating geometric center to origin.*

We might choose to shift the geometric center α_c of an N -point list $\{x_\ell\}$ (arranged columnar in X) to the origin; [312] [142]

$$\alpha = \alpha_c \triangleq Xb_c \triangleq \frac{1}{N}X\mathbf{1} \in \mathcal{P} \subseteq \mathcal{A} \quad (888)$$

where \mathcal{A} represents the list's affine hull. If we were to associate a point-mass m_ℓ with each of the points x_ℓ in the list, then their *center of mass* (or *gravity*) would be $(\sum x_\ell m_\ell) / \sum m_\ell$. The geometric center is the same as the center of mass under the assumption of uniform mass density across points. [194] The geometric center always lies in the convex hull \mathcal{P} of the list; *id est*, $\alpha_c \in \mathcal{P}$ because $b_c^T \mathbf{1} = 1$ and $b_c \succeq 0$.^{5.22} Subtracting the geometric center from every list member,

$$X - \alpha_c \mathbf{1}^T = X - \frac{1}{N}X\mathbf{1}\mathbf{1}^T = X(I - \frac{1}{N}\mathbf{1}\mathbf{1}^T) = XV \in \mathbb{R}^{n \times N} \quad (889)$$

where V is the geometric centering matrix (831). So we have (*confer* (808))

$$\mathbf{D}(X) = \mathbf{D}(XV) = \delta(V^T X^T X V) \mathbf{1}^T + \mathbf{1} \delta(V^T X^T X V)^T - 2V^T X^T X V \in \mathbb{EDM}^N \quad (890)$$

□

5.5.1.1 Gram-form invariance

Following from (890) and the linear Gram-form EDM operator (820):

$$\mathbf{D}(G) = \mathbf{D}(VGV) = \delta(VGV) \mathbf{1}^T + \mathbf{1} \delta(VGV)^T - 2VGV \in \mathbb{EDM}^N \quad (891)$$

The Gram-form consequently exhibits invariance to translation by a *doublet* (§B.2) $u\mathbf{1}^T + \mathbf{1}u^T$;

$$\mathbf{D}(G) = \mathbf{D}(G - (u\mathbf{1}^T + \mathbf{1}u^T)) \quad (892)$$

because, for any $u \in \mathbb{R}^N$, $\mathbf{D}(u\mathbf{1}^T + \mathbf{1}u^T) = \mathbf{0}$. The collection of all such doublets forms the nullspace (908) to the operator; the *translation-invariant subspace* $\mathbb{S}_c^{N\perp}$ (1887) of the symmetric matrices \mathbb{S}^N . This means matrix G can belong to an expanse more broad than a positive semidefinite cone; *id est*, $G \in \mathbb{S}_+^N - \mathbb{S}_c^{N\perp}$. So explains Gram matrix sufficiency in EDM definition (820).^{5.23}

^{5.22}Any b from $\alpha = Xb$ chosen such that $b^T \mathbf{1} = 1$, more generally, makes an auxiliary V -matrix. (§B.4.5)

^{5.23}A constraint $G\mathbf{1} = \mathbf{0}$ would prevent excursion into the translation-invariant subspace (numerical unboundedness).

5.5.2 Rotation/Reflection

Rotation of the list $X \in \mathbb{R}^{n \times N}$ about some arbitrary point $\alpha \in \mathbb{R}^n$, or reflection through some affine subset containing α , can be accomplished via $Q(X - \alpha \mathbf{1}^T)$ where Q is an orthogonal matrix (§B.5).

We rightfully expect

$$\mathbf{D}(Q(X - \alpha \mathbf{1}^T)) = \mathbf{D}(QX - \beta \mathbf{1}^T) = \mathbf{D}(QX) = \mathbf{D}(X) \quad (893)$$

Because list-form $\mathbf{D}(X)$ is translation invariant, we may safely ignore offset and consider only the impact of matrices that premultiply X . Interpoint distances are unaffected by rotation or reflection; we say, EDM D is *rotation/reflection invariant*. Proof follows from the fact, $Q^T = Q^{-1} \Rightarrow X^T Q^T Q X = X^T X$. So (893) follows directly from (808).

The class of premultiplying matrices for which interpoint distances are unaffected is a little more broad than orthogonal matrices. Looking at EDM definition (808), it appears that any matrix Q_p such that

$$X^T Q_p^T Q_p X = X^T X \quad (894)$$

will have the property

$$\mathbf{D}(Q_p X) = \mathbf{D}(X) \quad (895)$$

An example is skinny $Q_p \in \mathbb{R}^{m \times n}$ ($m > n$) having orthonormal columns. We call such a matrix *orthonormal*.

5.5.2.0.1 Example. Reflection prevention and quadrature rotation.

Consider the EDM graph in Figure 107(b) representing known distance between vertices (Figure 107(a)) of a tilted-square diamond in \mathbb{R}^2 . Suppose some geometrical optimization problem were posed where isometric transformation is feasible excepting reflection, and where rotation must be quantized so that only *quadrature* rotations are feasible; only multiples of $\pi/2$.

In two dimensions, a counterclockwise rotation of any vector about the origin by angle θ is prescribed by the orthogonal matrix

$$Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \quad (896)$$




Figure 107: (a) Four points in quadrature in two dimensions about their geometric center. (b) Complete EDM graph of diamond vertices. (c) Quadrature rotation of a Euclidean body in \mathbb{R}^2 first requires a shroud that is the smallest Cartesian square containing it.

whereas reflection of any point through a hyperplane containing the origin

$$\partial\mathcal{H} = \left\{ x \in \mathbb{R}^2 \mid \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}^\top x = 0 \right\} \tag{897}$$

is accomplished via multiplication with symmetric orthogonal matrix (§B.5.2)

$$R = \begin{bmatrix} \sin(\theta)^2 - \cos(\theta)^2 & -2 \sin(\theta) \cos(\theta) \\ -2 \sin(\theta) \cos(\theta) & \cos(\theta)^2 - \sin(\theta)^2 \end{bmatrix} \tag{898}$$

Rotation matrix Q is characterized by identical diagonal entries and by antidiagonal entries equal but opposite in sign, whereas reflection matrix R is characterized in the reverse sense.

Assign the diamond vertices $\{x_\ell \in \mathbb{R}^2, \ell = 1 \dots 4\}$ to columns of a matrix

$$X = [x_1 \ x_2 \ x_3 \ x_4] \in \mathbb{R}^{2 \times 4} \tag{69}$$

Our scheme to prevent reflection enforces a rotation matrix characteristic upon the coordinates of adjacent points themselves: First shift the geometric center of X to the origin; for geometric centering matrix $V \in \mathbb{S}^4$ (§5.5.1.0.1), define

$$Y \triangleq XV \in \mathbb{R}^{2 \times 4} \tag{899}$$

To maintain relative quadrature between points (Figure 107(a)) and to prevent reflection, it is sufficient that all interpoint distances be specified and that adjacencies $Y(:, 1:2)$, $Y(:, 2:3)$, and $Y(:, 3:4)$ be proportional to 2×2 rotation matrices; any clockwise rotation would ascribe a reflection matrix characteristic. Counterclockwise rotation is thereby enforced by constraining equality among diagonal and antidiagonal entries as prescribed by (896);

$$Y(:, 1:3) = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} Y(:, 2:4) \quad (900)$$

Quadrature quantization of rotation can be regarded as a constraint on tilt of the smallest Cartesian square containing the diamond as in Figure 107(c). Our scheme to quantize rotation requires that all square vertices be described by vectors whose entries are nonnegative when the square is translated anywhere interior to the nonnegative orthant. We capture the four square vertices as columns of a product YC where

$$C = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix} \quad (901)$$

Then, assuming a unit-square shroud, the affine constraint

$$YC + \begin{bmatrix} 1/2 \\ 1/2 \end{bmatrix} \mathbf{1}^T \geq \mathbf{0} \quad (902)$$

quantizes rotation, as desired. \square

5.5.2.1 Inner-product form invariance

Likewise, $\mathbf{D}(\Theta)$ (873) is rotation/reflection invariant;

$$\mathbf{D}(Q_p \Theta) = \mathbf{D}(Q \Theta) = \mathbf{D}(\Theta) \quad (903)$$

so (894) and (895) similarly apply.

5.5.3 Invariance conclusion

In the making of an EDM, absolute rotation, reflection, and translation information is lost. Given an EDM, reconstruction of point position (§5.12, the list X) can be guaranteed correct only in affine dimension r and relative position. Given a noiseless complete EDM, this isometric reconstruction is unique in so far as every realization of a corresponding list X is *congruent*:

5.6 Injectivity of \mathbf{D} & unique reconstruction

Injectivity implies uniqueness of isometric reconstruction; hence, we endeavor to demonstrate it.

EDM operators list-form $\mathbf{D}(X)$ (808), Gram-form $\mathbf{D}(G)$ (820), and inner-product form $\mathbf{D}(\Theta)$ (873) are many-to-one surjections (§5.5) onto the same range; the EDM cone (§6): (*confer*(821) (910))

$$\begin{aligned} \text{EDM}^N &= \{\mathbf{D}(X) : \mathbb{R}^{N-1 \times N} \rightarrow \mathbb{S}_h^N \mid X \in \mathbb{R}^{N-1 \times N}\} \\ &= \{\mathbf{D}(G) : \mathbb{S}^N \rightarrow \mathbb{S}_h^N \mid G \in \mathbb{S}_+^N - \mathbb{S}_c^{N\perp}\} \\ &= \{\mathbf{D}(\Theta) : \mathbb{R}^{N-1 \times N-1} \rightarrow \mathbb{S}_h^N \mid \Theta \in \mathbb{R}^{N-1 \times N-1}\} \end{aligned} \quad (904)$$

where (§5.5.1.1)

$$\mathbb{S}_c^{N\perp} = \{u\mathbf{1}^T + \mathbf{1}u^T \mid u \in \mathbb{R}^N\} \subseteq \mathbb{S}^N \quad (1887)$$

5.6.1 Gram-form bijectivity

Because linear Gram-form EDM operator

$$\mathbf{D}(G) = \delta(G)\mathbf{1}^T + \mathbf{1}\delta(G)^T - 2G \quad (820)$$

has no nullspace [75, §A.1] on the geometric center subspace^{5.24} (§E.7.2.0.2)

$$\begin{aligned} \mathbb{S}_c^N &\triangleq \{G \in \mathbb{S}^N \mid G\mathbf{1} = \mathbf{0}\} \quad (1885) \\ &= \{G \in \mathbb{S}^N \mid \mathcal{N}(G) \supseteq \mathbf{1}\} = \{G \in \mathbb{S}^N \mid \mathcal{R}(G) \subseteq \mathcal{N}(\mathbf{1}^T)\} \\ &= \{VYV \mid Y \in \mathbb{S}^N\} \subset \mathbb{S}^N \quad (1886) \\ &\equiv \{V_{\mathcal{N}}AV_{\mathcal{N}}^T \mid A \in \mathbb{S}^{N-1}\} \end{aligned} \quad (905)$$

then $\mathbf{D}(G)$ on that subspace is injective.

^{5.24}The equivalence \equiv in (905) follows from the fact: Given $B = VYV = V_{\mathcal{N}}AV_{\mathcal{N}}^T \in \mathbb{S}_c^N$ with only matrix $A \in \mathbb{S}^{N-1}$ unknown, then $V_{\mathcal{N}}^\dagger BV_{\mathcal{N}}^{\dagger T} = A$ or $V_{\mathcal{N}}^\dagger YV_{\mathcal{N}}^{\dagger T} = A$.




Figure 108: Orthogonal complements in \mathbb{S}^N abstractly oriented in isometrically isomorphic $\mathbb{R}^{N(N+1)/2}$. Case $N=2$ accurately illustrated in \mathbb{R}^3 . Orthogonal projection of basis for $\mathbb{S}_h^{N\perp}$ on $\mathbb{S}_c^{N\perp}$ yields another basis for $\mathbb{S}_c^{N\perp}$. (Basis vectors for $\mathbb{S}_c^{N\perp}$ are illustrated lying in a plane orthogonal to \mathbb{S}_c^N in this dimension. Basis vectors for each \perp space outnumber those for its respective orthogonal complement; such is not the case in higher dimension.)

To prove injectivity of $\mathbf{D}(G)$ on \mathbb{S}_c^N : Any matrix $Y \in \mathbb{S}^N$ can be decomposed into orthogonal components in \mathbb{S}^N ;

$$Y = VYV + (Y - VYV) \quad (906)$$

where $VYV \in \mathbb{S}_c^N$ and $Y - VYV \in \mathbb{S}_c^{N\perp}$ (1887). Because of translation invariance (§5.5.1.1) and linearity, $\mathbf{D}(Y - VYV) = \mathbf{0}$ hence $\mathcal{N}(\mathbf{D}) \supseteq \mathbb{S}_c^{N\perp}$. It remains only to show

$$\mathbf{D}(VYV) = \mathbf{0} \Leftrightarrow VYV = \mathbf{0} \quad (907)$$

($\Leftrightarrow Y = u\mathbf{1}^T + \mathbf{1}u^T$ for some $u \in \mathbb{R}^N$). $\mathbf{D}(VYV)$ will vanish whenever $2VYV = \delta(VYV)\mathbf{1}^T + \mathbf{1}\delta(VYV)^T$. But this implies $\mathcal{R}(\mathbf{1})$ (§B.2) were a subset of $\mathcal{R}(VYV)$, which is contradictory. Thus we have

$$\mathcal{N}(\mathbf{D}) = \{Y \mid \mathbf{D}(Y) = \mathbf{0}\} = \{Y \mid VYV = \mathbf{0}\} = \mathbb{S}_c^{N\perp} \quad (908)$$

◆

Since $G\mathbf{1} = \mathbf{0} \Leftrightarrow X\mathbf{1} = \mathbf{0}$ (829) simply means list X is geometrically centered at the origin, and because the Gram-form EDM operator \mathbf{D} is translation invariant and $\mathcal{N}(\mathbf{D})$ is the translation-invariant subspace $\mathbb{S}_c^{N\perp}$, then EDM definition $\mathbf{D}(G)$ (904) on^{5.25} (confer §6.6.1, §6.7.1, §A.7.4.0.1)

$$\mathbb{S}_c^N \cap \mathbb{S}_+^N = \{VYV \succeq 0 \mid Y \in \mathbb{S}^N\} \equiv \{V_{\mathcal{N}}AV_{\mathcal{N}}^T \mid A \in \mathbb{S}_+^{N-1}\} \subset \mathbb{S}^N \quad (909)$$

must be surjective onto \mathbb{EDM}^N ; (confer (821))

$$\mathbb{EDM}^N = \{\mathbf{D}(G) \mid G \in \mathbb{S}_c^N \cap \mathbb{S}_+^N\} \quad (910)$$

5.6.1.1 Gram-form operator \mathbf{D} inversion

Define the linear *geometric centering operator* \mathbf{V} ; (confer (830))

$$\mathbf{V}(D) : \mathbb{S}^N \rightarrow \mathbb{S}^N \triangleq -VDV\frac{1}{2} \quad (911)$$

[78, §4.3]^{5.26} This orthogonal projector \mathbf{V} has no nullspace on

$$\mathbb{S}_h^N = \text{aff } \mathbb{EDM}^N \quad (1164)$$

^{5.25}Equivalence \equiv in (909) follows from the fact: Given $B = VYV = V_{\mathcal{N}}AV_{\mathcal{N}}^T \in \mathbb{S}_+^N$ with only matrix A unknown, then $V_{\mathcal{N}}^\dagger BV_{\mathcal{N}}^{\dagger T} = A$ and $A \in \mathbb{S}_+^{N-1}$ must be positive semidefinite by positive semidefiniteness of B and Corollary A.3.1.0.5.

^{5.26}Critchley cites Torgerson (1958) [309, ch.11, §2] for a history and derivation of (911).

because the projection of $-D/2$ on \mathbb{S}_c^N (1885) can be $\mathbf{0}$ if and only if $D \in \mathbb{S}_c^{N\perp}$; but $\mathbb{S}_c^{N\perp} \cap \mathbb{S}_h^N = \mathbf{0}$ (Figure 108). Projector \mathbf{V} on \mathbb{S}_h^N is therefore injective hence uniquely invertible. Further, $-V\mathbb{S}_h^N V/2$ is equivalent to the geometric center subspace \mathbb{S}_c^N in the ambient space of symmetric matrices; a surjection,

$$\mathbb{S}_c^N = \mathbf{V}(\mathbb{S}^N) = \mathbf{V}(\mathbb{S}_h^N \oplus \mathbb{S}_h^{N\perp}) = \mathbf{V}(\mathbb{S}_h^N) \quad (912)$$

because (66)

$$\mathbf{V}(\mathbb{S}_h^N) \supseteq \mathbf{V}(\mathbb{S}_h^{N\perp}) = \mathbf{V}(\delta^2(\mathbb{S}^N)) \quad (913)$$

Because $\mathbf{D}(G)$ on \mathbb{S}_c^N is injective, and $\text{aff } \mathbf{D}(\mathbf{V}(\text{EDM}^N)) = \mathbf{D}(\mathbf{V}(\text{aff EDM}^N))$ by property (119) of the affine hull, we find for $D \in \mathbb{S}_h^N$ (confer (834))

$$\mathbf{D}(-VDV\frac{1}{2}) = \delta(-VDV\frac{1}{2})\mathbf{1}^T + \mathbf{1}\delta(-VDV\frac{1}{2})^T - 2(-VDV\frac{1}{2}) \quad (914)$$

id est,

$$D = \mathbf{D}(\mathbf{V}(D)) \quad (915)$$

$$-VDV = \mathbf{V}(\mathbf{D}(-VDV)) \quad (916)$$

or

$$\mathbb{S}_h^N = \mathbf{D}(\mathbf{V}(\mathbb{S}_h^N)) \quad (917)$$

$$-V\mathbb{S}_h^N V = \mathbf{V}(\mathbf{D}(-V\mathbb{S}_h^N V)) \quad (918)$$

These operators \mathbf{V} and \mathbf{D} are mutual inverses.

The Gram-form $\mathbf{D}(\mathbb{S}_c^N)$ (820) is equivalent to \mathbb{S}_h^N ;

$$\mathbf{D}(\mathbb{S}_c^N) = \mathbf{D}(\mathbf{V}(\mathbb{S}_h^N \oplus \mathbb{S}_h^{N\perp})) = \mathbb{S}_h^N + \mathbf{D}(\mathbf{V}(\mathbb{S}_h^{N\perp})) = \mathbb{S}_h^N \quad (919)$$

because $\mathbb{S}_h^N \supseteq \mathbf{D}(\mathbf{V}(\mathbb{S}_h^{N\perp}))$. In summary, for the Gram-form we have the isomorphisms [79, §2] [78, p.76, p.107] [6, §2.1]^{5.27} [5, §2] [7, §18.2.1] [1, §2.1]

$$\mathbb{S}_h^N = \mathbf{D}(\mathbb{S}_c^N) \quad (920)$$

$$\mathbb{S}_c^N = \mathbf{V}(\mathbb{S}_h^N) \quad (921)$$

and from the bijectivity results in §5.6.1,

$$\text{EDM}^N = \mathbf{D}(\mathbb{S}_c^N \cap \mathbb{S}_+^N) \quad (922)$$

$$\mathbb{S}_c^N \cap \mathbb{S}_+^N = \mathbf{V}(\text{EDM}^N) \quad (923)$$

^{5.27}In [6, p.6, line 20], delete sentence: *Since G is also ... not a singleton set.*
[6, p.10, line 11] $x_3=2$ (not 1).

5.6.2 Inner-product form bijectivity

The Gram-form EDM operator $\mathbf{D}(G) = \delta(G)\mathbf{1}^T + \mathbf{1}\delta(G)^T - 2G$ (820) is an injective map, for example, on the domain that is the subspace of symmetric matrices having all zeros in the first row and column

$$\begin{aligned} \mathbb{S}_1^N &\triangleq \{G \in \mathbb{S}^N \mid Ge_1 = \mathbf{0}\} \\ &= \left\{ \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} Y \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} \mid Y \in \mathbb{S}^N \right\} \end{aligned} \quad (1889)$$

because it obviously has no nullspace there. Since $Ge_1 = \mathbf{0} \Leftrightarrow Xe_1 = \mathbf{0}$ (822) means the first point in the list X resides at the origin, then $\mathbf{D}(G)$ on $\mathbb{S}_1^N \cap \mathbb{S}_+^N$ must be surjective onto \mathbb{EDM}^N .

Substituting $\Theta^T\Theta \leftarrow -V_{\mathcal{N}}^T D V_{\mathcal{N}}$ (885) into inner-product form EDM definition $\mathbf{D}(\Theta)$ (873), it may be further decomposed: (*confer* (828))

$$\mathbf{D}(D) = \begin{bmatrix} 0 \\ \delta(-V_{\mathcal{N}}^T D V_{\mathcal{N}}) \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & \delta(-V_{\mathcal{N}}^T D V_{\mathcal{N}})^T \end{bmatrix} - 2 \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & -V_{\mathcal{N}}^T D V_{\mathcal{N}} \end{bmatrix} \quad (924)$$

This linear operator \mathbf{D} is another flavor of inner-product form and an injective map of the EDM cone onto itself. Yet when its domain is instead the entire symmetric hollow subspace $\mathbb{S}_h^N = \text{aff } \mathbb{EDM}^N$, $\mathbf{D}(D)$ becomes an injective map onto that same subspace. Proof follows directly from the fact: linear \mathbf{D} has no nullspace [75, §A.1] on $\mathbb{S}_h^N = \text{aff } \mathbf{D}(\mathbb{EDM}^N) = \mathbf{D}(\text{aff } \mathbb{EDM}^N)$ (119).

5.6.2.1 Inversion of $\mathbf{D}(-V_{\mathcal{N}}^T D V_{\mathcal{N}})$

Injectivity of $\mathbf{D}(D)$ suggests inversion of (*confer* (825))

$$\mathbf{V}_{\mathcal{N}}(D) : \mathbb{S}^N \rightarrow \mathbb{S}^{N-1} \triangleq -V_{\mathcal{N}}^T D V_{\mathcal{N}} \quad (925)$$

a linear surjective^{5.28} mapping onto \mathbb{S}^{N-1} having nullspace^{5.29} $\mathbb{S}_c^{N\perp}$;

$$\mathbf{V}_{\mathcal{N}}(\mathbb{S}_h^N) = \mathbb{S}^{N-1} \quad (926)$$

^{5.28}Surjectivity of $\mathbf{V}_{\mathcal{N}}(D)$ is demonstrated via the Gram-form EDM operator $\mathbf{D}(G)$: Since $\mathbb{S}_h^N = \mathbf{D}(\mathbb{S}_c^N)$ (919), then for any $Y \in \mathbb{S}^{N-1}$, $-V_{\mathcal{N}}^T \mathbf{D}(V_{\mathcal{N}}^{\dagger T} Y V_{\mathcal{N}}^{\dagger} / 2) V_{\mathcal{N}} = Y$.

^{5.29} $\mathcal{N}(\mathbf{V}_{\mathcal{N}}) \supseteq \mathbb{S}_c^{N\perp}$ is apparent. There exists a linear mapping

$$T(\mathbf{V}_{\mathcal{N}}(D)) \triangleq V_{\mathcal{N}}^{\dagger T} \mathbf{V}_{\mathcal{N}}(D) V_{\mathcal{N}}^{\dagger} = -V D V \frac{1}{2} = \mathbf{V}(D)$$

injective on domain \mathbb{S}_h^N because $\mathbb{S}_c^{N\perp} \cap \mathbb{S}_h^N = \mathbf{0}$. Revising the argument of this inner-product form (924), we get another flavor

$$\mathbf{D}(-V_{\mathcal{N}}^T D V_{\mathcal{N}}) = \begin{bmatrix} 0 \\ \delta(-V_{\mathcal{N}}^T D V_{\mathcal{N}}) \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & \delta(-V_{\mathcal{N}}^T D V_{\mathcal{N}})^T \end{bmatrix} - 2 \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & -V_{\mathcal{N}}^T D V_{\mathcal{N}} \end{bmatrix} \quad (927)$$

and we obtain mutual inversion of operators $\mathbf{V}_{\mathcal{N}}$ and \mathbf{D} , for $D \in \mathbb{S}_h^N$

$$D = \mathbf{D}(\mathbf{V}_{\mathcal{N}}(D)) \quad (928)$$

$$-V_{\mathcal{N}}^T D V_{\mathcal{N}} = \mathbf{V}_{\mathcal{N}}(\mathbf{D}(-V_{\mathcal{N}}^T D V_{\mathcal{N}})) \quad (929)$$

or

$$\mathbb{S}_h^N = \mathbf{D}(\mathbf{V}_{\mathcal{N}}(\mathbb{S}_h^N)) \quad (930)$$

$$-V_{\mathcal{N}}^T \mathbb{S}_h^N V_{\mathcal{N}} = \mathbf{V}_{\mathcal{N}}(\mathbf{D}(-V_{\mathcal{N}}^T \mathbb{S}_h^N V_{\mathcal{N}})) \quad (931)$$

Substituting $\Theta^T \Theta \leftarrow \Phi$ into inner-product form EDM definition (873), any EDM may be expressed by the new flavor

$$\begin{aligned} \mathbf{D}(\Phi) &\triangleq \begin{bmatrix} 0 \\ \delta(\Phi) \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & \delta(\Phi)^T \end{bmatrix} - 2 \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \Phi \end{bmatrix} \in \mathbb{EDM}^N \\ &\Leftrightarrow \\ &\Phi \succeq 0 \end{aligned} \quad (932)$$

where this \mathbf{D} is a linear surjective operator onto \mathbb{EDM}^N by definition, injective because it has no nullspace on domain \mathbb{S}_+^{N-1} . More broadly, $\text{aff } \mathbf{D}(\mathbb{S}_+^{N-1}) = \mathbf{D}(\text{aff } \mathbb{S}_+^{N-1})$ (119),

$$\begin{aligned} \mathbb{S}_h^N &= \mathbf{D}(\mathbb{S}^{N-1}) \\ \mathbb{S}^{N-1} &= \mathbf{V}_{\mathcal{N}}(\mathbb{S}_h^N) \end{aligned} \quad (933)$$

demonstrably isomorphisms, and by bijectivity of this inner-product form:

$$\mathbb{EDM}^N = \mathbf{D}(\mathbb{S}_+^{N-1}) \quad (934)$$

$$\mathbb{S}_+^{N-1} = \mathbf{V}_{\mathcal{N}}(\mathbb{EDM}^N) \quad (935)$$

such that

$$\mathcal{N}(T(\mathbf{V}_{\mathcal{N}})) = \mathcal{N}(\mathbf{V}) \supseteq \mathcal{N}(\mathbf{V}_{\mathcal{N}}) \supseteq \mathbb{S}_c^{N\perp} = \mathcal{N}(\mathbf{V})$$

where the equality $\mathbb{S}_c^{N\perp} = \mathcal{N}(\mathbf{V})$ is known (§E.7.2.0.2). \blacklozenge

5.7 Embedding in affine hull

The affine hull \mathcal{A} (71) of a point list $\{x_\ell\}$ (arranged columnar in $X \in \mathbb{R}^{n \times N}$ (69)) is identical to the affine hull of that polyhedron \mathcal{P} (79) formed from all convex combinations of the x_ℓ ; [53, §2] [271, §17]

$$\mathcal{A} = \text{aff } X = \text{aff } \mathcal{P} \quad (936)$$

Comparing hull definitions (71) and (79), it becomes obvious that the x_ℓ and their convex hull \mathcal{P} are embedded in their unique affine hull \mathcal{A} ;

$$\mathcal{A} \supseteq \mathcal{P} \supseteq \{x_\ell\} \quad (937)$$

Recall: affine dimension r is a lower bound on embedding, equal to dimension of the subspace parallel to that nonempty affine set \mathcal{A} in which the points are embedded. (§2.3.1) We define dimension of the convex hull \mathcal{P} to be the same as dimension r of the affine hull \mathcal{A} [271, §2], but r is not necessarily equal to rank of X (956).

For the particular example illustrated in Figure 95, \mathcal{P} is the triangle in union with its relative interior while its three vertices constitute the entire list X . Affine hull \mathcal{A} is the unique plane that contains the triangle, so affine dimension $r = 2$ in that example while rank of X is 3. Were there only two points in Figure 95, then the affine hull would instead be the unique line passing through them; r would become 1 while rank would then be 2.

5.7.1 Determining affine dimension

Knowledge of affine dimension r becomes important because we lose any absolute offset common to all the generating x_ℓ in \mathbb{R}^n when reconstructing convex polyhedra given only distance information. (§5.5.1) To calculate r , we first remove any offset that serves to increase dimensionality of the subspace required to contain polyhedron \mathcal{P} ; subtracting any $\alpha \in \mathcal{A}$ in the affine hull from every list member will work,

$$X - \alpha \mathbf{1}^T \quad (938)$$

translating \mathcal{A} to the origin:^{5.30}

$$\mathcal{A} - \alpha = \text{aff}(X - \alpha \mathbf{1}^T) = \text{aff}(X) - \alpha \quad (939)$$

$$\mathcal{P} - \alpha = \text{conv}(X - \alpha \mathbf{1}^T) = \text{conv}(X) - \alpha \quad (940)$$

^{5.30}The manipulation of hull functions aff and conv follows from their definitions.

Because (936) and (937) translate,

$$\mathbb{R}^n \supseteq \mathcal{A} - \alpha = \text{aff}(X - \alpha \mathbf{1}^T) = \text{aff}(\mathcal{P} - \alpha) \supseteq \mathcal{P} - \alpha \supseteq \{x_\ell - \alpha\} \quad (941)$$

where from the previous relations it is easily shown

$$\text{aff}(\mathcal{P} - \alpha) = \text{aff}(\mathcal{P}) - \alpha \quad (942)$$

Translating \mathcal{A} neither changes its dimension or the dimension of the embedded polyhedron \mathcal{P} ; (70)

$$r \triangleq \dim \mathcal{A} = \dim(\mathcal{A} - \alpha) \triangleq \dim(\mathcal{P} - \alpha) = \dim \mathcal{P} \quad (943)$$

For any $\alpha \in \mathbb{R}^n$, (939)-(943) remain true. [271, p.4, p.12] Yet when $\alpha \in \mathcal{A}$, the affine set $\mathcal{A} - \alpha$ becomes a unique subspace of \mathbb{R}^n in which the $\{x_\ell - \alpha\}$ and their convex hull $\mathcal{P} - \alpha$ are embedded (941), and whose dimension is more easily calculated.

5.7.1.0.1 Example. *Translating first list-member to origin.*

Subtracting the first member $\alpha \triangleq x_1$ from every list member will translate their affine hull \mathcal{A} and their convex hull \mathcal{P} and, in particular, $x_1 \in \mathcal{P} \subseteq \mathcal{A}$ to the origin in \mathbb{R}^n ; *videlicet*,

$$X - x_1 \mathbf{1}^T = X - X e_1 \mathbf{1}^T = X(I - e_1 \mathbf{1}^T) = X \begin{bmatrix} \mathbf{0} & \sqrt{2} V_{\mathcal{N}} \end{bmatrix} \in \mathbb{R}^{n \times N} \quad (944)$$

where $V_{\mathcal{N}}$ is defined in (814), and e_1 in (824). Applying (941) to (944),

$$\mathbb{R}^n \supseteq \mathcal{R}(XV_{\mathcal{N}}) = \mathcal{A} - x_1 = \text{aff}(X - x_1 \mathbf{1}^T) = \text{aff}(\mathcal{P} - x_1) \supseteq \mathcal{P} - x_1 \ni \mathbf{0} \quad (945)$$

where $XV_{\mathcal{N}} \in \mathbb{R}^{n \times N-1}$. Hence

$$r = \dim \mathcal{R}(XV_{\mathcal{N}}) \quad (946)$$

□

Since shifting the geometric center to the origin (§5.5.1.0.1) translates the affine hull to the origin as well, then it must also be true

$$r = \dim \mathcal{R}(XV) \quad (947)$$

For any matrix whose range is $\mathcal{R}(V) = \mathcal{N}(\mathbf{1}^\top)$ we get the same result; *e.g.*,

$$r = \dim \mathcal{R}(XV_{\mathcal{N}}^{\dagger\top}) \quad (948)$$

because

$$\mathcal{R}(XV) = \{Xz \mid z \in \mathcal{N}(\mathbf{1}^\top)\} \quad (949)$$

and $\mathcal{R}(V) = \mathcal{R}(V_{\mathcal{N}}) = \mathcal{R}(V_{\mathcal{N}}^{\dagger\top})$ (§E). These auxiliary matrices (§B.4.2) are more closely related;

$$V = V_{\mathcal{N}}V_{\mathcal{N}}^{\dagger} \quad (1553)$$

5.7.1.1 Affine dimension r versus rank

Now, suppose D is an EDM as defined by

$$\mathbf{D}(X) \triangleq \delta(X^\top X)\mathbf{1}^\top + \mathbf{1}\delta(X^\top X)^\top - 2X^\top X \in \text{EDM}^N \quad (808)$$

and we premultiply by $-V_{\mathcal{N}}^\top$ and postmultiply by $V_{\mathcal{N}}$. Then because $V_{\mathcal{N}}^\top \mathbf{1} = \mathbf{0}$ (815), it is always true that

$$-V_{\mathcal{N}}^\top D V_{\mathcal{N}} = 2V_{\mathcal{N}}^\top X^\top X V_{\mathcal{N}} = 2V_{\mathcal{N}}^\top G V_{\mathcal{N}} \in \mathbb{S}^{N-1} \quad (950)$$

where G is a Gram matrix. Similarly pre- and postmultiplying by V (*confer* (830))

$$-VDV = 2VX^\top XV = 2VGV \in \mathbb{S}^N \quad (951)$$

always holds because $V\mathbf{1} = \mathbf{0}$ (1543). Likewise, multiplying inner-product form EDM definition (873), it always holds:

$$-V_{\mathcal{N}}^\top D V_{\mathcal{N}} = \Theta^\top \Theta \in \mathbb{S}^{N-1} \quad (877)$$

For any matrix A , $\text{rank } A^\top A = \text{rank } A = \text{rank } A^\top$. [180, §0.4]^{5.31} So, by (949), affine dimension

$$\begin{aligned} r &= \text{rank } XV = \text{rank } XV_{\mathcal{N}} = \text{rank } XV_{\mathcal{N}}^{\dagger\top} = \text{rank } \Theta \\ &= \text{rank } VDV = \text{rank } VGV = \text{rank } V_{\mathcal{N}}^\top D V_{\mathcal{N}} = \text{rank } V_{\mathcal{N}}^\top G V_{\mathcal{N}} \end{aligned} \quad (952)$$

^{5.31}For $A \in \mathbb{R}^{m \times n}$, $\mathcal{N}(A^\top A) = \mathcal{N}(A)$. [294, §3.3]

By conservation of dimension, (§A.7.3.0.1)

$$r + \dim \mathcal{N}(V_{\mathcal{N}}^T D V_{\mathcal{N}}) = N - 1 \quad (953)$$

$$r + \dim \mathcal{N}(V D V) = N \quad (954)$$

For $D \in \text{EDM}^N$

$$-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succ 0 \Leftrightarrow r = N - 1 \quad (955)$$

but $-V D V \not\succeq 0$. The general fact^{5.32} (confer (841))

$$r \leq \min\{n, N - 1\} \quad (956)$$

is evident from (944) but can be visualized in the example illustrated in Figure 95. There we imagine a vector from the origin to each point in the list. Those three vectors are linearly independent in \mathbb{R}^3 , but affine dimension r is 2 because the three points lie in a plane. When that plane is translated to the origin, it becomes the only subspace of dimension $r=2$ that can contain the translated triangular polyhedron.

5.7.2 Précis

We collect expressions for affine dimension: for list $X \in \mathbb{R}^{n \times N}$ and Gram matrix $G \in \mathbb{S}_+^N$

$$\begin{aligned}
 r &\stackrel{\Delta}{=} \dim(\mathcal{P} - \alpha) = \dim \mathcal{P} = \dim \text{conv } X \\
 &= \dim(\mathcal{A} - \alpha) = \dim \mathcal{A} = \dim \text{aff } X \\
 &= \text{rank}(X - x_1 \mathbf{1}^T) = \text{rank}(X - \alpha_c \mathbf{1}^T) \\
 &= \text{rank } \Theta \quad (875) \\
 &= \text{rank } X V_{\mathcal{N}} = \text{rank } X V = \text{rank } X V_{\mathcal{N}}^{\dagger T} \\
 &= \text{rank } X, \quad X e_1 = \mathbf{0} \quad \text{or} \quad X \mathbf{1} = \mathbf{0} \\
 &= \text{rank } V_{\mathcal{N}}^T G V_{\mathcal{N}} = \text{rank } V G V = \text{rank } V_{\mathcal{N}}^{\dagger} G V_{\mathcal{N}} \\
 &= \text{rank } G, \quad G e_1 = \mathbf{0} \quad (825) \quad \text{or} \quad G \mathbf{1} = \mathbf{0} \quad (830) \\
 &= \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} = \text{rank } V D V = \text{rank } V_{\mathcal{N}}^{\dagger} D V_{\mathcal{N}} = \text{rank } V_{\mathcal{N}} (V_{\mathcal{N}}^T D V_{\mathcal{N}}) V_{\mathcal{N}}^T \\
 &= \text{rank } \Lambda \quad (1043) \\
 &= N - 1 - \dim \mathcal{N} \left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \right) = \text{rank} \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} - 2 \quad (964)
 \end{aligned}
 \left. \vphantom{\begin{aligned} r &\stackrel{\Delta}{=} \dim(\mathcal{P} - \alpha) = \dim \mathcal{P} = \dim \text{conv } X \\ &= \dim(\mathcal{A} - \alpha) = \dim \mathcal{A} = \dim \text{aff } X \\ &= \text{rank}(X - x_1 \mathbf{1}^T) = \text{rank}(X - \alpha_c \mathbf{1}^T) \\ &= \text{rank } \Theta \quad (875) \\ &= \text{rank } X V_{\mathcal{N}} = \text{rank } X V = \text{rank } X V_{\mathcal{N}}^{\dagger T} \\ &= \text{rank } X, \quad X e_1 = \mathbf{0} \quad \text{or} \quad X \mathbf{1} = \mathbf{0} \\ &= \text{rank } V_{\mathcal{N}}^T G V_{\mathcal{N}} = \text{rank } V G V = \text{rank } V_{\mathcal{N}}^{\dagger} G V_{\mathcal{N}} \\ &= \text{rank } G, \quad G e_1 = \mathbf{0} \quad (825) \quad \text{or} \quad G \mathbf{1} = \mathbf{0} \quad (830) \\ &= \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} = \text{rank } V D V = \text{rank } V_{\mathcal{N}}^{\dagger} D V_{\mathcal{N}} = \text{rank } V_{\mathcal{N}} (V_{\mathcal{N}}^T D V_{\mathcal{N}}) V_{\mathcal{N}}^T \\ &= \text{rank } \Lambda \quad (1043) \\ &= N - 1 - \dim \mathcal{N} \left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \right) = \text{rank} \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} - 2 \quad (964) \end{aligned}} \right\} D \in \text{EDM}^N$$

^{5.32} $\text{rank } X \leq \min\{n, N\}$

5.7.3 Eigenvalues of $-VDV$ versus $-V_{\mathcal{N}}^{\dagger}DV_{\mathcal{N}}$

Suppose for $D \in \text{EDM}^N$ we are given eigenvectors $v_i \in \mathbb{R}^N$ of $-VDV$ and corresponding eigenvalues $\lambda \in \mathbb{R}^N$ so that

$$-VDVv_i = \lambda_i v_i, \quad i = 1 \dots N \quad (958)$$

From these we can determine the eigenvectors and eigenvalues of $-V_{\mathcal{N}}^{\dagger}DV_{\mathcal{N}}$: Define

$$\nu_i \triangleq V_{\mathcal{N}}^{\dagger} v_i, \quad \lambda_i \neq 0 \quad (959)$$

Then we have:

$$-VDV_{\mathcal{N}}V_{\mathcal{N}}^{\dagger}v_i = \lambda_i v_i \quad (960)$$

$$-V_{\mathcal{N}}^{\dagger}VDV_{\mathcal{N}}\nu_i = \lambda_i V_{\mathcal{N}}^{\dagger}v_i \quad (961)$$

$$-V_{\mathcal{N}}^{\dagger}DV_{\mathcal{N}}\nu_i = \lambda_i \nu_i \quad (962)$$

the eigenvectors of $-V_{\mathcal{N}}^{\dagger}DV_{\mathcal{N}}$ are given by (959) while its corresponding nonzero eigenvalues are identical to those of $-VDV$ although $-V_{\mathcal{N}}^{\dagger}DV_{\mathcal{N}}$ is not necessarily positive semidefinite. In contrast, $-V_{\mathcal{N}}^{\text{T}}DV_{\mathcal{N}}$ is positive semidefinite but its nonzero eigenvalues are generally different.

5.7.3.0.1 Theorem. *EDM rank versus affine dimension r .*
[142, §3] [163, §3] [141, §3] For $D \in \text{EDM}^N$ (confer (1119))

1. $r = \text{rank}(D) - 1 \Leftrightarrow \mathbf{1}^{\text{T}}D^{\dagger}\mathbf{1} \neq 0$

Points constituting a list X generating the polyhedron corresponding to D lie on the relative boundary of an r -dimensional *circumhypersphere* having

$$\begin{aligned} \text{diameter} &= \sqrt{2} (\mathbf{1}^{\text{T}}D^{\dagger}\mathbf{1})^{-1/2} \\ \text{circumcenter} &= \frac{XD^{\dagger}\mathbf{1}}{\mathbf{1}^{\text{T}}D^{\dagger}\mathbf{1}} \end{aligned} \quad (963)$$

2. $r = \text{rank}(D) - 2 \Leftrightarrow \mathbf{1}^{\text{T}}D^{\dagger}\mathbf{1} = 0$

There can be no circumhypersphere whose relative boundary contains a generating list for the corresponding polyhedron.

3. In *Cayley-Menger form* [96, §6.2] [77, §3.3] [44, §40] (§5.11.2),

$$r = N - 1 - \dim \mathcal{N} \left(\begin{bmatrix} 0 & \mathbf{1}^{\text{T}} \\ \mathbf{1} & -D \end{bmatrix} \right) = \text{rank} \begin{bmatrix} 0 & \mathbf{1}^{\text{T}} \\ \mathbf{1} & -D \end{bmatrix} - 2 \quad (964)$$

Circumhyperspheres exist for $r < \text{rank}(D) - 2$. [308, §7] \diamond

For all practical purposes, (956)

$$\max\{0, \text{rank}(D) - 2\} \leq r \leq \min\{n, N - 1\} \quad (965)$$

5.8 Euclidean metric *versus* matrix criteria

5.8.1 Nonnegativity property 1

When $D = [d_{ij}]$ is an EDM (808), then it is apparent from (950)

$$2V_{\mathcal{N}}^T X^T X V_{\mathcal{N}} = -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \quad (966)$$

because for any matrix A , $A^T A \succeq 0$.^{5.33} We claim nonnegativity of the d_{ij} is enforced primarily by the matrix inequality (966); *id est*,

$$\left. \begin{array}{l} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \\ D \in \mathbb{S}_h^N \end{array} \right\} \Rightarrow d_{ij} \geq 0, \quad i \neq j \quad (967)$$

(The matrix inequality to enforce strict positivity differs by a stroke of the pen. (970))

We now support our claim: If any matrix $A \in \mathbb{R}^{m \times m}$ is positive semidefinite, then its main diagonal $\delta(A) \in \mathbb{R}^m$ must have all nonnegative entries. [139, §4.2] Given $D \in \mathbb{S}_h^N$

$$\begin{aligned} & -V_{\mathcal{N}}^T D V_{\mathcal{N}} = \\ & \left[\begin{array}{cccc} d_{12} & \frac{1}{2}(d_{12} + d_{13} - d_{23}) & \frac{1}{2}(d_{1,i+1} + d_{1,j+1} - d_{i+1,j+1}) & \cdots & \frac{1}{2}(d_{12} + d_{1N} - d_{2N}) \\ \frac{1}{2}(d_{12} + d_{13} - d_{23}) & d_{13} & \frac{1}{2}(d_{1,i+1} + d_{1,j+1} - d_{i+1,j+1}) & \cdots & \frac{1}{2}(d_{13} + d_{1N} - d_{3N}) \\ \frac{1}{2}(d_{1,j+1} + d_{1,i+1} - d_{j+1,i+1}) & \frac{1}{2}(d_{1,j+1} + d_{1,i+1} - d_{j+1,i+1}) & d_{1,i+1} & \ddots & \frac{1}{2}(d_{14} + d_{1N} - d_{4N}) \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ \frac{1}{2}(d_{12} + d_{1N} - d_{2N}) & \frac{1}{2}(d_{13} + d_{1N} - d_{3N}) & \frac{1}{2}(d_{14} + d_{1N} - d_{4N}) & \cdots & d_{1N} \end{array} \right] \\ & = \frac{1}{2}(\mathbf{1}D_{1,2:N} + D_{2:N,1}\mathbf{1}^T - D_{2:N,2:N}) \in \mathbb{S}^{N-1} \quad (968) \end{aligned}$$

^{5.33}For $A \in \mathbb{R}^{m \times n}$, $A^T A \succeq 0 \Leftrightarrow y^T A^T A y = \|Ay\|^2 \geq 0$ for all $\|y\| = 1$. When A is full-rank skinny-or-square, $A^T A \succ 0$.

where row, column indices $i, j \in \{1 \dots N-1\}$. [276] It follows:

$$\left. \begin{array}{l} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \\ D \in \mathbb{S}_h^N \end{array} \right\} \Rightarrow \delta(-V_{\mathcal{N}}^T D V_{\mathcal{N}}) = \begin{bmatrix} d_{12} \\ d_{13} \\ \vdots \\ d_{1N} \end{bmatrix} \succeq 0 \quad (969)$$

Multiplication of $V_{\mathcal{N}}$ by any permutation matrix Ξ has null effect on its range and nullspace. In other words, any permutation of the rows or columns of $V_{\mathcal{N}}$ produces a basis for $\mathcal{N}(\mathbf{1}^T)$; *id est*, $\mathcal{R}(\Xi_r V_{\mathcal{N}}) = \mathcal{R}(V_{\mathcal{N}} \Xi_c) = \mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^T)$. Hence, $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \Leftrightarrow -V_{\mathcal{N}}^T \Xi_r^T D \Xi_r V_{\mathcal{N}} \succeq 0$ ($\Leftrightarrow -\Xi_c^T V_{\mathcal{N}}^T D V_{\mathcal{N}} \Xi_c \succeq 0$). Various permutation matrices^{5.34} will sift the remaining d_{ij} similarly to (969) thereby proving their nonnegativity. Hence $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$ is a sufficient test for the first property (§5.2) of the Euclidean metric, nonnegativity. \blacklozenge

When affine dimension r equals 1, in particular, nonnegativity symmetry and hollowness become necessary and sufficient criteria satisfying matrix inequality (966). (§6.6.0.0.1)

5.8.1.1 Strict positivity

Should we require the points in \mathbb{R}^n to be distinct, then entries of D off the main diagonal must be strictly positive $\{d_{ij} > 0, i \neq j\}$ and only those entries along the main diagonal of D are 0. By similar argument, the strict matrix inequality is a sufficient test for strict positivity of Euclidean distance-square;

$$\left. \begin{array}{l} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succ 0 \\ D \in \mathbb{S}_h^N \end{array} \right\} \Rightarrow d_{ij} > 0, \quad i \neq j \quad (970)$$

^{5.34}The rule of thumb is: If $\Xi_r(i, 1) = 1$, then $\delta(-V_{\mathcal{N}}^T \Xi_r^T D \Xi_r V_{\mathcal{N}}) \in \mathbb{R}^{N-1}$ is some permutation of the i^{th} row or column of D excepting the 0 entry from the main diagonal.

5.8.2 Triangle inequality property 4

In light of Kreyszig's observation [201, §1.1, prob.15] that properties 2 through 4 of the Euclidean metric (§5.2) together imply nonnegativity property 1,

$$2\sqrt{d_{jk}} = \sqrt{d_{jk}} + \sqrt{d_{kj}} \geq \sqrt{d_{jj}} = 0, \quad j \neq k \quad (971)$$

nonnegativity criterion (967) suggests that the matrix inequality $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$ might somehow take on the role of triangle inequality; *id est*,

$$\left. \begin{array}{l} \delta(D) = \mathbf{0} \\ D^T = D \\ -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \end{array} \right\} \Rightarrow \sqrt{d_{ij}} \leq \sqrt{d_{ik}} + \sqrt{d_{kj}}, \quad i \neq j \neq k \quad (972)$$

We now show that is indeed the case: Let T be the *leading principal submatrix* in \mathbb{S}^2 of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ (upper left 2×2 submatrix from (968));

$$T \triangleq \begin{bmatrix} d_{12} & \frac{1}{2}(d_{12} + d_{13} - d_{23}) \\ \frac{1}{2}(d_{12} + d_{13} - d_{23}) & d_{13} \end{bmatrix} \quad (973)$$

Submatrix T must be positive (semi)definite whenever $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ is. (§A.3.1.0.4, §5.8.3) Now we have,

$$\begin{aligned} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 &\Rightarrow T \succeq 0 \Leftrightarrow \lambda_1 \geq \lambda_2 \geq 0 \\ -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succ 0 &\Rightarrow T \succ 0 \Leftrightarrow \lambda_1 > \lambda_2 > 0 \end{aligned} \quad (974)$$

where λ_1 and λ_2 are the eigenvalues of T , real due only to symmetry of T :

$$\begin{aligned} \lambda_1 &= \frac{1}{2} \left(d_{12} + d_{13} + \sqrt{d_{23}^2 - 2(d_{12} + d_{13})d_{23} + 2(d_{12}^2 + d_{13}^2)} \right) \in \mathbb{R} \\ \lambda_2 &= \frac{1}{2} \left(d_{12} + d_{13} - \sqrt{d_{23}^2 - 2(d_{12} + d_{13})d_{23} + 2(d_{12}^2 + d_{13}^2)} \right) \in \mathbb{R} \end{aligned} \quad (975)$$

Nonnegativity of eigenvalue λ_1 is guaranteed by only nonnegativity of the d_{ij} which in turn is guaranteed by matrix inequality (967). Inequality between the eigenvalues in (974) follows from only realness of the d_{ij} . Since λ_1 always equals or exceeds λ_2 , conditions for the positive (semi)definiteness of submatrix T can be completely determined by examining λ_2 the smaller of its two eigenvalues. A triangle inequality is made apparent when we express

T eigenvalue nonnegativity in terms of D matrix entries; *videlicet*,

$$\begin{aligned} T \succeq 0 &\Leftrightarrow \det T = \lambda_1 \lambda_2 \geq 0, \quad d_{12}, d_{13} \geq 0 & \text{(c)} \\ &\Leftrightarrow \\ &\lambda_2 \geq 0 & \text{(b)} \\ &\Leftrightarrow \\ |\sqrt{d_{12}} - \sqrt{d_{23}}| &\leq \sqrt{d_{13}} \leq \sqrt{d_{12}} + \sqrt{d_{23}} & \text{(a)} \end{aligned} \tag{976}$$

Triangle inequality (976a) (*confer* (871) (988)), in terms of three rooted entries from D , is equivalent to metric property 4

$$\begin{aligned} \sqrt{d_{13}} &\leq \sqrt{d_{12}} + \sqrt{d_{23}} \\ \sqrt{d_{23}} &\leq \sqrt{d_{12}} + \sqrt{d_{13}} \\ \sqrt{d_{12}} &\leq \sqrt{d_{13}} + \sqrt{d_{23}} \end{aligned} \tag{977}$$

for the corresponding points x_1, x_2, x_3 from some length- N list.^{5.35}

5.8.2.1 Comment

Given D whose dimension N equals or exceeds 3, there are $N!/(3!(N-3)!)$ distinct triangle inequalities in total like (871) that must be satisfied, of which each d_{ij} is involved in $N-2$, and each point x_i is in $(N-1)!/(2!(N-1-2)!)$. We have so far revealed only one of those triangle inequalities; namely, (976a) that came from T (973). Yet we claim if $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$ then all triangle inequalities will be satisfied simultaneously;

$$|\sqrt{d_{ik}} - \sqrt{d_{kj}}| \leq \sqrt{d_{ij}} \leq \sqrt{d_{ik}} + \sqrt{d_{kj}}, \quad i < k < j \tag{978}$$

(There are no more.) To verify our claim, we must prove the matrix inequality $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$ to be a sufficient test of all the triangle inequalities; more efficient, we mention, for larger N :

^{5.35}Accounting for symmetry property 3, the fourth metric property demands three inequalities be satisfied per one of type (976a). The first of those inequalities in (977) is self evident from (976a), while the two remaining follow from the left-hand side of (976a) and the fact for scalars, $|a| \leq b \Leftrightarrow a \leq b$ and $-a \leq b$.

5.8.2.1.1 Shore. The columns of $\Xi_r V_{\mathcal{N}} \Xi_c$ hold a basis for $\mathcal{N}(\mathbf{1}^T)$ when Ξ_r and Ξ_c are permutation matrices. In other words, any permutation of the rows or columns of $V_{\mathcal{N}}$ leaves its range and nullspace unchanged; *id est*, $\mathcal{R}(\Xi_r V_{\mathcal{N}} \Xi_c) = \mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^T)$ (815). Hence, two distinct matrix inequalities can be equivalent tests of the positive semidefiniteness of D on $\mathcal{R}(V_{\mathcal{N}})$; *id est*, $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \Leftrightarrow -(\Xi_r V_{\mathcal{N}} \Xi_c)^T D (\Xi_r V_{\mathcal{N}} \Xi_c) \succeq 0$. By properly choosing permutation matrices,^{5.36} the leading principal submatrix $T_{\Xi} \in \mathbb{S}^2$ of $-(\Xi_r V_{\mathcal{N}} \Xi_c)^T D (\Xi_r V_{\mathcal{N}} \Xi_c)$ may be loaded with the entries of D needed to test any particular triangle inequality (similarly to (968)-(976)). Because all the triangle inequalities can be individually tested using a test equivalent to the lone matrix inequality $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$, it logically follows that the lone matrix inequality tests all those triangle inequalities simultaneously. We conclude that $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$ is a sufficient test for the fourth property of the Euclidean metric, triangle inequality. \blacklozenge

5.8.2.2 Strict triangle inequality

Without exception, all the inequalities in (976) and (977) can be made strict while their corresponding implications remain true. The then strict inequality (976a) or (977) may be interpreted as a *strict triangle inequality* under which collinear arrangement of points is not allowed. [198, §24/6, p.322] Hence by similar reasoning, $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succ 0$ is a sufficient test of all the strict triangle inequalities; *id est*,

$$\left. \begin{array}{l} \delta(D) = \mathbf{0} \\ D^T = D \\ -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succ 0 \end{array} \right\} \Rightarrow \sqrt{d_{ij}} < \sqrt{d_{ik}} + \sqrt{d_{kj}}, \quad i \neq j \neq k \quad (979)$$

5.8.3 $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ nesting

From (973) observe that $T = -V_{\mathcal{N}}^T D V_{\mathcal{N}}|_{N \leftarrow 3}$. In fact, for $D \in \text{EDM}^N$, the leading principal submatrices of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ form a nested sequence (by inclusion) whose members are individually positive semidefinite [139] [180] [294] and have the same form as T ; *videlicet*,^{5.37}

^{5.36}To individually test triangle inequality $|\sqrt{d_{ik}} - \sqrt{d_{kj}}| \leq \sqrt{d_{ij}} \leq \sqrt{d_{ik}} + \sqrt{d_{kj}}$ for particular i, k, j , set $\Xi_r(i, 1) = \Xi_r(k, 2) = \Xi_r(j, 3) = 1$ and $\Xi_c = I$.

^{5.37} $-V D V|_{N \leftarrow 1} = 0 \in \mathbb{S}_+^0$ (§B.4.1)

$$-V_{\mathcal{N}}^T DV_{\mathcal{N}}|_{N \leftarrow 1} = [\emptyset] \quad (\text{o})$$

$$-V_{\mathcal{N}}^T DV_{\mathcal{N}}|_{N \leftarrow 2} = [d_{12}] \in \mathbb{S}_+ \quad (\text{a})$$

$$-V_{\mathcal{N}}^T DV_{\mathcal{N}}|_{N \leftarrow 3} = \begin{bmatrix} d_{12} & \frac{1}{2}(d_{12} + d_{13} - d_{23}) \\ \frac{1}{2}(d_{12} + d_{13} - d_{23}) & d_{13} \end{bmatrix} = T \in \mathbb{S}_+^2 \quad (\text{b})$$

$$-V_{\mathcal{N}}^T DV_{\mathcal{N}}|_{N \leftarrow 4} = \begin{bmatrix} d_{12} & \frac{1}{2}(d_{12} + d_{13} - d_{23}) & \frac{1}{2}(d_{12} + d_{14} - d_{24}) \\ \frac{1}{2}(d_{12} + d_{13} - d_{23}) & d_{13} & \frac{1}{2}(d_{13} + d_{14} - d_{34}) \\ \frac{1}{2}(d_{12} + d_{14} - d_{24}) & \frac{1}{2}(d_{13} + d_{14} - d_{34}) & d_{14} \end{bmatrix} \quad (\text{c})$$

$$\vdots$$

$$-V_{\mathcal{N}}^T DV_{\mathcal{N}}|_{N \leftarrow i} = \begin{bmatrix} -V_{\mathcal{N}}^T DV_{\mathcal{N}}|_{N \leftarrow i-1} & \nu(i) \\ \nu(i)^T & d_{1i} \end{bmatrix} \in \mathbb{S}_+^{i-1} \quad (\text{d})$$

$$\vdots$$

$$-V_{\mathcal{N}}^T DV_{\mathcal{N}} = \begin{bmatrix} -V_{\mathcal{N}}^T DV_{\mathcal{N}}|_{N \leftarrow N-1} & \nu(N) \\ \nu(N)^T & d_{1N} \end{bmatrix} \in \mathbb{S}_+^{N-1} \quad (\text{e})$$

(980)

where

$$\nu(i) \triangleq \frac{1}{2} \begin{bmatrix} d_{12} + d_{1i} - d_{2i} \\ d_{13} + d_{1i} - d_{3i} \\ \vdots \\ d_{1,i-1} + d_{1i} - d_{i-1,i} \end{bmatrix} \in \mathbb{R}^{i-2}, \quad i > 2 \quad (981)$$

Hence, the leading principal submatrices of EDM D must also be EDMs. ^{5.38}

Bordered symmetric matrices in the form (980d) are known to have *intertwined* [294, §6.4] (or *interlaced* [180, §4.3] [291, §IV.4.1]) eigenvalues; (*confer* §5.11.1) that means, for the particular submatrices (980a) and (980b),

^{5.38}In fact, each and every principal submatrix of an EDM D is another EDM. [207, §4.1]

$$\lambda_2 \leq d_{12} \leq \lambda_1 \quad (982)$$

where d_{12} is the eigenvalue of submatrix (980a) and λ_1, λ_2 are the eigenvalues of T (980b) (973). Intertwining in (982) predicts that should d_{12} become 0, then λ_2 must go to 0.^{5.39} The eigenvalues are similarly intertwined for submatrices (980b) and (980c);

$$\gamma_3 \leq \lambda_2 \leq \gamma_2 \leq \lambda_1 \leq \gamma_1 \quad (983)$$

where $\gamma_1, \gamma_2, \gamma_3$ are the eigenvalues of submatrix (980c). Intertwining likewise predicts that should λ_2 become 0 (a possibility revealed in §5.8.3.1), then γ_3 must go to 0. Combining results so far for $N=2, 3, 4$: (982) (983)

$$\gamma_3 \leq \lambda_2 \leq d_{12} \leq \lambda_1 \leq \gamma_1 \quad (984)$$

The preceding logic extends by induction through the remaining members of the sequence (980).

5.8.3.1 Tightening the triangle inequality

Now we apply Schur complement from §A.4 to tighten the triangle inequality from (972) in case: cardinality $N=4$. We find that the gains by doing so are modest. From (980) we identify:

$$\begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \triangleq -V_N^T D V_N |_{N \leftarrow 4} \quad (985)$$

$$A \triangleq T = -V_N^T D V_N |_{N \leftarrow 3} \quad (986)$$

both positive semidefinite by assumption, where $B = \nu(4)$ (981), and $C = d_{14}$. Using nonstrict CC^\dagger -form (1420), $C \succeq 0$ by assumption (§5.8.1) and $CC^\dagger = I$. So by the *positive semidefinite ordering of eigenvalues theorem* (§A.3.1.0.1),

$$-V_N^T D V_N |_{N \leftarrow 4} \succeq 0 \Leftrightarrow T \succeq d_{14}^{-1} \nu(4) \nu(4)^T \Rightarrow \begin{cases} \lambda_1 \geq d_{14}^{-1} \|\nu(4)\|^2 \\ \lambda_2 \geq 0 \end{cases} \quad (987)$$

where $\{d_{14}^{-1} \|\nu(4)\|^2, 0\}$ are the eigenvalues of $d_{14}^{-1} \nu(4) \nu(4)^T$ while λ_1, λ_2 are the eigenvalues of T .

^{5.39}If d_{12} were 0, eigenvalue λ_2 becomes 0 (975) because d_{13} must then be equal to d_{23} ; *id est*, $d_{12} = 0 \Leftrightarrow x_1 = x_2$. (§5.4)

5.8.3.1.1 Example. *Small completion problem, II.*

Applying the inequality for λ_1 in (987) to the *small completion problem* on page 352 Figure 96, the lower bound on $\sqrt{d_{14}}$ (1.236 in (801)) is tightened to 1.289. The correct value of $\sqrt{d_{14}}$ to three significant figures is 1.414.

□

5.8.4 Affine dimension reduction in two dimensions

(confer §5.14.4) The leading principal 2×2 submatrix T of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ has largest eigenvalue λ_1 (975) which is a convex function of D .^{5.40} λ_1 can never be 0 unless $d_{12} = d_{13} = d_{23} = 0$. Eigenvalue λ_1 can never be negative while the d_{ij} are nonnegative. The remaining eigenvalue λ_2 is a concave function of D that becomes 0 only at the upper and lower bounds of inequality (976a) and its equivalent forms: (confer (978))

$$|\sqrt{d_{12}} - \sqrt{d_{23}}| \leq \sqrt{d_{13}} \leq \sqrt{d_{12}} + \sqrt{d_{23}} \quad (\text{a})$$

$$|\sqrt{d_{12}} - \sqrt{d_{13}}| \leq \sqrt{d_{23}} \leq \sqrt{d_{12}} + \sqrt{d_{13}} \quad (\text{b}) \quad (988)$$

$$|\sqrt{d_{13}} - \sqrt{d_{23}}| \leq \sqrt{d_{12}} \leq \sqrt{d_{13}} + \sqrt{d_{23}} \quad (\text{c})$$

In between those bounds, λ_2 is strictly positive; otherwise, it would be negative but prevented by the condition $T \succeq 0$.

When λ_2 becomes 0, it means triangle Δ_{123} has collapsed to a line segment; a potential reduction in affine dimension r . The same logic is valid for any particular principal 2×2 submatrix of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$, hence applicable to other triangles.

^{5.40}The largest eigenvalue of any symmetric matrix is always a convex function of its entries, while the smallest eigenvalue is always concave. [53, exmp.3.10] In our particular case, say $\underline{d} \triangleq \begin{bmatrix} d_{12} \\ d_{13} \\ d_{23} \end{bmatrix} \in \mathbb{R}^3$. Then the Hessian (1649) $\nabla^2 \lambda_1(\underline{d}) \succeq 0$ certifies convexity whereas $\nabla^2 \lambda_2(\underline{d}) \preceq 0$ certifies concavity. Each Hessian has rank equal to 1. The respective gradients $\nabla \lambda_1(\underline{d})$ and $\nabla \lambda_2(\underline{d})$ are nowhere $\mathbf{0}$.

5.9 Bridge: Convex polyhedra to EDMs

The criteria for the existence of an EDM include, by definition (808) (873), the properties imposed upon its entries d_{ij} by the Euclidean metric. From §5.8.1 and §5.8.2, we know there is a relationship of matrix criteria to those properties. Here is a snapshot of what we are sure: for $i, j, k \in \{1 \dots N\}$ (confer §5.2)

$$\left. \begin{aligned} \sqrt{d_{ij}} &\geq 0, \quad i \neq j \\ \sqrt{d_{ij}} &= 0, \quad i = j \\ \sqrt{d_{ij}} &= \sqrt{d_{ji}} \\ \sqrt{d_{ij}} &\leq \sqrt{d_{ik}} + \sqrt{d_{kj}}, \quad i \neq j \neq k \end{aligned} \right\} \Leftrightarrow \begin{aligned} -V_{\mathcal{N}}^T D V_{\mathcal{N}} &\succeq 0 \\ \delta(D) &= \mathbf{0} \\ D^T &= D \end{aligned} \quad (989)$$

all implied by $D \in \text{EDM}^N$. In words, these four Euclidean metric properties are necessary conditions for D to be a distance matrix. At the moment, we have no converse. As of concern in §5.3, we have yet to establish metric requirements beyond the four Euclidean metric properties that would allow D to be certified an EDM or might facilitate polyhedron or list reconstruction from an incomplete EDM. We deal with this problem in §5.14. Our present goal is to establish *ab initio* the necessary and sufficient matrix criteria that will subsume all the Euclidean metric properties and any further requirements^{5.41} for all $N > 1$ (§5.8.3); *id est*,

$$\left. \begin{aligned} -V_{\mathcal{N}}^T D V_{\mathcal{N}} &\succeq 0 \\ D &\in \mathbb{S}_h^N \end{aligned} \right\} \Leftrightarrow D \in \text{EDM}^N \quad (827)$$

or for EDM definition (882),

$$\left. \begin{aligned} \Omega &\succeq 0 \\ \sqrt{\delta(d)} &\succeq 0 \end{aligned} \right\} \Leftrightarrow D = \mathbf{D}(\Omega, d) \in \text{EDM}^N \quad (990)$$

^{5.41}In 1935, Schoenberg [276, (1)] first extolled matrix product $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ (968) (predicated on symmetry and self-distance) specifically incorporating $V_{\mathcal{N}}$, albeit algebraically. He showed: nonnegativity $-y^T V_{\mathcal{N}}^T D V_{\mathcal{N}} y \geq 0$, for all $y \in \mathbb{R}^{N-1}$, is necessary and sufficient for D to be an EDM. Gower [141, §3] remarks how surprising it is that such a fundamental property of Euclidean geometry was obtained so late.




Figure 109: Elliptope \mathcal{E}^3 in isometrically isomorphic \mathbb{R}^6 (projected on \mathbb{R}^3) is a convex body that appears to possess some kind of symmetry in this dimension; it resembles a malformed pillow in the shape of a bulging tetrahedron. Elliptope relative boundary is not *smooth* and comprises all set members (991) having at least one 0 eigenvalue. [210, §2.1] This elliptope has an infinity of vertices, but there are only four vertices corresponding to a rank-1 matrix. Those yy^T , evident in the illustration, have binary vector $y \in \mathbb{R}^3$ with entries in $\{\pm 1\}$.




Figure 110: Elliptope \mathcal{E}^2 in isometrically isomorphic \mathbb{R}^3 is a line segment illustrated interior to positive semidefinite cone \mathbb{S}_+^2 (Figure 39).

5.9.1 Geometric arguments

5.9.1.0.1 Definition. *Elliptope:* [210] [207, §2.3] [96, §31.5] a unique bounded immutable convex Euclidean body in \mathbb{S}^n ; intersection of positive semidefinite cone \mathbb{S}_+^n with that set of n hyperplanes defined by unity main diagonal;

$$\mathcal{E}^n \triangleq \mathbb{S}_+^n \cap \{\Phi \in \mathbb{S}^n \mid \delta(\Phi) = \mathbf{1}\} \quad (991)$$

a. k. a, the set of all *correlation matrices* of dimension

$$\dim \mathcal{E}^n = n(n-1)/2 \text{ in } \mathbb{R}^{n(n+1)/2} \quad (992)$$

An elliptope \mathcal{E}^n is not a polyhedron, in general, but has some polyhedral faces and an infinity of vertices.^{5.42} Of those, 2^{n-1} vertices (some extreme points of the elliptope) are extreme directions yy^T of the positive semidefinite cone, where the entries of vector $y \in \mathbb{R}^n$ belong to $\{\pm 1\}$ and exercise every combination. Each of the remaining vertices has rank belonging to the set $\{k > 0 \mid k(k+1)/2 \leq n\}$. Each and every face of an elliptope is exposed.

△

^{5.42}Laurent defines vertex distinctly from the sense herein (§2.6.1.0.1); she defines *vertex* as a point with full-dimensional (nonempty interior) normal cone (§E.10.3.2.1). Her definition excludes point C in Figure 29, for example.

In fact, any positive semidefinite matrix whose entries belong to $\{\pm 1\}$ is a correlation matrix whose rank must be one: (As there are few equivalent conditions for rank constraints, this statement is rather important as a device for relaxing an integer, combinatorial, or Boolean problem.)

5.9.1.0.2 Theorem. *Some elliptope vertices.* [108, §2.1.1]

For $Y \in \mathbb{S}^n$, $y \in \mathbb{R}^n$, and all $i, j \in \{1 \dots n\}$ (confer §2.3.1.0.2)

$$Y \succeq 0, \quad Y_{ij} \in \{\pm 1\} \quad \Leftrightarrow \quad Y = yy^T, \quad y_i \in \{\pm 1\} \quad (993)$$

◇

The elliptope for dimension $n = 2$ is a line segment in isometrically isomorphic $\mathbb{R}^{n(n+1)/2}$ (Figure 110). Obviously, $\text{cone}(\mathcal{E}^n) \neq \mathbb{S}_+^n$. The elliptope for dimension $n = 3$ is realized in Figure 109.

5.9.1.0.3 Lemma. *Hypersphere.* [16, §4] (confer bullet p.362)

Matrix $A = [A_{ij}] \in \mathbb{S}^N$ belongs to the elliptope in \mathbb{S}^N iff there exist N points p on the boundary of a hypersphere in $\mathbb{R}^{\text{rank } A}$ having radius 1 such that

$$\|p_i - p_j\| = \sqrt{2}\sqrt{1 - A_{ij}}, \quad i, j = 1 \dots N \quad (994)$$

◇

There is a similar theorem for Euclidean distance matrices:

We derive matrix criteria for D to be an EDM, validating (827) using simple geometry; distance to the polyhedron formed by the convex hull of a list of points (69) in Euclidean space \mathbb{R}^n .

5.9.1.0.4 EDM assertion.

D is a Euclidean distance matrix if and only if $D \in \mathbb{S}_h^N$ and distances-square from the origin

$$\{\|p(y)\|^2 = -y^T V_N^T D V_N y \mid y \in \mathcal{S} - \beta\} \quad (995)$$

correspond to points p in some bounded convex polyhedron

$$\mathcal{P} - \alpha = \{p(y) \mid y \in \mathcal{S} - \beta\} \quad (996)$$

having N or fewer vertices embedded in an r -dimensional subspace $\mathcal{A} - \alpha$ of \mathbb{R}^n , where $\alpha \in \mathcal{A} = \text{aff } \mathcal{P}$ and where the domain of linear surjection $p(y)$ is the unit simplex $\mathcal{S} \subset \mathbb{R}_+^{N-1}$ shifted such that its vertex at the origin is translated to $-\beta$ in \mathbb{R}^{N-1} . When $\beta = 0$, then $\alpha = x_1$. ◇

In terms of $V_{\mathcal{N}}$, the unit simplex (266) in \mathbb{R}^{N-1} has an equivalent representation:

$$\mathcal{S} = \{s \in \mathbb{R}^{N-1} \mid \sqrt{2}V_{\mathcal{N}}s \succeq -e_1\} \quad (997)$$

where e_1 is as in (824). Incidental to the *EDM assertion*, shifting the unit-simplex domain in \mathbb{R}^{N-1} translates the polyhedron \mathcal{P} in \mathbb{R}^n . Indeed, there is a map from vertices of the unit simplex to members of the list generating \mathcal{P} ;

$$p : \mathbb{R}^{N-1} \rightarrow \mathbb{R}^n$$

$$p \left(\left\{ \begin{array}{c} -\beta \\ e_1 - \beta \\ e_2 - \beta \\ \vdots \\ e_{N-1} - \beta \end{array} \right\} \right) = \left\{ \begin{array}{c} x_1 - \alpha \\ x_2 - \alpha \\ x_3 - \alpha \\ \vdots \\ x_N - \alpha \end{array} \right\} \quad (998)$$

5.9.1.0.5 Proof. EDM assertion.

(\Rightarrow) We demonstrate that if D is an EDM, then each distance-square $\|p(y)\|^2$ described by (995) corresponds to a point p in some embedded polyhedron $\mathcal{P} - \alpha$. Assume D is indeed an EDM; *id est*, D can be made from some list X of N unknown points in Euclidean space \mathbb{R}^n ; $D = \mathbf{D}(X)$ for $X \in \mathbb{R}^{n \times N}$ as in (808). Since D is translation invariant (§5.5.1), we may shift the affine hull \mathcal{A} of those unknown points to the origin as in (938). Then take any point p in their convex hull (79);

$$\mathcal{P} - \alpha = \{p = (X - Xb\mathbf{1}^T)a \mid a^T\mathbf{1} = 1, a \succeq 0\} \quad (999)$$

where $\alpha = Xb \in \mathcal{A} \Leftrightarrow b^T\mathbf{1} = 1$. Solutions to $a^T\mathbf{1} = 1$ are:^{5.43}

$$a \in \{e_1 + \sqrt{2}V_{\mathcal{N}}s \mid s \in \mathbb{R}^{N-1}\} \quad (1000)$$

where e_1 is as in (824). Similarly, $b = e_1 + \sqrt{2}V_{\mathcal{N}}\beta$.

$$\begin{aligned} \mathcal{P} - \alpha &= \{p = X(I - (e_1 + \sqrt{2}V_{\mathcal{N}}\beta)\mathbf{1}^T)(e_1 + \sqrt{2}V_{\mathcal{N}}s) \mid \sqrt{2}V_{\mathcal{N}}s \succeq -e_1\} \\ &= \{p = X\sqrt{2}V_{\mathcal{N}}(s - \beta) \mid \sqrt{2}V_{\mathcal{N}}s \succeq -e_1\} \end{aligned} \quad (1001)$$

^{5.43}Since $\mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^T)$ and $\mathcal{N}(\mathbf{1}^T) \perp \mathcal{R}(\mathbf{1})$, then over all $s \in \mathbb{R}^{N-1}$, $V_{\mathcal{N}}s$ is a hyperplane through the origin orthogonal to $\mathbf{1}$. Thus the solutions $\{a\}$ constitute a hyperplane orthogonal to the vector $\mathbf{1}$, and offset from the origin in \mathbb{R}^N by any particular solution; in this case, $a = e_1$.

that describes the domain of $p(s)$ as the unit simplex

$$\mathcal{S} = \{s \mid \sqrt{2}V_{\mathcal{N}}s \succeq -e_1\} \subset \mathbb{R}_+^{N-1} \quad (997)$$

Making the substitution $s - \beta \leftarrow y$

$$\mathcal{P} - \alpha = \{p = X\sqrt{2}V_{\mathcal{N}}y \mid y \in \mathcal{S} - \beta\} \quad (1002)$$

Point p belongs to a convex polyhedron $\mathcal{P} - \alpha$ embedded in an r -dimensional subspace of \mathbb{R}^n because the convex hull of any list forms a polyhedron, and because the translated affine hull $\mathcal{A} - \alpha$ contains the translated polyhedron $\mathcal{P} - \alpha$ (941) and the origin (when $\alpha \in \mathcal{A}$), and because \mathcal{A} has dimension r by definition (943). Now, any distance-square from the origin to the polyhedron $\mathcal{P} - \alpha$ can be formulated

$$\{p^{\text{T}}p = \|p\|^2 = 2y^{\text{T}}V_{\mathcal{N}}^{\text{T}}X^{\text{T}}XV_{\mathcal{N}}y \mid y \in \mathcal{S} - \beta\} \quad (1003)$$

Applying (950) to (1003) we get (995).

(\Leftarrow) To validate the *EDM assertion* in the reverse direction, we prove: If each distance-square $\|p(y)\|^2$ (995) on the shifted unit-simplex $\mathcal{S} - \beta \subset \mathbb{R}^{N-1}$ corresponds to a point $p(y)$ in some embedded polyhedron $\mathcal{P} - \alpha$, then D is an EDM. The r -dimensional subspace $\mathcal{A} - \alpha \subseteq \mathbb{R}^n$ is spanned by

$$p(\mathcal{S} - \beta) = \mathcal{P} - \alpha \quad (1004)$$

because $\mathcal{A} - \alpha = \text{aff}(\mathcal{P} - \alpha) \supseteq \mathcal{P} - \alpha$ (941). So, outside the domain $\mathcal{S} - \beta$ of linear surjection $p(y)$, the simplex complement $\setminus \mathcal{S} - \beta \subset \mathbb{R}^{N-1}$ must contain the domain of the distance-square $\|p(y)\|^2 = p(y)^{\text{T}}p(y)$ to remaining points in the subspace $\mathcal{A} - \alpha$; *id est*, to the polyhedron's relative exterior $\setminus \mathcal{P} - \alpha$. For $\|p(y)\|^2$ to be nonnegative on the entire subspace $\mathcal{A} - \alpha$, $-V_{\mathcal{N}}^{\text{T}}DV_{\mathcal{N}}$ must be positive semidefinite and is assumed symmetric;^{5.44}

$$-V_{\mathcal{N}}^{\text{T}}DV_{\mathcal{N}} \stackrel{\Delta}{=} \Theta_{\text{p}}^{\text{T}}\Theta_{\text{p}} \quad (1005)$$

where^{5.45} $\Theta_{\text{p}} \in \mathbb{R}^{m \times N-1}$ for some $m \geq r$. Because $p(\mathcal{S} - \beta)$ is a convex polyhedron, it is necessarily a set of linear combinations of points from some

^{5.44}The antisymmetric part $(-V_{\mathcal{N}}^{\text{T}}DV_{\mathcal{N}} - (-V_{\mathcal{N}}^{\text{T}}DV_{\mathcal{N}})^{\text{T}})/2$ is annihilated by $\|p(y)\|^2$. By the same reasoning, any positive (semi)definite matrix A is generally assumed symmetric because only the symmetric part $(A + A^{\text{T}})/2$ survives the test $y^{\text{T}}Ay \geq 0$. [180, §7.1]

^{5.45} $A^{\text{T}} = A \succeq 0 \Leftrightarrow A = R^{\text{T}}R$ for some real matrix R . [294, §6.3]

length- N list because every convex polyhedron having N or fewer vertices can be generated that way (§2.12.2). Equivalent to (995) are

$$\{p^T p \mid p \in \mathcal{P} - \alpha\} = \{p^T p = y^T \Theta_p^T \Theta_p y \mid y \in \mathcal{S} - \beta\} \quad (1006)$$

Because $p \in \mathcal{P} - \alpha$ may be found by factoring (1006), the list Θ_p is found by factoring (1005). A unique EDM can be made from that list using inner-product form definition $\mathbf{D}(\Theta)|_{\Theta=\Theta_p}$ (873). That EDM will be identical to D if $\delta(D)=\mathbf{0}$, by injectivity of \mathbf{D} (924). \blacklozenge

5.9.2 Necessity and sufficiency

From (966) we learned that matrix inequality $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$ is a necessary test for D to be an EDM. In §5.9.1, the connection between convex polyhedra and EDMs was pronounced by the *EDM assertion*; the matrix inequality together with $D \in \mathbb{S}_h^N$ became a sufficient test when the *EDM assertion* demanded that every bounded convex polyhedron have a corresponding EDM. For all $N > 1$ (§5.8.3), the matrix criteria for the existence of an EDM in (827), (990), and (803) are therefore necessary and sufficient and subsume all the Euclidean metric properties and further requirements.

5.9.3 Example revisited

Now we apply the necessary and sufficient EDM criteria (827) to an earlier problem.

5.9.3.0.1 Example. *Small completion problem, III.* (confer §5.8.3.1.1) Continuing Example 5.3.0.0.2 pertaining to Figure 96 where $N=4$, distance-square d_{14} is ascertainable from the matrix inequality $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$. Because all distances in (800) are known except $\sqrt{d_{14}}$, we may simply calculate the smallest eigenvalue of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ over a range of d_{14} as in Figure 111. We observe a unique value of d_{14} satisfying (827) where the abscissa is tangent to the hypograph of the smallest eigenvalue. Since the smallest eigenvalue of a symmetric matrix is known to be a concave function (§5.8.4), we calculate its second partial derivative with respect to d_{14} evaluated at 2 and find $-1/3$. We conclude there are no other satisfying values of d_{14} . Further, that value of d_{14} does not meet an upper or lower bound of a triangle inequality like (978), so neither does it cause the collapse

of any triangle. Because the smallest eigenvalue is 0, affine dimension r of any point list corresponding to D cannot exceed $N-2$. (§5.7.1.1) \square

5.10 EDM-entry composition

Laurent [207, §2.3] applies results from Schoenberg (1938) [277] to show certain nonlinear compositions of individual EDM entries yield EDMs; in particular,

$$\begin{aligned} D \in \text{EDM}^N &\Leftrightarrow [1 - e^{-\alpha d_{ij}}] \in \text{EDM}^N \quad \forall \alpha > 0 \\ &\Leftrightarrow [e^{-\alpha d_{ij}}] \in \mathcal{E}^N \quad \forall \alpha > 0 \end{aligned} \quad (1007)$$

where $D = [d_{ij}]$ and \mathcal{E}^N is the ellipsope (991).

5.10.0.0.1 Proof. (Laurent, 2003) [277] (confer [201])

Lemma 2.1. from *A Tour d'Horizon ... on Completion Problems*. [207] The following assertions are equivalent: for $D = [d_{ij}, i, j = 1 \dots N] \in \mathbb{S}_h^N$ and \mathcal{E}^N the ellipsope in \mathbb{S}^N (§5.9.1.0.1),

- (i) $D \in \text{EDM}^N$
- (ii) $e^{-\alpha D} \triangleq [e^{-\alpha d_{ij}}] \in \mathcal{E}^N$ for all $\alpha > 0$
- (iii) $\mathbf{1}\mathbf{1}^T - e^{-\alpha D} \triangleq [1 - e^{-\alpha d_{ij}}] \in \text{EDM}^N$ for all $\alpha > 0$ \diamond

- 1) Equivalence of Lemma 2.1 (i) (ii) is stated in Schoenberg's Theorem 1 [277, p.527].
- 2) (ii) \Rightarrow (iii) can be seen from the statement in the beginning of section 3, saying that a distance space embeds in L_2 iff some associated matrix is PSD. We reformulate it:

Let $d = (d_{ij})_{i,j=0,1\dots N}$ be a distance space on $N+1$ points (*i.e.*, symmetric hollow matrix of order $N+1$) and let $p = (p_{ij})_{i,j=1\dots N}$ be the symmetric matrix of order N related by:

$$(A) \quad 2p_{ij} = d_{0i} + d_{0j} - d_{ij} \quad \text{for } i, j = 1 \dots N$$

or equivalently

$$(B) \quad d_{0i} = p_{ii}, \quad d_{ij} = p_{ii} + p_{jj} - 2p_{ij} \quad \text{for } i, j = 1 \dots N$$




Figure 111: Smallest eigenvalue of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ makes it a PSD matrix for only one value of d_{14} : 2.




Figure 112: Some entrywise EDM compositions: **(a)** $\alpha = 2$. Concave nondecreasing in d_{ij} . **(b)** Trajectory convergence in α for $d_{ij} = 2$.

Then d embeds in L_2 iff p is a positive semidefinite matrix iff d is of negative type (second half page 525/top of page 526 in [277]).

For the implication from (ii) to (iii), set: $p = e^{-\alpha d}$ and define d' from p using (B) above. Then d' is a distance space on $N+1$ points that embeds in L_2 . Thus its subspace of N points also embeds in L_2 and is precisely $1 - e^{-\alpha d}$.

Note that (iii) \Rightarrow (ii) cannot be read immediately from this argument since (iii) involves the subdistance of d' on N points (and not the full d' on $N+1$ points).

- 3) Show (iii) \Rightarrow (i) by using the series expansion of the function $1 - e^{-\alpha d}$: the constant term cancels, α factors out; there remains a summation of d plus a multiple of α . Letting α go to 0 gives the result.

This is not explicitly written in Schoenberg, but he also uses such an argument; expansion of the exponential function then $\alpha \rightarrow 0$ (first proof on [277, p.526]). \blacklozenge

Schoenberg's results [277, §6, thm.5] (confer [201, p.108-109]) also suggest certain finite positive roots of EDM entries produce EDMs; specifically,

$$D \in \text{EDM}^N \Leftrightarrow [d_{ij}^{1/\alpha}] \in \text{EDM}^N \quad \forall \alpha > 1 \quad (1008)$$

The special case $\alpha = 2$ is of interest because it corresponds to absolute distance; *e.g.*,

$$D \in \text{EDM}^N \Rightarrow \sqrt[D]{} \in \text{EDM}^N \quad (1009)$$

Assuming that points constituting a corresponding list X are distinct (970), then it follows: for $D \in \mathbb{S}_h^N$

$$\lim_{\alpha \rightarrow \infty} [d_{ij}^{1/\alpha}] = \lim_{\alpha \rightarrow \infty} [1 - e^{-\alpha d_{ij}}] = -E \triangleq \mathbf{1}\mathbf{1}^T - I \quad (1010)$$

Negative elementary matrix $-E$ (§B.3) is relatively interior to the EDM cone (§6.6) and terminal to the respective trajectories (1007) and (1008) as functions of α . Both trajectories are confined to the EDM cone; in engineering terms, the EDM cone is an *invariant set* [274] to either trajectory. Further, if D is not an EDM but for some particular α_p it becomes an EDM, then for all greater values of α it remains an EDM.

These preliminary findings lead one to speculate whether any concave nondecreasing composition of individual EDM entries d_{ij} on \mathbb{R}_+ will produce another EDM; *e.g.*, given EDM D , empirical evidence suggests that for each fixed $\alpha \geq 1$ [*sic*] the composition $[\log_2(1 + d_{ij}^{1/\alpha})]$ is also an EDM. Figure 112 illustrates that composition's concavity in d_{ij} together with functions from (1007) and (1008).

5.10.0.0.2 Exercise. *Taxicab distance matrix as EDM.*

Determine whether taxicab distance matrices $(\mathbf{D}_1(X))$ in Example 3.2.0.0.2 are all numerically equivalent to EDMs. Explain why or why not. ▼

5.10.1 EDM by elliptope

For some $\kappa \in \mathbb{R}_+$ and $C \in \mathbb{S}_+^N$ in the elliptope \mathcal{E}^N (§5.9.1.0.1), Alfakih asserts any given EDM D is expressible [8] [96, §31.5]

$$D = \kappa(\mathbf{1}\mathbf{1}^T - C) \in \mathbb{EDM}^N \quad (1011)$$

This expression exhibits nonlinear combination of variables κ and C . We therefore propose a different expression requiring redefinition of the elliptope (991) by scalar parametrization;

$$\mathcal{E}_t^n \triangleq \mathbb{S}_+^n \cap \{\Phi \in \mathbb{S}^n \mid \delta(\Phi) = t\mathbf{1}\} \quad (1012)$$

where, of course, $\mathcal{E}^n = \mathcal{E}_1^n$. Then any given EDM D is expressible

$$D = t\mathbf{1}\mathbf{1}^T - \mathfrak{E} \in \mathbb{EDM}^N \quad (1013)$$

which is linear in variables $t \in \mathbb{R}_+$ and $\mathfrak{E} \in \mathcal{E}_t^N$.

5.11 EDM indefiniteness

By the known result in §A.7.2 regarding a 0-valued entry on the main diagonal of a symmetric positive semidefinite matrix, there can be no positive nor negative semidefinite EDM except the $\mathbf{0}$ matrix because $\mathbb{EDM}^N \subseteq \mathbb{S}_h^N$ (807) and

$$\mathbb{S}_h^N \cap \mathbb{S}_+^N = \mathbf{0} \quad (1014)$$

the origin. So when $D \in \mathbb{EDM}^N$, there can be no factorization $D = A^T A$ nor $-D = A^T A$. [294, §6.3] Hence eigenvalues of an EDM are neither all nonnegative or all nonpositive; an EDM is indefinite and possibly invertible.

5.11.1 EDM eigenvalues, congruence transformation

For any symmetric $-D$, we can characterize its eigenvalues by congruence transformation: [294, §6.3]

$$-W^T D W = - \begin{bmatrix} V_N^T \\ \mathbf{1}^T \end{bmatrix} D \begin{bmatrix} V_N & \mathbf{1} \end{bmatrix} = - \begin{bmatrix} V_N^T D V_N & V_N^T D \mathbf{1} \\ \mathbf{1}^T D V_N & \mathbf{1}^T D \mathbf{1} \end{bmatrix} \in \mathbb{S}^N \quad (1015)$$

Because

$$W \triangleq \begin{bmatrix} V_N & \mathbf{1} \end{bmatrix} \in \mathbb{R}^{N \times N} \quad (1016)$$

is full-rank, then (1425)

$$\text{inertia}(-D) = \text{inertia}(-W^T D W) \quad (1017)$$

the congruence (1015) has the same number of positive, zero, and negative eigenvalues as $-D$. Further, if we denote by $\{\gamma_i, i=1 \dots N-1\}$ the eigenvalues of $-V_N^T D V_N$ and denote eigenvalues of the congruence $-W^T D W$ by $\{\zeta_i, i=1 \dots N\}$ and if we arrange each respective set of eigenvalues in nonincreasing order, then by theory of *interlacing eigenvalues for bordered symmetric matrices* [180, §4.3] [294, §6.4] [291, §IV.4.1]

$$\zeta_N \leq \gamma_{N-1} \leq \zeta_{N-1} \leq \gamma_{N-2} \leq \dots \leq \gamma_2 \leq \zeta_2 \leq \gamma_1 \leq \zeta_1 \quad (1018)$$

When $D \in \text{EDM}^N$, then $\gamma_i \geq 0 \forall i$ (1363) because $-V_N^T D V_N \succeq 0$ as we know. That means the congruence must have $N-1$ nonnegative eigenvalues; $\zeta_i \geq 0, i=1 \dots N-1$. The remaining eigenvalue ζ_N cannot be nonnegative because then $-D$ would be positive semidefinite, an impossibility; so $\zeta_N < 0$. By congruence, nontrivial $-D$ must therefore have exactly one negative eigenvalue;^{5.46} [96, §2.4.5]

^{5.46}All the entries of the corresponding eigenvector must have the same sign with respect to each other [78, p.116] because that eigenvector is the *Perron vector* corresponding to the spectral radius; [180, §8.2.6] the predominant characteristic of negative [*sic*] matrices.

$$D \in \text{EDM}^N \Rightarrow \begin{cases} \lambda(-D)_i \geq 0, & i=1 \dots N-1 \\ \left(\sum_{i=1}^N \lambda(-D)_i = 0 \right) \\ D \in \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N} \end{cases} \quad (1019)$$

where the $\lambda(-D)_i$ are nonincreasingly ordered eigenvalues of $-D$ whose sum must be 0 only because $\text{tr } D = 0$ [294, §5.1]. The eigenvalue summation condition, therefore, can be considered redundant. Even so, all these conditions are insufficient to determine whether some given $H \in \mathbb{S}_h^N$ is an EDM, as shown by counter-example.^{5.47}

5.11.1.0.1 Exercise. Spectral inequality.

Prove whether it holds: for $D = [d_{ij}] \in \text{EDM}^N$

$$\lambda(-D)_1 \geq d_{ij} \geq \lambda(-D)_{N-1} \quad \forall i \neq j \quad (1020)$$

▼

Terminology: a *spectral cone* is a convex cone containing all *eigenspectra* [201, p.365] [291, p.26] corresponding to some set of matrices. Any positive semidefinite matrix, for example, possesses a vector of nonnegative eigenvalues corresponding to an eigenspectrum contained in a spectral cone that is a nonnegative orthant.

5.11.2 Spectral cones

Denoting the eigenvalues of Cayley-Menger matrix $\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \in \mathbb{S}^{N+1}$ by

$$\lambda\left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix}\right) \in \mathbb{R}^{N+1} \quad (1021)$$

^{5.47}When $N=3$, for example, the symmetric hollow matrix

$$H = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 5 \\ 1 & 5 & 0 \end{bmatrix} \in \mathbb{S}_h^3 \cap \mathbb{R}_+^{3 \times 3}$$

is not an EDM, although $\lambda(-H) = [5 \ 0.3723 \ -5.3723]^T$ conforms to (1019).

we have the Cayley-Menger form (§5.7.3.0.1) of necessary and sufficient conditions for $D \in \text{EDM}^N$ from the literature: [163, §3]^{5.48} [65, §3] [96, §6.2] (confer (827) (803))

$$D \in \text{EDM}^N \Leftrightarrow \left\{ \begin{array}{l} \lambda \left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \right)_i \geq 0, \quad i = 1 \dots N \\ D \in \mathbb{S}_h^N \end{array} \right\} \Leftrightarrow \left\{ \begin{array}{l} -V_N^T D V_N \succeq 0 \\ D \in \mathbb{S}_h^N \end{array} \right\} \quad (1022)$$

These conditions say the Cayley-Menger form has one and only one negative eigenvalue. When D is an EDM, eigenvalues $\lambda \left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \right)$ belong to that particular orthant in \mathbb{R}^{N+1} having the $N+1^{\text{th}}$ coordinate as sole negative coordinate^{5.49}:

$$\begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} = \text{cone} \{e_1, e_2, \dots, e_N, -e_{N+1}\} \quad (1023)$$

5.11.2.1 Cayley-Menger *versus* Schoenberg

Connection to the Schoenberg criterion (827) is made when the Cayley-Menger form is further partitioned:

$$\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} = \begin{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} & \begin{bmatrix} \mathbf{1}^T \\ -D_{1,2:N} \end{bmatrix} \\ \begin{bmatrix} \mathbf{1} & -D_{2:N,1} \end{bmatrix} & -D_{2:N,2:N} \end{bmatrix} \quad (1024)$$

Matrix $D \in \mathbb{S}_h^N$ is an EDM if and only if the Schur complement of $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ (§A.4) in this partition is positive semidefinite; [16, §1] [192, §3] *id est*,

^{5.48}Recall: for $D \in \mathbb{S}_h^N$, $-V_N^T D V_N \succeq 0$ subsumes nonnegativity property 1 (§5.8.1).

^{5.49}Empirically, all except one entry of the corresponding eigenvector have the same sign with respect to each other.

(confer (968))

$$\begin{aligned}
& D \in \text{EDM}^N \\
& \Leftrightarrow \\
& -D_{2:N,2:N} - [\mathbf{1} \quad -D_{2:N,1}] \begin{bmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{bmatrix} \begin{bmatrix} \mathbf{1}^T \\ -D_{1,2:N} \end{bmatrix} = -2V_N^T D V_N \succeq 0 \quad (1025) \\
& \text{and} \\
& D \in \mathbb{S}_h^N
\end{aligned}$$

Positive semidefiniteness of that Schur complement insures nonnegativity ($D \in \mathbb{R}_+^{N \times N}$, §5.8.1), whereas *complementary inertia* (1427) insures existence of that lone negative eigenvalue of the Cayley-Menger form.

Now we apply results from chapter 2 with regard to polyhedral cones and their duals.

5.11.2.2 Ordered eigenspectra

Conditions (1022) specify eigenvalue membership to the smallest pointed polyhedral *spectral cone* for $\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -\text{EDM}^N \end{bmatrix}$:

$$\begin{aligned}
\mathcal{K}_\lambda & \triangleq \{\zeta \in \mathbb{R}^{N+1} \mid \zeta_1 \geq \zeta_2 \geq \cdots \geq \zeta_N \geq 0 \geq \zeta_{N+1}, \mathbf{1}^T \zeta = 0\} \\
& = \mathcal{K}_{\mathcal{M}} \cap \begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} \cap \partial \mathcal{H} \\
& = \lambda \left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -\text{EDM}^N \end{bmatrix} \right)
\end{aligned} \quad (1026)$$

where

$$\partial \mathcal{H} \triangleq \{\zeta \in \mathbb{R}^{N+1} \mid \mathbf{1}^T \zeta = 0\} \quad (1027)$$

is a hyperplane through the origin, and $\mathcal{K}_{\mathcal{M}}$ is the monotone cone (§2.13.9.4.3, implying ordered eigenspectra) which has nonempty interior but is not pointed;

$$\mathcal{K}_{\mathcal{M}} = \{\zeta \in \mathbb{R}^{N+1} \mid \zeta_1 \geq \zeta_2 \geq \cdots \geq \zeta_{N+1}\} \quad (396)$$

$$\mathcal{K}_{\mathcal{M}}^* = \{[e_1 - e_2 \quad e_2 - e_3 \quad \cdots \quad e_N - e_{N+1}] a \mid a \succeq 0\} \subset \mathbb{R}^{N+1} \quad (397)$$

So because of the hyperplane,

$$\dim \text{aff } \mathcal{K}_\lambda = \dim \partial \mathcal{H} = N \quad (1028)$$

indicating \mathcal{K}_λ has empty interior. Defining

$$A \triangleq \begin{bmatrix} e_1^T - e_2^T \\ e_2^T - e_3^T \\ \vdots \\ e_N^T - e_{N+1}^T \end{bmatrix} \in \mathbb{R}^{N \times N+1}, \quad B \triangleq \begin{bmatrix} e_1^T \\ e_2^T \\ \vdots \\ e_N^T \\ -e_{N+1}^T \end{bmatrix} \in \mathbb{R}^{N+1 \times N+1} \quad (1029)$$

we have the halfspace-description:

$$\mathcal{K}_\lambda = \{\zeta \in \mathbb{R}^{N+1} \mid A\zeta \succeq 0, B\zeta \succeq 0, \mathbf{1}^T \zeta = 0\} \quad (1030)$$

From this and (404) we get a vertex-description for a pointed spectral cone having empty interior:

$$\mathcal{K}_\lambda = \left\{ V_{\mathcal{N}} \left(\begin{bmatrix} \hat{A} \\ \hat{B} \end{bmatrix} V_{\mathcal{N}} \right)^\dagger b \mid b \succeq 0 \right\} \quad (1031)$$

where $V_{\mathcal{N}} \in \mathbb{R}^{N+1 \times N}$, and where [sic]

$$\hat{B} = e_N^T \in \mathbb{R}^{1 \times N+1} \quad (1032)$$

and

$$\hat{A} = \begin{bmatrix} e_1^T - e_2^T \\ e_2^T - e_3^T \\ \vdots \\ e_{N-1}^T - e_N^T \end{bmatrix} \in \mathbb{R}^{N-1 \times N+1} \quad (1033)$$

hold those rows of A and B corresponding to conically independent rows (§2.10) in $\begin{bmatrix} A \\ B \end{bmatrix} V_{\mathcal{N}}$.

Conditions (1022) can be equivalently restated in terms of a spectral cone for Euclidean distance matrices:

$$D \in \text{EDM}^N \Leftrightarrow \begin{cases} \lambda \left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \right) \in \mathcal{K}_{\mathcal{M}} \cap \begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} \cap \partial \mathcal{H} \\ D \in \mathbb{S}_h^N \end{cases} \quad (1034)$$

Vertex-description of the dual spectral cone is, (285)

$$\begin{aligned} \mathcal{K}_\lambda^* &= \overline{\mathcal{K}_\mathcal{M}^* + \begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} + \partial\mathcal{H}^*} \subseteq \mathbb{R}^{N+1} \\ &= \{ [A^\top \ B^\top \ \mathbf{1} \ -\mathbf{1}] b \mid b \succeq 0 \} = \left\{ \begin{bmatrix} \hat{A}^\top & \hat{B}^\top & \mathbf{1} & -\mathbf{1} \end{bmatrix} a \mid a \succeq 0 \right\} \end{aligned} \quad (1035)$$

From (1031) and (405) we get a halfspace-description:

$$\mathcal{K}_\lambda^* = \{ y \in \mathbb{R}^{N+1} \mid (V_N^\top [\hat{A}^\top \ \hat{B}^\top])^\dagger V_N^\top y \succeq 0 \} \quad (1036)$$

This polyhedral dual spectral cone \mathcal{K}_λ^* is closed, convex, has nonempty interior because \mathcal{K}_λ is pointed, but is not pointed because \mathcal{K}_λ has empty interior.

5.11.2.3 Unordered eigenspectra

Spectral cones are not unique; eigenspectra ordering can be rendered benign within a cone by presorting a vector of eigenvalues into nonincreasing order.^{5.50} Then things simplify: Conditions (1022) now specify eigenvalue membership to the spectral cone

$$\begin{aligned} \lambda \left(\begin{bmatrix} 0 & \mathbf{1}^\top \\ \mathbf{1} & -\text{EDM}^N \end{bmatrix} \right) &= \begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \\ &= \{ \zeta \in \mathbb{R}^{N+1} \mid B\zeta \succeq 0, \mathbf{1}^\top \zeta = 0 \} \end{aligned} \quad (1037)$$

where B is defined in (1029), and $\partial\mathcal{H}$ in (1027). From (404) we get a vertex-description for a pointed spectral cone having empty interior:

$$\begin{aligned} \lambda \left(\begin{bmatrix} 0 & \mathbf{1}^\top \\ \mathbf{1} & -\text{EDM}^N \end{bmatrix} \right) &= \{ V_N (\tilde{B} V_N)^\dagger b \mid b \succeq 0 \} \\ &= \left\{ \begin{bmatrix} I \\ -\mathbf{1}^\top \end{bmatrix} b \mid b \succeq 0 \right\} \end{aligned} \quad (1038)$$

^{5.50}Eigenspectra ordering (represented by a cone having monotone description such as (1026)) becomes benign in (1247), for example, where projection of a given presorted vector on the nonnegative orthant in a subspace is equivalent to its projection on the monotone nonnegative cone in that same subspace; equivalence is a consequence of presorting.

where $V_{\mathcal{N}} \in \mathbb{R}^{N+1 \times N}$ and

$$\tilde{B} \triangleq \begin{bmatrix} e_1^T \\ e_2^T \\ \vdots \\ e_N^T \end{bmatrix} \in \mathbb{R}^{N \times N+1} \quad (1039)$$

holds only those rows of B corresponding to conically independent rows in $BV_{\mathcal{N}}$.

For presorted eigenvalues, (1022) can be equivalently restated

$$D \in \text{EDM}^N \Leftrightarrow \begin{cases} \lambda\left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix}\right) \in \begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \\ D \in \mathbb{S}_h^N \end{cases} \quad (1040)$$

Vertex-description of the dual spectral cone is, (285)

$$\begin{aligned} \lambda\left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -\text{EDM}^N \end{bmatrix}\right)^* &= \begin{bmatrix} \mathbb{R}_+^N \\ \mathbb{R}_- \end{bmatrix} + \partial\mathcal{H}^* \subseteq \mathbb{R}^{N+1} \\ &= \{ [B^T \ \mathbf{1} \ -\mathbf{1}] b \mid b \succeq 0 \} = \left\{ \begin{bmatrix} \tilde{B}^T & \mathbf{1} & -\mathbf{1} \end{bmatrix} a \mid a \succeq 0 \right\} \end{aligned} \quad (1041)$$

From (405) we get a halfspace-description:

$$\begin{aligned} \lambda\left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -\text{EDM}^N \end{bmatrix}\right)^* &= \{y \in \mathbb{R}^{N+1} \mid (V_{\mathcal{N}}^T \tilde{B}^T)^\dagger V_{\mathcal{N}}^T y \succeq 0\} \\ &= \{y \in \mathbb{R}^{N+1} \mid [I \ -\mathbf{1}] y \succeq 0\} \end{aligned} \quad (1042)$$

This polyhedral dual spectral cone is closed, convex, has nonempty interior but is not pointed. (Notice that any nonincreasingly ordered eigenspectrum belongs to this dual spectral cone.)

5.11.2.4 Dual cone *versus* dual spectral cone

An open question regards the relationship of convex cones and their duals to the corresponding spectral cones and their duals. A positive semidefinite cone, for example, is self-dual. Both the nonnegative orthant and the monotone nonnegative cone are spectral cones for it. When we consider the nonnegative orthant, then that spectral cone for the self-dual positive semidefinite cone is also self-dual.

5.12 List reconstruction

The traditional term *metric multidimensional scaling*^{5.51} [227] [90] [313] [88] [230] [78] refers to any reconstruction of a list $X \in \mathbb{R}^{n \times N}$ in Euclidean space from interpoint distance information, possibly incomplete (§6.4), ordinal (§5.13.2), or specified perhaps only by bounding-constraints (§5.4.2.2.8) [311]. Techniques for reconstruction are essentially methods for optimally embedding an unknown list of points, corresponding to given Euclidean distance data, in an affine subset of desired or minimum dimension. The oldest known precursor is called *principal component analysis* [144] which analyzes the correlation matrix (§5.9.1.0.1); [46, §22] a. k. a, *Karhunen–Loève transform* in the digital signal processing literature.

Isometric reconstruction (§5.5.3) of point list X is best performed by eigen decomposition of a Gram matrix; for then, numerical errors of factorization are easily spotted in the eigenvalues: Now we consider how rotation/reflection and translation invariance factor into a reconstruction.

5.12.1 x_1 at the origin. V_N

At the stage of reconstruction, we have $D \in \mathbb{EDM}^N$ and we wish to find a generating list (§2.3.2) for $\mathcal{P} - \alpha$ by factoring positive semidefinite $-V_N^T D V_N$ (1005) as suggested in §5.9.1.0.5. One way to factor $-V_N^T D V_N$ is via diagonalization of symmetric matrices; [294, §5.6] [180] (§A.5.2, §A.3)

$$-V_N^T D V_N \triangleq Q \Lambda Q^T \quad (1043)$$

$$Q \Lambda Q^T \succeq 0 \Leftrightarrow \Lambda \succeq 0 \quad (1044)$$

where $Q \in \mathbb{R}^{N-1 \times N-1}$ is an orthogonal matrix containing eigenvectors while $\Lambda \in \mathbb{S}^{N-1}$ is a diagonal matrix containing corresponding nonnegative eigenvalues ordered by nonincreasing value. From the diagonalization, identify the list using (950);

$$-V_N^T D V_N = 2V_N^T X^T X V_N \triangleq Q \sqrt{\Lambda} Q_p^T Q_p \sqrt{\Lambda} Q^T \quad (1045)$$

^{5.51}Scaling [309] means making a scale, i.e., a numerical representation of qualitative data. If the scale is multidimensional, it's multidimensional scaling. — Jan de Leeuw
When data comprises measurable distances, then reconstruction is termed *metric multidimensional scaling*. In one dimension, N coordinates in X define the scale.

where $\sqrt{\Lambda} Q_p^T Q_p \sqrt{\Lambda} \triangleq \Lambda = \sqrt{\Lambda} \sqrt{\Lambda}$ and where $Q_p \in \mathbb{R}^{n \times N-1}$ is unknown as is its dimension n . Rotation/reflection is accounted for by Q_p yet only its first r columns are necessarily orthonormal.^{5.52} Assuming membership to the unit simplex $y \in \mathcal{S}$ (1002), then point $p = X\sqrt{2}V_N y = Q_p \sqrt{\Lambda} Q^T y$ in \mathbb{R}^n belongs to the translated polyhedron

$$\mathcal{P} - x_1 \quad (1046)$$

whose generating list constitutes the columns of (944)

$$\begin{aligned} [\mathbf{0} \quad X\sqrt{2}V_N] &= [\mathbf{0} \quad Q_p \sqrt{\Lambda} Q^T] \in \mathbb{R}^{n \times N} \\ &= [\mathbf{0} \quad x_2 - x_1 \quad x_3 - x_1 \quad \cdots \quad x_N - x_1] \end{aligned} \quad (1047)$$

The scaled auxiliary matrix V_N represents that translation. A simple choice for Q_p has n set to $N-1$; *id est*, $Q_p = I$. Ideally, each member of the generating list has at most r nonzero entries; r being, affine dimension

$$\text{rank } V_N^T D V_N = \text{rank } Q_p \sqrt{\Lambda} Q^T = \text{rank } \Lambda = r \quad (1048)$$

Each member then has at least $N-1-r$ zeros in its higher-dimensional coordinates because $r \leq N-1$. (956) To truncate those zeros, choose n equal to affine dimension which is the smallest n possible because XV_N has rank $r \leq n$ (952).^{5.53} In that case, the simplest choice for Q_p is $[I \quad \mathbf{0}]$ having dimensions $r \times N-1$.

We may wish to verify the list (1047) found from the diagonalization of $-V_N^T D V_N$. Because of rotation/reflection and translation invariance (§5.5), EDM D can be uniquely made from that list by calculating: (808)

^{5.52}Recall r signifies affine dimension. Q_p is not necessarily an orthogonal matrix. Q_p is constrained such that only its first r columns are necessarily orthonormal because there are only r nonzero eigenvalues in Λ when $-V_N^T D V_N$ has rank r (§5.7.1.1). Remaining columns of Q_p are arbitrary.

^{5.53}If we write $Q^T = \begin{bmatrix} q_1^T \\ \vdots \\ q_{N-1}^T \end{bmatrix}$ as rowwise eigenvectors, $\Lambda = \begin{bmatrix} \lambda_1 & & & \mathbf{0} \\ & \ddots & & \\ & & \lambda_r & \\ \mathbf{0} & & & \ddots & 0 \\ & & & & \ddots & 0 \end{bmatrix}$

in terms of eigenvalues, and $Q_p = [q_{p_1} \cdots q_{p_{N-1}}]$ as column vectors, then $Q_p \sqrt{\Lambda} Q^T = \sum_{i=1}^r \sqrt{\lambda_i} q_{p_i} q_i^T$ is a sum of r linearly independent rank-one matrices (§B.1.1). Hence the summation has rank r .

$$\mathbf{D}(X) = \mathbf{D}(X[\mathbf{0} \quad \sqrt{2}V_{\mathcal{N}}]) = \mathbf{D}(Q_p[\mathbf{0} \quad \sqrt{\Lambda}Q^T]) = \mathbf{D}([\mathbf{0} \quad \sqrt{\Lambda}Q^T]) \quad (1049)$$

This suggests a way to find EDM D given $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$; (*confer*(928))

$$D = \begin{bmatrix} 0 \\ \delta(-V_{\mathcal{N}}^T D V_{\mathcal{N}}) \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & \delta(-V_{\mathcal{N}}^T D V_{\mathcal{N}})^T \end{bmatrix} - 2 \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & -V_{\mathcal{N}}^T D V_{\mathcal{N}} \end{bmatrix} \quad (828)$$

5.12.2 0 geometric center. V

Alternatively, we may perform reconstruction using instead the auxiliary matrix V (§B.4.1), corresponding to the polyhedron

$$\mathcal{P} - \alpha_c \quad (1050)$$

whose geometric center has been translated to the origin. Redimensioning diagonalization factors $Q, \Lambda \in \mathbb{R}^{N \times N}$ and unknown $Q_p \in \mathbb{R}^{n \times N}$, (951)

$$-VDV = 2VX^T X V \triangleq Q\sqrt{\Lambda}Q_p^T Q_p \sqrt{\Lambda}Q^T \triangleq Q\Lambda Q^T \quad (1051)$$

where the geometrically centered generating list constitutes (*confer*(1047))

$$\begin{aligned} XV &= \frac{1}{\sqrt{2}} Q_p \sqrt{\Lambda} Q^T \in \mathbb{R}^{n \times N} \\ &= [x_1 - \frac{1}{N}X\mathbf{1} \quad x_2 - \frac{1}{N}X\mathbf{1} \quad x_3 - \frac{1}{N}X\mathbf{1} \quad \cdots \quad x_N - \frac{1}{N}X\mathbf{1}] \end{aligned} \quad (1052)$$

where $\alpha_c = \frac{1}{N}X\mathbf{1}$. (§5.5.1.0.1) The simplest choice for Q_p is $[I \quad \mathbf{0}] \in \mathbb{R}^{r \times N}$.

Now EDM D can be uniquely made from the list found, by calculating: (808)

$$\mathbf{D}(X) = \mathbf{D}(XV) = \mathbf{D}\left(\frac{1}{\sqrt{2}} Q_p \sqrt{\Lambda} Q^T\right) = \mathbf{D}(\sqrt{\Lambda} Q^T) \frac{1}{2} \quad (1053)$$

This EDM is, of course, identical to (1049). Similarly to (828), from $-VDV$ we can find EDM D ; (*confer*(915))

$$D = \delta(-VDV \frac{1}{2}) \mathbf{1}^T + \mathbf{1} \delta(-VDV \frac{1}{2})^T - 2(-VDV \frac{1}{2}) \quad (834)$$




Figure 113: Map of United States of America showing some state boundaries and the Great Lakes. All plots made by connecting 5020 points. Any difference in scale in (a) through (d) is artifact of plotting routine.

- (a) Shows original map made from decimated (latitude, longitude) data.
- (b) Original map data rotated (freehand) to highlight curvature of Earth.
- (c) Map isometrically reconstructed from an EDM (from distance only).
- (d) Same reconstructed map illustrating curvature.
- (e)(f) Two views of one isotonic reconstruction (from comparative distance); problem (1062) with no sort constraint $\Pi \underline{d}$ (and no hidden line removal).

5.13 Reconstruction examples

5.13.1 Isometric reconstruction

5.13.1.0.1 Example. *Map of the USA.*

The most fundamental application of EDMs is to reconstruct relative point position given only interpoint distance information. Drawing a map of the United States is a good illustration of isometric reconstruction from complete distance data. We obtained latitude and longitude information for the coast, border, states, and Great Lakes from the [usalo atlas data file](#) within the MATLAB Mapping Toolbox; the conversion to Cartesian coordinates (x, y, z) via:

$$\begin{aligned}\phi &\triangleq \pi/2 - \text{latitude} \\ \theta &\triangleq \text{longitude} \\ x &= \sin(\phi) \cos(\theta) \\ y &= \sin(\phi) \sin(\theta) \\ z &= \cos(\phi)\end{aligned}\tag{1054}$$

We used 64% of the available map data to calculate EDM D from $N = 5020$ points. The original (decimated) data and its isometric reconstruction via (1045) are shown in Figure 113(a)-(d). MATLAB code is on [Wikimization](#). The eigenvalues computed for (1043) are

$$\lambda(-V_N^T D V_N) = [199.8 \ 152.3 \ 2.465 \ 0 \ 0 \ 0 \ \dots]^T \tag{1055}$$

The 0 eigenvalues have absolute numerical error on the order of $2\text{E-}13$; meaning, the EDM data indicates three dimensions ($r = 3$) are required for reconstruction to nearly machine precision. \square

5.13.2 Isotonic reconstruction

Sometimes only comparative information about distance is known (Earth is closer to the Moon than it is to the Sun). Suppose, for example, the EDM D for three points is unknown:

$$D = [d_{ij}] = \begin{bmatrix} 0 & d_{12} & d_{13} \\ d_{12} & 0 & d_{23} \\ d_{13} & d_{23} & 0 \end{bmatrix} \in \mathbb{S}_h^{\mathbf{3}} \quad (797)$$

but the comparative data is available:

$$d_{13} \geq d_{23} \geq d_{12} \quad (1056)$$

With the vectorization $\underline{d} = [d_{12} \ d_{13} \ d_{23}]^T \in \mathbb{R}^{\mathbf{3}}$, we express the comparative distance relationship as the nonincreasing sorting

$$\Pi \underline{d} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} d_{12} \\ d_{13} \\ d_{23} \end{bmatrix} = \begin{bmatrix} d_{13} \\ d_{23} \\ d_{12} \end{bmatrix} \in \mathcal{K}_{\mathcal{M}+} \quad (1057)$$

where Π is a given permutation matrix expressing the known sorting action on the entries of unknown EDM D , and $\mathcal{K}_{\mathcal{M}+}$ is the monotone nonnegative cone (§2.13.9.4.2)

$$\mathcal{K}_{\mathcal{M}+} \triangleq \{z \mid z_1 \geq z_2 \geq \cdots \geq z_{N(N-1)/2} \geq 0\} \subseteq \mathbb{R}_+^{N(N-1)/2} \quad (389)$$

where $N(N-1)/2 = 3$ for the present example. From the sorted vectorization (1057) we create the *sort-index matrix*

$$O = \begin{bmatrix} 0 & 1^2 & 3^2 \\ 1^2 & 0 & 2^2 \\ 3^2 & 2^2 & 0 \end{bmatrix} \in \mathbb{S}_h^{\mathbf{3}} \cap \mathbb{R}_+^{\mathbf{3} \times \mathbf{3}} \quad (1058)$$

generally defined

$$O_{ij} \triangleq k^2 \mid d_{ij} = (\Xi \Pi \underline{d})_k, \quad j \neq i \quad (1059)$$

where Ξ is a permutation matrix (1619) completely reversing order of vector entries.

Replacing EDM data with indices-square of a nonincreasing sorting like this is, of course, a heuristic we invented and may be regarded as a nonlinear introduction of much noise into the Euclidean distance matrix. For large data sets, this heuristic makes an otherwise intense problem computationally tractable; we see an example in relaxed problem (1063).




Figure 114: Largest ten eigenvalues, of $-V_N^T O V_N$ for map of USA, sorted by nonincreasing value.

Any process of reconstruction that leaves comparative distance information intact is called *ordinal multidimensional scaling* or *isotonic reconstruction*. Beyond rotation, reflection, and translation error, (§5.5) list reconstruction by isotonic reconstruction is subject to error in absolute scale (*dilation*) and distance ratio. Yet Borg & Groenen argue: [46, §2.2] reconstruction from complete comparative distance information for a large number of points is as highly constrained as reconstruction from an EDM; the larger the number, the better.

5.13.2.1 Isotonic map of the USA

To test Borg & Groenen's conjecture, suppose we make a complete sort-index matrix $O \in \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N}$ for the map of the USA and then substitute O in place of EDM D in the reconstruction process of §5.12. Whereas EDM D returned only three significant eigenvalues (1055), the sort-index matrix O is generally not an EDM (certainly not an EDM with corresponding affine dimension 3) so returns many more. The eigenvalues, calculated with absolute numerical error approximately $5\text{E-}7$, are plotted in Figure 114: (In the code on [Wikimization](#), matrix O is normalized by $(N(N-1)/2)^2$.)

$$\lambda(-V_N^T O V_N) = [880.1 \ 463.9 \ 186.1 \ 46.20 \ 17.12 \ 9.625 \ 8.257 \ 1.701 \ 0.7128 \ 0.6460 \ \dots]^T$$

(1060)

The extra eigenvalues indicate that affine dimension corresponding to an EDM near O is likely to exceed 3. To realize the map, we must simultaneously reduce that dimensionality and find an EDM D closest to O in some sense (a problem explored more in §7) while maintaining the known comparative distance relationship; *e.g.*, given permutation matrix Π expressing the known sorting action on the entries \underline{d} of unknown $D \in \mathbb{S}_h^N$, (67)

$$\underline{d} \triangleq \frac{1}{\sqrt{2}} \text{dvec } D = \begin{bmatrix} d_{12} \\ d_{13} \\ d_{23} \\ d_{14} \\ d_{24} \\ d_{34} \\ \vdots \\ d_{N-1,N} \end{bmatrix} \in \mathbb{R}^{N(N-1)/2} \quad (1061)$$

we can make the sort-index matrix O input to the optimization problem

$$\begin{aligned} & \underset{D}{\text{minimize}} && \| -V_{\mathcal{N}}^T(D - O)V_{\mathcal{N}} \|_{\text{F}} \\ & \text{subject to} && \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} \leq 3 \\ & && \Pi \underline{d} \in \mathcal{K}_{\mathcal{M}+} \\ & && D \in \text{EDM}^N \end{aligned} \quad (1062)$$

that finds the EDM D (corresponding to affine dimension not exceeding 3 in isomorphic $\text{dvec } \text{EDM}^N \cap \Pi^T \mathcal{K}_{\mathcal{M}+}$) closest to O in the sense of Schoenberg (827).

Analytical solution to this problem, ignoring the sort constraint $\Pi \underline{d} \in \mathcal{K}_{\mathcal{M}+}$, is known [313]: we get the convex optimization [*sic*] (§7.1)

$$\begin{aligned} & \underset{D}{\text{minimize}} && \| -V_{\mathcal{N}}^T(D - O)V_{\mathcal{N}} \|_{\text{F}} \\ & \text{subject to} && \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} \leq 3 \\ & && D \in \text{EDM}^N \end{aligned} \quad (1063)$$

Only the three largest nonnegative eigenvalues in (1060) need be retained to make list (1047); the rest are discarded. The reconstruction from EDM D found in this manner is plotted in Figure 113(e)(f). MATLAB code is on [Wikimization](#). From these plots it becomes obvious that inclusion of the sort constraint is necessary for isotonic reconstruction.

That sort constraint demands: any optimal solution D^* must possess the known comparative distance relationship that produces the original ordinal distance data O (1059). Ignoring the sort constraint, apparently, violates it. Yet even more remarkable is how much the map reconstructed using only ordinal data still resembles the original map of the USA after suffering the many violations produced by solving relaxed problem (1063). This suggests the simple reconstruction techniques of §5.12 are robust to a significant amount of noise.

5.13.2.2 Isotonic solution with sort constraint

Because problems involving rank are generally difficult, we will partition (1062) into two problems we know how to solve and then alternate their solution until convergence:

$$\begin{aligned} & \underset{D}{\text{minimize}} && \| -V_{\mathcal{N}}^T(D - O)V_{\mathcal{N}} \|_F \\ & \text{subject to} && \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} \leq 3 \\ & && D \in \mathbb{EDM}^N \end{aligned} \quad (\text{a}) \quad (1064)$$

$$\begin{aligned} & \underset{\sigma}{\text{minimize}} && \| \sigma - \Pi \underline{d} \| \\ & \text{subject to} && \sigma \in \mathcal{K}_{\mathcal{M}+} \end{aligned} \quad (\text{b})$$

where the sort-index matrix O (a given constant in (a)) becomes an implicit vector variable \underline{o}_i solving the i^{th} instance of (1064b)

$$\underline{o}_i \triangleq \Pi^T \sigma^* = \frac{1}{\sqrt{2}} \text{dvec } O_i \in \mathbb{R}^{N(N-1)/2}, \quad i \in \{1, 2, 3, \dots\} \quad (1065)$$

As mentioned in discussion of relaxed problem (1063), a closed-form solution to problem (1064a) exists. Only the first iteration of (1064a) sees the original sort-index matrix O whose entries are nonnegative whole numbers; *id est*, $O_0 = O \in \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N}$ (1059). Subsequent iterations i take the previous solution of (1064b) as input

$$O_i = \text{dvec}^{-1}(\sqrt{2} \underline{o}_i) \in \mathbb{S}^N \quad (1066)$$

real successors to the sort-index matrix O .

New problem (1064b) finds the unique minimum-distance projection of $\Pi \underline{d}$ on the monotone nonnegative cone $\mathcal{K}_{\mathcal{M}+}$. By defining

$$Y^{\dagger T} \triangleq [e_1 - e_2 \quad e_2 - e_3 \quad e_3 - e_4 \quad \cdots \quad e_m] \in \mathbb{R}^{m \times m} \quad (390)$$

where $m \triangleq N(N-1)/2$, we may rewrite (1064b) as an equivalent *quadratic program*; a convex optimization problem [53, §4] in terms of the halfspace-description of $\mathcal{K}_{\mathcal{M}+}$:

$$\begin{aligned} & \underset{\sigma}{\text{minimize}} && (\sigma - \Pi \underline{d})^T (\sigma - \Pi \underline{d}) \\ & \text{subject to} && Y^\dagger \sigma \succeq 0 \end{aligned} \quad (1067)$$

This quadratic program can be converted to a semidefinite program via Schur-form (§3.1.7.2); we get the equivalent problem

$$\begin{aligned} & \underset{t \in \mathbb{R}, \sigma}{\text{minimize}} && t \\ & \text{subject to} && \begin{bmatrix} tI & \sigma - \Pi \underline{d} \\ (\sigma - \Pi \underline{d})^T & 1 \end{bmatrix} \succeq 0 \\ & && Y^\dagger \sigma \succeq 0 \end{aligned} \quad (1068)$$

5.13.2.3 Convergence

In §E.10 we discuss convergence of alternating projection on intersecting convex sets in a Euclidean vector space; convergence to a point in their intersection. Here the situation is different for two reasons:

Firstly, sets of positive semidefinite matrices having an upper bound on rank are generally not convex. Yet in §7.1.4.0.1 we prove (1064a) is equivalent to a projection of nonincreasingly ordered eigenvalues on a subset of the nonnegative orthant:

$$\begin{aligned} & \underset{D}{\text{minimize}} && \|-V_{\mathcal{N}}^T(D - O)V_{\mathcal{N}}\|_F \\ & \text{subject to} && \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} \leq 3 \\ & && D \in \text{EDM}^N \end{aligned} \quad \equiv \quad \begin{aligned} & \underset{\Upsilon}{\text{minimize}} && \|\Upsilon - \Lambda\|_F \\ & \text{subject to} && \delta(\Upsilon) \in \begin{bmatrix} \mathbb{R}_+^3 \\ \mathbf{0} \end{bmatrix} \end{aligned} \quad (1069)$$

where $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \triangleq U \Upsilon U^T \in \mathbb{S}^{N-1}$ and $-V_{\mathcal{N}}^T O V_{\mathcal{N}} \triangleq Q \Lambda Q^T \in \mathbb{S}^{N-1}$ are ordered diagonalizations (§A.5). It so happens: optimal orthogonal U^* always equals Q given. Linear operator $T(A) = U^{*T} A U^*$, acting on square matrix A , is a bijective isometry because the Frobenius norm is orthogonally invariant (42). This isometric isomorphism T thus maps a nonconvex problem to a convex one that preserves distance.

Secondly, the second half (1064b) of the *alternation* takes place in a different vector space; \mathbb{S}_h^N (*versus* \mathbb{S}^{N-1}). From §5.6 we know these two vector spaces are related by an isomorphism, $\mathbb{S}^{N-1} = \mathbf{V}_{\mathcal{N}}(\mathbb{S}_h^N)$ (933), but not by an isometry.

We have, therefore, no guarantee from theory of alternating projection that the alternation (1064) converges to a point, in the set of all EDMs corresponding to affine dimension not in excess of 3, belonging to $\text{dvec } \mathbb{EDM}^N \cap \Pi^T \mathcal{K}_{\mathcal{M}_+}$.

5.13.2.4 Interlude

We have not implemented the second half (1067) of alternation (1064) for USA map data because memory-demands exceed the capability of our 32-bit laptop computer.

5.13.2.4.1 Exercise. *Convergence of isotonic solution by alternation.* Empirically demonstrate convergence, discussed in §5.13.2.3, on a smaller data set. ▼

It would be remiss not to mention another method of solution to this isotonic reconstruction problem: Once again we assume only comparative distance data like (1056) is available. Given known set of indices \mathcal{I}

$$\begin{aligned} & \underset{D}{\text{minimize}} && \text{rank } V D V \\ & \text{subject to} && d_{ij} \leq d_{kl} \leq d_{mn} \quad \forall (i, j, k, l, m, n) \in \mathcal{I} \\ & && D \in \mathbb{EDM}^N \end{aligned} \tag{1070}$$

this problem minimizes affine dimension while finding an EDM whose entries satisfy known comparative relationships. Suitable rank heuristics are discussed in §4.4.1 and §7.2.2 that will transform this to a convex optimization problem.

Using contemporary computers, even with a rank heuristic in place of the objective function, this problem formulation is more difficult to compute than the relaxed counterpart problem (1063). That is because there exist efficient algorithms to compute a selected few eigenvalues and eigenvectors from a very large matrix. Regardless, it is important to recognize: the optimal solution set for this problem (1070) is practically always different from the optimal solution set for its counterpart, problem (1062).

5.14 Fifth property of Euclidean metric

We continue now with the question raised in §5.3 regarding the necessity for at least one requirement more than the four properties of the Euclidean metric (§5.2) to certify realizability of a bounded convex polyhedron or to reconstruct a generating list for it from incomplete distance information. There we saw that the four Euclidean metric properties are necessary for $D \in \text{EDM}^N$ in the case $N = 3$, but become insufficient when cardinality N exceeds 3 (regardless of affine dimension).

5.14.1 Recapitulate

In the particular case $N = 3$, $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$ (974) and $D \in \mathbb{S}_h^3$ are necessary and sufficient conditions for D to be an EDM. By (976), triangle inequality is then the only Euclidean condition bounding the necessarily nonnegative d_{ij} ; and those bounds are tight. That means the first four properties of the Euclidean metric are necessary and sufficient conditions for D to be an EDM in the case $N = 3$; for $i, j \in \{1, 2, 3\}$

$$\begin{aligned} \sqrt{d_{ij}} &\geq 0, \quad i \neq j \\ \sqrt{d_{ij}} &= 0, \quad i = j \\ \sqrt{d_{ij}} &= \sqrt{d_{ji}} \\ \sqrt{d_{ij}} &\leq \sqrt{d_{ik}} + \sqrt{d_{kj}}, \quad i \neq j \neq k \end{aligned} \quad \Leftrightarrow \quad \begin{aligned} -V_{\mathcal{N}}^T D V_{\mathcal{N}} &\succeq 0 \\ D &\in \mathbb{S}_h^3 \end{aligned} \quad \Leftrightarrow \quad D \in \text{EDM}^3 \quad (1071)$$

Yet those four properties become insufficient when $N > 3$.

5.14.2 Derivation of the fifth

Correspondence between the triangle inequality and the EDM was developed in §5.8.2 where a triangle inequality (976a) was revealed within the leading principal 2×2 submatrix of $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ when positive semidefinite. Our choice of the leading principal submatrix was arbitrary; actually, a unique triangle inequality like (871) corresponds to any one of the $(N-1)!/(2!(N-1-2)!)$ principal 2×2 submatrices.^{5.54} Assuming $D \in \mathbb{S}_h^4$ and $-V_{\mathcal{N}}^T D V_{\mathcal{N}} \in \mathbb{S}^3$, then by the *positive (semi)definite principal submatrices*

^{5.54}There are fewer principal 2×2 submatrices in $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ than there are triangles made by four or more points because there are $N!/(3!(N-3)!)$ triangles made by point triples. The triangles corresponding to those submatrices all have vertex x_1 . (confer §5.8.2.1)

theorem (§A.3.1.0.4) it is sufficient to prove all d_{ij} are nonnegative, all triangle inequalities are satisfied, and $\det(-V_{\mathcal{N}}^T D V_{\mathcal{N}})$ is nonnegative. When $N=4$, in other words, that nonnegative determinant becomes the fifth and last Euclidean metric requirement for $D \in \mathbb{EDM}^N$. We now endeavor to ascribe geometric meaning to it.

5.14.2.1 Nonnegative determinant

By (877) when $D \in \mathbb{EDM}^4$, $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ is equal to inner product (872),

$$\Theta^T \Theta = \begin{bmatrix} d_{12} & \sqrt{d_{12}d_{13}} \cos \theta_{213} & \sqrt{d_{12}d_{14}} \cos \theta_{214} \\ \sqrt{d_{12}d_{13}} \cos \theta_{213} & d_{13} & \sqrt{d_{13}d_{14}} \cos \theta_{314} \\ \sqrt{d_{12}d_{14}} \cos \theta_{214} & \sqrt{d_{13}d_{14}} \cos \theta_{314} & d_{14} \end{bmatrix} \quad (1072)$$

Because Euclidean space is an inner-product space, the more concise inner-product form of the determinant is admitted;

$$\det(\Theta^T \Theta) = -d_{12}d_{13}d_{14}(\cos(\theta_{213})^2 + \cos(\theta_{214})^2 + \cos(\theta_{314})^2 - 2 \cos \theta_{213} \cos \theta_{214} \cos \theta_{314} - 1) \quad (1073)$$

The determinant is nonnegative if and only if

$$\begin{aligned} \cos \theta_{214} \cos \theta_{314} - \sqrt{\sin(\theta_{214})^2 \sin(\theta_{314})^2} &\leq \cos \theta_{213} \leq \cos \theta_{214} \cos \theta_{314} + \sqrt{\sin(\theta_{214})^2 \sin(\theta_{314})^2} \\ &\Leftrightarrow \\ \cos \theta_{213} \cos \theta_{314} - \sqrt{\sin(\theta_{213})^2 \sin(\theta_{314})^2} &\leq \cos \theta_{214} \leq \cos \theta_{213} \cos \theta_{314} + \sqrt{\sin(\theta_{213})^2 \sin(\theta_{314})^2} \\ &\Leftrightarrow \\ \cos \theta_{213} \cos \theta_{214} - \sqrt{\sin(\theta_{213})^2 \sin(\theta_{214})^2} &\leq \cos \theta_{314} \leq \cos \theta_{213} \cos \theta_{214} + \sqrt{\sin(\theta_{213})^2 \sin(\theta_{214})^2} \end{aligned} \quad (1074)$$

which simplifies, for $0 \leq \theta_{i1\ell}, \theta_{\ell 1j}, \theta_{i1j} \leq \pi$ and all $i \neq j \neq \ell \in \{2, 3, 4\}$, to

$$\cos(\theta_{i1\ell} + \theta_{\ell 1j}) \leq \cos \theta_{i1j} \leq \cos(\theta_{i1\ell} - \theta_{\ell 1j}) \quad (1075)$$

Analogously to triangle inequality (988), the determinant is 0 upon equality on either side of (1075) which is tight. Inequality (1075) can be equivalently written linearly as a triangle inequality between relative angles [352, §1.4];

$$\begin{aligned} |\theta_{i1\ell} - \theta_{\ell 1j}| &\leq \theta_{i1j} \leq \theta_{i1\ell} + \theta_{\ell 1j} \\ \theta_{i1\ell} + \theta_{\ell 1j} + \theta_{i1j} &\leq 2\pi \\ 0 &\leq \theta_{i1\ell}, \theta_{\ell 1j}, \theta_{i1j} \leq \pi \end{aligned} \quad (1076)$$




Figure 115: The *relative-angle inequality tetrahedron* (1077) bounding \mathbb{EDM}^4 is regular; drawn in entirety. Each angle θ (869) must belong to this solid to be realizable.

Generalizing this:

5.14.2.1.1 Fifth property of Euclidean metric - restatement.

Relative-angle inequality. (confer §5.3.1.0.1) [43] [44, p.17, p.107] [207, §3.1] Augmenting the four fundamental Euclidean metric properties in \mathbb{R}^n , for all $i, j, \ell \neq k \in \{1 \dots N\}$, $i < j < \ell$, and for $N \geq 4$ distinct points $\{x_k\}$, the inequalities

$$\begin{aligned}
 |\theta_{ik\ell} - \theta_{\ell kj}| &\leq \theta_{ikj} \leq \theta_{ik\ell} + \theta_{\ell kj} & (a) \\
 \theta_{ik\ell} + \theta_{\ell kj} + \theta_{ikj} &\leq 2\pi & (b) \\
 0 \leq \theta_{ik\ell}, \theta_{\ell kj}, \theta_{ikj} &\leq \pi & (c)
 \end{aligned} \tag{1077}$$

where $\theta_{ikj} = \theta_{jki}$ is the angle between vectors at vertex x_k (as defined in (869) and illustrated in Figure 97), must be satisfied at each point x_k regardless of affine dimension. \diamond

Because point labelling is arbitrary, this fifth Euclidean metric requirement must apply to each of the N points as though each were in turn labelled x_1 ; hence the new index k in (1077). Just as the triangle inequality is the ultimate test for realizability of only three points, the relative-angle inequality is the ultimate test for only four. For four distinct points, the triangle inequality remains a necessary although penultimate test; (§5.4.3)

$$\begin{aligned} \text{Four Euclidean metric properties (§5.2).} & \Leftrightarrow -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \\ \text{Angle } \theta \text{ inequality (802) or (1077).} & \Leftrightarrow D \in \mathbb{S}_h^4 \Leftrightarrow D = \mathbf{D}(\Theta) \in \mathbb{EDM}^4 \end{aligned} \quad (1078)$$

The relative-angle inequality, for this case, is illustrated in Figure 115.

5.14.2.2 Beyond the fifth metric property

When cardinality N exceeds 4, the first four properties of the Euclidean metric and the relative-angle inequality together become insufficient conditions for realizability. In other words, the four Euclidean metric properties and relative-angle inequality remain necessary but become a sufficient test only for positive semidefiniteness of all the principal 3×3 submatrices [*sic*] in $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$. Relative-angle inequality can be considered the ultimate test only for realizability at each vertex x_k of each and every purported tetrahedron constituting a hyperdimensional body.

When $N=5$ in particular, relative-angle inequality becomes the penultimate Euclidean metric requirement while nonnegativity of then unwieldy $\det(\Theta^T \Theta)$ corresponds (by the *positive (semi)definite principal submatrices theorem* in §A.3.1.0.4) to the sixth and last Euclidean metric requirement. Together these six tests become necessary and sufficient, and so on.

Yet for all values of N , only assuming nonnegative d_{ij} , relative-angle matrix inequality in (990) is necessary and sufficient to certify realizability; (§5.4.3.1)

$$\begin{aligned} \text{Euclidean metric property 1 (§5.2).} & \Leftrightarrow -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \\ \text{Angle matrix inequality } \Omega \succeq 0 \text{ (878).} & \Leftrightarrow D \in \mathbb{S}_h^N \Leftrightarrow D = \mathbf{D}(\Omega, d) \in \mathbb{EDM}^N \end{aligned} \quad (1079)$$

Like matrix criteria (803), (827), and (990), the relative-angle matrix inequality and nonnegativity property subsume all the Euclidean metric properties and further requirements.

5.14.3 Path not followed

As a means to test for realizability of four or more points, an intuitively appealing way to augment the four Euclidean metric properties is to recognize generalizations of the triangle inequality: In the case $N=4$, the three-dimensional analogue to triangle & distance is tetrahedron & facet-area, while in the case $N=5$ the four-dimensional analogue is polychoron & facet-volume, *ad infinitum*. For N points, $N+1$ metric properties are required.

5.14.3.1 $N = 4$

Each of the four facets of a general tetrahedron is a triangle and its relative interior. Suppose we identify each facet of the tetrahedron by its area-squared: c_1, c_2, c_3, c_4 . Then analogous to metric property 4, we may write a tight^{5.55} area inequality for the facets

$$\sqrt{c_i} \leq \sqrt{c_j} + \sqrt{c_k} + \sqrt{c_\ell}, \quad i \neq j \neq k \neq \ell \in \{1, 2, 3, 4\} \quad (1080)$$

which is a generalized “triangle” inequality [201, §1.1] that follows from

$$\sqrt{c_i} = \sqrt{c_j} \cos \varphi_{ij} + \sqrt{c_k} \cos \varphi_{ik} + \sqrt{c_\ell} \cos \varphi_{i\ell} \quad (1081)$$

[214] [333, *Law of Cosines*] where φ_{ij} is the dihedral angle at the common edge between triangular facets i and j .

If D is the EDM corresponding to the whole tetrahedron, then area-squared of the i^{th} triangular facet has a convenient formula in terms of $D_i \in \mathbb{EDM}^{N-1}$ the EDM corresponding to that particular facet: From the *Cayley-Menger determinant*^{5.56} for simplices, [333] [114] [141, §4] [77, §3.3] the i^{th} facet area-squared for $i \in \{1 \dots N\}$ is (§A.4.1)

^{5.55}The upper bound is met when all angles in (1081) are simultaneously 0; that occurs, for example, if one point is relatively interior to the convex hull of the three remaining.

^{5.56}whose foremost characteristic is: the determinant vanishes if and only if affine dimension does not equal penultimate cardinality; *id est*, $\det \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} = 0 \Leftrightarrow r < N-1$ where D is any EDM (§5.7.3.0.1). Otherwise, the determinant is negative.

$$c_i = \frac{-1}{2^{N-2}(N-2)!^2} \det \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D_i \end{bmatrix} \quad (1082)$$

$$= \frac{(-1)^N}{2^{N-2}(N-2)!^2} \det D_i (\mathbf{1}^T D_i^{-1} \mathbf{1}) \quad (1083)$$

$$= \frac{(-1)^N}{2^{N-2}(N-2)!^2} \mathbf{1}^T \text{cof}(D_i)^T \mathbf{1} \quad (1084)$$

where D_i is the i^{th} principal $(N-1) \times (N-1)$ submatrix^{5.57} of $D \in \text{EDM}^N$, and $\text{cof}(D_i)$ is the $(N-1) \times (N-1)$ matrix of *cofactors* [294, §4] corresponding to D_i . The number of principal 3×3 submatrices in D is, of course, equal to the number of triangular facets in the tetrahedron; four ($N!/(3!(N-3)!)$) when $N=4$.

5.14.3.1.1 Exercise. *Sufficiency conditions for an EDM of four points.* Triangle inequality (property 4) and area inequality (1080) are conditions necessary for D to be an EDM. Prove their sufficiency in conjunction with the remaining three Euclidean metric properties. ▼

5.14.3.2 $N = 5$

Moving to the next level, we might encounter a Euclidean body called *polychoron*: a bounded polyhedron in four dimensions.^{5.58} The polychoron has five ($N!/(4!(N-4)!)$) facets, each of them a general tetrahedron whose volume-squared c_i is calculated using the same formula; (1082) where D is the EDM corresponding to the polychoron, and D_i is the EDM corresponding to the i^{th} facet (the principal 4×4 submatrix of $D \in \text{EDM}^N$ corresponding to the i^{th} tetrahedron). The analogue to triangle & distance is now polychoron & facet-volume. We could then write another generalized “triangle” inequality like (1080) but in terms of facet volume; [338, §IV]

$$\sqrt{c_i} \leq \sqrt{c_j} + \sqrt{c_k} + \sqrt{c_\ell} + \sqrt{c_m}, \quad i \neq j \neq k \neq \ell \neq m \in \{1 \dots 5\} \quad (1085)$$

^{5.57}Every principal submatrix of an EDM remains an EDM. [207, §4.1]

^{5.58}The simplest polychoron is called a pentatope [333]; a regular simplex hence convex. (A *pentahedron* is a three-dimensional body having five vertices.)




Figure 116: Length of one-dimensional face a equals height $h=a=1$ of this nonsimplicial pyramid in \mathbb{R}^3 with square base inscribed in a circle of radius R centered at the origin. [333, *Pyramid*]

5.14.3.2.1 Exercise. *Sufficiency for an EDM of five points.*

For $N=5$, triangle (distance) inequality (§5.2), area inequality (1080), and volume inequality (1085) are conditions necessary for D to be an EDM. Prove their sufficiency. ▼

5.14.3.3 Volume of simplices

There is no known formula for the volume of a bounded general convex polyhedron expressed either by halfspace or vertex-description. [350, §2.1] [248, p.173] [205] [152] [153] Volume is a concept germane to \mathbb{R}^3 ; in higher dimensions it is called *content*. Applying the *EDM assertion* (§5.9.1.0.4) and a result from [53, p.407], a general nonempty simplex (§2.12.3) in \mathbb{R}^{N-1} corresponding to an EDM $D \in \mathbb{S}_h^N$ has content

$$\sqrt{c} = \text{content}(\mathcal{S}) \sqrt{\det(-V_N^T D V_N)} \quad (1086)$$

where the content-squared of the unit simplex $\mathcal{S} \subset \mathbb{R}^{N-1}$ is proportional to its Cayley-Menger determinant;

$$\text{content}(\mathcal{S})^2 = \frac{-1}{2^{N-1}(N-1)!^2} \det \begin{bmatrix} 0 & & & & \mathbf{1}^T \\ \mathbf{1} & -\mathbf{D}([\mathbf{0} & e_1 & e_2 & \cdots & e_{N-1}]) \end{bmatrix} \quad (1087)$$

where $e_i \in \mathbb{R}^{N-1}$ and the EDM operator used is $\mathbf{D}(X)$ (808).

5.14.3.3.1 Example. *Pyramid.*

A formula for volume of a pyramid is known;^{5.59} it is $\frac{1}{3}$ the product of its base area with its height. [198] The pyramid in Figure 116 has volume $\frac{1}{3}$. To find its volume using EDMs, we must first decompose the pyramid into simplicial parts. Slicing it in half along the plane containing the line segments corresponding to radius R and height h we find the vertices of one simplex,

$$X = \begin{bmatrix} 1/2 & 1/2 & -1/2 & 0 \\ 1/2 & -1/2 & -1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \in \mathbb{R}^{n \times N} \quad (1088)$$

where $N = n + 1$ for any nonempty simplex in \mathbb{R}^n . The volume of this simplex is half that of the entire pyramid; *id est*, $\sqrt{c} = \frac{1}{6}$ found by evaluating (1086). \square

With that, we conclude digression of path.

5.14.4 Affine dimension reduction in three dimensions

(confer §5.8.4) The determinant of any $M \times M$ matrix is equal to the product of its M eigenvalues. [294, §5.1] When $N = 4$ and $\det(\Theta^T \Theta)$ is 0, that means one or more eigenvalues of $\Theta^T \Theta \in \mathbb{R}^{3 \times 3}$ are 0. The determinant will go to 0 whenever equality is attained on either side of (802), (1077a), or (1077b), meaning that a tetrahedron has collapsed to a lower affine dimension; *id est*, $r = \text{rank } \Theta^T \Theta = \text{rank } \Theta$ is reduced below $N - 1$ exactly by the number of 0 eigenvalues (§5.7.1.1).

In solving completion problems of any size N where one or more entries of an EDM are unknown, therefore, dimension r of the affine hull required to contain the unknown points is potentially reduced by selecting distances to attain equality in (802) or (1077a) or (1077b).

^{5.59}Pyramid volume is independent of the paramount vertex position as long as its height remains constant.

5.14.4.1 *Exemplum redux*

We now apply the *fifth Euclidean metric property* to an earlier problem:

5.14.4.1.1 Example. *Small completion problem, IV.* (confer §5.9.3.0.1) Returning again to Example 5.3.0.0.2 that pertains to Figure 96 where $N=4$, distance-square d_{14} is ascertainable from the fifth Euclidean metric property. Because all distances in (800) are known except $\sqrt{d_{14}}$, then $\cos \theta_{123}=0$ and $\theta_{324}=0$ result from identity (869). Applying (802),

$$\begin{aligned} \cos(\theta_{123} + \theta_{324}) &\leq \cos \theta_{124} \leq \cos(\theta_{123} - \theta_{324}) \\ 0 &\leq \cos \theta_{124} \leq 0 \end{aligned} \tag{1089}$$

It follows again from (869) that d_{14} can only be 2. As explained in this subsection, affine dimension r cannot exceed $N-2$ because equality is attained in (1089). \square

Chapter 6

Cone of distance matrices

For $N > 3$, the cone of EDMs is no longer a circular cone and the geometry becomes complicated. . .

–Hayden, Wells, Liu, & Tarazaga (1991) [164, §3]

In the subspace of symmetric matrices \mathbb{S}^N , we know that the convex cone of Euclidean distance matrices \mathbb{EDM}^N (the EDM cone) does not intersect the positive semidefinite cone \mathbb{S}_+^N (PSD cone) except at the origin, their only vertex; there can be no positive nor negative semidefinite EDM. (1014) [207]

$$\mathbb{EDM}^N \cap \mathbb{S}_+^N = \mathbf{0} \quad (1090)$$

Even so, the two convex cones can be related. In §6.8.1 we prove the equality

$$\mathbb{EDM}^N = \mathbb{S}_h^N \cap (\mathbb{S}_c^{N\perp} - \mathbb{S}_+^N) \quad (1183)$$

a resemblance to EDM definition (808) where

$$\mathbb{S}_h^N \triangleq \{A \in \mathbb{S}^N \mid \delta(A) = \mathbf{0}\} \quad (60)$$

is the symmetric hollow subspace (§2.2.3) and where

$$\mathbb{S}_c^{N\perp} = \{u\mathbf{1}^T + \mathbf{1}u^T \mid u \in \mathbb{R}^N\} \quad (1887)$$

is the orthogonal complement of the geometric center subspace (§E.7.2.0.2)

$$\mathbb{S}_c^N \triangleq \{Y \in \mathbb{S}^N \mid Y\mathbf{1} = \mathbf{0}\} \quad (1885)$$

6.0.1 gravity

Equality (1183) is equally important as the known isomorphisms (922) (923) (934) (935) relating the EDM cone \mathbb{EDM}^N to positive semidefinite cone \mathbb{S}_+^{N-1} (§5.6.2.1) or to an $N(N-1)/2$ -dimensional face of \mathbb{S}_+^N (§5.6.1.1).^{6.1} But those isomorphisms have never led to this equality relating whole cones \mathbb{EDM}^N and \mathbb{S}_+^N .

Equality (1183) is not obvious from the various EDM definitions such as (808) or (1109) because inclusion must be proved algebraically in order to establish equality; $\mathbb{EDM}^N \supseteq \mathbb{S}_h^N \cap (\mathbb{S}_c^{N\perp} - \mathbb{S}_+^N)$. We will instead prove (1183) using purely geometric methods.

6.0.2 highlight

In §6.8.1.7 we show: the Schoenberg criterion for discriminating Euclidean distance matrices

$$D \in \mathbb{EDM}^N \Leftrightarrow \begin{cases} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \in \mathbb{S}_+^{N-1} \\ D \in \mathbb{S}_h^N \end{cases} \quad (827)$$

is a discretized membership relation (§2.13.4, dual generalized inequalities) between the EDM cone and its ordinary dual.

^{6.1}Because both positive semidefinite cones are frequently in play, dimension is explicit.

6.1 Defining EDM cone

We invoke a popular matrix criterion to illustrate correspondence between the EDM and PSD cones belonging to the ambient space of symmetric matrices:

$$D \in \mathbb{EDM}^N \Leftrightarrow \begin{cases} -VDV \in \mathbb{S}_+^N \\ D \in \mathbb{S}_h^N \end{cases} \quad (832)$$

where $V \in \mathbb{S}^N$ is the geometric centering matrix (§B.4). The set of all EDMs of dimension $N \times N$ forms a closed convex cone \mathbb{EDM}^N because any pair of EDMs satisfies the definition of a convex cone (158); *videlicet*, for each and every $\zeta_1, \zeta_2 \geq 0$ (§A.3.1.0.2)

$$\begin{aligned} \zeta_1 VD_1V + \zeta_2 VD_2V \succeq 0 \\ \zeta_1 D_1 + \zeta_2 D_2 \in \mathbb{S}_h^N \end{aligned} \Leftrightarrow \begin{aligned} VD_1V \succeq 0, \quad VD_2V \succeq 0 \\ D_1 \in \mathbb{S}_h^N, \quad D_2 \in \mathbb{S}_h^N \end{aligned} \quad (1091)$$

and convex cones are invariant to inverse linear transformation [271, p.22].

6.1.0.0.1 Definition. Cone of Euclidean distance matrices.

In the subspace of symmetric matrices, the set of all Euclidean distance matrices forms a unique immutable pointed closed convex cone called the *EDM cone*: for $N > 0$

$$\begin{aligned} \mathbb{EDM}^N &\triangleq \{D \in \mathbb{S}_h^N \mid -VDV \in \mathbb{S}_+^N\} \\ &= \bigcap_{z \in \mathcal{N}(\mathbf{1}^T)} \{D \in \mathbb{S}^N \mid \langle zz^T, -D \rangle \geq 0, \delta(D) = \mathbf{0}\} \end{aligned} \quad (1092)$$

The EDM cone in isomorphic $\mathbb{R}^{N(N+1)/2}$ [*sic*] is the intersection of an infinite number (when $N > 2$) of halfspaces about the origin and a finite number of hyperplanes through the origin in vectorized variable $D = [d_{ij}]$. Hence \mathbb{EDM}^N has empty interior with respect to \mathbb{S}^N because it is confined to the symmetric hollow subspace \mathbb{S}_h^N . The EDM cone relative interior comprises

$$\begin{aligned} \text{rel int } \mathbb{EDM}^N &= \bigcap_{z \in \mathcal{N}(\mathbf{1}^T)} \{D \in \mathbb{S}^N \mid \langle zz^T, -D \rangle > 0, \delta(D) = \mathbf{0}\} \\ &= \{D \in \mathbb{EDM}^N \mid \text{rank}(VDV) = N - 1\} \end{aligned} \quad (1093)$$

while its relative boundary comprises

$$\begin{aligned} \text{rel } \partial \mathbb{EDM}^N &= \{D \in \mathbb{EDM}^N \mid \langle zz^T, -D \rangle = 0 \text{ for some } z \in \mathcal{N}(\mathbf{1}^T)\} \\ &= \{D \in \mathbb{EDM}^N \mid \text{rank}(VDV) < N - 1\} \end{aligned} \quad (1094)$$

△




Figure 117: Relative boundary (tiled) of EDM cone EDM^3 drawn truncated in isometrically isomorphic subspace \mathbb{R}^3 . **(a)** EDM cone drawn in usual distance-square coordinates d_{ij} . View is from interior toward origin. Unlike positive semidefinite cone, EDM cone is not self-dual, neither is it proper in ambient symmetric subspace (dual EDM cone for this example belongs to isomorphic \mathbb{R}^6). **(b)** Drawn in its natural coordinates $\sqrt{d_{ij}}$ (absolute distance), cone remains convex (*confer* §5.10); intersection of three halfspaces (977) whose partial boundaries each contain origin. Cone geometry becomes “complicated” (nonpolyhedral) in higher dimension. [164, §3] **(c)** Two coordinate systems artificially superimposed. Coordinate transformation from d_{ij} to $\sqrt{d_{ij}}$ appears a topological contraction. **(d)** Sitting on its vertex $\mathbf{0}$, pointed EDM^3 is a circular cone having axis of revolution $\text{dvec}(-E) = \text{dvec}(\mathbf{1}\mathbf{1}^T - I)$ (1010) (67). Rounded vertex is plot artifact.

This cone is more easily visualized in the isomorphic vector subspace $\mathbb{R}^{N(N-1)/2}$ corresponding to \mathbb{S}_h^N :

In the case $N = 1$ point, the EDM cone is the origin in \mathbb{R}^0 .

In the case $N = 2$, the EDM cone is the nonnegative real line in \mathbb{R} ; a halfline in a subspace of the realization in Figure 125.

The EDM cone in the case $N = 3$ is a circular cone in \mathbb{R}^3 illustrated in Figure 117(a)(d); rather, the set of all matrices

$$D = \begin{bmatrix} 0 & d_{12} & d_{13} \\ d_{12} & 0 & d_{23} \\ d_{13} & d_{23} & 0 \end{bmatrix} \in \mathbb{EDM}^3 \quad (1095)$$

makes a circular cone in this dimension. In this case, the first four Euclidean metric properties are necessary and sufficient tests to certify realizability of triangles; (1071). Thus triangle inequality property 4 describes three halfspaces (977) whose intersection makes a polyhedral cone in \mathbb{R}^3 of realizable $\sqrt{d_{ij}}$ (absolute distance); an isomorphic subspace representation of the set of all EDMs D in the natural coordinates

$$\sqrt{\circ}D \triangleq \begin{bmatrix} 0 & \sqrt{d_{12}} & \sqrt{d_{13}} \\ \sqrt{d_{12}} & 0 & \sqrt{d_{23}} \\ \sqrt{d_{13}} & \sqrt{d_{23}} & 0 \end{bmatrix} \quad (1096)$$

illustrated in Figure 117(b).

6.2 Polyhedral bounds

The convex cone of EDMs is nonpolyhedral in d_{ij} for $N > 2$; *e.g.*, Figure 117(a). Still we found necessary and sufficient bounding polyhedral relations consistent with EDM cones for cardinality $N = 1, 2, 3, 4$:

- $N = 3$. Transforming distance-square coordinates d_{ij} by taking their positive square root provides the polyhedral cone in Figure 117(b); polyhedral because an intersection of three halfspaces in natural coordinates $\sqrt{d_{ij}}$ is provided by triangle inequalities (977). This polyhedral cone implicitly encompasses necessary and sufficient metric properties: nonnegativity, self-distance, symmetry, and triangle inequality.




Figure 118: (a) In isometrically isomorphic subspace \mathbb{R}^3 , intersection of \mathbb{EDM}^3 with hyperplane $\partial\mathcal{H}$ representing one fixed symmetric entry $d_{23} = \kappa$ (both drawn truncated, rounded vertex is artifact of plot). EDMs in this dimension corresponding to affine dimension 1 comprise relative boundary of EDM cone (§6.6). Since intersection illustrated includes a nontrivial subset of cone's relative boundary, then it is apparent there exist infinitely many EDM completions corresponding to affine dimension 1. In this dimension it is impossible to represent a unique nonzero completion corresponding to affine dimension 1, for example, using a single hyperplane because any hyperplane supporting relative boundary at a particular point Γ contains an entire ray $\{\zeta\Gamma \mid \zeta \geq 0\}$ belonging to $\text{rel } \partial\mathbb{EDM}^3$ by Lemma 2.8.0.0.1. (b) $d_{13} = \kappa$. (c) $d_{12} = \kappa$.

$N = 4$. Relative-angle inequality (1077) together with four Euclidean metric properties are necessary and sufficient tests for realizability of tetrahedra. (1078) Albeit relative angles θ_{ikj} (869) are nonlinear functions of the d_{ij} , relative-angle inequality provides a regular tetrahedron in \mathbb{R}^3 [sic] (Figure 115) bounding angles θ_{ikj} at vertex x_k consistently with EDM^4 .^{6.2}

Yet were we to employ the procedure outlined in §5.14.3 for making generalized triangle inequalities, then we would find all the necessary and sufficient d_{ij} -transformations for generating bounding polyhedra consistent with EDMs of any higher dimension ($N > 3$).

6.3 $\sqrt{\text{EDM}}$ cone is not convex

For some applications, like a molecular conformation problem (Figure 3, Figure 106) or multidimensional scaling [87] [314], absolute distance $\sqrt{d_{ij}}$ is the preferred variable. Taking square root of the entries in all EDMs D of dimension N , we get another cone but not a convex cone when $N > 3$ (Figure 117(b)): [78, §4.5.2]

$$\sqrt{\text{EDM}^N} \triangleq \{\sqrt[3]{D} \mid D \in \text{EDM}^N\} \quad (1097)$$

where $\sqrt[3]{D}$ is defined like (1096). It is a cone simply because any cone is completely constituted by rays emanating from the origin: (§2.7) Any given ray $\{\zeta \Gamma \in \mathbb{R}^{N(N-1)/2} \mid \zeta \geq 0\}$ remains a ray under entrywise square root: $\{\sqrt{\zeta} \Gamma \in \mathbb{R}^{N(N-1)/2} \mid \zeta \geq 0\}$. It is already established that

$$D \in \text{EDM}^N \Rightarrow \sqrt[3]{D} \in \text{EDM}^N \quad (1009)$$

But because of how $\sqrt{\text{EDM}^N}$ is defined, it is obvious that (confer §5.10)

$$D \in \text{EDM}^N \Leftrightarrow \sqrt[3]{D} \in \sqrt{\text{EDM}^N} \quad (1098)$$

Were $\sqrt{\text{EDM}^N}$ convex, then given $\sqrt[3]{D_1}, \sqrt[3]{D_2} \in \sqrt{\text{EDM}^N}$ we would expect their conic combination $\sqrt[3]{D_1} + \sqrt[3]{D_2}$ to be a member of $\sqrt{\text{EDM}^N}$.

^{6.2}Still, property-4 triangle inequalities (977) corresponding to each principal 3×3 submatrix of $-V_N^T D V_N$ demand that the corresponding $\sqrt{d_{ij}}$ belong to a polyhedral cone like that in Figure 117(b).

That is easily proven false by counter-example via (1098), for then $(\sqrt[N]{D_1} + \sqrt[N]{D_2}) \circ (\sqrt[N]{D_1} + \sqrt[N]{D_2})$ would need to be a member of EDM^N .

Notwithstanding,

$$\sqrt{\text{EDM}^N} \subseteq \text{EDM}^N \quad (1099)$$

by (1009) (Figure 117), and we learn how to transform a nonconvex *proximity problem* in the natural coordinates $\sqrt{d_{ij}}$ to a convex optimization in §7.2.1.

6.4 A geometry of completion

Intriguing is the question of whether the list in $X \in \mathbb{R}^{n \times N}$ (69) may be reconstructed given an incomplete noiseless EDM, and under what circumstances reconstruction is unique. [1] [3] [4] [5] [7] [16] [183] [192] [206] [207] [208]

If one or more entries of a particular EDM are fixed, then geometric interpretation of the feasible set of completions is the intersection of the EDM cone EDM^N in isomorphic subspace $\mathbb{R}^{N(N-1)/2}$ with as many hyperplanes as there are fixed symmetric entries. (Depicted in Figure 118(a) is an intersection of the EDM cone EDM^3 with a single hyperplane representing the set of all EDMs having one fixed symmetric entry.) Assuming a nonempty intersection, then the number of completions is generally infinite, and those corresponding to particular affine dimension $r < N - 1$ belong to some generally nonconvex subset of that intersection (*confer* §2.9.2.6.2) that can be as small as a point. Indeed, Trosset remarks: [312, §1] *It is not known how to proceed if one wishes to restrict the dimension of the Euclidean space in which the configuration of points may be constructed.*

6.4.0.0.1 Example. *Diffusing, decompacting, unfurling, unfolding.* [332] A process minimizing affine dimension (§2.1.5) of certain kinds of Euclidean *manifold* by topological transformation can be posed as a completion problem (*confer* §E.10.2.1.2). Weinberger & Saul, who originated the technique, specify an applicable manifold in three dimensions by analogy to an ordinary sheet of paper (*confer* §2.1.6); imagine, we find it deformed from flatness in some way introducing neither holes, tears, or self-intersections. [332, §2.2] The physical process is intuitively described as unfurling, unfolding, or *unraveling*. In particular instances, the process is a sort of




Figure 119: Neighborhood graph (dashed) with dimensionless EDM subgraph completion (solid) superimposed (but not covering dashed). Local view of a few dense samples \circ from relative interior of some arbitrary Euclidean manifold whose affine dimension appears two-dimensional in this neighborhood. All line segments measure absolute distance. Dashed line segments help visually locate nearest neighbors; suggesting, best number of nearest neighbors can be greater than value of embedding dimension after topological transformation. (*confer* [187, §2]) Solid line segments represent completion of EDM subgraph from available distance data for an arbitrarily chosen sample and its nearest neighbors. Each distance from EDM subgraph becomes distance-square in corresponding EDM submatrix.

flattening by stretching until taut (but not by crushing); *e.g.*, unfurling a three-dimensional Euclidean body resembling a billowy national flag reduces that manifold's affine dimension to $r=2$.

Data input to the proposed process originates from distances between neighboring relatively dense samples of a given manifold. Figure 119 realizes a densely sampled neighborhood; called, *neighborhood graph*. Essentially, the algorithmic process preserves local isometry between *nearest neighbors* allowing distant neighbors to excuse expansively by “maximizing variance” (Figure 5). The common number of nearest neighbors to each sample is a data-dependent algorithmic parameter whose minimum value connects the graph. The *dimensionless EDM subgraph* between each sample and its nearest neighbors is completed from available data and included as input; one such EDM subgraph completion is drawn superimposed upon the neighborhood graph in Figure 119.^{6.3} The consequent dimensionless EDM graph comprising all the subgraphs is incomplete, in general, because the neighbor number is relatively small; incomplete even though it is a superset of the neighborhood graph. Remaining distances (those not graphed at all) are squared then made variables within the algorithm; it is this variability that admits unfurling.

To demonstrate, consider untying the *trefoil knot* drawn in Figure 120(a). A corresponding Euclidean distance matrix $D = [d_{ij}, i, j = 1 \dots N]$ employing only 2 nearest neighbors is banded having the incomplete form

$$D = \begin{bmatrix} 0 & \check{d}_{12} & \check{d}_{13} & ? & \cdots & ? & \check{d}_{1,N-1} & \check{d}_{1N} \\ \check{d}_{12} & 0 & \check{d}_{23} & \check{d}_{24} & \ddots & ? & ? & \check{d}_{2N} \\ \check{d}_{13} & \check{d}_{23} & 0 & \check{d}_{34} & \ddots & ? & ? & ? \\ ? & \check{d}_{24} & \check{d}_{34} & 0 & \ddots & \ddots & ? & ? \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & ? \\ ? & ? & ? & \ddots & \ddots & 0 & \check{d}_{N-2,N-1} & \check{d}_{N-2,N} \\ \check{d}_{1,N-1} & ? & ? & ? & \ddots & \check{d}_{N-2,N-1} & 0 & \check{d}_{N-1,N} \\ \check{d}_{1N} & \check{d}_{2N} & ? & ? & ? & \check{d}_{N-2,N} & \check{d}_{N-1,N} & 0 \end{bmatrix} \quad (1100)$$

where \check{d}_{ij} denotes a given fixed distance-square. The unfurling algorithm can be expressed as an optimization problem; constrained distance-square

^{6.3}Local reconstruction of point position from the EDM submatrix corresponding to a complete dimensionless EDM subgraph is unique to within an isometry (§5.6, §5.12).




Figure 120: (a) *Trefoil knot* in \mathbb{R}^3 from Weinberger & Saul [332]. (b) Topological transformation algorithm employing 4 nearest neighbors and $N = 539$ samples reduces affine dimension of knot to $r = 2$. Choosing instead 2 nearest neighbors would make this embedding more circular.

maximization:

$$\begin{aligned}
& \underset{D}{\text{maximize}} && \langle -V, D \rangle \\
& \text{subject to} && \langle D, e_i e_j^T + e_j e_i^T \rangle_{\frac{1}{2}} = \check{d}_{ij} \quad \forall (i, j) \in \mathcal{I} \\
& && \text{rank}(VDV) = 2 \\
& && D \in \text{EDM}^N
\end{aligned} \tag{1101}$$

where $e_i \in \mathbb{R}^N$ is the i^{th} member of the standard basis, where set \mathcal{I} indexes the given distance-square data like that in (1100), where $V \in \mathbb{R}^{N \times N}$ is the geometric centering matrix (§B.4.1), and where

$$\langle -V, D \rangle = \text{tr}(-VDV) = 2 \text{tr} G = \frac{1}{N} \sum_{i,j} d_{ij} \tag{833}$$

where G is the Gram matrix producing D assuming $G\mathbf{1} = \mathbf{0}$.

If the (rank) constraint on affine dimension is ignored, then problem (1101) becomes convex, a corresponding solution D^* can be found, and a nearest rank-2 solution is then had by ordered eigen decomposition of $-VD^*V$ followed by *spectral projection* (§7.1.3) on $\begin{bmatrix} \mathbb{R}_+^2 \\ \mathbf{0} \end{bmatrix} \subset \mathbb{R}^N$. This two-step process is necessarily suboptimal. Yet because the decomposition for the trefoil knot reveals only two dominant eigenvalues, the spectral projection is nearly benign. Such a reconstruction of point position (§5.12) utilizing 4 nearest neighbors is drawn in Figure 120(b); a low-dimensional embedding of the trefoil knot.

This problem (1101) can, of course, be written equivalently in terms of Gram matrix G , facilitated by (839); *videlicet*, for Φ_{ij} as in (806)

$$\begin{aligned}
& \underset{G \in \mathbb{S}_c^N}{\text{maximize}} && \langle I, G \rangle \\
& \text{subject to} && \langle G, \Phi_{ij} \rangle = \check{d}_{ij} \quad \forall (i, j) \in \mathcal{I} \\
& && \text{rank } G = 2 \\
& && G \succeq 0
\end{aligned} \tag{1102}$$

The advantage to converting EDM to Gram is: Gram matrix G is a bridge between point list X and EDM D ; constraints on any or all of these three variables may now be introduced. (Example 5.4.2.2.4) Confinement to the geometric center subspace \mathbb{S}_c^N (implicit constraint $G\mathbf{1} = \mathbf{0}$) keeps G independent of its translation-invariant subspace $\mathbb{S}_c^{N\perp}$ (§5.5.1.1, Figure 127) so as not to become numerically unbounded.




Figure 121: *Trefoil ribbon*; courtesy, Kilian Weinberger. Same topological transformation algorithm with 5 nearest neighbors and $N = 1617$ samples.

The problem dual to maximum variance unfolding problem (1102) (less the Gram rank constraint) has been called the *fastest mixing Markov process*. That dual has simple interpretations in graph and circuit theory and in mechanical and thermal systems, explored in [302], and direct application to quick calculation of *PageRank* by search engines [203, §4]. Optimal Gram rank turns out to be tightly bounded above by minimum multiplicity of the second smallest eigenvalue of a dual optimal variable. \square

6.5 EDM definition in $\mathbf{1}\mathbf{1}^T$

Any EDM D corresponding to affine dimension r has representation

$$\mathbf{D}(V_{\mathcal{X}}) \triangleq \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)\mathbf{1}^T + \mathbf{1}\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)^T - 2V_{\mathcal{X}}V_{\mathcal{X}}^T \in \text{EDM}^N \quad (1103)$$

where $\mathcal{R}(V_{\mathcal{X}} \in \mathbb{R}^{N \times r}) \subseteq \mathcal{N}(\mathbf{1}^T) = \mathbf{1}^\perp$,

$$V_{\mathcal{X}}^T V_{\mathcal{X}} = \delta^2(V_{\mathcal{X}}^T V_{\mathcal{X}}) \quad \text{and} \quad V_{\mathcal{X}} \text{ is full-rank with orthogonal columns.} \quad (1104)$$

Equation (1103) is simply the standard EDM definition (808) with a centered list X as in (890); Gram matrix $X^T X$ has been replaced with the subcompact singular value decomposition (§A.6.2)^{6.4}

$$V_{\mathcal{X}}V_{\mathcal{X}}^T \equiv V^T X^T X V \in \mathbb{S}_c^N \cap \mathbb{S}_+^N \quad (1105)$$

This means: inner product $V_{\mathcal{X}}^T V_{\mathcal{X}}$ is an $r \times r$ diagonal matrix Σ of nonzero singular values.

Vector $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)$ may be decomposed into complementary parts by projecting it on orthogonal subspaces $\mathbf{1}^\perp$ and $\mathcal{R}(\mathbf{1})$: namely,

$$P_{\mathbf{1}^\perp}(\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)) = V\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \quad (1106)$$

$$P_{\mathbf{1}}(\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)) = \frac{1}{N}\mathbf{1}\mathbf{1}^T\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \quad (1107)$$

Of course

$$\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) = V\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) + \frac{1}{N}\mathbf{1}\mathbf{1}^T\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \quad (1108)$$

^{6.4}Subcompact SVD: $V_{\mathcal{X}}V_{\mathcal{X}}^T \triangleq Q\sqrt{\Sigma}\sqrt{\Sigma}Q^T \equiv V^T X^T X V$. So $V_{\mathcal{X}}^T$ is not necessarily XV (§5.5.1.0.1), although affine dimension $r = \text{rank}(V_{\mathcal{X}}^T) = \text{rank}(XV)$. (947)

by (831). Substituting this into EDM definition (1103), we get the Hayden, Wells, Liu, & Tarazaga EDM formula [164, §2]

$$\mathbf{D}(V_{\mathcal{X}}, y) \triangleq y\mathbf{1}^T + \mathbf{1}y^T + \frac{\lambda}{N}\mathbf{1}\mathbf{1}^T - 2V_{\mathcal{X}}V_{\mathcal{X}}^T \in \text{EDM}^N \quad (1109)$$

where

$$\lambda \triangleq 2\|V_{\mathcal{X}}\|_{\mathbb{F}}^2 = \mathbf{1}^T \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \mathbf{1} \quad \text{and} \quad y \triangleq \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) - \frac{\lambda}{2N}\mathbf{1} = V\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \quad (1110)$$

and $y = \mathbf{0}$ if and only if $\mathbf{1}$ is an eigenvector of EDM D . Scalar λ becomes an eigenvalue when corresponding eigenvector $\mathbf{1}$ exists.^{6.5}

Then the particular dyad sum from (1109)

$$y\mathbf{1}^T + \mathbf{1}y^T + \frac{\lambda}{N}\mathbf{1}\mathbf{1}^T \in \mathbb{S}_c^{N\perp} \quad (1111)$$

must belong to the orthogonal complement of the geometric center subspace (p.675), whereas $V_{\mathcal{X}}V_{\mathcal{X}}^T \in \mathbb{S}_c^N \cap \mathbb{S}_+^N$ (1105) belongs to the positive semidefinite cone in the geometric center subspace.

Proof. We validate eigenvector $\mathbf{1}$ and eigenvalue λ .

(\Rightarrow) Suppose $\mathbf{1}$ is an eigenvector of EDM D . Then because

$$V_{\mathcal{X}}^T \mathbf{1} = \mathbf{0} \quad (1112)$$

it follows

$$\begin{aligned} D\mathbf{1} &= \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)\mathbf{1}^T\mathbf{1} + \mathbf{1}\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)^T\mathbf{1} = N\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) + \|V_{\mathcal{X}}\|_{\mathbb{F}}^2\mathbf{1} \\ &\Rightarrow \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \propto \mathbf{1} \end{aligned} \quad (1113)$$

For some $\kappa \in \mathbb{R}_+$

$$\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)^T\mathbf{1} = N\kappa = \text{tr}(V_{\mathcal{X}}^T V_{\mathcal{X}}) = \|V_{\mathcal{X}}\|_{\mathbb{F}}^2 \Rightarrow \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) = \frac{1}{N}\|V_{\mathcal{X}}\|_{\mathbb{F}}^2\mathbf{1} \quad (1114)$$

so $y = \mathbf{0}$.

(\Leftarrow) Now suppose $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) = \frac{\lambda}{2N}\mathbf{1}$; *id est*, $y = \mathbf{0}$. Then

$$D = \frac{\lambda}{N}\mathbf{1}\mathbf{1}^T - 2V_{\mathcal{X}}V_{\mathcal{X}}^T \in \text{EDM}^N \quad (1115)$$

$\mathbf{1}$ is an eigenvector with corresponding eigenvalue λ . \blacklozenge

^{6.5} *e.g.*, when $X = I$ in EDM definition (808).




Figure 122: Example of $V_{\mathcal{X}}$ selection to make an EDM corresponding to cardinality $N=3$ and affine dimension $r=1$; $V_{\mathcal{X}}$ is a vector in nullspace $\mathcal{N}(\mathbf{1}^T) \subset \mathbb{R}^3$. Nullspace of $\mathbf{1}^T$ is hyperplane in \mathbb{R}^3 (drawn truncated) having normal $\mathbf{1}$. Vector $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)$ may or may not be in plane spanned by $\{\mathbf{1}, V_{\mathcal{X}}\}$, but belongs to nonnegative orthant which is strictly supported by $\mathcal{N}(\mathbf{1}^T)$.

6.5.1 Range of EDM D

From §B.1.1 pertaining to linear independence of dyad sums: If the transpose halves of all the dyads in the sum (1103)^{6.6} make a linearly independent set, then the nontranspose halves constitute a basis for the range of EDM D . Saying this mathematically: For $D \in \mathbb{EDM}^N$

$$\begin{aligned} \mathcal{R}(D) = \mathcal{R}([\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \quad \mathbf{1} \quad V_{\mathcal{X}}]) &\Leftrightarrow \text{rank}([\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \quad \mathbf{1} \quad V_{\mathcal{X}}]) = 2 + r \\ \mathcal{R}(D) = \mathcal{R}([\mathbf{1} \quad V_{\mathcal{X}}]) &\Leftrightarrow \text{otherwise} \end{aligned} \quad (1116)$$

To prove this, we need that condition under which the rank equality is satisfied: We know $\mathcal{R}(V_{\mathcal{X}}) \perp \mathbf{1}$, but what is the relative geometric orientation of $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)$? $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \succeq 0$ because $V_{\mathcal{X}}V_{\mathcal{X}}^T \succeq 0$, and $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \propto \mathbf{1}$ remains possible (1113); this means $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) \notin \mathcal{N}(\mathbf{1}^T)$ simply because it has no negative entries. (Figure 122) If the projection of $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)$ on $\mathcal{N}(\mathbf{1}^T)$ does not belong to $\mathcal{R}(V_{\mathcal{X}})$, then that is a necessary and sufficient condition for linear independence (l.i.) of $\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)$ with respect to $\mathcal{R}([\mathbf{1} \quad V_{\mathcal{X}}])$; *id est*,

$$\begin{aligned} V\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) &\neq V_{\mathcal{X}}a \quad \text{for any } a \in \mathbb{R}^r \\ (I - \frac{1}{N}\mathbf{1}\mathbf{1}^T)\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) &\neq V_{\mathcal{X}}a \\ \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) - \frac{1}{N}\|V_{\mathcal{X}}\|_{\text{F}}^2\mathbf{1} &\neq V_{\mathcal{X}}a \\ \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T) - \frac{\lambda}{2N}\mathbf{1} = y &\neq V_{\mathcal{X}}a \Leftrightarrow \{\mathbf{1}, \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T), V_{\mathcal{X}}\} \text{ is l.i.} \end{aligned} \quad (1117)$$

On the other hand when this condition is violated (when (1110) $y = V_{\mathcal{X}}a_p$ for some particular $a \in \mathbb{R}^r$), then from (1109) we have

$$\begin{aligned} \mathcal{R}(D = y\mathbf{1}^T + \mathbf{1}y^T + \frac{\lambda}{N}\mathbf{1}\mathbf{1}^T - 2V_{\mathcal{X}}V_{\mathcal{X}}^T) &= \mathcal{R}((V_{\mathcal{X}}a_p + \frac{\lambda}{N}\mathbf{1})\mathbf{1}^T + (\mathbf{1}a_p^T - 2V_{\mathcal{X}})V_{\mathcal{X}}^T) \\ &= \mathcal{R}([V_{\mathcal{X}}a_p + \frac{\lambda}{N}\mathbf{1} \quad \mathbf{1}a_p^T - 2V_{\mathcal{X}}]) \\ &= \mathcal{R}([\mathbf{1} \quad V_{\mathcal{X}}]) \end{aligned} \quad (1118)$$

An example of such a violation is (1115) where, in particular, $a_p = \mathbf{0}$. \blacklozenge

Then a statement parallel to (1116) is, for $D \in \mathbb{EDM}^N$ (Theorem 5.7.3.0.1)

$$\begin{aligned} \text{rank}(D) = r + 2 &\Leftrightarrow y \notin \mathcal{R}(V_{\mathcal{X}}) \quad (\Leftrightarrow \mathbf{1}^T D \mathbf{1} = 0) \\ \text{rank}(D) = r + 1 &\Leftrightarrow y \in \mathcal{R}(V_{\mathcal{X}}) \quad (\Leftrightarrow \mathbf{1}^T D \mathbf{1} \neq 0) \end{aligned} \quad (1119)$$

^{6.6}Identifying columns $V_{\mathcal{X}} \triangleq [v_1 \cdots v_r]$, then $V_{\mathcal{X}}V_{\mathcal{X}}^T = \sum_i v_i v_i^T$ is also a sum of dyads.

6.5.2 Boundary constituents of EDM cone

Expression (1103) has utility in forming the set of all EDMs corresponding to affine dimension r :

$$\begin{aligned} & \{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) = r\} \\ &= \{\mathbf{D}(V_{\mathcal{X}}) \mid V_{\mathcal{X}} \in \mathbb{R}^{N \times r}, \text{rank } V_{\mathcal{X}} = r, V_{\mathcal{X}}^T V_{\mathcal{X}} = \delta^2(V_{\mathcal{X}}^T V_{\mathcal{X}}), \mathcal{R}(V_{\mathcal{X}}) \subseteq \mathcal{N}(\mathbf{1}^T)\} \end{aligned} \quad (1120)$$

whereas $\{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) \leq r\}$ is the closure of this same set;

$$\{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) \leq r\} = \overline{\{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) = r\}} \quad (1121)$$

For example,

$$\begin{aligned} \text{rel } \partial \mathbf{EDM}^N &= \{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) < N-1\} \\ &= \bigcup_{r=0}^{N-2} \{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) = r\} \end{aligned} \quad (1122)$$

None of these are necessarily convex sets, although

$$\begin{aligned} \mathbf{EDM}^N &= \bigcup_{r=0}^{N-1} \{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) = r\} \\ &= \overline{\{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) = N-1\}} \\ \text{rel int } \mathbf{EDM}^N &= \{D \in \mathbf{EDM}^N \mid \text{rank}(VDV) = N-1\} \end{aligned} \quad (1123)$$

are pointed convex cones.

When cardinality $N=3$ and affine dimension $r=2$, for example, the relative interior $\text{rel int } \mathbf{EDM}^3$ is realized via (1120). (§6.6)

When $N=3$ and $r=1$, the relative boundary of the EDM cone $\text{dvec rel } \partial \mathbf{EDM}^3$ is realized in isomorphic \mathbb{R}^3 as in Figure 117(d). This figure could be constructed via (1121) by spiraling vector $V_{\mathcal{X}}$ tightly about the origin in $\mathcal{N}(\mathbf{1}^T)$; as can be imagined with aid of Figure 122. Vectors close to the origin in $\mathcal{N}(\mathbf{1}^T)$ are correspondingly close to the origin in \mathbf{EDM}^N . As vector $V_{\mathcal{X}}$ orbits the origin in $\mathcal{N}(\mathbf{1}^T)$, the corresponding EDM orbits the axis of revolution while remaining on the boundary of the circular cone $\text{dvec rel } \partial \mathbf{EDM}^3$. (Figure 123)




Figure 123: (a) Vector $V_{\mathcal{X}}$ from Figure 122 spirals in $\mathcal{N}(\mathbf{1}^T) \subset \mathbb{R}^3$ decaying toward origin. (Spiral is two-dimensional in vector space \mathbb{R}^3 .) (b) Corresponding trajectory $\mathbf{D}(V_{\mathcal{X}})$ on EDM cone relative boundary creates a vortex also decaying toward origin. There are two complete orbits on EDM cone boundary about axis of revolution for every single revolution of $V_{\mathcal{X}}$ about origin. (Vortex is three-dimensional in isometrically isomorphic \mathbb{R}^3 .)

6.5.3 Faces of EDM cone

Like the positive semidefinite cone, EDM cone faces are EDM cones.

6.5.3.0.1 Exercise. *Isomorphic faces.*

Prove that in high cardinality N , any set of EDMs made via (1120) or (1121) with particular affine dimension r is isomorphic with any set admitting the same affine dimension but made in lower cardinality. ▼

6.5.3.1 Smallest face containing an EDM

Now suppose we are given a particular EDM $\mathbf{D}(V_{\mathcal{X}_p}) \in \mathbb{EDM}^N$ corresponding to affine dimension r and parametrized by $V_{\mathcal{X}_p}$ in (1103). The EDM cone's smallest face that contains $\mathbf{D}(V_{\mathcal{X}_p})$ is

$$\begin{aligned} & \mathcal{F}(\mathbb{EDM}^N \ni \mathbf{D}(V_{\mathcal{X}_p})) \\ &= \overline{\{\mathbf{D}(V_{\mathcal{X}}) \mid V_{\mathcal{X}} \in \mathbb{R}^{N \times r}, \text{rank } V_{\mathcal{X}} = r, V_{\mathcal{X}}^T V_{\mathcal{X}} = \delta^2(V_{\mathcal{X}}^T V_{\mathcal{X}}), \mathcal{R}(V_{\mathcal{X}}) \subseteq \mathcal{R}(V_{\mathcal{X}_p})\}} \\ & \simeq \mathbb{EDM}^{r+1} \end{aligned} \quad (1124)$$

which is isomorphic^{6.7} with the convex cone \mathbb{EDM}^{r+1} , hence of dimension

$$\dim \mathcal{F}(\mathbb{EDM}^N \ni \mathbf{D}(V_{\mathcal{X}_p})) = (r+1)r/2 \quad (1125)$$

in isomorphic $\mathbb{R}^{N(N-1)/2}$. Not all dimensions are represented; *e.g.*, the EDM cone has no two-dimensional faces.

When cardinality $N=4$ and affine dimension $r=2$ so that $\mathcal{R}(V_{\mathcal{X}_p})$ is any two-dimensional subspace of three-dimensional $\mathcal{N}(\mathbf{1}^T)$ in \mathbb{R}^4 , for example, then the corresponding face of \mathbb{EDM}^4 is isometrically isomorphic with: (1121)

$$\mathbb{EDM}^3 = \{D \in \mathbb{EDM}^3 \mid \text{rank}(VDV) \leq 2\} \simeq \mathcal{F}(\mathbb{EDM}^4 \ni \mathbf{D}(V_{\mathcal{X}_p})) \quad (1126)$$

Each two-dimensional subspace of $\mathcal{N}(\mathbf{1}^T)$ corresponds to another three-dimensional face.

Because each and every principal submatrix of an EDM in \mathbb{EDM}^N (§5.14.3) is another EDM [207, §4.1], for example, then each principal submatrix belongs to a particular face of \mathbb{EDM}^N .

^{6.7}The fact that the smallest face is isomorphic with another EDM cone (perhaps smaller than \mathbb{EDM}^N) is implicit in [164, §2].

6.5.3.2 Extreme directions of EDM cone

In particular, extreme directions (§2.8.1) of \mathbf{EDM}^N correspond to affine dimension $r = 1$ and are simply represented: for any particular cardinality $N \geq 2$ (§2.8.2) and each and every nonzero vector z in $\mathcal{N}(\mathbf{1}^T)$

$$\begin{aligned}\Gamma &\triangleq (z \circ z)\mathbf{1}^T + \mathbf{1}(z \circ z)^T - 2zz^T \in \mathbf{EDM}^N \\ &= \delta(zz^T)\mathbf{1}^T + \mathbf{1}\delta(zz^T)^T - 2zz^T\end{aligned}\quad (1127)$$

is an extreme direction corresponding to a one-dimensional face of the EDM cone \mathbf{EDM}^N that is a ray in isomorphic subspace $\mathbb{R}^{N(N-1)/2}$.

Proving this would exercise the fundamental definition (169) of extreme direction. Here is a sketch: Any EDM may be represented

$$\mathbf{D}(V_{\mathcal{X}}) \triangleq \delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)\mathbf{1}^T + \mathbf{1}\delta(V_{\mathcal{X}}V_{\mathcal{X}}^T)^T - 2V_{\mathcal{X}}V_{\mathcal{X}}^T \in \mathbf{EDM}^N \quad (1103)$$

where matrix $V_{\mathcal{X}}$ (1104) has orthogonal columns. For the same reason (1380) that zz^T is an extreme direction of the positive semidefinite cone (§2.9.2.4) for any particular nonzero vector z , there is no conic combination of distinct EDMs (each conically independent of Γ (§2.10)) equal to Γ . ■

6.5.3.2.1 Example. Biorthogonal expansion of an EDM.

(confer §2.13.7.1.1) When matrix D belongs to the EDM cone, nonnegative coordinates for biorthogonal expansion are the eigenvalues $\lambda \in \mathbb{R}^N$ of $-VDV_{\frac{1}{2}}$: For any $D \in \mathbb{S}_h^N$ it holds

$$D = \delta(-VDV_{\frac{1}{2}})\mathbf{1}^T + \mathbf{1}\delta(-VDV_{\frac{1}{2}})^T - 2(-VDV_{\frac{1}{2}}) \quad (914)$$

By diagonalization $-VDV_{\frac{1}{2}} \triangleq Q\Lambda Q^T \in \mathbb{S}_c^N$ (§A.5.2) we may write

$$\begin{aligned}D &= \delta\left(\sum_{i=1}^N \lambda_i q_i q_i^T\right)\mathbf{1}^T + \mathbf{1}\delta\left(\sum_{i=1}^N \lambda_i q_i q_i^T\right)^T - 2\sum_{i=1}^N \lambda_i q_i q_i^T \\ &= \sum_{i=1}^N \lambda_i (\delta(q_i q_i^T)\mathbf{1}^T + \mathbf{1}\delta(q_i q_i^T)^T - 2q_i q_i^T)\end{aligned}\quad (1128)$$

where q_i is the i^{th} eigenvector of $-VDV_{\frac{1}{2}}$ arranged columnar in orthogonal matrix

$$Q = [q_1 \ q_2 \ \cdots \ q_N] \in \mathbb{R}^{N \times N} \quad (360)$$

and where $\{\delta(q_i q_i^T) \mathbf{1}^T + \mathbf{1} \delta(q_i q_i^T)^T - 2q_i q_i^T, i=1 \dots N\}$ are extreme directions of some pointed polyhedral cone $\mathcal{K} \subset \mathbb{S}_h^N$ and extreme directions of \mathbb{EDM}^N . Invertibility of (1128)

$$\begin{aligned} -VDV\frac{1}{2} &= -V \sum_{i=1}^N \lambda_i (\delta(q_i q_i^T) \mathbf{1}^T + \mathbf{1} \delta(q_i q_i^T)^T - 2q_i q_i^T) V\frac{1}{2} \\ &= \sum_{i=1}^N \lambda_i q_i q_i^T \end{aligned} \quad (1129)$$

implies linear independence of those extreme directions. Then the biorthogonal expansion is expressed

$$\text{dvec } D = YY^\dagger \text{dvec } D = Y \lambda (-VDV\frac{1}{2}) \quad (1130)$$

where

$$Y \triangleq [\text{dvec}(\delta(q_i q_i^T) \mathbf{1}^T + \mathbf{1} \delta(q_i q_i^T)^T - 2q_i q_i^T), i=1 \dots N] \in \mathbb{R}^{N(N-1)/2 \times N} \quad (1131)$$

When D belongs to the EDM cone in the subspace of symmetric hollow matrices, unique coordinates $Y^\dagger \text{dvec } D$ for this biorthogonal expansion must be the nonnegative eigenvalues λ of $-VDV\frac{1}{2}$. This means D simultaneously belongs to the EDM cone and to the pointed polyhedral cone $\text{dvec } \mathcal{K} = \text{cone}(Y)$. \square

6.5.3.3 Open question

Result (1125) is analogous to that for the positive semidefinite cone (205), although the question remains open whether all faces of \mathbb{EDM}^N (whose dimension is less than dimension of the cone) are exposed like they are for the positive semidefinite cone.^{6.8} (§2.9.2.3) [308]

6.6 Correspondence to PSD cone \mathbb{S}_+^{N-1}

Hayden, Wells, Liu, & Tarazaga [164, §2] assert one-to-one correspondence of EDMs with positive semidefinite matrices in the symmetric subspace. Because $\text{rank}(VDV) \leq N-1$ (§5.7.1.1), that PSD cone corresponding to

^{6.8}Elementary example of a face not exposed is given in Figure 29 and Figure 38.

the EDM cone can only be \mathbb{S}_+^{N-1} . [7, §18.2.1] To clearly demonstrate this correspondence, we invoke inner-product form EDM definition

$$\begin{aligned} \mathbf{D}(\Phi) &\triangleq \begin{bmatrix} 0 \\ \delta(\Phi) \end{bmatrix} \mathbf{1}^T + \mathbf{1} \begin{bmatrix} 0 & \delta(\Phi)^T \end{bmatrix} - 2 \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \Phi \end{bmatrix} \in \mathbb{EDM}^N \\ &\Leftrightarrow \\ &\Phi \succeq 0 \end{aligned} \quad (932)$$

Then the EDM cone may be expressed

$$\mathbb{EDM}^N = \{\mathbf{D}(\Phi) \mid \Phi \in \mathbb{S}_+^{N-1}\} \quad (1132)$$

Hayden & Wells' assertion can therefore be equivalently stated in terms of an inner-product form EDM operator

$$\mathbf{D}(\mathbb{S}_+^{N-1}) = \mathbb{EDM}^N \quad (934)$$

$$\mathbf{V}_{\mathcal{N}}(\mathbb{EDM}^N) = \mathbb{S}_+^{N-1} \quad (935)$$

identity (935) holding because $\mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^T)$ (815), linear functions $\mathbf{D}(\Phi)$ and $\mathbf{V}_{\mathcal{N}}(D) = -V_{\mathcal{N}}^T D V_{\mathcal{N}}$ (§5.6.2.1) being mutually inverse.

In terms of affine dimension r , Hayden & Wells claim particular correspondence between PSD and EDM cones:

$r = N - 1$: Symmetric hollow matrices $-D$ positive definite on $\mathcal{N}(\mathbf{1}^T)$ correspond to points relatively interior to the EDM cone.

$r < N - 1$: Symmetric hollow matrices $-D$ positive semidefinite on $\mathcal{N}(\mathbf{1}^T)$, where $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ has at least one 0 eigenvalue, correspond to points on the relative boundary of the EDM cone.

$r = 1$: Symmetric hollow nonnegative matrices rank-one on $\mathcal{N}(\mathbf{1}^T)$ correspond to extreme directions (1127) of the EDM cone; *id est*, for some nonzero vector u (§A.3.1.0.7)

$$\left. \begin{array}{l} \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} = 1 \\ D \in \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N} \end{array} \right\} \Leftrightarrow \begin{array}{l} D \in \mathbb{EDM}^N \\ D \text{ is an extreme direction} \end{array} \Leftrightarrow \left\{ \begin{array}{l} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \equiv uu^T \\ D \in \mathbb{S}_h^N \end{array} \right. \quad (1133)$$

6.6.0.0.1 Proof. Case $r = 1$ is easily proved: From the nonnegativity development in §5.8.1, extreme direction (1127), and Schoenberg criterion (827), we need show only sufficiency; *id est*, prove

$$\left. \begin{array}{l} \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} = 1 \\ D \in \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N} \end{array} \right\} \Rightarrow \begin{array}{l} D \in \mathbb{EDM}^N \\ D \text{ is an extreme direction} \end{array}$$

Any symmetric matrix D satisfying the rank condition must have the form, for $z, q \in \mathbb{R}^N$ and nonzero $z \in \mathcal{N}(\mathbf{1}^T)$,

$$D = \pm(\mathbf{1}q^T + q\mathbf{1}^T - 2zz^T) \quad (1134)$$

because (§5.6.2.1, confer §E.7.2.0.2)

$$\mathcal{N}(\mathbf{V}_{\mathcal{N}}(D)) = \{\mathbf{1}q^T + q\mathbf{1}^T \mid q \in \mathbb{R}^N\} \subseteq \mathbb{S}^N \quad (1135)$$

Hollowness demands $q = \delta(zz^T)$ while nonnegativity demands choice of positive sign in (1134). Matrix D thus takes the form of an extreme direction (1127) of the EDM cone. \blacklozenge

The foregoing proof is not extensible in rank: An EDM with corresponding affine dimension r has the general form, for $\{z_i \in \mathcal{N}(\mathbf{1}^T), i = 1 \dots r\}$ an independent set,

$$D = \mathbf{1}\delta\left(\sum_{i=1}^r z_i z_i^T\right)^T + \delta\left(\sum_{i=1}^r z_i z_i^T\right)\mathbf{1}^T - 2\sum_{i=1}^r z_i z_i^T \in \mathbb{EDM}^N \quad (1136)$$

The EDM so defined relies principally on the sum $\sum z_i z_i^T$ having positive summand coefficients ($\Leftrightarrow -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0$)^{6.9}. Then it is easy to find a sum incorporating negative coefficients while meeting rank, nonnegativity, and symmetric hollowness conditions but not positive semidefiniteness on subspace $\mathcal{R}(V_{\mathcal{N}})$; *e.g.*, from page 422,

$$-V \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 5 \\ 1 & 5 & 0 \end{bmatrix} V \frac{1}{2} = z_1 z_1^T - z_2 z_2^T \quad (1137)$$

^{6.9} (\Leftrightarrow) For $a_i \in \mathbb{R}^{N-1}$, let $z_i = V_{\mathcal{N}}^{\dagger T} a_i$.

6.6.0.0.2 Example. *Extreme rays versus rays on the boundary.*

The EDM $D = \begin{bmatrix} 0 & 1 & 4 \\ 1 & 0 & 1 \\ 4 & 1 & 0 \end{bmatrix}$ is an extreme direction of \mathbb{EDM}^3 where

$u = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ in (1133). Because $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ has eigenvalues $\{0, 5\}$, the ray whose direction is D also lies on the relative boundary of \mathbb{EDM}^3 .

In exception, EDM $D = \kappa \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, for any particular $\kappa > 0$, is an extreme direction of \mathbb{EDM}^2 but $-V_{\mathcal{N}}^T D V_{\mathcal{N}}$ has only one eigenvalue: $\{\kappa\}$. Because \mathbb{EDM}^2 is a ray whose relative boundary (§2.6.1.3.1) is the origin, this conventional boundary does not include D which belongs to the relative interior in this dimension. (§2.7.0.0.1) \square

6.6.1 Gram-form correspondence to \mathbb{S}_+^{N-1}

With respect to $\mathbf{D}(G) = \delta(G)\mathbf{1}^T + \mathbf{1}\delta(G)^T - 2G$ (820) the linear Gram-form EDM operator, results in §5.6.1 provide [1, §2.6]

$$\mathbb{EDM}^N = \mathbf{D}(\mathbf{V}(\mathbb{EDM}^N)) \equiv \mathbf{D}(V_{\mathcal{N}}\mathbb{S}_+^{N-1}V_{\mathcal{N}}^T) \quad (1138)$$

$$V_{\mathcal{N}}\mathbb{S}_+^{N-1}V_{\mathcal{N}}^T \equiv \mathbf{V}(\mathbf{D}(V_{\mathcal{N}}\mathbb{S}_+^{N-1}V_{\mathcal{N}}^T)) = \mathbf{V}(\mathbb{EDM}^N) \stackrel{\Delta}{=} -V\mathbb{EDM}^N V \frac{1}{2} = \mathbb{S}_c^N \cap \mathbb{S}_+^N \quad (1139)$$

a one-to-one correspondence between \mathbb{EDM}^N and \mathbb{S}_+^{N-1} .

6.6.2 EDM cone by ellipotope


(confer §5.10.1) Defining the ellipotope parametrized by scalar $t > 0$

$$\mathcal{E}_t^N = \mathbb{S}_+^N \cap \{\Phi \in \mathbb{S}^N \mid \delta(\Phi) = t\mathbf{1}\} \quad (1012)$$

then following Alfakih [8] we have

$$\mathbb{EDM}^N = \overline{\text{cone}\{\mathbf{1}\mathbf{1}^T - \mathcal{E}_1^N\}} = \overline{\{t(\mathbf{1}\mathbf{1}^T - \mathcal{E}_1^N) \mid t \geq 0\}} \quad (1140)$$

Identification $\mathcal{E}^N = \mathcal{E}_1^N$ equates the standard ellipotope (§5.9.1.0.1, Figure 109) to our parametrized ellipotope.



$$\mathbb{EDM}^N = \overline{\text{cone}\{\mathbf{1}\mathbf{1}^T - \mathcal{E}^N\}} = \overline{\{t(\mathbf{1}\mathbf{1}^T - \mathcal{E}^N) \mid t \geq 0\}} \quad (1140)$$

Figure 124: Three views of translated negated ellipsope $\mathbf{1}\mathbf{1}^T - \mathcal{E}_1^3$ (confer Figure 109) shrouded by truncated EDM cone. Fractal on EDM cone relative boundary is numerical artifact belonging to intersection with ellipsope relative boundary. The fractal is trying to convey existence of a neighborhood about the origin where the translated ellipsope boundary and EDM cone boundary intersect.

6.6.2.0.1 Expository. Define $\mathcal{T}_{\mathcal{E}}(\mathbf{1}\mathbf{1}^T)$ to be the *tangent cone* to the ellipsope \mathcal{E} at point $\mathbf{1}\mathbf{1}^T$; *id est*,

$$\mathcal{T}_{\mathcal{E}}(\mathbf{1}\mathbf{1}^T) \triangleq \overline{\{t(\mathcal{E} - \mathbf{1}\mathbf{1}^T) \mid t \geq 0\}} \quad (1141)$$

The normal cone $\mathcal{K}_{\mathcal{E}}^{\perp}(\mathbf{1}\mathbf{1}^T)$ to the ellipsope at $\mathbf{1}\mathbf{1}^T$ is a closed convex cone defined (§E.10.3.2.1, Figure 153)

$$\mathcal{K}_{\mathcal{E}}^{\perp}(\mathbf{1}\mathbf{1}^T) \triangleq \{B \mid \langle B, \Phi - \mathbf{1}\mathbf{1}^T \rangle \leq 0, \Phi \in \mathcal{E}\} \quad (1142)$$

The *polar cone* of any set \mathcal{K} is the closed convex cone (*confer*(270))

$$\mathcal{K}^{\circ} \triangleq \{B \mid \langle B, A \rangle \leq 0, \text{ for all } A \in \mathcal{K}\} \quad (1143)$$

The normal cone is well known to be the polar of the tangent cone,

$$\mathcal{K}_{\mathcal{E}}^{\perp}(\mathbf{1}\mathbf{1}^T) = \mathcal{T}_{\mathcal{E}}(\mathbf{1}\mathbf{1}^T)^{\circ} \quad (1144)$$

and *vice versa*; [177, §A.5.2.4]

$$\mathcal{K}_{\mathcal{E}}^{\perp}(\mathbf{1}\mathbf{1}^T)^{\circ} = \mathcal{T}_{\mathcal{E}}(\mathbf{1}\mathbf{1}^T) \quad (1145)$$

From Deza & Laurent [96, p.535] we have the EDM cone

$$\text{EDM} = -\mathcal{T}_{\mathcal{E}}(\mathbf{1}\mathbf{1}^T) \quad (1146)$$

The polar EDM cone is also expressible in terms of the ellipsope. From (1144) we have

$$\text{EDM}^{\circ} = -\mathcal{K}_{\mathcal{E}}^{\perp}(\mathbf{1}\mathbf{1}^T) \quad (1147)$$

★

In §5.10.1 we proposed the expression for EDM D

$$D = t\mathbf{1}\mathbf{1}^T - \mathfrak{E} \in \text{EDM}^N \quad (1013)$$

where $t \in \mathbb{R}_+$ and \mathfrak{E} belongs to the parametrized ellipsope \mathcal{E}_t^N . We further propose, for any particular $t > 0$

$$\text{EDM}^N = \overline{\text{cone}\{t\mathbf{1}\mathbf{1}^T - \mathcal{E}_t^N\}} \quad (1148)$$

Proof. Pending. ◆

Relationship of the translated negated ellipsope with the EDM cone is illustrated in Figure 124. We speculate

$$\text{EDM}^N = \overline{\lim_{t \rightarrow \infty} t\mathbf{1}\mathbf{1}^T - \mathcal{E}_t^N} \quad (1149)$$

6.7 Vectorization & projection interpretation

In §E.7.2.0.2 we learn: $-VDV$ can be interpreted as orthogonal projection [5, §2] of vectorized $-D \in \mathbb{S}_h^N$ on the subspace of geometrically centered symmetric matrices

$$\begin{aligned} \mathbb{S}_c^N &= \{G \in \mathbb{S}^N \mid G\mathbf{1} = \mathbf{0}\} & (1885) \\ &= \{G \in \mathbb{S}^N \mid \mathcal{N}(G) \supseteq \mathbf{1}\} = \{G \in \mathbb{S}^N \mid \mathcal{R}(G) \subseteq \mathcal{N}(\mathbf{1}^T)\} & (905) \\ &= \{VYV \mid Y \in \mathbb{S}^N\} \subset \mathbb{S}^N & (1886) \\ &\equiv \{V_{\mathcal{N}}AV_{\mathcal{N}}^T \mid A \in \mathbb{S}^{N-1}\} \end{aligned}$$

because elementary auxiliary matrix V is an orthogonal projector (§B.4.1). Yet there is another useful projection interpretation:

Revising the fundamental matrix criterion for membership to the EDM cone (803),^{6.10}

$$\left. \begin{aligned} \langle zz^T, -D \rangle \geq 0 \quad \forall zz^T \mid \mathbf{1}\mathbf{1}^T zz^T = \mathbf{0} \\ D \in \mathbb{S}_h^N \end{aligned} \right\} \Leftrightarrow D \in \text{EDM}^N \quad (1150)$$

this is equivalent, of course, to the Schoenberg criterion

$$\left. \begin{aligned} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \\ D \in \mathbb{S}_h^N \end{aligned} \right\} \Leftrightarrow D \in \text{EDM}^N \quad (827)$$

because $\mathcal{N}(\mathbf{1}\mathbf{1}^T) = \mathcal{R}(V_{\mathcal{N}})$. When $D \in \text{EDM}^N$, correspondence (1150) means $-z^T D z$ is proportional to a nonnegative coefficient of orthogonal projection (§E.6.4.2, Figure 126) of $-D$ in isometrically isomorphic $\mathbb{R}^{N(N+1)/2}$ on the range of each and every vectorized (§2.2.2.1) symmetric dyad (§B.1) in the nullspace of $\mathbf{1}\mathbf{1}^T$; *id est*, on each and every member of

$$\begin{aligned} \mathcal{T} &\triangleq \{\text{svec}(zz^T) \mid z \in \mathcal{N}(\mathbf{1}\mathbf{1}^T) = \mathcal{R}(V_{\mathcal{N}})\} \subset \text{svec} \partial \mathbb{S}_+^N \\ &= \{\text{svec}(V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T) \mid v \in \mathbb{R}^{N-1}\} \end{aligned} \quad (1151)$$

whose dimension is

$$\dim \mathcal{T} = N(N-1)/2 \quad (1152)$$

^{6.10} $\mathcal{N}(\mathbf{1}\mathbf{1}^T) = \mathcal{N}(\mathbf{1}^T)$ and $\mathcal{R}(zz^T) = \mathcal{R}(z)$

The set of all symmetric dyads $\{zz^T \mid z \in \mathbb{R}^N\}$ constitute the extreme directions of the positive semidefinite cone (§2.8.1, §2.9) \mathbb{S}_+^N , hence lie on its boundary. Yet only those dyads in $\mathcal{R}(V_{\mathcal{N}})$ are included in the test (1150), thus only a subset \mathcal{T} of all vectorized extreme directions of \mathbb{S}_+^N is observed.

In the particularly simple case $D \in \mathbb{EDM}^2 = \{D \in \mathbb{S}_h^2 \mid d_{12} \geq 0\}$, for example, only one extreme direction of the PSD cone is involved:

$$zz^T = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} \quad (1153)$$

Any nonnegative scaling of vectorized zz^T belongs to the set \mathcal{T} illustrated in Figure 125 and Figure 126.

6.7.1 Face of PSD cone \mathbb{S}_+^N containing V

In any case, set \mathcal{T} (1151) constitutes the vectorized extreme directions of an $N(N-1)/2$ -dimensional face of the PSD cone \mathbb{S}_+^N containing auxiliary matrix V ; a face isomorphic with $\mathbb{S}_+^{N-1} = \mathbb{S}_+^{\text{rank } V}$ (§2.9.2.3).


To show this, we must first find the smallest face that contains auxiliary matrix V and then determine its extreme directions. From (204),

$$\begin{aligned} \mathcal{F}(\mathbb{S}_+^N \ni V) &= \{W \in \mathbb{S}_+^N \mid \mathcal{N}(W) \supseteq \mathcal{N}(V)\} = \{W \in \mathbb{S}_+^N \mid \mathcal{N}(W) \supseteq \mathbf{1}\} \\ &= \{VYV \succeq 0 \mid Y \in \mathbb{S}^N\} \equiv \{V_{\mathcal{N}} B V_{\mathcal{N}}^T \mid B \in \mathbb{S}_+^{N-1}\} \\ &\simeq \mathbb{S}_+^{\text{rank } V} = -V_{\mathcal{N}}^T \mathbb{EDM}^N V_{\mathcal{N}} \end{aligned} \quad (1154)$$

where the equivalence \equiv is from §5.6.1 while isomorphic equality \simeq with transformed EDM cone is from (935). Projector V belongs to $\mathcal{F}(\mathbb{S}_+^N \ni V)$ because $V_{\mathcal{N}} V_{\mathcal{N}}^\dagger V_{\mathcal{N}}^{\dagger T} V_{\mathcal{N}}^T = V$. (§B.4.3) Each and every rank-one matrix belonging to this face is therefore of the form:


$$V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1} \quad (1155)$$

Because $\mathcal{F}(\mathbb{S}_+^N \ni V)$ is isomorphic with a positive semidefinite cone \mathbb{S}_+^{N-1} , then \mathcal{T} constitutes the vectorized extreme directions of \mathcal{F} , the origin constitutes the extreme points of \mathcal{F} , and auxiliary matrix V is some convex combination of those extreme points and directions by the *extremes theorem* (§2.8.1.1.1). \blacklozenge



$$\mathcal{T} \triangleq \left\{ \text{svec}(zz^T) \mid z \in \mathcal{N}(\mathbf{1}\mathbf{1}^T) = \kappa \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \kappa \in \mathbb{R} \right\} \subset \text{svec } \partial \mathbb{S}_+^2$$

Figure 125: Truncated boundary of positive semidefinite cone \mathbb{S}_+^2 in isometrically isomorphic \mathbb{R}^3 (via svec (50)) is, in this dimension, constituted solely by its extreme directions. Truncated cone of Euclidean distance matrices \mathbb{EDM}^2 in isometrically isomorphic subspace \mathbb{R} . Relative boundary of EDM cone is constituted solely by matrix $\mathbf{0}$. Halfline $\mathcal{T} = \{\kappa^2[1 \ -\sqrt{2} \ 1]^T \mid \kappa \in \mathbb{R}\}$ on PSD cone boundary depicts that lone extreme ray (1153) on which orthogonal projection of $-D$ must be positive semidefinite if D is to belong to \mathbb{EDM}^2 . $\text{aff cone } \mathcal{T} = \text{svec } \mathbb{S}_c^2$. (1158) Dual EDM cone is halfspace in \mathbb{R}^3 whose bounding hyperplane has inward-normal $\text{svec } \mathbb{EDM}^2$.



Projection of vectorized $-D$ on range of vectorized zz^T :

$$P_{\text{svec } zz^T}(\text{svec}(-D)) = \frac{\langle zz^T, -D \rangle}{\langle zz^T, zz^T \rangle} zz^T$$

$$D \in \text{EDM}^N \Leftrightarrow \begin{cases} \langle zz^T, -D \rangle \geq 0 \quad \forall zz^T \mid \mathbf{1}\mathbf{1}^T zz^T = \mathbf{0} \\ D \in \mathbb{S}_h^N \end{cases} \quad (1150)$$

Figure 126: Given-matrix D is assumed to belong to symmetric hollow subspace \mathbb{S}_h^2 ; a line in this dimension. Negating D , we find its polar along \mathbb{S}_h^2 . Set \mathcal{T} (1151) has only one ray member in this dimension; not orthogonal to \mathbb{S}_h^2 . Orthogonal projection of $-D$ on \mathcal{T} (indicated by half dot) has nonnegative projection coefficient. Matrix D must therefore be an EDM.

In fact, the smallest face that contains auxiliary matrix V of the PSD cone \mathbb{S}_+^N is the intersection with the geometric center subspace (1885) (1886);

$$\begin{aligned}\mathcal{F}(\mathbb{S}_+^N \ni V) &= \text{cone}\{V_{\mathcal{N}}vv^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1}\} \\ &= \mathbb{S}_c^N \cap \mathbb{S}_+^N \\ &\equiv \{X \succeq 0 \mid \langle X, \mathbf{1}\mathbf{1}^T \rangle = 0\} \quad (1491)\end{aligned}\tag{1156}$$

In isometrically isomorphic $\mathbb{R}^{N(N+1)/2}$

$$\text{svec } \mathcal{F}(\mathbb{S}_+^N \ni V) = \text{cone } \mathcal{T} \tag{1157}$$

related to \mathbb{S}_c^N by

$$\text{aff cone } \mathcal{T} = \text{svec } \mathbb{S}_c^N \tag{1158}$$

6.7.2 EDM criteria in $\mathbf{1}\mathbf{1}^T$

(confer §6.5) Laurent specifies an ellipsope trajectory condition for EDM cone membership: [207, §2.3]

$$D \in \text{EDM}^N \Leftrightarrow [1 - e^{-\alpha d_{ij}}] \in \text{EDM}^N \quad \forall \alpha > 0 \quad (1007)$$

From the parametrized ellipsope \mathcal{E}_t^N in §6.6.2 we propose

$$D \in \text{EDM}^N \Leftrightarrow \left. \begin{array}{l} t \in \mathbb{R}_+ \\ \mathfrak{E} \in \mathcal{E}_t^N \end{array} \right\} \ni D = t\mathbf{1}\mathbf{1}^T - \mathfrak{E} \tag{1159}$$

Chabrillac & Crouzeix [65, §4] prove a different criterion they attribute to Finsler (1937) [125]. We apply it to EDMs: for $D \in \mathbb{S}_h^N$ (955)

$$\begin{aligned}-V_{\mathcal{N}}^T D V_{\mathcal{N}} \succ 0 &\Leftrightarrow \exists \kappa > 0 \ni -D + \kappa \mathbf{1}\mathbf{1}^T \succ 0 \\ &\Leftrightarrow\end{aligned}\tag{1160}$$

$D \in \text{EDM}^N$ with corresponding affine dimension $r = N - 1$

This *Finsler criterion* has geometric interpretation in terms of the vectorization & projection already discussed in connection with (1150). With reference to Figure 125, the offset $\mathbf{1}\mathbf{1}^T$ is simply a direction orthogonal to \mathcal{T} in isomorphic \mathbb{R}^3 . Intuitively, translation of $-D$ in direction $\mathbf{1}\mathbf{1}^T$ is like orthogonal projection on \mathcal{T} in so far as similar information can be obtained.

When the Finsler criterion (1160) is applied despite lower affine dimension, the constant κ can go to infinity making the test $-D + \kappa \mathbf{1}\mathbf{1}^T \succeq 0$ impractical for numerical computation. Chabrillac & Crouzeix invent a criterion for the semidefinite case, but is no more practical: for $D \in \mathbb{S}_h^N$

$$D \in \mathbb{EDM}^N \Leftrightarrow \quad (1161)$$

$\exists \kappa_p > 0 \ni \forall \kappa \geq \kappa_p, -D - \kappa \mathbf{1}\mathbf{1}^T$ [sic] has exactly one negative eigenvalue

6.8 Dual EDM cone

6.8.1 Ambient \mathbb{S}^N

We consider finding the ordinary dual EDM cone in ambient space \mathbb{S}^N where \mathbb{EDM}^N is pointed, closed, convex, but has empty interior. The set of all EDMs in \mathbb{S}^N is a closed convex cone because it is the intersection of halfspaces about the origin in vectorized variable D (§2.4.1.1.1, §2.7.2):

$$\mathbb{EDM}^N = \bigcap_{\substack{z \in \mathcal{N}(\mathbf{1}^T) \\ i=1 \dots N}} \{D \in \mathbb{S}^N \mid \langle e_i e_i^T, D \rangle \geq 0, \langle e_i e_i^T, D \rangle \leq 0, \langle z z^T, -D \rangle \geq 0\} \quad (1162)$$

By definition (270), dual cone \mathcal{K}^* comprises each and every vector inward-normal to a hyperplane supporting convex cone \mathcal{K} (§2.4.2.6.1) or bounding a halfspace containing \mathcal{K} . The dual EDM cone in the ambient space of symmetric matrices is therefore expressible as the aggregate of every conic combination of inward-normals from (1162):

$$\begin{aligned} \mathbb{EDM}^{N*} &= \text{cone}\{e_i e_i^T, -e_j e_j^T \mid i, j = 1 \dots N\} - \text{cone}\{z z^T \mid \mathbf{1}\mathbf{1}^T z z^T = \mathbf{0}\} \\ &= \left\{ \sum_{i=1}^N \zeta_i e_i e_i^T - \sum_{j=1}^N \xi_j e_j e_j^T \mid \zeta_i, \xi_j \geq 0 \right\} - \text{cone}\{z z^T \mid \mathbf{1}\mathbf{1}^T z z^T = \mathbf{0}\} \\ &= \{\delta(u) \mid u \in \mathbb{R}^N\} - \text{cone}\{V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1}, (\|v\| = 1)\} \subset \mathbb{S}^N \\ &= \{\delta^2(Y) - V_{\mathcal{N}} \Psi V_{\mathcal{N}}^T \mid Y \in \mathbb{S}^N, \Psi \in \mathbb{S}_+^{N-1}\} \quad (1163) \end{aligned}$$

The EDM cone is not self-dual in ambient \mathbb{S}^N because its affine hull belongs to a proper subspace

$$\text{aff } \mathbb{EDM}^N = \mathbb{S}_h^N \quad (1164)$$

The ordinary dual EDM cone cannot, therefore, be pointed. (§2.13.1.1)

When $N = 1$, the EDM cone is the point at the origin in \mathbb{R} . Auxiliary matrix $V_{\mathcal{N}}$ is empty $[\emptyset]$, and dual cone \mathbb{EDM}^* is the real line.

When $N = 2$, the EDM cone is a nonnegative real line in isometrically isomorphic \mathbb{R}^3 ; there \mathbb{S}_h^2 is a real line containing the EDM cone. Dual cone \mathbb{EDM}^{2*} is the particular halfspace in \mathbb{R}^3 whose boundary has inward-normal \mathbb{EDM}^2 . Diagonal matrices $\{\delta(u)\}$ in (1163) are represented by a hyperplane through the origin $\{\underline{d} \mid [0 \ 1 \ 0] \underline{d} = 0\}$ while the term $\text{cone}\{V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T\}$ is represented by the halfline \mathcal{T} in Figure 125 belonging to the positive semidefinite cone boundary. The dual EDM cone is formed by translating the hyperplane along the negative semidefinite halfline $-\mathcal{T}$; the union of each and every translation. (confer §2.10.2.0.1)

When cardinality N exceeds 2, the dual EDM cone can no longer be polyhedral simply because the EDM cone cannot. (§2.13.1.1)

6.8.1.1 EDM cone and its dual in ambient \mathbb{S}^N

Consider the two convex cones

$$\begin{aligned} \mathcal{K}_1 &\triangleq \mathbb{S}_h^N \\ \mathcal{K}_2 &\triangleq \bigcap_{y \in \mathcal{N}(\mathbf{1}^T)} \{A \in \mathbb{S}^N \mid \langle y y^T, -A \rangle \geq 0\} \\ &= \{A \in \mathbb{S}^N \mid -z^T V A V z \geq 0 \quad \forall z z^T (\succeq 0)\} \\ &= \{A \in \mathbb{S}^N \mid -V A V \succeq 0\} \end{aligned} \quad (1165)$$

so

$$\mathcal{K}_1 \cap \mathcal{K}_2 = \mathbb{EDM}^N \quad (1166)$$

Dual cone $\mathcal{K}_1^* = \mathbb{S}_h^{N\perp} \subseteq \mathbb{S}^N$ (66) is the subspace of diagonal matrices. From (1163) via (285),

$$\mathcal{K}_2^* = -\text{cone}\{V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1}\} \subset \mathbb{S}^N \quad (1167)$$

Gaffke & Mathar [128, §5.3] observe that projection on \mathcal{K}_1 and \mathcal{K}_2 have simple closed forms: Projection on subspace \mathcal{K}_1 is easily performed by symmetrization and zeroing the main diagonal or *vice versa*, while projection of $H \in \mathbb{S}^N$ on \mathcal{K}_2 is

$$P_{\mathcal{K}_2} H = H - P_{\mathbb{S}_+^N}(V H V) \quad (1168)$$

Proof. First, we observe membership of $H - P_{\mathbb{S}_+^N}(VHV)$ to \mathcal{K}_2 because

$$P_{\mathbb{S}_+^N}(VHV) - H = \left(P_{\mathbb{S}_+^N}(VHV) - VHV \right) + (VHV - H) \quad (1169)$$

The term $P_{\mathbb{S}_+^N}(VHV) - VHV$ necessarily belongs to the (dual) positive semidefinite cone by Theorem E.9.2.0.1. $V^2 = V$, hence

$$-V \left(H - P_{\mathbb{S}_+^N}(VHV) \right) V \succeq 0 \quad (1170)$$

by Corollary A.3.1.0.5.

Next, we require

$$\langle P_{\mathcal{K}_2}H - H, P_{\mathcal{K}_2}H \rangle = 0 \quad (1171)$$

Expanding,

$$\langle -P_{\mathbb{S}_+^N}(VHV), H - P_{\mathbb{S}_+^N}(VHV) \rangle = 0 \quad (1172)$$

$$\langle P_{\mathbb{S}_+^N}(VHV), (P_{\mathbb{S}_+^N}(VHV) - VHV) + (VHV - H) \rangle = 0 \quad (1173)$$

$$\langle P_{\mathbb{S}_+^N}(VHV), (VHV - H) \rangle = 0 \quad (1174)$$

Product VHV belongs to the geometric center subspace; (§E.7.2.0.2)

$$VHV \in \mathbb{S}_c^N = \{Y \in \mathbb{S}^N \mid \mathcal{N}(Y) \supseteq \mathbf{1}\} \quad (1175)$$

Diagonalize $VHV \stackrel{\Delta}{=} Q\Lambda Q^T$ (§A.5) whose nullspace is spanned by the eigenvectors corresponding to 0 eigenvalues by Theorem A.7.3.0.1. Projection of VHV on the PSD cone (§7.1) simply zeroes negative eigenvalues in diagonal matrix Λ . Then

$$\mathcal{N}(P_{\mathbb{S}_+^N}(VHV)) \supseteq \mathcal{N}(VHV) (\supseteq \mathcal{N}(V)) \quad (1176)$$

from which it follows:

$$P_{\mathbb{S}_+^N}(VHV) \in \mathbb{S}_c^N \quad (1177)$$

so $P_{\mathbb{S}_+^N}(VHV) \perp (VHV - H)$ because $VHV - H \in \mathbb{S}_c^{N\perp}$.

Finally, we must have $P_{\mathcal{K}_2}H - H = -P_{\mathbb{S}_+^N}(VHV) \in \mathcal{K}_2^*$. From §6.7.1 we know dual cone $\mathcal{K}_2^* = -\mathcal{F}(\mathbb{S}_+^N \ni V)$ is the negative of the positive semidefinite cone's smallest face that contains auxiliary matrix V . Thus $P_{\mathbb{S}_+^N}(VHV) \in \mathcal{F}(\mathbb{S}_+^N \ni V) \Leftrightarrow \mathcal{N}(P_{\mathbb{S}_+^N}(VHV)) \supseteq \mathcal{N}(V)$ (§2.9.2.3) which was already established in (1176). \blacklozenge

$$\text{EDM}^2 = \mathbb{S}_h^2 \cap (\mathbb{S}_c^{2\perp} - \mathbb{S}_+^2)$$




Figure 127: Orthogonal complement $\mathbb{S}_c^{2\perp}$ (1887) (§B.2) of geometric center subspace (a plane in isometrically isomorphic \mathbb{R}^3 ; drawn is a tiled fragment) apparently supporting positive semidefinite cone. (Rounded vertex is artifact of plot.) Line $\text{svec } \mathbb{S}_c^2 = \text{aff cone } \mathcal{T}$ (1158) intersects $\text{svec } \partial \mathbb{S}_+^2$, also drawn in Figure 125; it runs along PSD cone boundary. (confer Figure 108)

$$\text{EDM}^2 = \mathbb{S}_h^2 \cap (\mathbb{S}_c^{2\perp} - \mathbb{S}_+^2)$$




Figure 128: EDM cone construction in isometrically isomorphic \mathbb{R}^3 by adding polar PSD cone to $\text{svec } \mathbb{S}_c^{2\perp}$. Difference $\text{svec}(\mathbb{S}_c^{2\perp} - \mathbb{S}_+^2)$ is halfspace partially bounded by $\text{svec } \mathbb{S}_c^{2\perp}$. EDM cone is nonnegative halfline along $\text{svec } \mathbb{S}_h^2$ in this dimension.

From the results in §E.7.2.0.2, we know matrix product VHV is the orthogonal projection of $H \in \mathbb{S}^N$ on the geometric center subspace \mathbb{S}_c^N . Thus the projection product

$$P_{\mathcal{K}_2}H = H - P_{\mathbb{S}_+^N}P_{\mathbb{S}_c^N}H \quad (1178)$$

6.8.1.1.1 Lemma. *Projection on PSD cone \cap geometric center subspace.*

$$P_{\mathbb{S}_+^N \cap \mathbb{S}_c^N} = P_{\mathbb{S}_+^N}P_{\mathbb{S}_c^N} \quad (1179)$$

◇

Proof. For each and every $H \in \mathbb{S}^N$, projection of $P_{\mathbb{S}_c^N}H$ on the positive semidefinite cone remains in the geometric center subspace

$$\mathbb{S}_c^N = \{G \in \mathbb{S}^N \mid G\mathbf{1} = \mathbf{0}\} \quad (1885)$$

$$= \{G \in \mathbb{S}^N \mid \mathcal{N}(G) \supseteq \mathbf{1}\} = \{G \in \mathbb{S}^N \mid \mathcal{R}(G) \subseteq \mathcal{N}(\mathbf{1}^T)\} \quad (905)$$

$$= \{VYV \mid Y \in \mathbb{S}^N\} \subset \mathbb{S}^N \quad (1886)$$

That is because: eigenvectors of $P_{\mathbb{S}_c^N}H$ corresponding to 0 eigenvalues span its nullspace $\mathcal{N}(P_{\mathbb{S}_c^N}H)$. (§A.7.3.0.1) To project $P_{\mathbb{S}_c^N}H$ on the positive semidefinite cone, its negative eigenvalues are zeroed. (§7.1.2) The nullspace is thereby expanded while eigenvectors originally spanning $\mathcal{N}(P_{\mathbb{S}_c^N}H)$ remain intact. Because the geometric center subspace is invariant to projection on the PSD cone, then the rule for projection on a convex set in a subspace governs (§E.9.5, projectors do not commute) and statement (1179) follows directly. ◆

From the lemma it follows

$$\{P_{\mathbb{S}_+^N}P_{\mathbb{S}_c^N}H \mid H \in \mathbb{S}^N\} = \{P_{\mathbb{S}_+^N \cap \mathbb{S}_c^N}H \mid H \in \mathbb{S}^N\} \quad (1180)$$

Then from (1912)

$$-(\mathbb{S}_c^N \cap \mathbb{S}_+^N)^* = \{H - P_{\mathbb{S}_+^N}P_{\mathbb{S}_c^N}H \mid H \in \mathbb{S}^N\} \quad (1181)$$

From (285) we get closure of a vector sum

$$\mathcal{K}_2 = -(\mathbb{S}_c^N \cap \mathbb{S}_+^N)^* = \mathbb{S}_c^{N\perp} - \mathbb{S}_+^N \quad (1182)$$

therefore the equality

$$\text{EDM}^N = \mathcal{K}_1 \cap \mathcal{K}_2 = \mathbb{S}_h^N \cap (\mathbb{S}_c^{N\perp} - \mathbb{S}_+^N) \quad (1183)$$

whose veracity is intuitively evident, in hindsight, [78, p.109] from the most fundamental EDM definition (808). Formula (1183) is not a matrix criterion for membership to the EDM cone, it is not an EDM definition, and it is not an equivalence between EDM operators nor an isomorphism. Rather, it is a recipe for constructing the EDM cone whole from large Euclidean bodies: the positive semidefinite cone, orthogonal complement of the geometric center subspace, and symmetric hollow subspace. A realization of this construction in low dimension is illustrated in Figure 127 and Figure 128.

The dual EDM cone follows directly from (1183) by standard properties of cones (§2.13.1.1):

$$\mathbb{EDM}^{N*} = \overline{\mathcal{K}_1^* + \mathcal{K}_2^*} = \mathbb{S}_h^{N\perp} - \mathbb{S}_c^N \cap \mathbb{S}_+^N \quad (1184)$$

which bears strong resemblance to (1163).

6.8.1.2 nonnegative orthant contains \mathbb{EDM}^N

That \mathbb{EDM}^N is a proper subset of the nonnegative orthant is not obvious from (1183). We wish to verify

$$\mathbb{EDM}^N = \mathbb{S}_h^N \cap (\mathbb{S}_c^{N\perp} - \mathbb{S}_+^N) \subset \mathbb{R}_+^{N \times N} \quad (1185)$$

While there are many ways to prove this, it is sufficient to show that all entries of the extreme directions of \mathbb{EDM}^N must be nonnegative; *id est*, for any particular nonzero vector $z = [z_i, i=1 \dots N] \in \mathcal{N}(\mathbf{1}^T)$ (§6.5.3.2),

$$\delta(zz^T)\mathbf{1}^T + \mathbf{1}\delta(zz^T)^T - 2zz^T \geq \mathbf{0} \quad (1186)$$

where the inequality denotes entrywise comparison. The inequality holds because the i, j^{th} entry of an extreme direction is squared: $(z_i - z_j)^2$.

We observe that the dyad $2zz^T \in \mathbb{S}_+^N$ belongs to the positive semidefinite cone, the doublet

$$\delta(zz^T)\mathbf{1}^T + \mathbf{1}\delta(zz^T)^T \in \mathbb{S}_c^{N\perp} \quad (1187)$$

to the orthogonal complement (1887) of the geometric center subspace, while their difference is a member of the symmetric hollow subspace \mathbb{S}_h^N .

Here is an algebraic method to prove nonnegativity: Suppose we are given $A \in \mathbb{S}_c^{N\perp}$ and $B = [B_{ij}] \in \mathbb{S}_+^N$ and $A - B \in \mathbb{S}_h^N$. Then we have, for some vector u , $A = u\mathbf{1}^T + \mathbf{1}u^T = [A_{ij}] = [u_i + u_j]$ and $\delta(B) = \delta(A) = 2u$. Positive semidefiniteness of B requires nonnegativity $A - B \geq \mathbf{0}$ because

$$(e_i - e_j)^T B (e_i - e_j) = (B_{ii} - B_{ij}) - (B_{ji} - B_{jj}) = 2(u_i + u_j) - 2B_{ij} \geq 0 \quad (1188)$$

6.8.1.3 Dual Euclidean distance matrix criterion

Conditions necessary for membership of a matrix $D^* \in \mathbb{S}^N$ to the dual EDM cone \mathbb{EDM}^{N^*} may be derived from (1163): $D^* \in \mathbb{EDM}^{N^*} \Rightarrow D^* = \delta(y) - V_{\mathcal{N}} A V_{\mathcal{N}}^T$ for some vector y and positive semidefinite matrix $A \in \mathbb{S}_+^{N-1}$. This in turn implies $\delta(D^* \mathbf{1}) = \delta(y)$. Then, for $D^* \in \mathbb{S}^N$

$$D^* \in \mathbb{EDM}^{N^*} \Leftrightarrow \delta(D^* \mathbf{1}) - D^* \succeq 0 \quad (1189)$$

where, for any symmetric matrix D^*

$$\delta(D^* \mathbf{1}) - D^* \in \mathbb{S}_c^N \quad (1190)$$

To show sufficiency of the matrix criterion in (1189), recall Gram-form EDM operator

$$\mathbf{D}(G) = \delta(G) \mathbf{1}^T + \mathbf{1} \delta(G)^T - 2G \quad (820)$$

where Gram matrix G is positive semidefinite by definition, and recall the self-adjointness property of the main-diagonal linear operator δ (§A.1):

$$\langle D, D^* \rangle = \langle \delta(G) \mathbf{1}^T + \mathbf{1} \delta(G)^T - 2G, D^* \rangle = \langle G, \delta(D^* \mathbf{1}) - D^* \rangle 2 \quad (839)$$

Assuming $\langle G, \delta(D^* \mathbf{1}) - D^* \rangle \geq 0$ (1395), then we have known membership relation (§2.13.2.0.1)

$$D^* \in \mathbb{EDM}^{N^*} \Leftrightarrow \langle D, D^* \rangle \geq 0 \quad \forall D \in \mathbb{EDM}^N \quad (1191)$$

◆

Elegance of this matrix criterion (1189) for membership to the dual EDM cone is the lack of any other assumptions except D^* be symmetric. (Recall: Schoenberg criterion (827) for membership to the EDM cone requires membership to the symmetric hollow subspace.)

Linear Gram-form EDM operator (820) has adjoint, for $Y \in \mathbb{S}^N$

$$\mathbf{D}^T(Y) \triangleq (\delta(Y \mathbf{1}) - Y) 2 \quad (1192)$$

Then we have: [78, p.111]

$$\mathbb{EDM}^{N^*} = \{Y \in \mathbb{S}^N \mid \delta(Y \mathbf{1}) - Y \succeq 0\} \quad (1193)$$

the dual EDM cone expressed in terms of the adjoint operator. A dual EDM cone determined this way is illustrated in Figure 130.

6.8.1.3.1 Exercise. *Dual EDM spectral cone.*

Find a spectral cone as in §5.11.2 corresponding to \mathbb{EDM}^{N^*} . ▼

6.8.1.4 Nonorthogonal components of dual EDM

Now we tie construct (1184) for the dual EDM cone together with the matrix criterion (1189) for dual EDM cone membership. For any $D^* \in \mathbb{S}^N$ it is obvious:

$$\delta(D^* \mathbf{1}) \in \mathbb{S}_h^{N\perp} \quad (1194)$$

any diagonal matrix belongs to the subspace of diagonal matrices (61). We know when $D^* \in \mathbb{EDM}^{N^*}$

$$\delta(D^* \mathbf{1}) - D^* \in \mathbb{S}_c^N \cap \mathbb{S}_+^N \quad (1195)$$

this adjoint expression (1192) belongs to that face (1156) of the positive semidefinite cone \mathbb{S}_+^N in the geometric center subspace. Any nonzero dual EDM

$$D^* = \delta(D^* \mathbf{1}) - (\delta(D^* \mathbf{1}) - D^*) \in \mathbb{S}_h^{N\perp} \ominus \mathbb{S}_c^N \cap \mathbb{S}_+^N = \mathbb{EDM}^{N^*} \quad (1196)$$

can therefore be expressed as the difference of two linearly independent nonorthogonal components (Figure 108, Figure 129).

6.8.1.5 Affine dimension complementarity

From §6.8.1.3 we have, for some $A \in \mathbb{S}_+^{N-1}$ (*confer* (1195))

$$\delta(D^* \mathbf{1}) - D^* = V_{\mathcal{N}} A V_{\mathcal{N}}^T \in \mathbb{S}_c^N \cap \mathbb{S}_+^N \quad (1197)$$

if and only if D^* belongs to the dual EDM cone. Call $\text{rank}(V_{\mathcal{N}} A V_{\mathcal{N}}^T)$ *dual affine dimension*. Empirically, we find a complementary relationship in affine dimension between the projection of some arbitrary symmetric matrix H on the polar EDM cone, $\mathbb{EDM}^{N^\circ} = -\mathbb{EDM}^{N^*}$, and its projection on the EDM cone; *id est*, the optimal solution of ^{6.11}

$$\begin{aligned} & \underset{D^\circ \in \mathbb{S}^N}{\text{minimize}} && \|D^\circ - H\|_F \\ & \text{subject to} && D^\circ - \delta(D^\circ \mathbf{1}) \succeq 0 \end{aligned} \quad (1198)$$

^{6.11}This dual projection can be solved quickly (without semidefinite programming) via

$$D^\circ = \delta(D^\circ \mathbf{1}) + (D^\circ - \delta(D^\circ \mathbf{1})) \in \mathbb{S}_h^{N\perp} \oplus \mathbb{S}_c^N \cap \mathbb{S}_+^N = \mathbf{EDM}^{N^\circ}$$




Figure 129: [Hand-drawn abstraction](#) of polar EDM cone (drawn truncated). Any member D° of polar EDM cone can be decomposed into two linearly independent nonorthogonal components: $\delta(D^\circ \mathbf{1})$ and $D^\circ - \delta(D^\circ \mathbf{1})$.

has dual affine dimension complementary to affine dimension corresponding to the optimal solution of

$$\begin{aligned} & \underset{D \in \mathbb{S}_h^N}{\text{minimize}} && \|D - H\|_{\text{F}} \\ & \text{subject to} && -V_{\mathcal{N}}^{\text{T}} D V_{\mathcal{N}} \succeq 0 \end{aligned} \quad (1199)$$

Precisely,

$$\text{rank}(D^{\circ\star} - \delta(D^{\circ\star} \mathbf{1})) + \text{rank}(V_{\mathcal{N}}^{\text{T}} D^{\star} V_{\mathcal{N}}) = N - 1 \quad (1200)$$

and $\text{rank}(D^{\circ\star} - \delta(D^{\circ\star} \mathbf{1})) \leq N - 1$ because vector $\mathbf{1}$ is always in the nullspace of rank's argument. This is similar to the known result for projection on the self-dual positive semidefinite cone and its polar:

$$\text{rank } P_{-\mathbb{S}_+^N} H + \text{rank } P_{\mathbb{S}_+^N} H = N \quad (1201)$$

When low affine dimension is a desirable result of projection on the EDM cone, projection on the polar EDM cone should be performed instead. Convex polar problem (1198) can be solved for $D^{\circ\star}$ by transforming to an equivalent Schur-form semidefinite program (§3.1.7.2). Interior-point methods for numerically solving semidefinite programs tend to produce high-rank solutions. (§4.1.1.1) Then $D^{\star} = H - D^{\circ\star} \in \text{EDM}^N$ by Corollary E.9.2.2.1, and D^{\star} will tend to have low affine dimension. This approach breaks when attempting projection on a cone subset discriminated by affine dimension or rank, because then we have no complementarity relation like (1200) or (1201) (§7.1.4.1).

Lemma 6.8.1.1.1; rewriting,

$$\begin{aligned} & \underset{D^{\circ} \in \mathbb{S}^N}{\text{minimize}} && \|(D^{\circ} - \delta(D^{\circ} \mathbf{1})) - (H - \delta(D^{\circ} \mathbf{1}))\|_{\text{F}} \\ & \text{subject to} && D^{\circ} - \delta(D^{\circ} \mathbf{1}) \succeq 0 \end{aligned}$$

which is the projection of affinely transformed optimal solution $H - \delta(D^{\circ\star} \mathbf{1})$ on $\mathbb{S}_c^N \cap \mathbb{S}_+^N$;

$$D^{\circ\star} - \delta(D^{\circ\star} \mathbf{1}) = P_{\mathbb{S}_+^N} P_{\mathbb{S}_c^N} (H - \delta(D^{\circ\star} \mathbf{1}))$$

Foreknowledge of an optimal solution $D^{\circ\star}$ as argument to projection suggests recursion.

6.8.1.6 EDM cone duality

In §5.6.1.1, via Gram-form EDM operator

$$\mathbf{D}(G) = \delta(G)\mathbf{1}^T + \mathbf{1}\delta(G)^T - 2G \in \mathbb{EDM}^N \iff G \succeq 0 \quad (820)$$

we established clear connection between the EDM cone and that face (1156) of positive semidefinite cone \mathbb{S}_+^N in the geometric center subspace:

$$\mathbb{EDM}^N = \mathbf{D}(\mathbb{S}_c^N \cap \mathbb{S}_+^N) \quad (922)$$

$$\mathbf{V}(\mathbb{EDM}^N) = \mathbb{S}_c^N \cap \mathbb{S}_+^N \quad (923)$$

where

$$\mathbf{V}(D) = -VDV\frac{1}{2} \quad (911)$$

In §5.6.1 we established

$$\mathbb{S}_c^N \cap \mathbb{S}_+^N = V_{\mathcal{N}}\mathbb{S}_+^{N-1}V_{\mathcal{N}}^T \quad (909)$$

Then from (1189), (1197), and (1163) we can deduce

$$\delta(\mathbb{EDM}^{N*}\mathbf{1}) - \mathbb{EDM}^{N*} = V_{\mathcal{N}}\mathbb{S}_+^{N-1}V_{\mathcal{N}}^T = \mathbb{S}_c^N \cap \mathbb{S}_+^N \quad (1202)$$

which, by (922) and (923), means the EDM cone can be related to the dual EDM cone by an equality:

$$\mathbb{EDM}^N = \mathbf{D}\left(\delta(\mathbb{EDM}^{N*}\mathbf{1}) - \mathbb{EDM}^{N*}\right) \quad (1203)$$

$$\mathbf{V}(\mathbb{EDM}^N) = \delta(\mathbb{EDM}^{N*}\mathbf{1}) - \mathbb{EDM}^{N*} \quad (1204)$$

This means projection $-\mathbf{V}(\mathbb{EDM}^N)$ of the EDM cone on the geometric center subspace \mathbb{S}_c^N (§E.7.2.0.2) is a linear transformation of the dual EDM cone: $\mathbb{EDM}^{N*} - \delta(\mathbb{EDM}^{N*}\mathbf{1})$. Secondly, it means the EDM cone is not self-dual.

6.8.1.7 Schoenberg criterion is discretized membership relation

We show the Schoenberg criterion

$$\left. \begin{array}{l} -V_{\mathcal{N}}^T D V_{\mathcal{N}} \in \mathbb{S}_+^{N-1} \\ D \in \mathbb{S}_h^N \end{array} \right\} \Leftrightarrow D \in \text{EDM}^N \quad (827)$$

to be a discretized membership relation (§2.13.4) between a closed convex cone \mathcal{K} and its dual \mathcal{K}^* like

$$\langle y, x \rangle \geq 0 \text{ for all } y \in \mathcal{G}(\mathcal{K}^*) \Leftrightarrow x \in \mathcal{K} \quad (331)$$

where $\mathcal{G}(\mathcal{K}^*)$ is any set of generators whose conic hull constructs closed convex dual cone \mathcal{K}^* :

The Schoenberg criterion is the same as

$$\left. \begin{array}{l} \langle zz^T, -D \rangle \geq 0 \quad \forall zz^T \mid \mathbf{1}\mathbf{1}^T zz^T = \mathbf{0} \\ D \in \mathbb{S}_h^N \end{array} \right\} \Leftrightarrow D \in \text{EDM}^N \quad (1150)$$

which, by (1151), is the same as

$$\left. \begin{array}{l} \langle zz^T, -D \rangle \geq 0 \quad \forall zz^T \in \{V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1}\} \\ D \in \mathbb{S}_h^N \end{array} \right\} \Leftrightarrow D \in \text{EDM}^N \quad (1205)$$

where the zz^T constitute a set of generators \mathcal{G} for the positive semidefinite cone's smallest face $\mathcal{F}(\mathbb{S}_+^N \ni V)$ (§6.7.1) that contains auxiliary matrix V . From the aggregate in (1163) we get the ordinary membership relation, assuming only $D \in \mathbb{S}^N$ [177, p.58]

$$\langle D^*, D \rangle \geq 0 \quad \forall D^* \in \text{EDM}^{N^*} \Leftrightarrow D \in \text{EDM}^N \quad (1206)$$

$$\langle D^*, D \rangle \geq 0 \quad \forall D^* \in \{\delta(u) \mid u \in \mathbb{R}^N\} - \text{cone}\{V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1}\} \Leftrightarrow D \in \text{EDM}^N$$

Discretization (331) yields:

$$\langle D^*, D \rangle \geq 0 \quad \forall D^* \in \{e_i e_i^T, -e_j e_j^T, -V_{\mathcal{N}} v v^T V_{\mathcal{N}}^T \mid i, j = 1 \dots N, v \in \mathbb{R}^{N-1}\} \Leftrightarrow D \in \text{EDM}^N \quad (1207)$$

Because $\langle \{\delta(u) \mid u \in \mathbb{R}^N\}, D \rangle \geq 0 \Leftrightarrow D \in \mathbb{S}_h^N$, we can restrict observation to the symmetric hollow subspace without loss of generality. Then for $D \in \mathbb{S}_h^N$

$$\langle D^*, D \rangle \geq 0 \quad \forall D^* \in \{-V_{\mathcal{N}}vv^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1}\} \Leftrightarrow D \in \text{EDM}^N \quad (1208)$$

this discretized membership relation becomes (1205); identical to the Schoenberg criterion.

Hitherto a correspondence between the EDM cone and a face of a PSD cone, the Schoenberg criterion is now accurately interpreted as a discretized membership relation between the EDM cone and its ordinary dual.

6.8.2 Ambient \mathbb{S}_h^N

When instead we consider the ambient space of symmetric hollow matrices (1164), then still we find the EDM cone is not self-dual for $N > 2$. The simplest way to prove this is as follows:

Given a set of generators $\mathcal{G} = \{\Gamma\}$ (1127) for the pointed closed convex EDM cone, the *discretized membership theorem* in §2.13.4.2.1 asserts that members of the dual EDM cone in the ambient space of symmetric hollow matrices can be discerned via discretized membership relation:

$$\begin{aligned} \text{EDM}^{N*} \cap \mathbb{S}_h^N &\triangleq \{D^* \in \mathbb{S}_h^N \mid \langle \Gamma, D^* \rangle \geq 0 \quad \forall \Gamma \in \mathcal{G}(\text{EDM}^N)\} & (1209) \\ &= \{D^* \in \mathbb{S}_h^N \mid \langle \delta(zz^T)\mathbf{1}^T + \mathbf{1}\delta(zz^T)^T - 2zz^T, D^* \rangle \geq 0 \quad \forall z \in \mathcal{N}(\mathbf{1}^T)\} \\ &= \{D^* \in \mathbb{S}_h^N \mid \langle \mathbf{1}\delta(zz^T)^T - zz^T, D^* \rangle \geq 0 \quad \forall z \in \mathcal{N}(\mathbf{1}^T)\} \end{aligned}$$

By comparison

$$\text{EDM}^N = \{D \in \mathbb{S}_h^N \mid \langle -zz^T, D \rangle \geq 0 \quad \forall z \in \mathcal{N}(\mathbf{1}^T)\} \quad (1210)$$

the term $\delta(zz^T)^T D^* \mathbf{1}$ foils any hope of self-duality in ambient \mathbb{S}_h^N . \blacklozenge

To find the dual EDM cone in ambient \mathbb{S}_h^N per §2.13.9.4 we prune the aggregate in (1163) describing the ordinary dual EDM cone, removing any member having nonzero main diagonal:

$$\begin{aligned} \text{EDM}^{N*} \cap \mathbb{S}_h^N &= \text{cone}\{\delta^2(V_{\mathcal{N}}vv^T V_{\mathcal{N}}^T) - V_{\mathcal{N}}vv^T V_{\mathcal{N}}^T \mid v \in \mathbb{R}^{N-1}\} \\ &= \{\delta^2(V_{\mathcal{N}}\Psi V_{\mathcal{N}}^T) - V_{\mathcal{N}}\Psi V_{\mathcal{N}}^T \mid \Psi \in \mathbb{S}_+^{N-1}\} \end{aligned} \tag{1211}$$

When $N=1$, the EDM cone and its dual in ambient \mathbb{S}_h each comprise the origin in isomorphic \mathbb{R}^0 ; thus, self-dual in this dimension. (*confer*(96))

When $N=2$, the EDM cone is the nonnegative real line in isomorphic \mathbb{R} . (Figure 125) EDM^{2*} in \mathbb{S}_h^2 is identical, thus self-dual in this dimension. This result is in agreement with (1209), verified directly: for all $\kappa \in \mathbb{R}$, $z = \kappa \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ and $\delta(zz^T) = \kappa^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \Rightarrow d_{12}^* \geq 0$.

The first case adverse to self-duality $N=3$ may be deduced from Figure 117; the EDM cone is a circular cone in isomorphic \mathbb{R}^3 corresponding to no rotation of the Lorentz cone (161) (the self-dual circular cone). Figure 130 illustrates the EDM cone and its dual in ambient \mathbb{S}_h^3 ; no longer self-dual.


6.9 Theorem of the alternative

In §2.13.2.1.1 we showed how alternative systems of generalized inequality can be derived from closed convex cones and their duals. This section is, therefore, a fitting postscript to the discussion of the dual EDM cone.

6.9.0.0.1 Theorem. *EDM alternative.* [142, §1]
 Given $D \in \mathbb{S}_h^N$

$$\begin{aligned} D &\in \text{EDM}^N \\ &\text{or in the alternative} \\ \exists z \text{ such that } &\begin{cases} \mathbf{1}^T z = 1 \\ Dz = \mathbf{0} \end{cases} \end{aligned} \tag{1212}$$

In words, either $\mathcal{N}(D)$ intersects hyperplane $\{z \mid \mathbf{1}^T z = 1\}$ or D is an EDM; the alternatives are incompatible. \diamond



$$D^* \in EDM^{N^*} \Leftrightarrow \delta(D^* \mathbf{1}) - D^* \succeq 0 \quad (1189)$$

Figure 130: Ordinary dual EDM cone projected on S_h^3 shrouds EDM^3 ; drawn tiled in isometrically isomorphic \mathbb{R}^3 . (It so happens: intersection $EDM^{N^*} \cap S_h^N$ (§2.13.9.3) is identical to projection of dual EDM cone on S_h^N .)

When D is an EDM [229, §2]

$$\mathcal{N}(D) \subset \mathcal{N}(\mathbf{1}^T) = \{z \mid \mathbf{1}^T z = 0\} \quad (1213)$$

Because [142, §2] (§E.0.1)

$$\begin{aligned} DD^\dagger \mathbf{1} &= \mathbf{1} \\ \mathbf{1}^T D^\dagger D &= \mathbf{1}^T \end{aligned} \quad (1214)$$

then

$$\mathcal{R}(\mathbf{1}) \subset \mathcal{R}(D) \quad (1215)$$

6.10 Postscript

We provided an equality (1183) relating the convex cone of Euclidean distance matrices to the convex cone of positive semidefinite matrices. Projection on a positive semidefinite cone, constrained by an upper bound on rank, is easy and well known; [112] simply, a matter of truncating a list of eigenvalues. Projection on a positive semidefinite cone with such a rank constraint is, in fact, a convex optimization problem. (§7.1.4)

In the past, it was difficult to project on the EDM cone under a constraint on rank or affine dimension. A surrogate method was to invoke the Schoenberg criterion (827) and then project on a positive semidefinite cone under a rank constraint bounding affine dimension from above. But a solution acquired that way is necessarily suboptimal.

In §7.3.3 we present a method for projecting directly on the EDM cone under a constraint on rank or affine dimension.

Chapter 7

Proximity problems

In summary, we find that the solution to problem [(1225.3) p.505] is difficult and depends on the dimension of the space as the geometry of the cone of EDMs becomes more complex.

–Hayden, Wells, Liu, & Tarazaga (1991) [164, §3]

A problem common to various sciences is to find the Euclidean distance matrix (EDM) $D \in \mathbb{EDM}^N$ closest in some sense to a given complete matrix of measurements H under a constraint on affine dimension $0 \leq r \leq N-1$ (§2.3.1, §5.7.1.1); rather, r is bounded above by desired affine dimension ρ .

7.0.1 Measurement matrix H

Ideally, we want a given matrix of measurements $H \in \mathbb{R}^{N \times N}$ to conform with the first three Euclidean metric properties (§5.2); to belong to the intersection of the orthant of nonnegative matrices $\mathbb{R}_+^{N \times N}$ with the symmetric hollow subspace \mathbb{S}_h^N (§2.2.3.0.1). Geometrically, we want H to belong to the polyhedral cone (§2.12.1.0.1)

$$\mathcal{K} \triangleq \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N} \quad (1216)$$

Yet in practice, H can possess significant measurement uncertainty (noise).

Sometimes realization of an optimization problem demands that its input, the given matrix H , possess some particular characteristics; perhaps symmetry and hollowness or nonnegativity. When that H given does not possess the desired properties, then we must impose them upon H prior to optimization:

- When *measurement matrix* H is not symmetric or hollow, taking its symmetric hollow part is equivalent to orthogonal projection on the symmetric hollow subspace \mathbb{S}_h^N .
- When measurements of distance in H are negative, zeroing negative entries effects unique minimum-distance projection on the orthant of nonnegative matrices $\mathbb{R}_+^{N \times N}$ in isomorphic \mathbb{R}^{N^2} (§E.9.2.2.3).

7.0.1.1 Order of imposition

Since convex cone \mathcal{K} (1216) is the intersection of an orthant with a subspace, we want to project on that subset of the orthant belonging to the subspace; on the nonnegative orthant in the symmetric hollow subspace that is, in fact, the intersection. For that reason alone, unique minimum-distance projection of H on \mathcal{K} (that member of \mathcal{K} closest to H in isomorphic \mathbb{R}^{N^2} in the Euclidean sense) can be attained by first taking its symmetric hollow part, and only then clipping negative entries of the result to 0; *id est*, there is only one correct *order of projection*, in general, on an orthant intersecting a subspace:

- project on the subspace, then project the result on the orthant in that subspace. (confer §E.9.5)

In contrast, order of projection on an intersection of subspaces is arbitrary.

That order-of-projection rule applies more generally, of course, to the intersection of any convex set \mathcal{C} with any subspace. Consider the *proximity problem*^{7.1} over convex feasible set $\mathbb{S}_h^N \cap \mathcal{C}$ given nonsymmetric nonhollow $H \in \mathbb{R}^{N \times N}$:

$$\begin{aligned} & \underset{B \in \mathbb{S}_h^N}{\text{minimize}} && \|B - H\|_{\text{F}}^2 \\ & \text{subject to} && B \in \mathcal{C} \end{aligned} \tag{1217}$$

a convex optimization problem. Because the symmetric hollow subspace is orthogonal to the antisymmetric antihollow subspace (§2.2.3), then for $B \in \mathbb{S}_h^N$

$$\text{tr}\left(B^T\left(\frac{1}{2}(H - H^T) + \delta^2(H)\right)\right) = 0 \tag{1218}$$

so the objective function is equivalent to

$$\|B - H\|_{\text{F}}^2 \equiv \left\|B - \left(\frac{1}{2}(H + H^T) - \delta^2(H)\right)\right\|_{\text{F}}^2 + \left\|\frac{1}{2}(H - H^T) + \delta^2(H)\right\|_{\text{F}}^2 \tag{1219}$$

This means the antisymmetric antihollow part of given matrix H would be ignored by minimization with respect to symmetric hollow variable B under the Frobenius norm; *id est*, minimization proceeds as though given the symmetric hollow part of H .

This action of the Frobenius norm (1219) is effectively a Euclidean projection (minimum-distance projection) of H on the symmetric hollow subspace \mathbb{S}_h^N prior to minimization. Thus minimization proceeds inherently following the correct order for projection on $\mathbb{S}_h^N \cap \mathcal{C}$. Therefore we may either assume $H \in \mathbb{S}_h^N$, or take its symmetric hollow part prior to optimization.

^{7.1}There are two equivalent interpretations of projection (§E.9): one finds a set normal, the other, minimum distance between a point and a set. Here we realize the latter view.




Figure 131: Pseudo-Venn diagram: The EDM cone belongs to the intersection of the symmetric hollow subspace with the nonnegative orthant; $\text{EDM}^N \subseteq \mathcal{K}$ (807). EDM^N cannot exist outside \mathbb{S}_h^N , but $\mathbb{R}_+^{N \times N}$ does.

7.0.1.2 Egregious input error under nonnegativity demand

More pertinent to the optimization problems presented herein where

$$\mathcal{C} \triangleq \text{EDM}^N \subseteq \mathcal{K} = \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N} \quad (1220)$$

then should some particular realization of a proximity problem demand input H be nonnegative, and were we only to zero negative entries of a nonsymmetric nonhollow input H prior to optimization, then the ensuing projection on EDM^N would be guaranteed incorrect (out of order).

Now comes a surprising fact: Even were we to correctly follow the order-of-projection rule and provide $H \in \mathcal{K}$ prior to optimization, then the ensuing projection on EDM^N will be incorrect whenever input H has negative entries and some proximity problem demands nonnegative input H .




Figure 132: Pseudo-Venn diagram from Figure 131 showing elbow placed in path of projection of H on $\text{EDM}^N \subset \mathbb{S}_h^N$ by an optimization problem demanding nonnegative input matrix H . The first two line segments leading away from H result from correct order-of-projection required to provide nonnegative H prior to optimization. Were H nonnegative, then its projection on \mathbb{S}_h^N would instead belong to \mathcal{K} ; making the elbow disappear. (confer Figure 144)

This is best understood referring to Figure 131: Suppose nonnegative input H is demanded, and then the problem realization correctly projects its input first on \mathbb{S}_h^N and then directly on $\mathcal{C} = \text{EDM}^N$. That demand for nonnegativity effectively requires imposition of \mathcal{K} on input H prior to optimization so as to obtain correct order of projection (on \mathbb{S}_h^N first). Yet such an imposition prior to projection on EDM^N generally introduces an *elbow* into the path of projection (illustrated in Figure 132) caused by the technique itself; that being, a particular proximity problem realization requiring nonnegative input.

Any procedure for imposition of nonnegativity on input H can only be incorrect in this circumstance. There is no resolution unless input H is guaranteed nonnegative with no tinkering. Otherwise, we have no choice but to employ a different problem realization; one not demanding nonnegative input.

7.0.2 Lower bound

Most of the problems we encounter in this chapter have the general form:

$$\begin{aligned} & \underset{B}{\text{minimize}} && \|B - A\|_F \\ & \text{subject to} && B \in \mathcal{C} \end{aligned} \tag{1221}$$

where $A \in \mathbb{R}^{m \times n}$ is given data. This particular objective denotes Euclidean projection (§E) of vectorized matrix A on the set \mathcal{C} which may or may not be convex. When \mathcal{C} is convex, then the projection is unique minimum-distance because the Frobenius norm when squared is a strictly convex function of variable B and because the optimal solution is the same regardless of the square (463). When \mathcal{C} is a subspace, then the direction of projection is orthogonal to \mathcal{C} .

Denoting by $A \triangleq U_A \Sigma_A Q_A^T$ and $B \triangleq U_B \Sigma_B Q_B^T$ their full singular value decompositions (whose singular values are always nonincreasingly ordered (§A.6)), there exists a tight lower bound on the objective over the manifold of orthogonal matrices;

$$\|\Sigma_B - \Sigma_A\|_F \leq \inf_{U_A, U_B, Q_A, Q_B} \|B - A\|_F \tag{1222}$$

This least lower bound holds more generally for any orthogonally invariant norm on $\mathbb{R}^{m \times n}$ (§2.2.1) including the Frobenius and *spectral norm* [291, §II.3]. [180, §7.4.51]

7.0.3 Problem approach

Problems traditionally posed in terms of point position $\{x_i \in \mathbb{R}^n, i = 1 \dots N\}$ such as

$$\underset{\{x_i\}}{\text{minimize}} \sum_{i,j \in \mathcal{I}} (\|x_i - x_j\| - h_{ij})^2 \quad (1223)$$

or

$$\underset{\{x_i\}}{\text{minimize}} \sum_{i,j \in \mathcal{I}} (\|x_i - x_j\|^2 - h_{ij})^2 \quad (1224)$$

(where \mathcal{I} is an abstract set of indices and h_{ij} is given data) are everywhere converted herein to the distance-square variable D or to Gram matrix G ; the Gram matrix acting as bridge between position and distance. (That conversion is performed regardless of whether known data is complete.) Then the techniques of chapter 5 or chapter 6 are applied to find relative or absolute position. This approach is taken because we prefer introduction of rank constraints into convex problems rather than searching a googol of local minima in nonconvex problems like (1223) or (1224) [89].

7.0.4 Three prevalent proximity problems

There are three statements of the closest-EDM problem prevalent in the literature, the multiplicity due primarily to choice of projection on the EDM *versus* positive semidefinite (PSD) cone and vacillation between the distance-square variable d_{ij} *versus* absolute distance $\sqrt{d_{ij}}$. In their most fundamental form, the three prevalent proximity problems are (1225.1), (1225.2), and (1225.3): [304] for $D \triangleq [d_{ij}]$ and $\sqrt[3]{D} \triangleq [\sqrt{d_{ij}}]$

$$\begin{array}{ll} \underset{D}{\text{minimize}} & \|-V(D - H)V\|_{\text{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & D \in \text{EDM}^N \end{array} \quad \begin{array}{ll} \underset{\sqrt[3]{D}}{\text{minimize}} & \|\sqrt[3]{D} - H\|_{\text{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & \sqrt[3]{D} \in \sqrt{\text{EDM}^N} \end{array} \quad (2) \quad (1225)$$

$$\begin{array}{ll} \underset{D}{\text{minimize}} & \|D - H\|_{\text{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & D \in \text{EDM}^N \end{array} \quad \begin{array}{ll} \underset{\sqrt[3]{D}}{\text{minimize}} & \|-V(\sqrt[3]{D} - H)V\|_{\text{F}}^2 \\ \text{subject to} & \text{rank } VDV \leq \rho \\ & \sqrt[3]{D} \in \sqrt{\text{EDM}^N} \end{array} \quad (4)$$

where we have made explicit an imposed upper bound ρ on affine dimension

$$r = \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} = \text{rank } V D V \quad (957)$$

that is benign when $\rho = N - 1$ or H were realizable with $r \leq \rho$. Problems (1225.2) and (1225.3) are Euclidean projections of a vectorized matrix H on an EDM cone (§6.3), whereas problems (1225.1) and (1225.4) are Euclidean projections of a vectorized matrix $-VHV$ on a PSD cone.^{7.2} Problem (1225.4) is not posed in the literature because it has limited theoretical foundation.^{7.3}

Analytical solution to (1225.1) is known in closed form for any bound ρ although, as the problem is stated, it is a convex optimization only in the case $\rho = N - 1$. We show in §7.1.4 how (1225.1) becomes a convex optimization problem for any ρ when transformed to the spectral domain. When expressed as a function of point list in a matrix X as in (1223), problem (1225.2) is a variant of what is known in statistics literature as the *stress problem*. [46, p.34] [87] [314] Problems (1225.2) and (1225.3) are convex optimization problems in D for the case $\rho = N - 1$, wherein (1225.3) becomes equivalent to (1224). Even with the rank constraint removed from (1225.2), we will see that the convex problem remaining inherently minimizes affine dimension.

Generally speaking, each problem in (1225) produces a different result because there is no isometry relating them. Of the various auxiliary V -matrices (§B.4), the geometric centering matrix V (831) appears in the literature most often although $V_{\mathcal{N}}$ (814) is the auxiliary matrix naturally consequent to Schoenberg's seminal exposition [276]. Substitution of any auxiliary matrix or its pseudoinverse into these problems produces another valid problem.

Substitution of $V_{\mathcal{N}}^T$ for left-hand V in (1225.1), in particular, produces a different result because

$$\begin{aligned} & \underset{D}{\text{minimize}} && \| -V(D - H)V \|_{\text{F}}^2 \\ & \text{subject to} && D \in \text{EDM}^N \end{aligned} \quad (1226)$$

^{7.2}Because $-VHV$ is orthogonal projection of $-H$ on the geometric center subspace \mathbb{S}_c^N (§E.7.2.0.2), problems (1225.1) and (1225.4) may be interpreted as oblique (nonminimum distance) projections of $-H$ on a positive semidefinite cone.

^{7.3} $D \in \text{EDM}^N \Rightarrow \sqrt[N]{D} \in \text{EDM}^N, -V\sqrt[N]{D}V \in \mathbb{S}_+^N$ (§5.10)

finds D to attain Euclidean distance of vectorized $-VHV$ to the positive semidefinite cone in ambient isometrically isomorphic $\mathbb{R}^{N(N+1)/2}$, whereas

$$\begin{aligned} & \underset{D}{\text{minimize}} \quad \|-V_{\mathcal{N}}^{\text{T}}(D - H)V_{\mathcal{N}}\|_{\mathbb{F}}^2 \\ & \text{subject to} \quad D \in \text{EDM}^N \end{aligned} \tag{1227}$$

attains Euclidean distance of vectorized $-V_{\mathcal{N}}^{\text{T}}HV_{\mathcal{N}}$ to the positive semidefinite cone in isometrically isomorphic subspace $\mathbb{R}^{N(N-1)/2}$; quite different projections^{7.4} regardless of whether affine dimension is constrained. But substitution of auxiliary matrix $V_{\mathcal{W}}^{\text{T}}$ (§B.4.3) or $V_{\mathcal{N}}^{\dagger}$ yields the same result as (1225.1) because $V = V_{\mathcal{W}}V_{\mathcal{W}}^{\text{T}} = V_{\mathcal{N}}V_{\mathcal{N}}^{\dagger}$; *id est*,

$$\begin{aligned} \|-V(D - H)V\|_{\mathbb{F}}^2 &= \|-V_{\mathcal{W}}V_{\mathcal{W}}^{\text{T}}(D - H)V_{\mathcal{W}}V_{\mathcal{W}}^{\text{T}}\|_{\mathbb{F}}^2 = \|-V_{\mathcal{W}}^{\text{T}}(D - H)V_{\mathcal{W}}\|_{\mathbb{F}}^2 \\ &= \|-V_{\mathcal{N}}V_{\mathcal{N}}^{\dagger}(D - H)V_{\mathcal{N}}V_{\mathcal{N}}^{\dagger}\|_{\mathbb{F}}^2 = \|-V_{\mathcal{N}}^{\dagger}(D - H)V_{\mathcal{N}}\|_{\mathbb{F}}^2 \end{aligned} \tag{1228}$$

We see no compelling reason to prefer one particular auxiliary V -matrix over another. Each has its own coherent interpretations; *e.g.*, §5.4.2, §6.7. Neither can we say any particular problem formulation produces generally better results than another.^{7.5}

7.1 First prevalent problem: Projection on PSD cone

This first problem

$$\left. \begin{aligned} & \underset{D}{\text{minimize}} \quad \|-V_{\mathcal{N}}^{\text{T}}(D - H)V_{\mathcal{N}}\|_{\mathbb{F}}^2 \\ & \text{subject to} \quad \text{rank } V_{\mathcal{N}}^{\text{T}}DV_{\mathcal{N}} \leq \rho \\ & \quad \quad \quad D \in \text{EDM}^N \end{aligned} \right\} \text{ Problem 1} \tag{1229}$$

^{7.4}The isomorphism $T(Y) = V_{\mathcal{N}}^{\dagger \text{T}}YV_{\mathcal{N}}^{\dagger}$ onto $\mathbb{S}_c^N = \{VXV \mid X \in \mathbb{S}^N\}$ relates the map in (1227) to that in (1226), but is not an isometry.

^{7.5}All four problem formulations (1225) produce identical results when affine dimension r , implicit to a realizable measurement matrix H , does not exceed desired affine dimension ρ ; because, the optimal objective value will vanish ($\|\cdot\| = 0$).

poses a Euclidean projection of $-V_N^T H V_N$ in subspace \mathbb{S}^{N-1} on a generally nonconvex subset (when $\rho < N-1$) of the positive semidefinite cone boundary $\partial\mathbb{S}_+^{N-1}$ whose elemental matrices have rank no greater than desired affine dimension ρ (§5.7.1.1). Problem 1 finds the closest EDM D in the sense of Schoenberg. (827) [276] As it is stated, this optimization problem is convex only when desired affine dimension is largest $\rho = N-1$ although its analytical solution is known [227, thm.14.4.2] for all nonnegative $\rho \leq N-1$.^{7.6}

We assume only that the given measurement matrix H is symmetric,^{7.7}

$$H \in \mathbb{S}^N \quad (1230)$$

Arranging the eigenvalues λ_i of $-V_N^T H V_N$ in nonincreasing order for all i , $\lambda_i \geq \lambda_{i+1}$ with v_i the corresponding i^{th} eigenvector, then an optimal solution to Problem 1 is [313, §2]

$$-V_N^T D^* V_N = \sum_{i=1}^{\rho} \max\{0, \lambda_i\} v_i v_i^T \quad (1231)$$

where

$$-V_N^T H V_N \triangleq \sum_{i=1}^{N-1} \lambda_i v_i v_i^T \in \mathbb{S}^{N-1} \quad (1232)$$

is an eigenvalue decomposition and

$$D^* \in \mathbb{EDM}^N \quad (1233)$$

is an optimal Euclidean distance matrix.

In §7.1.4 we show how to transform Problem 1 to a convex optimization problem for any ρ .

^{7.6} being first pronounced in the context of multidimensional scaling by Mardia [226] in 1978 who attributes the generic result (§7.1.2) to Eckart & Young, 1936 [112].

^{7.7}Projection in Problem 1 is on a rank ρ subset of the positive semidefinite cone \mathbb{S}_+^{N-1} (§2.9.2.1) in the subspace of symmetric matrices \mathbb{S}^{N-1} . It is wrong here to zero the main diagonal of given H because first projecting H on the symmetric hollow subspace places an elbow in the path of projection in Problem 1. (Figure 132)

7.1.1 Closest-EDM Problem 1, convex case

7.1.1.0.1 Proof. *Solution (1231), convex case.*

When desired affine dimension is unconstrained, $\rho = N - 1$, the rank function disappears from (1229) leaving a convex optimization problem; a simple unique minimum-distance projection on the positive semidefinite cone \mathbb{S}_+^{N-1} : *videlicet*

$$\begin{aligned} & \underset{D \in \mathbb{S}_h^N}{\text{minimize}} && \| -V_{\mathcal{N}}^T(D - H)V_{\mathcal{N}} \|_{\mathbb{F}}^2 \\ & \text{subject to} && -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \end{aligned} \quad (1234)$$

by (827). Because

$$\mathbb{S}^{N-1} = -V_{\mathcal{N}}^T \mathbb{S}_h^N V_{\mathcal{N}} \quad (926)$$

then the necessary and sufficient conditions for projection in isometrically isomorphic $\mathbb{R}^{N(N-1)/2}$ on the self-dual (341) positive semidefinite cone \mathbb{S}_+^{N-1} are:^{7.8} (§E.9.2.0.1) (1488) (*confer*(1919))

$$\begin{aligned} & -V_{\mathcal{N}}^T D^* V_{\mathcal{N}} \succeq 0 \\ & -V_{\mathcal{N}}^T D^* V_{\mathcal{N}} (-V_{\mathcal{N}}^T D^* V_{\mathcal{N}} + V_{\mathcal{N}}^T H V_{\mathcal{N}}) = \mathbf{0} \\ & -V_{\mathcal{N}}^T D^* V_{\mathcal{N}} + V_{\mathcal{N}}^T H V_{\mathcal{N}} \succeq 0 \end{aligned} \quad (1235)$$

Symmetric $-V_{\mathcal{N}}^T H V_{\mathcal{N}}$ is diagonalizable hence decomposable in terms of its eigenvectors v and eigenvalues λ as in (1232). Therefore (*confer*(1231))

$$-V_{\mathcal{N}}^T D^* V_{\mathcal{N}} = \sum_{i=1}^{N-1} \max\{0, \lambda_i\} v_i v_i^T \quad (1236)$$

satisfies (1235), optimally solving (1234). To see that, recall: these eigenvectors constitute an orthogonal set and

$$-V_{\mathcal{N}}^T D^* V_{\mathcal{N}} + V_{\mathcal{N}}^T H V_{\mathcal{N}} = - \sum_{i=1}^{N-1} \min\{0, \lambda_i\} v_i v_i^T \quad (1237)$$

◆

^{7.8}These conditions for projection on a convex cone are identical to the Karush-Kuhn-Tucker (KKT) optimality conditions [246, p.328] for problem (1234).

7.1.2 Generic problem

Prior to determination of D^* , analytical solution (1231) to Problem 1 is equivalent to solution of a generic rank-constrained projection problem: Given desired affine dimension ρ and

$$A \triangleq -V_{\mathcal{N}}^T H V_{\mathcal{N}} = \sum_{i=1}^{N-1} \lambda_i v_i v_i^T \in \mathbb{S}^{N-1} \quad (1232)$$

Euclidean projection on a rank ρ subset of a PSD cone (on a generally nonconvex subset of the PSD cone boundary $\partial\mathbb{S}_+^{N-1}$ when $\rho < N-1$)

$$\left. \begin{array}{l} \text{minimize}_{B \in \mathbb{S}^{N-1}} \|B - A\|_{\text{F}}^2 \\ \text{subject to } \text{rank } B \leq \rho \\ B \succeq 0 \end{array} \right\} \text{Generic 1} \quad (1238)$$

has well known optimal solution (Eckart & Young) [112]

$$B^* \triangleq -V_{\mathcal{N}}^T D^* V_{\mathcal{N}} = \sum_{i=1}^{\rho} \max\{0, \lambda_i\} v_i v_i^T \in \mathbb{S}^{N-1} \quad (1231)$$

Once optimal B^* is found, the technique of §5.12 can be used to determine a uniquely corresponding optimal Euclidean distance matrix D^* ; a unique correspondence by injectivity arguments in §5.6.2.

7.1.2.1 Projection on rank ρ subset of PSD cone

Because Problem 1 is the same as

$$\left. \begin{array}{l} \text{minimize}_{D \in \mathbb{S}_h^N} \|-V_{\mathcal{N}}^T (D - H) V_{\mathcal{N}}\|_{\text{F}}^2 \\ \text{subject to } \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} \leq \rho \\ -V_{\mathcal{N}}^T D V_{\mathcal{N}} \succeq 0 \end{array} \right\} \quad (1239)$$

and because (926) provides invertible mapping to the generic problem, then Problem 1 is truly a Euclidean projection of vectorized $-V_{\mathcal{N}}^T H V_{\mathcal{N}}$ on that generally nonconvex subset of symmetric matrices (belonging to the positive semidefinite cone \mathbb{S}_+^{N-1}) having rank no greater than desired affine

dimension ρ ; ^{7.9} called rank ρ subset: (199) (233)

$$\mathbb{S}_+^{N-1} \setminus \mathbb{S}_+^{N-1}(\rho + 1) = \{X \in \mathbb{S}_+^{N-1} \mid \text{rank } X \leq \rho\} \quad (238)$$

7.1.3 Choice of spectral cone

Spectral projection substitutes projection on a polyhedral cone, containing a complete set of eigenspectra, in place of projection on a convex set of diagonalizable matrices; *e.g.*, (1251). In this section we develop a method of spectral projection for constraining rank of positive semidefinite matrices in a proximity problem like (1238). We will see why an orthant turns out to be the best choice of spectral cone, and why presorting is critical.

Define a nonlinear permutation-operator $\pi(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ that sorts its vector argument x into nonincreasing order.

7.1.3.0.1 Definition. Spectral projection.

Let R be an orthogonal matrix and Λ a nonincreasingly ordered diagonal matrix of eigenvalues. *Spectral projection* means unique minimum-distance projection of a rotated (R , §B.5.4) nonincreasingly ordered (π) vector (δ) of eigenvalues

$$\pi(\delta(R^T \Lambda R)) \quad (1240)$$

on a polyhedral cone containing all eigenspectra corresponding to a rank ρ subset of a positive semidefinite cone (§2.9.2.1) or the EDM cone (in Cayley-Menger form, §5.11.2.3). \triangle

In the simplest and most common case, projection on a positive semidefinite cone, orthogonal matrix R equals I (§7.1.4.0.1) and diagonal matrix Λ is ordered during diagonalization (§A.5.2). Then spectral projection simply means projection of $\delta(\Lambda)$ on a subset of the nonnegative orthant, as we shall now ascertain:

It is curious how nonconvex Problem 1 has such a simple analytical solution (1231). Although solution to generic problem (1238) is well known since 1936 [112], its equivalence was observed in 1997 [313, §2] to projection

^{7.9}Recall: affine dimension is a lower bound on embedding (§2.3.1), equal to dimension of the smallest affine set in which points from a list X corresponding to an EDM D can be embedded.

of an ordered vector of eigenvalues (in diagonal matrix Λ) on a subset of the monotone nonnegative cone (§2.13.9.4.2)

$$\mathcal{K}_{\mathcal{M}+} = \{v \mid v_1 \geq v_2 \geq \cdots \geq v_{N-1} \geq 0\} \subseteq \mathbb{R}_+^{N-1} \quad (389)$$

Of interest, momentarily, is only the smallest convex subset of the monotone nonnegative cone $\mathcal{K}_{\mathcal{M}+}$ containing every nonincreasingly ordered eigenspectrum corresponding to a rank ρ subset of the positive semidefinite cone \mathbb{S}_+^{N-1} ; *id est*,

$$\mathcal{K}_{\mathcal{M}+}^\rho \triangleq \{v \in \mathbb{R}^\rho \mid v_1 \geq v_2 \geq \cdots \geq v_\rho \geq 0\} \subseteq \mathbb{R}_+^\rho \quad (1241)$$

a pointed polyhedral cone, a ρ -dimensional convex subset of the monotone nonnegative cone $\mathcal{K}_{\mathcal{M}+} \subseteq \mathbb{R}_+^{N-1}$ having property, for λ denoting eigenspectra,

$$\begin{bmatrix} \mathcal{K}_{\mathcal{M}+}^\rho \\ \mathbf{0} \end{bmatrix} = \pi(\lambda(\text{rank } \rho \text{ subset})) \subseteq \mathcal{K}_{\mathcal{M}+}^{N-1} \triangleq \mathcal{K}_{\mathcal{M}+} \quad (1242)$$

For each and every elemental eigenspectrum

$$\gamma \in \lambda(\text{rank } \rho \text{ subset}) \subseteq \mathbb{R}_+^{N-1} \quad (1243)$$

of the rank ρ subset (ordered or unordered in λ), there is a nonlinear surjection $\pi(\gamma)$ onto $\mathcal{K}_{\mathcal{M}+}^\rho$.

7.1.3.0.2 Exercise. *Smallest spectral cone.*

Prove that there is no convex subset of $\mathcal{K}_{\mathcal{M}+}$ smaller than $\mathcal{K}_{\mathcal{M}+}^\rho$ containing every ordered eigenspectrum corresponding to the rank ρ subset of a positive semidefinite cone (§2.9.2.1). ▼

7.1.3.0.3 Proposition. (Hardy-Littlewood-Pólya) *Inequalities.* [158, §X] [48, §1.2] Any vectors σ and γ in \mathbb{R}^{N-1} satisfy a tight inequality

$$\pi(\sigma)^\top \pi(\gamma) \geq \sigma^\top \gamma \geq \pi(\sigma)^\top \Xi \pi(\gamma) \quad (1244)$$

where Ξ is the order-reversing permutation matrix defined in (1619), and permutator $\pi(\gamma)$ is a nonlinear function that sorts vector γ into nonincreasing order thereby providing the greatest upper bound and least lower bound with respect to every possible sorting. ◇

7.1.3.0.4 Corollary. *Monotone nonnegative sort.*

Any given vectors $\sigma, \gamma \in \mathbb{R}^{N-1}$ satisfy a tight Euclidean distance inequality

$$\|\pi(\sigma) - \pi(\gamma)\| \leq \|\sigma - \gamma\| \quad (1245)$$

where nonlinear function $\pi(\gamma)$ sorts vector γ into nonincreasing order thereby providing the least lower bound with respect to every possible sorting. \diamond

Given $\gamma \in \mathbb{R}^{N-1}$

$$\inf_{\sigma \in \mathbb{R}_+^{N-1}} \|\sigma - \gamma\| = \inf_{\sigma \in \mathbb{R}_+^{N-1}} \|\pi(\sigma) - \pi(\gamma)\| = \inf_{\sigma \in \mathbb{R}_+^{N-1}} \|\sigma - \pi(\gamma)\| = \inf_{\sigma \in \mathcal{K}_{\mathcal{M}+}} \|\sigma - \pi(\gamma)\| \quad (1246)$$

Yet for γ representing an arbitrary vector of eigenvalues, because

$$\inf_{\sigma \in \begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix}} \|\sigma - \gamma\|^2 \geq \inf_{\sigma \in \begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix}} \|\sigma - \pi(\gamma)\|^2 = \inf_{\sigma \in \begin{bmatrix} \mathcal{K}_{\mathcal{M}+}^\rho \\ \mathbf{0} \end{bmatrix}} \|\sigma - \pi(\gamma)\|^2 \quad (1247)$$

then projection of γ on the eigenspectra of a rank ρ subset can be tightened simply by presorting γ into nonincreasing order.

Proof. Simply because $\pi(\gamma)_{1:\rho} \succeq \pi(\gamma_{1:\rho})$

$$\begin{aligned} \inf_{\sigma \in \begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix}} \|\sigma - \gamma\|^2 &= \gamma_{\rho+1:N-1}^T \gamma_{\rho+1:N-1} + \inf_{\sigma \in \mathbb{R}_+^{N-1}} \|\sigma_{1:\rho} - \gamma_{1:\rho}\|^2 \\ &= \gamma^T \gamma + \inf_{\sigma \in \mathbb{R}_+^{N-1}} \sigma_{1:\rho}^T \sigma_{1:\rho} - 2\sigma_{1:\rho}^T \gamma_{1:\rho} \\ &\geq \gamma^T \gamma + \inf_{\sigma \in \mathbb{R}_+^{N-1}} \sigma_{1:\rho}^T \sigma_{1:\rho} - 2\sigma_{1:\rho}^T \pi(\gamma)_{1:\rho} \end{aligned} \quad (1248)$$

$$\inf_{\sigma \in \begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix}} \|\sigma - \gamma\|^2 \geq \inf_{\sigma \in \begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix}} \|\sigma - \pi(\gamma)\|^2$$

\blacklozenge

7.1.3.1 Orthant is best spectral cone for Problem 1

This means unique minimum-distance projection of γ on the nearest spectral member of the rank ρ subset is tantamount to presorting γ into nonincreasing order. Only then does unique spectral projection on a subset $\mathcal{K}_{\mathcal{M}^+}^\rho$ of the monotone nonnegative cone become equivalent to unique spectral projection on a subset \mathbb{R}_+^ρ of the nonnegative orthant (which is simpler); in other words, unique minimum-distance projection of sorted γ on the nonnegative orthant in a ρ -dimensional subspace of \mathbb{R}^N is indistinguishable from its projection on the subset $\mathcal{K}_{\mathcal{M}^+}^\rho$ of the monotone nonnegative cone in that same subspace.

7.1.4 Closest-EDM Problem 1, “nonconvex” case

Proof of solution (1231), for projection on a rank ρ subset of the positive semidefinite cone \mathbb{S}_+^{N-1} , can be algebraic in nature. [313, §2] Here we derive that known result but instead using a more geometric argument via spectral projection on a polyhedral cone (subsuming the proof in §7.1.1). In so doing, we demonstrate how nonconvex Problem 1 is transformed to a convex optimization:

7.1.4.0.1 Proof. Solution (1231), nonconvex case.

As explained in §7.1.2, we may instead work with the more facile generic problem (1238). With diagonalization of unknown

$$B \triangleq U\Upsilon U^T \in \mathbb{S}^{N-1} \quad (1249)$$

given desired affine dimension $0 \leq \rho \leq N-1$ and diagonalizable

$$A \triangleq Q\Lambda Q^T \in \mathbb{S}^{N-1} \quad (1250)$$

having eigenvalues in Λ arranged in nonincreasing order, by (42) the generic problem is equivalent to

$$\begin{array}{ll} \underset{B \in \mathbb{S}^{N-1}}{\text{minimize}} & \|B - A\|_{\mathbb{F}}^2 \\ \text{subject to} & \text{rank } B \leq \rho \\ & B \succeq 0 \end{array} \quad \equiv \quad \begin{array}{ll} \underset{R, \Upsilon}{\text{minimize}} & \|\Upsilon - R^T \Lambda R\|_{\mathbb{F}}^2 \\ \text{subject to} & \text{rank } \Upsilon \leq \rho \\ & \Upsilon \succeq 0 \\ & R^{-1} = R^T \end{array} \quad (1251)$$

where

$$R \triangleq Q^T U \in \mathbb{R}^{N-1 \times N-1} \quad (1252)$$

in U on the set of orthogonal matrices is a linear bijection. We propose solving (1251) by instead solving the problem sequence:

$$\begin{aligned} & \underset{\Upsilon}{\text{minimize}} && \|\Upsilon - R^T \Lambda R\|_{\mathbb{F}}^2 \\ & \text{subject to} && \text{rank } \Upsilon \leq \rho \\ & && \Upsilon \succeq 0 \end{aligned} \quad (\text{a}) \quad (1253)$$

$$\begin{aligned} & \underset{R}{\text{minimize}} && \|\Upsilon^* - R^T \Lambda R\|_{\mathbb{F}}^2 \\ & \text{subject to} && R^{-1} = R^T \end{aligned} \quad (\text{b})$$

Problem (1253a) is equivalent to: (1) orthogonal projection of $R^T \Lambda R$ on an $N-1$ -dimensional subspace of isometrically isomorphic $\mathbb{R}^{N(N-1)/2}$ containing $\delta(\Upsilon) \in \mathbb{R}_+^{N-1}$, (2) nonincreasingly ordering the result, (3) unique minimum-distance projection of the ordered result on $\begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix}$. (§E.9.5)

Projection on that $N-1$ -dimensional subspace amounts to zeroing $R^T \Lambda R$ at all entries off the main diagonal; thus, the equivalent sequence leading with a spectral projection:

$$\begin{aligned} & \underset{\Upsilon}{\text{minimize}} && \|\delta(\Upsilon) - \pi(\delta(R^T \Lambda R))\|^2 \\ & \text{subject to} && \delta(\Upsilon) \in \begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix} \end{aligned} \quad (\text{a}) \quad (1254)$$

$$\begin{aligned} & \underset{R}{\text{minimize}} && \|\Upsilon^* - R^T \Lambda R\|_{\mathbb{F}}^2 \\ & \text{subject to} && R^{-1} = R^T \end{aligned} \quad (\text{b})$$

Because any permutation matrix is an orthogonal matrix, it is always feasible that $\delta(R^T \Lambda R) \in \mathbb{R}^{N-1}$ be arranged in nonincreasing order; hence, the permutation operator π . Unique minimum-distance projection of vector $\pi(\delta(R^T \Lambda R))$ on the ρ -dimensional subset $\begin{bmatrix} \mathbb{R}_+^\rho \\ \mathbf{0} \end{bmatrix}$ of nonnegative orthant

\mathbb{R}_+^{N-1} requires: (§E.9.2.0.1)

$$\begin{aligned} \delta(\Upsilon^*)_{\rho+1:N-1} &= \mathbf{0} \\ \delta(\Upsilon^*) &\succeq 0 \\ \delta(\Upsilon^*)^T(\delta(\Upsilon^*) - \pi(\delta(R^T\Lambda R))) &= 0 \\ \delta(\Upsilon^*) - \pi(\delta(R^T\Lambda R)) &\succeq 0 \end{aligned} \tag{1255}$$

which are necessary and sufficient conditions. Any value Υ^* satisfying conditions (1255) is optimal for (1254a). So

$$\delta(\Upsilon^*)_i = \begin{cases} \max\{0, \pi(\delta(R^T\Lambda R))_i\}, & i=1 \dots \rho \\ 0, & i=\rho+1 \dots N-1 \end{cases} \tag{1256}$$

specifies an optimal solution. The lower bound on the objective with respect to R in (1254b) is tight; by (1222)

$$\| |\Upsilon^*| - |\Lambda| \|_F \leq \|\Upsilon^* - R^T\Lambda R\|_F \tag{1257}$$

where $|\cdot|$ denotes absolute entry-value. For selection of Υ^* as in (1256), this lower bound is attained when (*confer* §C.4.2.2)

$$R^* = I \tag{1258}$$

which is the known solution. \blacklozenge

7.1.4.1 Significance

Importance of this well-known [112] optimal solution (1231) for projection on a rank ρ subset of a positive semidefinite cone should not be dismissed:

- Problem 1, as stated, is generally nonconvex. This analytical solution at once encompasses projection on a rank ρ subset (238) of the positive semidefinite cone (generally, a nonconvex subset of its boundary) from either the exterior or interior of that cone.^{7.10} By problem transformation to the spectral domain, projection on a rank ρ subset becomes a convex optimization problem.

^{7.10}Projection on the boundary from the interior of a convex Euclidean body is generally a nonconvex problem.

- This solution is closed form.
- This solution is equivalent to projection on a polyhedral cone in the spectral domain (spectral projection, projection on a spectral cone §7.1.3.0.1); a necessary and sufficient condition (§A.3.1) for membership of a symmetric matrix to a rank ρ subset of a positive semidefinite cone (§2.9.2.1).
- Because $U^* = Q$, a minimum-distance projection on a rank ρ subset of the positive semidefinite cone is a positive semidefinite matrix orthogonal (in the Euclidean sense) to direction of projection.^{7.11}
- For the convex case problem, this solution is always unique. Otherwise, distinct eigenvalues (multiplicity 1) in Λ guarantees uniqueness of this solution by the reasoning in §A.5.0.1.^{7.12}

7.1.5 Problem 1 in spectral norm, convex case

When instead we pose the matrix 2-norm (spectral norm) in Problem 1 (1229) for the convex case $\rho = N - 1$, then the new problem

$$\begin{aligned} & \underset{D}{\text{minimize}} && \| -V_{\mathcal{N}}^T(D - H)V_{\mathcal{N}} \|_2 \\ & \text{subject to} && D \in \text{EDM}^N \end{aligned} \quad (1259)$$

is convex although its solution is not necessarily unique;^{7.13} giving rise to nonorthogonal projection (§E.1) on the positive semidefinite cone \mathbb{S}_+^{N-1} . Indeed, its solution set includes the Frobenius solution (1231) for the convex case whenever $-V_{\mathcal{N}}^T H V_{\mathcal{N}}$ is a normal matrix. [163, §1] [156] [53, §8.1.1] *Singular value problem* (1259) is equivalent to

$$\begin{aligned} & \underset{\mu, D}{\text{minimize}} && \mu \\ & \text{subject to} && -\mu I \preceq -V_{\mathcal{N}}^T(D - H)V_{\mathcal{N}} \preceq \mu I \\ & && D \in \text{EDM}^N \end{aligned} \quad (1260)$$

^{7.11}But Theorem E.9.2.0.1 for unique projection on a closed convex cone does not apply here because the direction of projection is not necessarily a member of the dual PSD cone. This occurs, for example, whenever positive eigenvalues are truncated.

^{7.12}Uncertainty of uniqueness prevents the erroneous conclusion that a rank ρ subset (199) were a convex body by the *Bunt-Motzkin theorem* (§E.9.0.0.1).

^{7.13}For each and every $|t| \leq 2$, for example, $\begin{bmatrix} 2 & 0 \\ 0 & t \end{bmatrix}$ has the same spectral-norm value.

by (1596) where

$$\mu^* = \max_i \{ |\lambda(-V_{\mathcal{N}}^T(D^* - H)V_{\mathcal{N}})_i|, \quad i = 1 \dots N-1 \} \in \mathbb{R}_+ \quad (1261)$$

the minimized largest absolute eigenvalue (due to matrix symmetry).

For lack of a unique solution here, we prefer the Frobenius rather than spectral norm.

7.2 Second prevalent problem: Projection on EDM cone in $\sqrt{d_{ij}}$

Let

$$\overset{\circ}{\sqrt{D}} \triangleq [\sqrt{d_{ij}}] \in \mathcal{K} = \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N} \quad (1262)$$

be an unknown matrix of absolute distance; *id est*,

$$D = [d_{ij}] \triangleq \overset{\circ}{\sqrt{D}} \circ \overset{\circ}{\sqrt{D}} \in \mathbb{EDM}^N \quad (1263)$$

where \circ denotes Hadamard product. The second prevalent proximity problem is a Euclidean projection (in the natural coordinates $\sqrt{d_{ij}}$) of matrix H on a nonconvex subset of the boundary of the nonconvex cone of Euclidean absolute-distance matrices $\text{rel } \partial \sqrt{\mathbb{EDM}^N}$: (§6.3, confer Figure 117(b))

$$\left. \begin{array}{l} \underset{\overset{\circ}{\sqrt{D}}}{\text{minimize}} \quad \|\overset{\circ}{\sqrt{D}} - H\|_{\text{F}}^2 \\ \text{subject to} \quad \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} \leq \rho \\ \quad \quad \quad \overset{\circ}{\sqrt{D}} \in \sqrt{\mathbb{EDM}^N} \end{array} \right\} \text{Problem 2} \quad (1264)$$

where

$$\sqrt{\mathbb{EDM}^N} = \{ \overset{\circ}{\sqrt{D}} \mid D \in \mathbb{EDM}^N \} \quad (1097)$$

This statement of the second proximity problem is considered difficult to solve because of the constraint on desired affine dimension ρ (§5.7.2) and because the objective function

$$\|\overset{\circ}{\sqrt{D}} - H\|_{\text{F}}^2 = \sum_{i,j} (\sqrt{d_{ij}} - h_{ij})^2 \quad (1265)$$

is expressed in the natural coordinates; projection on a doubly nonconvex set.

Our solution to this second problem prevalent in the literature requires measurement matrix H to be nonnegative;

$$H = [h_{ij}] \in \mathbb{R}_+^{N \times N} \quad (1266)$$

If the H matrix given has negative entries, then the technique of solution presented here becomes invalid. As explained in §7.0.1, projection of H on $\mathcal{K} = \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N}$ (1216) prior to application of this proposed solution is incorrect.

7.2.1 Convex case

When $\rho = N - 1$, the rank constraint vanishes and a convex problem that is equivalent to (1223) emerges.^{7.14}

$$\begin{aligned} \underset{\sqrt[D]{D}}{\text{minimize}} \quad & \|\sqrt[D]{D} - H\|_{\mathbb{F}}^2 & \Leftrightarrow & \underset{D}{\text{minimize}} \quad \sum_{i,j} d_{ij} - 2h_{ij}\sqrt{d_{ij}} + h_{ij}^2 \\ \text{subject to} \quad & \sqrt[D]{D} \in \sqrt{\text{EDM}^N} & & \text{subject to} \quad D \in \text{EDM}^N \end{aligned} \quad (1267)$$

For any fixed i and j , the argument of summation is a convex function of d_{ij} because (for nonnegative constant h_{ij}) the negative square root is convex in nonnegative d_{ij} and because $d_{ij} + h_{ij}^2$ is affine (convex). Because the sum of any number of convex functions in D remains convex [53, §3.2.1] and because the feasible set is convex in D , we have a convex optimization problem:

$$\begin{aligned} \underset{D}{\text{minimize}} \quad & \mathbf{1}^T(D - 2H \circ \sqrt[D]{D})\mathbf{1} + \|H\|_{\mathbb{F}}^2 \\ \text{subject to} \quad & D \in \text{EDM}^N \end{aligned} \quad (1268)$$

The objective function being a sum of strictly convex functions is, moreover, strictly convex in D on the nonnegative orthant. Existence of a unique solution D^* for this second prevalent problem depends upon nonnegativity of H and a convex feasible set (§3.1.2).^{7.15}

^{7.14} still thought to be a nonconvex problem as late as 1997 [314] even though discovered convex by de Leeuw in 1993. [87] [46, §13.6] Yet using methods from §3, it can be easily ascertained: $\|\sqrt[D]{D} - H\|_{\mathbb{F}}$ is not convex in D .

^{7.15}The transformed problem in variable D no longer describes Euclidean projection on an EDM cone. Otherwise we might erroneously conclude $\sqrt{\text{EDM}^N}$ were a convex body by the *Bunt-Motzkin theorem* (§E.9.0.0.1).

7.2.1.1 Equivalent semidefinite program, Problem 2, convex case

Convex problem (1267) is numerically solvable for its global minimum using an interior-point method [53, §11] [350] [256] [244] [341] [10] [126]. We translate (1267) to an equivalent semidefinite program (SDP) for a pedagogical reason made clear in §7.2.2.2 and because there exist readily available computer programs for numerical solution [146] [343] [344] [319] [31] [342].

Substituting a new matrix variable $Y \triangleq [y_{ij}] \in \mathbb{R}_+^{N \times N}$

$$h_{ij}\sqrt{d_{ij}} \leftarrow y_{ij} \quad (1269)$$

Boyd proposes: problem (1267) is equivalent to the semidefinite program

$$\begin{aligned} & \text{minimize} && \sum_{i,j} d_{ij} - 2y_{ij} + h_{ij}^2 \\ & \text{subject to} && \begin{bmatrix} d_{ij} & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0, \quad i, j = 1 \dots N \\ & && D \in \mathbb{EDM}^N \end{aligned} \quad (1270)$$

To see that, recall $d_{ij} \geq 0$ is implicit to $D \in \mathbb{EDM}^N$ (§5.8.1, (827)). So when $H \in \mathbb{R}_+^{N \times N}$ is nonnegative as assumed,

$$\begin{bmatrix} d_{ij} & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0 \Leftrightarrow h_{ij}\sqrt{d_{ij}} \geq \sqrt{y_{ij}^2} \quad (1271)$$

Minimization of the objective function implies maximization of y_{ij} that is bounded above. Hence nonnegativity of y_{ij} is implicit to (1270) and, as desired, $y_{ij} \rightarrow h_{ij}\sqrt{d_{ij}}$ as optimization proceeds. \blacklozenge

If the given matrix H is now assumed symmetric and nonnegative,

$$H = [h_{ij}] \in \mathbb{S}^N \cap \mathbb{R}_+^{N \times N} \quad (1272)$$

then $Y = H \circ \sqrt{\mathring{D}}$ must belong to $\mathcal{K} = \mathbb{S}_h^N \cap \mathbb{R}_+^{N \times N}$ (1216). Because $Y \in \mathbb{S}_h^N$ (§B.4.2 no.20), then

$$\|\sqrt{\mathring{D}} - H\|_{\mathbb{F}}^2 = \sum_{i,j} d_{ij} - 2y_{ij} + h_{ij}^2 = -N \operatorname{tr}(V(D - 2Y)V) + \|H\|_{\mathbb{F}}^2 \quad (1273)$$

So convex problem (1270) is equivalent to the semidefinite program

$$\begin{aligned}
& \underset{D, Y}{\text{minimize}} && -\text{tr}(V(D - 2Y)V) \\
& \text{subject to} && \begin{bmatrix} d_{ij} & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\
& && Y \in \mathbb{S}_h^N \\
& && D \in \mathbb{EDM}^N
\end{aligned} \tag{1274}$$

where the constants h_{ij}^2 and N have been dropped arbitrarily from the objective.

7.2.1.2 Gram-form semidefinite program, Problem 2, convex case

There is great advantage to expressing problem statement (1274) in Gram-form because Gram matrix G is a bidirectional bridge between point list X and distance matrix D ; *e.g.*, Example 5.4.2.2.4, Example 6.4.0.0.1. This way, problem convexity can be maintained while simultaneously constraining point list X , Gram matrix G , and distance matrix D at our discretion.

Convex problem (1274) may be equivalently written via linear bijective (§5.6.1) EDM operator $\mathbf{D}(G)$ (820);

$$\begin{aligned}
& \underset{G \in \mathbb{S}_c^N, Y \in \mathbb{S}_h^N}{\text{minimize}} && -\text{tr}(V(\mathbf{D}(G) - 2Y)V) \\
& \text{subject to} && \begin{bmatrix} \langle \Phi_{ij}, G \rangle & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\
& && G \succeq 0
\end{aligned} \tag{1275}$$

where distance-square $D = [d_{ij}] \in \mathbb{S}_h^N$ (804) is related to Gram matrix entries $G = [g_{ij}] \in \mathbb{S}_c^N \cap \mathbb{S}_+^N$ by

$$\begin{aligned}
d_{ij} &= g_{ii} + g_{jj} - 2g_{ij} \\
&= \langle \Phi_{ij}, G \rangle
\end{aligned} \tag{819}$$

where

$$\Phi_{ij} = (e_i - e_j)(e_i - e_j)^T \in \mathbb{S}_+^N \tag{806}$$

Confinement of G to the geometric center subspace provides numerical stability and no loss of generality (*confer* (1102)); implicit constraint $G\mathbf{1} = \mathbf{0}$ is otherwise unnecessary.

To include constraints on the list $X \in \mathbb{R}^{n \times N}$, we would first rewrite (1275)

$$\begin{aligned}
 & \underset{G \in \mathbb{S}_c^N, Y \in \mathbb{S}_h^N, X \in \mathbb{R}^{n \times N}}{\text{minimize}} && -\text{tr}(V(\mathbf{D}(G) - 2Y)V) \\
 & \text{subject to} && \begin{bmatrix} \langle \Phi_{ij}, G \rangle & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\
 & && \begin{bmatrix} I & X \\ X^T & G \end{bmatrix} \succeq 0 \\
 & && X \in \mathcal{C}
 \end{aligned} \tag{1276}$$

and then add the constraints, realized here in abstract membership to some convex set \mathcal{C} . This problem realization includes a convex relaxation of the nonconvex constraint $G = X^T X$ and, if desired, more constraints on G could be added. This technique is discussed in §5.4.2.2.4.

7.2.2 Minimization of affine dimension in Problem 2

When desired affine dimension ρ is diminished, the rank function becomes reinserted into problem (1270) that is then rendered difficult to solve because the feasible set $\{D, Y\}$ loses convexity in $\mathbb{S}_h^N \times \mathbb{R}^{N \times N}$. Indeed, the rank function is quasiconcave (§3.3) on the positive semidefinite cone; (§2.9.2.6.2) *id est*, its sublevel sets are not convex.

7.2.2.1 Rank minimization heuristic

A remedy developed in [232] [118] [119] [117] introduces convex envelope (cenv) of the quasiconcave rank function: (Figure 133)

7.2.2.1.1 Definition. *Convex envelope.* [176]

The convex envelope of a function $f: \mathcal{C} \rightarrow \mathbb{R}$ is defined as the largest convex function g such that $g \leq f$ on convex domain $\mathcal{C} \subseteq \mathbb{R}^n$. ^{7.16} \triangle

^{7.16} Provided $f \not\equiv +\infty$ and there exists an affine function $h \leq f$ on \mathbb{R}^n , then the convex envelope is equal to the convex conjugate (the *Legendre-Fenchel transform*) of the convex conjugate of f ; *id est*, the conjugate-conjugate function f^{**} . [177, §E.1]




Figure 133: Abstraction of convex envelope of rank function. Rank is a quasiconcave function on the positive semidefinite cone, but its convex envelope is the smallest convex function enveloping it. Vertical bar labelled \mathbf{g} illustrates a trace/rank gap; *id est*, rank found exceeds estimate (by 2).

- [118] [117] Convex envelope of rank function: for σ_i a singular value,

$$\text{cenv}(\text{rank } A) \text{ on } \{A \in \mathbb{R}^{m \times n} \mid \|A\|_2 \leq \kappa\} = \frac{1}{\kappa} \sum_i \sigma(A)_i \quad (1277)$$

$$\text{cenv}(\text{rank } A) \text{ on } \{A \in \mathbb{S}_+^n \mid \|A\|_2 \leq \kappa\} = \frac{1}{\kappa} \text{tr}(A) \quad (1278)$$

A properly scaled trace thus represents the best convex lower bound on rank for positive semidefinite matrices. The idea, then, is to substitute the convex envelope for rank of some variable $A \in \mathbb{S}_+^M$ (§A.6.5)

$$\text{rank } A \leftarrow \text{cenv}(\text{rank } A) \propto \text{tr } A = \sum_i \sigma(A)_i = \sum_i \lambda(A)_i = \|\lambda(A)\|_1 \quad (1279)$$

which is equivalent to the sum of all eigenvalues or singular values.

- [117] Convex envelope of the cardinality function is proportional to the 1-norm:

$$\text{cenv}(\text{card } x) \text{ on } \{x \in \mathbb{R}^n \mid \|x\|_\infty \leq \kappa\} = \frac{1}{\kappa} \|x\|_1 \quad (1280)$$

7.2.2.2 Applying trace rank-heuristic to Problem 2

Substituting rank envelope for rank function in Problem 2, for $D \in \text{EDM}^N$ (confer (957))

$$\text{cenv rank}(-V_{\mathcal{N}}^T D V_{\mathcal{N}}) = \text{cenv rank}(-V D V) \propto -\text{tr}(V D V) \quad (1281)$$

and for desired affine dimension $\rho \leq N-1$ and nonnegative H [sic] we get a convex optimization problem

$$\begin{aligned} & \underset{D}{\text{minimize}} && \|\sqrt{D} - H\|_{\text{F}}^2 \\ & \text{subject to} && -\text{tr}(V D V) \leq \kappa \rho \\ & && D \in \text{EDM}^N \end{aligned} \quad (1282)$$

where $\kappa \in \mathbb{R}_+$ is a constant determined by cut-and-try. The equivalent semidefinite program makes κ variable: for nonnegative and symmetric H

$$\begin{aligned} & \underset{D, Y, \kappa}{\text{minimize}} && \kappa \rho + 2 \text{tr}(V Y V) \\ & \text{subject to} && \begin{bmatrix} d_{ij} & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\ & && -\text{tr}(V D V) \leq \kappa \rho \\ & && Y \in \mathbb{S}_h^N \\ & && D \in \text{EDM}^N \end{aligned} \quad (1283)$$

which is the same as (1274), the problem with no explicit constraint on affine dimension. As the present problem is stated, the desired affine dimension ρ yields to the variable scale factor κ ; ρ is effectively ignored.

Yet this result is an illuminant for problem (1274) and it equivalents (all the way back to (1267)): When the given measurement matrix H is nonnegative and symmetric, finding the closest EDM D as in problem (1267), (1270), or (1274) implicitly entails minimization of affine dimension (confer §5.8.4, §5.14.4). Those non-rank-constrained problems are each inherently equivalent to $\text{cenv}(\text{rank})$ -minimization problem (1283), in other words, and their optimal solutions are unique because of the strictly convex objective function in (1267).

7.2.2.3 Rank-heuristic insight

Minimization of affine dimension by use of this trace rank-heuristic (1281) tends to find the list configuration of least energy; rather, it tends to optimize

compaction of the reconstruction by minimizing total distance. (833) It is best used where some physical equilibrium implies such an energy minimization; *e.g.*, [312, §5].

For this Problem 2, the trace rank-heuristic arose naturally in the objective in terms of V . We observe: V (in contrast to $V_{\mathcal{N}}^T$) spreads energy over all available distances (§B.4.2 no.20, *confer* no.22) although the rank function itself is insensitive to choice of auxiliary matrix.

7.2.2.4 Rank minimization heuristic beyond convex envelope

Fazel, Hindi, & Boyd [119] [346] [120] propose a rank heuristic more potent than trace (1279) for problems of rank minimization;

$$\text{rank } Y \leftarrow \log \det(Y + \varepsilon I) \quad (1284)$$

the concave surrogate function $\log \det$ in place of quasiconcave $\text{rank } Y$ (§2.9.2.6.2) when $Y \in \mathbb{S}_+^n$ is variable and where ε is a small positive constant. They propose minimization of the surrogate by substituting a sequence comprising infima of a linearized surrogate about the current estimate Y_i ; *id est*, from the first-order Taylor series expansion about Y_i on some open interval of $\|Y\|_2$ (§D.1.7)

$$\log \det(Y + \varepsilon I) \approx \log \det(Y_i + \varepsilon I) + \text{tr}((Y_i + \varepsilon I)^{-1}(Y - Y_i)) \quad (1285)$$

we make the surrogate sequence of infima over bounded convex feasible set \mathcal{C}

$$\arg \inf_{Y \in \mathcal{C}} \text{rank } Y \leftarrow \lim_{i \rightarrow \infty} Y_{i+1} \quad (1286)$$

where, for $i = 0 \dots$

$$Y_{i+1} = \arg \inf_{Y \in \mathcal{C}} \text{tr}((Y_i + \varepsilon I)^{-1}Y) \quad (1287)$$

Choosing $Y_0 = I$, the first step becomes equivalent to finding the infimum of $\text{tr } Y$; the trace rank-heuristic (1279). The intuition underlying (1287) is the new term in the argument of trace; specifically, $(Y_i + \varepsilon I)^{-1}$ weights Y so that relatively small eigenvalues of Y found by the infimum are made even smaller.

To see that, substitute the nonincreasingly ordered diagonalizations

$$Y_i + \varepsilon I \triangleq Q(\Lambda + \varepsilon I)Q^T \quad (\text{a}) \tag{1288}$$

$$Y \triangleq U\Upsilon U^T \quad (\text{b})$$

into (1287). Then from (1593) we have,

$$\begin{aligned} \inf_{\Upsilon \in U^* \mathcal{C} U^*} \delta((\Lambda + \varepsilon I)^{-1})^T \delta(\Upsilon) &= \inf_{\Upsilon \in U^* \mathcal{C} U} \inf_{R^T = R^{-1}} \text{tr}((\Lambda + \varepsilon I)^{-1} R^T \Upsilon R) \\ &\leq \inf_{Y \in \mathcal{C}} \text{tr}((Y_i + \varepsilon I)^{-1} Y) \end{aligned} \tag{1289}$$

where $R \triangleq Q^T U$ in U on the set of orthogonal matrices is a bijection. The role of ε is, therefore, to limit the maximum weight; the smallest entry on the main diagonal of Υ gets the largest weight. \blacklozenge

7.2.2.5 Applying log det rank-heuristic to Problem 2

When the log det rank-heuristic is inserted into Problem 2, problem (1283) becomes the problem sequence in i

$$\begin{aligned} &\underset{D, Y, \kappa}{\text{minimize}} && \kappa \rho + 2 \text{tr}(VYV) \\ &\text{subject to} && \begin{bmatrix} d_{jl} & y_{jl} \\ y_{jl} & h_{jl}^2 \end{bmatrix} \succeq 0, \quad l > j = 1 \dots N-1 \\ &&& -\text{tr}((-VD_iV + \varepsilon I)^{-1}VDV) \leq \kappa \rho \\ &&& Y \in \mathbb{S}_h^N \\ &&& D \in \text{EDM}^N \end{aligned} \tag{1290}$$

where $D_{i+1} \triangleq D^* \in \text{EDM}^N$, and $D_0 \triangleq \mathbf{1}\mathbf{1}^T - I$.

7.2.2.6 Tightening this log det rank-heuristic

Like the trace method, this log det technique for constraining rank offers no provision for meeting a predetermined upper bound ρ . Yet since the eigenvalues of the sum are simply determined, $\lambda(Y_i + \varepsilon I) = \delta(\Lambda + \varepsilon I)$, we may certainly force selected weights to ε^{-1} by manipulating diagonalization (1288a). Empirically we find this sometimes leads to better results, although affine dimension of a solution cannot be guaranteed.

7.2.2.7 Cumulative summary of rank heuristics

We have studied the *perturbation method* of rank reduction in §4.3, as well as the trace heuristic (convex envelope method §7.2.2.1.1) and log det heuristic in §7.2.2.4. There is another good contemporary method called LMIRank [252] based on alternating projection (§E.10) that does not solve the *ball packing* problem presented in §5.4.2.2.3.

7.2.2.7.1 Example. Rank regularization enforcing affine dimension.

We apply the *convex iteration* method from §4.4.1 to numerically solve an instance of Problem 2; a method empirically superior to the foregoing convex envelope and log det heuristics.

Unidimensional scaling, [89] a historically practical application of multidimensional scaling (§5.12), entails solution of an optimization problem having local minima whose multiplicity varies as the factorial of point-list cardinality. Geometrically, it means finding a list constrained to lie in one affine dimension. In terms of point list, the nonconvex problem is: given nonnegative symmetric matrix $H = [h_{ij}] \in \mathbb{S}^N \cap \mathbb{R}_+^{N \times N}$ (1272) whose entries h_{ij} are all known,

$$\underset{\{x_i \in \mathbb{R}\}}{\text{minimize}} \sum_{i,j=1}^N (|x_i - x_j| - h_{ij})^2 \quad (1223)$$

called a *raw stress* problem [46, p.34] which has an implicit constraint on dimensional embedding of points $\{x_i \in \mathbb{R}, i=1 \dots N\}$. This problem has proven NP-hard; *e.g.*, [64].

As always, we first transform variables to distance-square $D \in \mathbb{S}_h^N$; so we begin with convex problem (1274) on page 521

$$\begin{aligned} & \underset{D, Y}{\text{minimize}} \quad -\text{tr}(V(D - 2Y)V) \\ & \text{subject to} \quad \begin{bmatrix} d_{ij} & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\ & \quad Y \in \mathbb{S}_h^N \\ & \quad D \in \text{EDM}^N \\ & \quad \text{rank } V_{\mathcal{N}}^T D V_{\mathcal{N}} = 1 \end{aligned} \quad (1291)$$

that becomes equivalent to (1223) by making explicit the constraint on affine dimension via rank. The iteration is formed by moving the dimensional

constraint to the objective:

$$\begin{aligned}
 & \underset{D, Y}{\text{minimize}} && -\langle V(D - 2Y)V, I \rangle - w \langle V_{\mathcal{N}}^T D V_{\mathcal{N}}, W \rangle \\
 & \text{subject to} && \begin{bmatrix} d_{ij} & y_{ij} \\ y_{ij} & h_{ij}^2 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\
 & && Y \in \mathbb{S}_h^N \\
 & && D \in \text{EDM}^N
 \end{aligned} \tag{1292}$$

where w (≈ 10) is a positive scalar just large enough to make $\langle V_{\mathcal{N}}^T D V_{\mathcal{N}}, W \rangle$ vanish to within some numerical precision, and where direction matrix W is an optimal solution to semidefinite program (1592a)

$$\begin{aligned}
 & \underset{W}{\text{minimize}} && -\langle V_{\mathcal{N}}^T D^* V_{\mathcal{N}}, W \rangle \\
 & \text{subject to} && 0 \preceq W \preceq I \\
 & && \text{tr } W = N - 1
 \end{aligned} \tag{1293}$$

one of which is known in closed form. Semidefinite programs (1292) and (1293) are iterated until convergence in the sense defined on page 282. This iteration is not a projection method. Convex problem (1292) is neither a relaxation of unidimensional scaling problem (1291); instead, problem (1292) is a convex equivalent to (1291) at convergence of the iteration.

Jan de Leeuw provided us with some test data

$$H = \begin{bmatrix} 0.000000 & 5.235301 & 5.499274 & 6.404294 & 6.486829 & 6.263265 \\ 5.235301 & 0.000000 & 3.208028 & 5.840931 & 3.559010 & 5.353489 \\ 5.499274 & 3.208028 & 0.000000 & 5.679550 & 4.020339 & 5.239842 \\ 6.404294 & 5.840931 & 5.679550 & 0.000000 & 4.862884 & 4.543120 \\ 6.486829 & 3.559010 & 4.020339 & 4.862884 & 0.000000 & 4.618718 \\ 6.263265 & 5.353489 & 5.239842 & 4.543120 & 4.618718 & 0.000000 \end{bmatrix} \tag{1294}$$

and a globally optimal solution

$$\begin{aligned}
 X^* &= [-4.981494 \quad -2.121026 \quad -1.038738 \quad 4.555130 \quad 0.764096 \quad 2.822032] \\
 &= [\quad x_1^* \quad \quad x_2^* \quad \quad x_3^* \quad \quad x_4^* \quad \quad x_5^* \quad \quad x_6^*]
 \end{aligned} \tag{1295}$$

found by searching $6!$ local minima of (1223) [89]. By iterating convex problems (1292) and (1293) about twenty times (initial $W = \mathbf{0}$) we find the

global infimum 98.12812 of stress problem (1223), and by (1047) we find a corresponding one-dimensional point list that is a rigid transformation in \mathbb{R} of X^* .

Here we found the infimum to accuracy of the given data, but that ceases to hold as problem size increases. Because of machine numerical precision and an interior-point method of solution, we speculate, accuracy degrades quickly as problem size increases beyond this. \square

7.3 Third prevalent problem:

Projection on EDM cone in d_{ij}

Reformulating Problem 2 (p.518) in terms of EDM D changes the problem considerably:

$$\left. \begin{array}{l} \underset{D}{\text{minimize}} \quad \|D - H\|_{\text{F}}^2 \\ \text{subject to} \quad \text{rank } V_{\mathcal{N}}^{\text{T}} D V_{\mathcal{N}} \leq \rho \\ \quad \quad \quad D \in \text{EDM}^N \end{array} \right\} \text{ Problem 3} \quad (1296)$$

This third prevalent proximity problem is a Euclidean projection of given matrix H on a generally nonconvex subset ($\rho < N - 1$) of ∂EDM^N the boundary of the convex cone of Euclidean distance matrices relative to subspace \mathbb{S}_h^N (Figure 117(d)). Because coordinates of projection are distance-square and H presumably now holds distance-square measurements, numerical solution to Problem 3 is generally different than that of Problem 2.

For the moment, we need make no assumptions regarding measurement matrix H .

7.3.1 Convex case

$$\left. \begin{array}{l} \underset{D}{\text{minimize}} \quad \|D - H\|_{\text{F}}^2 \\ \text{subject to} \quad D \in \text{EDM}^N \end{array} \right\} \quad (1297)$$

When the rank constraint disappears (for $\rho = N - 1$), this third problem becomes obviously convex because the feasible set is then the entire EDM cone and because the objective function

$$\|D - H\|_{\text{F}}^2 = \sum_{i,j} (d_{ij} - h_{ij})^2 \quad (1298)$$

is a strictly convex quadratic in D ; ^{7.17}

$$\begin{aligned} & \text{minimize} && \sum_{i,j} d_{ij}^2 - 2h_{ij}d_{ij} + h_{ij}^2 \\ & \text{subject to} && D \in \text{EDM}^N \end{aligned} \quad (1299)$$

Optimal solution D^* is therefore unique, as expected, for this simple projection on the EDM cone equivalent to (1224).

7.3.1.1 Equivalent semidefinite program, Problem 3, convex case

In the past, this convex problem was solved numerically by means of alternating projection. (Example 7.3.1.1.1) [135] [128] [164, §1] We translate (1299) to an equivalent semidefinite program because we have a good solver:

Assume the given measurement matrix H to be nonnegative and symmetric; ^{7.18}

$$H = [h_{ij}] \in \mathbb{S}^N \cap \mathbb{R}_+^{N \times N} \quad (1272)$$

We then propose: Problem (1299) is equivalent to the semidefinite program, for

$$\partial \triangleq [d_{ij}^2] = D \circ D \quad (1300)$$

a matrix of distance-square squared,

$$\begin{aligned} & \text{minimize}_{\partial, D} && -\text{tr}(V(\partial - 2H \circ D)V) \\ & \text{subject to} && \begin{bmatrix} \partial_{ij} & d_{ij} \\ d_{ij} & 1 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\ & && D \in \text{EDM}^N \\ & && \partial \in \mathbb{S}_h^N \end{aligned} \quad (1301)$$

^{7.17}For nonzero $Y \in \mathbb{S}_h^N$ and some open interval of $t \in \mathbb{R}$ (§3.2.3.0.2, §D.2.3)

$$\frac{d^2}{dt^2} \|(D + tY) - H\|_{\text{F}}^2 = 2 \text{tr} Y^T Y > 0 \quad \blacklozenge$$

^{7.18}If that H given has negative entries, then the technique of solution presented here becomes invalid. Projection of H on \mathcal{K} (1216) prior to application of this proposed technique, as explained in §7.0.1, is incorrect.

where

$$\begin{bmatrix} \partial_{ij} & d_{ij} \\ d_{ij} & 1 \end{bmatrix} \succeq 0 \Leftrightarrow \partial_{ij} \geq d_{ij}^2 \quad (1302)$$

Symmetry of input H facilitates trace in the objective (§B.4.2 no.20), while its nonnegativity causes $\partial_{ij} \rightarrow d_{ij}^2$ as optimization proceeds.

7.3.1.1.1 Example. *Alternating projection on nearest EDM.*

By solving (1301) we confirm the result from an example given by Glunt, Hayden, *et alii* [135, §6] who found an analytical solution to convex optimization problem (1297) for particular cardinality $N=3$ by using the alternating projection method of von Neumann (§E.10):

$$H = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 9 \\ 1 & 9 & 0 \end{bmatrix}, \quad D^* = \begin{bmatrix} 0 & \frac{19}{9} & \frac{19}{9} \\ \frac{19}{9} & 0 & \frac{76}{9} \\ \frac{19}{9} & \frac{76}{9} & 0 \end{bmatrix} \quad (1303)$$

The original problem (1297) of projecting H on the EDM cone is transformed to an equivalent iterative sequence of projections on the two convex cones (1165) from §6.8.1.1. Using ordinary alternating projection, input H goes to D^* with an accuracy of four decimal places in about 17 iterations. Affine dimension corresponding to this optimal solution is $r=1$.

Obviation of semidefinite programming's computational expense is the principal advantage of this alternating projection technique. \square

7.3.1.2 Schur-form semidefinite program, Problem 3 convex case

Semidefinite program (1301) can be reformulated by moving the objective function in

$$\begin{aligned} & \underset{D}{\text{minimize}} && \|D - H\|_{\mathbb{F}}^2 \\ & \text{subject to} && D \in \text{EDM}^N \end{aligned} \quad (1297)$$

to the constraints. This makes an equivalent second-order cone program: for any measurement matrix H

$$\begin{aligned} & \underset{t \in \mathbb{R}, D}{\text{minimize}} && t \\ & \text{subject to} && \|D - H\|_{\mathbb{F}}^2 \leq t \\ & && D \in \text{EDM}^N \end{aligned} \quad (1304)$$

We can transform this problem to an equivalent Schur-form semidefinite program; (§3.1.7.2)

$$\begin{aligned} & \underset{t \in \mathbb{R}, D}{\text{minimize}} && t \\ & \text{subject to} && \begin{bmatrix} tI & \text{vec}(D - H) \\ \text{vec}(D - H)^T & 1 \end{bmatrix} \succeq 0 \\ & && D \in \text{EDM}^N \end{aligned} \quad (1305)$$

characterized by great sparsity and structure. The advantage of this SDP is lack of conditions on input H ; *e.g.*, negative entries would invalidate any solution provided by (1301). (§7.0.1.2)

7.3.1.3 Gram-form semidefinite program, Problem 3 convex case

Further, this problem statement may be equivalently written in terms of a Gram matrix via linear bijective (§5.6.1) EDM operator $\mathbf{D}(G)$ (820);

$$\begin{aligned} & \underset{G \in \mathbb{S}_c^N, t \in \mathbb{R}}{\text{minimize}} && t \\ & \text{subject to} && \begin{bmatrix} tI & \text{vec}(\mathbf{D}(G) - H) \\ \text{vec}(\mathbf{D}(G) - H)^T & 1 \end{bmatrix} \succeq 0 \\ & && G \succeq 0 \end{aligned} \quad (1306)$$

To include constraints on the list $X \in \mathbb{R}^{n \times N}$, we would rewrite this:

$$\begin{aligned} & \underset{G \in \mathbb{S}_c^N, t \in \mathbb{R}, X \in \mathbb{R}^{n \times N}}{\text{minimize}} && t \\ & \text{subject to} && \begin{bmatrix} tI & \text{vec}(\mathbf{D}(G) - H) \\ \text{vec}(\mathbf{D}(G) - H)^T & 1 \end{bmatrix} \succeq 0 \\ & && \begin{bmatrix} I & X \\ X^T & G \end{bmatrix} \succeq 0 \\ & && X \in \mathcal{C} \end{aligned} \quad (1307)$$

where \mathcal{C} is some abstract convex set. This technique is discussed in §5.4.2.2.4.

7.3.1.4 Dual interpretation, projection on EDM cone

From §E.9.1.1 we learn that projection on a convex set has a dual form. In the circumstance \mathcal{K} is a convex cone and point x exists exterior to the cone or on its boundary, distance to the nearest point Px in \mathcal{K} is found as the optimal value of the objective

$$\begin{aligned} \|x - Px\| = \underset{a}{\text{maximize}} \quad & a^T x \\ \text{subject to} \quad & \|a\| \leq 1 \\ & a \in \mathcal{K}^\circ \end{aligned} \quad (1903)$$

where \mathcal{K}° is the polar cone.

Applying this result to (1297), we get a convex optimization for any given symmetric matrix H exterior to or on the EDM cone boundary:

$$\begin{aligned} \underset{D}{\text{minimize}} \quad & \|D - H\|_F^2 \\ \text{subject to} \quad & D \in \text{EDM}^N \end{aligned} \equiv \begin{aligned} \underset{A^\circ}{\text{maximize}} \quad & \langle A^\circ, H \rangle \\ \text{subject to} \quad & \|A^\circ\|_F \leq 1 \\ & A^\circ \in \text{EDM}^{N^\circ} \end{aligned} \quad (1308)$$

Then from (1905) projection of H on cone EDM^N is

$$D^* = H - A^{\circ*} \langle A^{\circ*}, H \rangle \quad (1309)$$

Critchley proposed, instead, projection on the polar EDM cone in his 1980 thesis [78, p.113]: In that circumstance, by projection on the algebraic complement (§E.9.2.2.1),

$$D^* = A^* \langle A^*, H \rangle \quad (1310)$$

which is equal to (1309) when A^* solves

$$\begin{aligned} \underset{A}{\text{maximize}} \quad & \langle A, H \rangle \\ \text{subject to} \quad & \|A\|_F = 1 \\ & A \in \text{EDM}^N \end{aligned} \quad (1311)$$

This projection of symmetric H on polar cone EDM^{N° can be made a convex problem, of course, by relaxing the equality constraint ($\|A\|_F \leq 1$).

7.3.2 Minimization of affine dimension in Problem 3

When the desired affine dimension ρ is diminished, Problem 3 (1296) is difficult to solve [164, §3] because the feasible set in $\mathbb{R}^{N(N-1)/2}$ loses convexity. By substituting rank envelope (1281) into Problem 3, then for any given H we get a convex problem

$$\begin{aligned} & \underset{D}{\text{minimize}} && \|D - H\|_{\text{F}}^2 \\ & \text{subject to} && -\text{tr}(VDV) \leq \kappa \rho \\ & && D \in \text{EDM}^N \end{aligned} \tag{1312}$$

where $\kappa \in \mathbb{R}_+$ is a constant determined by cut-and-try. Given κ , problem (1312) is a convex optimization having unique solution in any desired affine dimension ρ ; an approximation to Euclidean projection on that nonconvex subset of the EDM cone containing EDMs with corresponding affine dimension no greater than ρ .

The SDP equivalent to (1312) does not move κ into the variables as on page 524: for nonnegative symmetric input H and distance-square squared variable ∂ as in (1300),

$$\begin{aligned} & \underset{\partial, D}{\text{minimize}} && -\text{tr}(V(\partial - 2H \circ D)V) \\ & \text{subject to} && \begin{bmatrix} \partial_{ij} & d_{ij} \\ d_{ij} & 1 \end{bmatrix} \succeq 0, \quad N \geq j > i = 1 \dots N-1 \\ & && -\text{tr}(VDV) \leq \kappa \rho \\ & && D \in \text{EDM}^N \\ & && \partial \in \mathbb{S}_h^N \end{aligned} \tag{1313}$$

That means we will not see equivalence of this $\text{cenv}(\text{rank})$ -minimization problem to the non-rank-constrained problems (1299) and (1301) like we saw for its counterpart (1283) in Problem 2.

Another approach to affine dimension minimization is to project instead on the polar EDM cone; discussed in §6.8.1.5.

7.3.3 Constrained affine dimension, Problem 3

When one desires affine dimension diminished further below what can be achieved via $\text{cenv}(\text{rank})$ -minimization as in (1313), spectral projection can be considered a natural means in light of its successful application to projection on a rank ρ subset of the positive semidefinite cone in §7.1.4.

Yet it is wrong here to zero eigenvalues of $-VDV$ or $-VGV$ or a variant to reduce affine dimension, because that particular method comes from projection on a positive semidefinite cone (1251); zeroing those eigenvalues here in Problem 3 would place an elbow in the projection path (Figure 132) thereby producing a result that is necessarily suboptimal. Problem 3 is instead a projection on the EDM cone whose associated spectral cone is considerably different. (§5.11.2.3) Proper choice of spectral cone is demanded by diagonalization of that variable argument to the objective:

7.3.3.1 Cayley-Menger form

We use Cayley-Menger composition of the Euclidean distance matrix to solve a problem that is the same as Problem 3 (1296): (§5.7.3.0.1)

$$\begin{aligned} & \underset{D}{\text{minimize}} \quad \left\| \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} - \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -H \end{bmatrix} \right\|_F^2 \\ & \text{subject to} \quad \text{rank} \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \leq \rho + 2 \\ & \quad \quad \quad D \in \text{EDM}^N \end{aligned} \tag{1314}$$

a projection of H on a generally nonconvex subset (when $\rho < N - 1$) of the Euclidean distance matrix cone boundary $\text{rel } \partial \text{EDM}^N$; *id est*, projection from the EDM cone interior or exterior on a subset of its relative boundary (§6.6, (1094)).

Rank of an optimal solution is intrinsically bounded above and below;

$$2 \leq \text{rank} \begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D^* \end{bmatrix} \leq \rho + 2 \leq N + 1 \tag{1315}$$

Our proposed strategy for low-rank solution is projection on that subset of a spectral cone $\lambda \left(\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -\text{EDM}^N \end{bmatrix} \right)$ (§5.11.2.3) corresponding to affine

dimension not in excess of that ρ desired; *id est*, spectral projection on

$$\begin{bmatrix} \mathbb{R}_+^{\rho+1} \\ \mathbf{0} \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \subset \mathbb{R}^{N+1} \quad (1316)$$

where

$$\partial\mathcal{H} = \{\lambda \in \mathbb{R}^{N+1} \mid \mathbf{1}^T \lambda = 0\} \quad (1027)$$

is a hyperplane through the origin. This pointed polyhedral cone (1316), to which membership subsumes the rank constraint, has empty interior.

Given desired affine dimension $0 \leq \rho \leq N-1$ and diagonalization (§A.5) of unknown EDM D

$$\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix} \triangleq U\Upsilon U^T \in \mathbb{S}_h^{N+1} \quad (1317)$$

and given symmetric H in diagonalization

$$\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -H \end{bmatrix} \triangleq Q\Lambda Q^T \in \mathbb{S}^{N+1} \quad (1318)$$

having eigenvalues arranged in nonincreasing order, then by (1040) problem (1314) is equivalent to

$$\begin{aligned} & \underset{\Upsilon, R}{\text{minimize}} && \|\delta(\Upsilon) - \pi(\delta(R^T \Lambda R))\|^2 \\ & \text{subject to} && \delta(\Upsilon) \in \begin{bmatrix} \mathbb{R}_+^{\rho+1} \\ \mathbf{0} \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \\ & && \delta(QR\Upsilon R^T Q^T) = \mathbf{0} \\ & && R^{-1} = R^T \end{aligned} \quad (1319)$$

where π is the permutation operator from §7.1.3 arranging its vector argument in nonincreasing order,^{7.19} where

$$R \triangleq Q^T U \in \mathbb{R}^{N+1 \times N+1} \quad (1320)$$

in U on the set of orthogonal matrices is a bijection, and where $\partial\mathcal{H}$ insures one negative eigenvalue. Hollowness constraint $\delta(QR\Upsilon R^T Q^T) = \mathbf{0}$ makes problem (1319) difficult by making the two variables dependent.

^{7.19}Recall, any permutation matrix is an orthogonal matrix.

Our plan is to instead divide problem (1319) into two and then iterate their solution:

$$\begin{aligned} & \underset{\Upsilon}{\text{minimize}} && \|\delta(\Upsilon) - \pi(\delta(R^T \Lambda R))\|^2 \\ & \text{subject to} && \delta(\Upsilon) \in \begin{bmatrix} \mathbb{R}_+^{\rho+1} \\ \mathbf{0} \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \end{aligned} \quad (\text{a}) \quad (1321)$$

$$\begin{aligned} & \underset{R}{\text{minimize}} && \|R \Upsilon^* R^T - \Lambda\|_{\mathbb{F}}^2 \\ & \text{subject to} && \delta(QR \Upsilon^* R^T Q^T) = \mathbf{0} \\ & && R^{-1} = R^T \end{aligned} \quad (\text{b})$$

Proof. We justify disappearance of the hollowness constraint in convex optimization problem (1321a): From the arguments in §7.1.3 with regard to π the permutation operator, cone membership constraint $\delta(\Upsilon) \in \begin{bmatrix} \mathbb{R}_+^{\rho+1} \\ \mathbf{0} \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H}$ from (1321a) is equivalent to

$$\delta(\Upsilon) \in \begin{bmatrix} \mathbb{R}_+^{\rho+1} \\ \mathbf{0} \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \cap \mathcal{K}_{\mathcal{M}} \quad (1322)$$

where $\mathcal{K}_{\mathcal{M}}$ is the monotone cone (§2.13.9.4.3). Membership of $\delta(\Upsilon)$ to the *polyhedral cone of majorization* (Theorem A.1.2.0.1)

$$\mathcal{K}_{\lambda\delta}^* = \partial\mathcal{H} \cap \mathcal{K}_{\mathcal{M}+}^* \quad (1339)$$

where $\mathcal{K}_{\mathcal{M}+}^*$ is the dual monotone nonnegative cone (§2.13.9.4.2), is a condition (in absence of a hollowness constraint) that would insure existence of a symmetric hollow matrix $\begin{bmatrix} 0 & \mathbf{1}^T \\ \mathbf{1} & -D \end{bmatrix}$. Curiously, intersection of

this feasible superset $\begin{bmatrix} \mathbb{R}_+^{\rho+1} \\ \mathbf{0} \\ \mathbb{R}_- \end{bmatrix} \cap \partial\mathcal{H} \cap \mathcal{K}_{\mathcal{M}}$ from (1322) with the cone of majorization $\mathcal{K}_{\lambda\delta}^*$ is a benign operation; *id est*,

$$\partial\mathcal{H} \cap \mathcal{K}_{\mathcal{M}+}^* \cap \mathcal{K}_{\mathcal{M}} = \partial\mathcal{H} \cap \mathcal{K}_{\mathcal{M}} \quad (1323)$$

verifiable by observing conic dependencies (§2.10.3) among the aggregate of halfspace-description normals. The cone membership constraint in (1321a) therefore inherently insures existence of a symmetric hollow matrix. ♦

Optimization (1321b) would be a Procrustes problem (§C.4) were it not for the hollowness constraint; it is, instead, a minimization over the intersection of the nonconvex manifold of orthogonal matrices with another nonconvex set in variable R specified by the hollowness constraint.

We solve problem (1321b) by a method introduced in §4.6.0.0.2: Define $R = [r_1 \cdots r_{N+1}] \in \mathbb{R}^{(N+1) \times (N+1)}$ and make the assignment

$$G = \begin{bmatrix} r_1 \\ \vdots \\ r_{N+1} \\ 1 \end{bmatrix} [r_1^T \cdots r_{N+1}^T \ 1] \in \mathbb{S}^{(N+1)^2+1} \quad (1324)$$

$$= \begin{bmatrix} R_{11} & \cdots & R_{1,N+1} & r_1 \\ \vdots & \ddots & \vdots & \vdots \\ R_{1,N+1}^T & & R_{N+1,N+1} & r_{N+1} \\ r_1^T & \cdots & r_{N+1}^T & 1 \end{bmatrix} \triangleq \begin{bmatrix} r_1 r_1^T & \cdots & r_1 r_{N+1}^T & r_1 \\ \vdots & \ddots & \vdots & \vdots \\ r_{N+1} r_1^T & & r_{N+1} r_{N+1}^T & r_{N+1} \\ r_1^T & \cdots & r_{N+1}^T & 1 \end{bmatrix}$$

where $R_{ij} \triangleq r_i r_j^T \in \mathbb{R}^{(N+1) \times (N+1)}$ and $\Upsilon_{ii}^* \in \mathbb{R}$. Since $R \Upsilon^* R^T = \sum_{i=1}^{N+1} \Upsilon_{ii}^* R_{ii}$ then problem (1321b) is equivalently expressed:

$$\begin{aligned} & \underset{R_{ij}, r_i}{\text{minimize}} && \left\| \sum_{i=1}^{N+1} \Upsilon_{ii}^* R_{ii} - \Lambda \right\|_{\text{F}}^2 \\ & \text{subject to} && \text{tr } R_{ii} = 1, && i = 1 \dots N+1 \\ & && \text{tr } R_{ij} = 0, && i < j = 2 \dots N+1 \\ & && G = \begin{bmatrix} R_{11} & \cdots & R_{1,N+1} & r_1 \\ \vdots & \ddots & \vdots & \vdots \\ R_{1,N+1}^T & & R_{N+1,N+1} & r_{N+1} \\ r_1^T & \cdots & r_{N+1}^T & 1 \end{bmatrix} (\succeq 0) && (1325) \\ & && \delta \left(Q \sum_{i=1}^{N+1} \Upsilon_{ii}^* R_{ii} Q^T \right) = \mathbf{0} \\ & && \text{rank } G = 1 \end{aligned}$$

The rank constraint is regularized by method of *convex iteration* developed in §4.4. Problem (1325) is partitioned into two convex problems:

$$\begin{aligned}
& \underset{R_{ij}, r_i}{\text{minimize}} && \left\| \sum_{i=1}^{N+1} \Upsilon_{ii}^* R_{ii} - \Lambda \right\|_{\text{F}}^2 + \langle G, W \rangle \\
& \text{subject to} && \text{tr } R_{ii} = 1, && i=1 \dots N+1 \\
& && \text{tr } R_{ij} = 0, && i < j = 2 \dots N+1 \\
& && G = \begin{bmatrix} R_{11} & \cdots & R_{1,N+1} & r_1 \\ \vdots & \ddots & \vdots & \vdots \\ R_{1,N+1}^T & & R_{N+1,N+1} & r_{N+1} \\ r_1^T & \cdots & r_{N+1}^T & 1 \end{bmatrix} \succeq 0 && (1326) \\
& && \delta \left(Q \sum_{i=1}^{N+1} \Upsilon_{ii}^* R_{ii} Q^T \right) = \mathbf{0}
\end{aligned}$$

and

$$\begin{aligned}
& \underset{W \in \mathbb{S}^{(N+1)^2+1}}{\text{minimize}} && \langle G^*, W \rangle \\
& \text{subject to} && 0 \preceq W \preceq I \\
& && \text{tr } W = (N+1)^2
\end{aligned} \tag{1327}$$

then iterated with convex problem (1321a) until a rank-1 G matrix is found and the objective of (1321a) is minimized. An optimal solution to (1327) is known in closed form (p.603). The hollowness constraint in (1326) may cause numerical infeasibility; in that case, it can be moved to the objective within an added weighted norm. Conditions for termination of the iteration would then comprise a vanishing norm of hollowness.

7.4 Conclusion

There has been little progress in spectral projection since the discovery by Eckart & Young in 1936 [112] leading to a formula for projection on a rank ρ subset of a positive semidefinite cone (§2.9.2.1). [127] The only closed-form spectral method presently available for solving proximity problems, having a constraint on rank, is based on their discovery (Problem 1, §7.1, §5.13).

One recourse is intentional misapplication of Eckart & Young's result by introducing spectral projection on a positive semidefinite cone into Problem 3 via $\mathbf{D}(G)$ (820), for example. [64] Since Problem 3 instead demands spectral

projection on the EDM cone, any solution acquired that way is necessarily suboptimal.

A second recourse is problem redesign: A presupposition to all proximity problems in this chapter is that matrix H is given. We considered H having various properties such as nonnegativity, symmetry, hollowness, or lack thereof. It was assumed that if H did not already belong to the EDM cone, then we wanted an EDM closest to H in some sense; *id est*, input-matrix H was assumed corrupted somehow. For practical problems, it withstands reason that such a proximity problem could instead be reformulated so that some or all entries of H were unknown but bounded above and below by known limits; the norm objective is thereby eliminated as in the development beginning on page 291. That redesign (*the art*, p.8), in terms of the Gram-matrix bridge between point-list X and EDM D , at once encompasses proximity and completion problems.

A third recourse is to apply the method of *convex iteration* just like we did in §7.2.2.7.1. This technique is applicable to any semidefinite problem requiring a rank constraint; it places a regularization term in the objective that enforces the rank constraint.

The importance and application of solving rank-constrained problems are enormous, a conclusion generally accepted *gratis* by the mathematics and engineering communities. *For example, one might be interested in the minimal order dynamic output feedback which stabilizes a given linear time invariant plant (this problem is considered to be among the most important open problems in control).* [233] Rank-constrained semidefinite programs arise in many vital feedback and control problems [151], optics [66], and communications [250] [224]. Rank-constrained problems also appear naturally in *combinatorial optimization*. (§4.6.0.0.8)

Appendix A

Linear algebra

A.1 Main-diagonal δ operator, λ , trace, vec

We introduce notation δ denoting the main-diagonal linear self-adjoint operator. When linear function δ operates on a square matrix $A \in \mathbb{R}^{N \times N}$, $\delta(A)$ returns a vector composed of all the entries from the main diagonal in the natural order;

$$\delta(A) \in \mathbb{R}^N \quad (1328)$$

Operating on a vector $y \in \mathbb{R}^N$, δ naturally returns a diagonal matrix;

$$\delta(y) \in \mathbb{S}^N \quad (1329)$$

Operating recursively on a vector $\Lambda \in \mathbb{R}^N$ or diagonal matrix $\Lambda \in \mathbb{S}^N$, $\delta(\delta(\Lambda))$ returns Λ itself;

$$\delta^2(\Lambda) \equiv \delta(\delta(\Lambda)) \triangleq \Lambda \quad (1330)$$

Defined in this manner, main-diagonal linear operator δ is *self-adjoint* [201, §3.10, §9.5-1];^{A.1} *videlicet*, (§2.2)

$$\delta(A)^T y = \langle \delta(A), y \rangle = \langle A, \delta(y) \rangle = \text{tr}(A^T \delta(y)) \quad (1331)$$

^{A.1}Linear operator $T : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^{M \times N}$ is self-adjoint when, for each and every $X_1, X_2 \in \mathbb{R}^{m \times n}$

$$\langle T(X_1), X_2 \rangle = \langle X_1, T(X_2) \rangle$$

A.1.1 Identities

This δ notation is efficient and unambiguous as illustrated in the following examples where $A \circ B$ denotes Hadamard product [180] [139, §1.1.4] of matrices of like size, \otimes Kronecker product [145] (§D.1.2.1), y a vector, X a matrix, e_i the i^{th} member of the standard basis for \mathbb{R}^n , \mathbb{S}_h^N the symmetric hollow subspace, σ a vector of (nonincreasingly) ordered singular values, and $\lambda(A)$ denotes a vector of nonincreasingly ordered eigenvalues of matrix A :

1. $\delta(A) = \delta(A^T)$
2. $\text{tr}(A) = \text{tr}(A^T) = \delta(A)^T \mathbf{1} = \langle I, A \rangle$
3. $\delta(cA) = c\delta(A)$ $c \in \mathbb{R}$
4. $\text{tr}(cA) = c \text{tr}(A) = c \mathbf{1}^T \lambda(A)$ $c \in \mathbb{R}$
5. $\text{vec}(cA) = c \text{vec}(A)$ $c \in \mathbb{R}$
6. $A \circ cB = cA \circ B$ $c \in \mathbb{R}$
7. $A \otimes cB = cA \otimes B$ $c \in \mathbb{R}$
8. $\delta(A + B) = \delta(A) + \delta(B)$
9. $\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B)$
10. $\text{vec}(A + B) = \text{vec}(A) + \text{vec}(B)$
11. $(A + B) \circ C = A \circ C + B \circ C$
 $A \circ (B + C) = A \circ B + A \circ C$
12. $(A + B) \otimes C = A \otimes C + B \otimes C$
 $A \otimes (B + C) = A \otimes B + A \otimes C$
13. $\text{sgn}(c) \lambda(|c|A) = c \lambda(A)$ $c \in \mathbb{R}$
14. $\text{sgn}(c) \sigma(|c|A) = c \sigma(A)$ $c \in \mathbb{R}$
15. $\text{tr}(c\sqrt{A^T A}) = c \text{tr} \sqrt{A^T A} = c \mathbf{1}^T \sigma(A)$ $c \in \mathbb{R}$
16. $\pi(\delta(A)) = \lambda(I \circ A)$ where π is the presorting function.
17. $\delta(AB) = (A \circ B^T) \mathbf{1} = (B^T \circ A) \mathbf{1}$
18. $\delta(AB)^T = \mathbf{1}^T (A^T \circ B) = \mathbf{1}^T (B \circ A^T)$

19. $\delta(uv^T) = \begin{bmatrix} u_1 v_1 \\ \vdots \\ u_N v_N \end{bmatrix} = u \circ v, \quad u, v \in \mathbb{R}^N$
20. $\text{tr}(A^T B) = \text{tr}(AB^T) = \text{tr}(BA^T) = \text{tr}(B^T A)$
 $= \mathbf{1}^T(A \circ B)\mathbf{1} = \mathbf{1}^T \delta(AB^T) = \delta(A^T B)^T \mathbf{1} = \delta(BA^T)^T \mathbf{1} = \delta(B^T A)^T \mathbf{1}$
21. $D = [d_{ij}] \in \mathbb{S}_h^N, \quad H = [h_{ij}] \in \mathbb{S}_h^N, \quad V = I - \frac{1}{N} \mathbf{1}\mathbf{1}^T \in \mathbb{S}^N$ (confer §B.4.2 no.20)
 $N \text{tr}(-V(D \circ H)V) = \text{tr}(D^T H) = \mathbf{1}^T(D \circ H)\mathbf{1} = \text{tr}(\mathbf{1}\mathbf{1}^T(D \circ H)) = \sum_{i,j} d_{ij} h_{ij}$
22. $\text{tr}(\Lambda A) = \delta(\Lambda)^T \delta(A), \quad \delta^2(\Lambda) \triangleq \Lambda \in \mathbb{S}^N$
23. $y^T B \delta(A) = \text{tr}(B \delta(A) y^T) = \text{tr}(\delta(B^T y) A) = \text{tr}(A \delta(B^T y))$
 $= \delta(A)^T B^T y = \text{tr}(y \delta(A)^T B^T) = \text{tr}(A^T \delta(B^T y)) = \text{tr}(\delta(B^T y) A^T)$
24. $\delta^2(A^T A) = \sum_i e_i e_i^T A^T A e_i e_i^T$
25. $\delta(\delta(A) \mathbf{1}^T) = \delta(\mathbf{1} \delta(A)^T) = \delta(A)$
26. $\delta(A \mathbf{1}) \mathbf{1} = \delta(A \mathbf{1}\mathbf{1}^T) = A \mathbf{1}, \quad \delta(y) \mathbf{1} = \delta(y \mathbf{1}^T) = y$
27. $\delta(I \mathbf{1}) = \delta(\mathbf{1}) = I$
28. $\delta(e_i e_j^T \mathbf{1}) = \delta(e_i) = e_i e_i^T$
29. For $\zeta = [\zeta_i] \in \mathbb{R}^k$ and $x = [x_i] \in \mathbb{R}^k, \quad \sum_i \zeta_i / x_i = \zeta^T \delta(x)^{-1} \mathbf{1}$
30. $\text{vec}(A \circ B) = \text{vec}(A) \circ \text{vec}(B) = \delta(\text{vec } A) \text{vec}(B)$
 $= \text{vec}(B) \circ \text{vec}(A) = \delta(\text{vec } B) \text{vec}(A)$ (36)(1675)
31. $\text{vec}(A X B) = (B^T \otimes A) \text{vec } X$ (not ^H)
32. $\text{vec}(B X A) = (A^T \otimes B) \text{vec } X$
33. $\text{tr}(A X B X^T) = \text{vec}(X)^T \text{vec}(A X B) = \text{vec}(X)^T (B^T \otimes A) \text{vec } X$ [145]
 $= \delta(\text{vec}(X) \text{vec}(X)^T (B^T \otimes A))^T \mathbf{1}$
34. $\text{tr}(A X^T B X) = \text{vec}(X)^T \text{vec}(B X A) = \text{vec}(X)^T (A^T \otimes B) \text{vec } X$
 $= \delta(\text{vec}(X) \text{vec}(X)^T (A^T \otimes B))^T \mathbf{1}$

35. For any permutation matrix Ξ and dimensionally compatible vector y or matrix A

$$\delta(\Xi y) = \Xi \delta(y) \Xi^T \quad (1332)$$

$$\delta(\Xi A \Xi^T) = \Xi \delta(A) \quad (1333)$$

So given any permutation matrix Ξ and any dimensionally compatible matrix B , for example,

$$\delta^2(B) = \Xi \delta^2(\Xi^T B \Xi) \Xi^T \quad (1334)$$

36. $A \otimes (B \otimes C) = (A \otimes B) \otimes C$

37. $(A \otimes B)(C \otimes D) = AC \otimes BD$

38. For A a vector, $(A \otimes B) = (A \otimes I)B$

39. For B a row vector, $(A \otimes B) = A(I \otimes B)$

40. $(A \otimes B)^T = A^T \otimes B^T$

41. $(A \otimes B)^{-1} = A^{-1} \otimes B^{-1}$

42. $\text{tr}(A \otimes B) = \text{tr} A \text{tr} B$

43. For $A \in \mathbb{R}^{m \times m}$, $B \in \mathbb{R}^{n \times n}$, $\det(A \otimes B) = \det^n(A) \det^m(B)$

44. There exist permutation matrices Ξ_1 and Ξ_2 such that [145, p.28]

$$A \otimes B = \Xi_1(B \otimes A)\Xi_2 \quad (1335)$$

45. For eigenvalues $\lambda(A) \in \mathbb{C}^n$ and eigenvectors $v(A) \in \mathbb{C}^{n \times n}$ such that $A = v \delta(\lambda) v^{-1} \in \mathbb{R}^{n \times n}$

$$\lambda(A \otimes B) = \lambda(A) \otimes \lambda(B), \quad v(A \otimes B) = v(A) \otimes v(B) \quad (1336)$$

46. Given *analytic function* [73] $f : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$, then $f(I \otimes A) = I \otimes f(A)$ and $f(A \otimes I) = f(A) \otimes I$ [145, p.28]

A.1.2 Majorization

A.1.2.0.1 Theorem. (Schur) *Majorization.* [352, §7.4] [180, §4.3] [181, §5.5] Let $\lambda \in \mathbb{R}^N$ denote a given vector of eigenvalues and let $\delta \in \mathbb{R}^N$ denote a given vector of main diagonal entries, both arranged in nonincreasing order. Then

$$\exists A \in \mathbb{S}^N \ni \lambda(A) = \lambda \text{ and } \delta(A) = \delta \iff \lambda - \delta \in \mathcal{K}_{\lambda\delta}^* \quad (1337)$$

and conversely

$$A \in \mathbb{S}^N \Rightarrow \lambda(A) - \delta(A) \in \mathcal{K}_{\lambda\delta}^* \quad (1338)$$

The difference belongs to the pointed polyhedral cone of majorization (empty interior, *confer* (284))

$$\mathcal{K}_{\lambda\delta}^* \triangleq \mathcal{K}_{\mathcal{M}^+}^* \cap \{\zeta \mathbf{1} \mid \zeta \in \mathbb{R}\}^* \quad (1339)$$

where $\mathcal{K}_{\mathcal{M}^+}^*$ is the dual monotone nonnegative cone (395), and where the dual of the line is a hyperplane; $\partial\mathcal{H} = \{\zeta \mathbf{1} \mid \zeta \in \mathbb{R}\}^* = \mathbf{1}^\perp$. \diamond

Majorization cone $\mathcal{K}_{\lambda\delta}^*$ is naturally consequent to the definition of majorization; *id est*, vector $y \in \mathbb{R}^N$ *majorizes* vector x if and only if

$$\sum_{i=1}^k x_i \leq \sum_{i=1}^k y_i \quad \forall 1 \leq k \leq N \quad (1340)$$

and

$$\mathbf{1}^\top x = \mathbf{1}^\top y \quad (1341)$$

Under these circumstances, rather, vector x is majorized by vector y .

In the particular circumstance $\delta(A) = \mathbf{0}$, we get:

A.1.2.0.2 Corollary. *Symmetric hollow majorization.*

Let $\lambda \in \mathbb{R}^N$ denote a given vector of eigenvalues arranged in nonincreasing order. Then

$$\exists A \in \mathbb{S}_h^N \ni \lambda(A) = \lambda \iff \lambda \in \mathcal{K}_{\lambda\delta}^* \quad (1342)$$

and conversely

$$A \in \mathbb{S}_h^N \Rightarrow \lambda(A) \in \mathcal{K}_{\lambda\delta}^* \quad (1343)$$

where $\mathcal{K}_{\lambda\delta}^*$ is defined in (1339). \diamond

A.2 Semidefiniteness: domain of test

The most fundamental necessary, sufficient, and definitive test for positive semidefiniteness of matrix $A \in \mathbb{R}^{n \times n}$ is: [181, §1]

$$x^T A x \geq 0 \text{ for each and every } x \in \mathbb{R}^n \text{ such that } \|x\| = 1 \quad (1344)$$

Traditionally, authors demand evaluation over broader domain; namely, over all $x \in \mathbb{R}^n$ which is sufficient but unnecessary. Indeed, that standard textbook requirement is far over-reaching because if $x^T A x$ is nonnegative for particular $x = x_p$, then it is nonnegative for any αx_p where $\alpha \in \mathbb{R}$. Thus, only normalized x in \mathbb{R}^n need be evaluated.

Many authors add the further requirement that the domain be complex; the broadest domain. By so doing, only *Hermitian matrices* ($A^H = A$ where superscript ^H denotes conjugate transpose)^{A.2} are admitted to the set of positive semidefinite matrices (1347); an unnecessary prohibitive condition.

A.2.1 Symmetry *versus* semidefiniteness

We call (1344) *the most fundamental test* of positive semidefiniteness. Yet some authors instead say, for real A and complex domain $\{x \in \mathbb{C}^n\}$, the complex test $x^H A x \geq 0$ is most fundamental. That complex broadening of the domain of test causes nonsymmetric real matrices to be excluded from the set of positive semidefinite matrices. Yet admitting nonsymmetric real matrices or not is a matter of preference^{A.3} unless that complex test is adopted, as we shall now explain.

Any real square matrix A has a representation in terms of its symmetric and antisymmetric parts; *id est*,

$$A = \frac{(A + A^T)}{2} + \frac{(A - A^T)}{2} \quad (47)$$

Because, for all real A , the antisymmetric part vanishes under real test,

$$x^T \frac{(A - A^T)}{2} x = 0 \quad (1345)$$

^{A.2}Hermitian symmetry is the complex analogue; the real part of a Hermitian matrix is symmetric while its imaginary part is antisymmetric. A Hermitian matrix has real eigenvalues and real main diagonal.

^{A.3}Golub & Van Loan [139, §4.2.2], for example, admit nonsymmetric real matrices.

only the symmetric part of A , $(A + A^T)/2$, has a role determining positive semidefiniteness. Hence the oft-made presumption that only symmetric matrices may be positive semidefinite is, of course, erroneous under (1344). Because eigenvalue-signs of a symmetric matrix translate unequivocally to its semidefiniteness, the eigenvalues that determine semidefiniteness are always those of the *symmetrized* matrix. (§A.3) For that reason, and because symmetric (or Hermitian) matrices must have real eigenvalues, the convention adopted in the literature is that semidefinite matrices are synonymous with symmetric semidefinite matrices. Certainly misleading under (1344), that presumption is typically bolstered with compelling examples from the physical sciences where symmetric matrices occur within the mathematical exposition of natural phenomena.^{A.4} [123, §52]

Perhaps a better explanation of this pervasive presumption of symmetry comes from Horn & Johnson [180, §7.1] whose perspective^{A.5} is the complex matrix, thus necessitating the complex domain of test throughout their treatise. They explain, if $A \in \mathbb{C}^{n \times n}$

*... and if $x^H A x$ is real for all $x \in \mathbb{C}^n$, then A is Hermitian.
Thus, the assumption that A is Hermitian is not necessary in the definition of positive definiteness. It is customary, however.*

Their comment is best explained by noting, the real part of $x^H A x$ comes from the Hermitian part $(A + A^H)/2$ of A ;

$$\operatorname{Re}(x^H A x) = x^H \frac{A + A^H}{2} x \quad (1346)$$

rather,

$$x^H A x \in \mathbb{R} \Leftrightarrow A^H = A \quad (1347)$$

because the imaginary part of $x^H A x$ comes from the anti-Hermitian part $(A - A^H)/2$;

$$\operatorname{Im}(x^H A x) = x^H \frac{A - A^H}{2} x \quad (1348)$$

that vanishes for nonzero x if and only if $A = A^H$. So the Hermitian symmetry assumption is unnecessary, according to Horn & Johnson, not

^{A.4}Symmetric matrices are certainly pervasive in the our chosen subject as well.

^{A.5}A totally complex perspective is not necessarily more advantageous. The positive semidefinite cone, for example, is not self-dual (§2.13.5) in the ambient space of Hermitian matrices. [174, §II]

because nonHermitian matrices could be regarded positive semidefinite, rather because nonHermitian (includes nonsymmetric real) matrices are not comparable on the real line under $x^H A x$. Yet that complex edifice is dismantled in the test of real matrices (1344) because the domain of test is no longer necessarily complex; meaning, $x^T A x$ will certainly always be real, regardless of symmetry, and so real A will always be comparable.

In summary, if we limit the domain of test to all x in \mathbb{R}^n as in (1344), then nonsymmetric real matrices are admitted to the realm of semidefinite matrices because they become comparable on the real line. One important exception occurs for rank-one matrices $\Psi = uv^T$ where u and v are real vectors: Ψ is positive semidefinite if and only if $\Psi = uu^T$. (§A.3.1.0.7)

We might choose to expand the domain of test to all x in \mathbb{C}^n so that only symmetric matrices would be comparable. The alternative to expanding domain of test is to assume all matrices of interest to be symmetric; that is commonly done, hence the synonymous relationship with semidefinite matrices.

A.2.1.0.1 Example. *Nonsymmetric positive definite product.*

Horn & Johnson assert and Zhang agrees:

If $A, B \in \mathbb{C}^{n \times n}$ are positive definite, then we know that the product AB is positive definite if and only if AB is Hermitian.
[180, §7.6, prob.10] [352, §6.2, §3.2]

Implicitly in their statement, A and B are assumed individually Hermitian and the domain of test is assumed complex.

We prove that assertion to be false for real matrices under (1344) that adopts a real domain of test.

$$A^T = A = \begin{bmatrix} 3 & 0 & -1 & 0 \\ 0 & 5 & 1 & 0 \\ -1 & 1 & 4 & 1 \\ 0 & 0 & 1 & 4 \end{bmatrix}, \quad \lambda(A) = \begin{bmatrix} 5.9 \\ 4.5 \\ 3.4 \\ 2.0 \end{bmatrix} \quad (1349)$$

$$B^T = B = \begin{bmatrix} 4 & 4 & -1 & -1 \\ 4 & 5 & 0 & 0 \\ -1 & 0 & 5 & 1 \\ -1 & 0 & 1 & 4 \end{bmatrix}, \quad \lambda(B) = \begin{bmatrix} 8.8 \\ 5.5 \\ 3.3 \\ 0.24 \end{bmatrix} \quad (1350)$$

$$(AB)^T \neq AB = \begin{bmatrix} 13 & 12 & -8 & -4 \\ 19 & 25 & 5 & 1 \\ -5 & 1 & 22 & 9 \\ -5 & 0 & 9 & 17 \end{bmatrix}, \quad \lambda(AB) = \begin{bmatrix} 36. \\ 29. \\ 10. \\ 0.72 \end{bmatrix} \quad (1351)$$

$$\frac{1}{2}(AB + (AB)^T) = \begin{bmatrix} 13 & 15.5 & -6.5 & -4.5 \\ 15.5 & 25 & 3 & 0.5 \\ -6.5 & 3 & 22 & 9 \\ -4.5 & 0.5 & 9 & 17 \end{bmatrix}, \quad \lambda\left(\frac{1}{2}(AB + (AB)^T)\right) = \begin{bmatrix} 36. \\ 30. \\ 10. \\ 0.014 \end{bmatrix} \quad (1352)$$

Whenever $A \in \mathbb{S}_+^n$ and $B \in \mathbb{S}_+^n$, then $\lambda(AB) = \lambda(\sqrt{A}B\sqrt{A})$ will always be a nonnegative vector by (1376) and Corollary A.3.1.0.5. Yet positive definiteness of product AB is certified instead by the nonnegative eigenvalues $\lambda(\frac{1}{2}(AB + (AB)^T))$ in (1352) (§A.3.1.0.1) despite the fact that AB is not symmetric.^{A.6} Horn & Johnson and Zhang resolve the anomaly by choosing to exclude nonsymmetric matrices and products; they do so by expanding the domain of test to \mathbb{C}^n . \square

A.3 Proper statements of positive semidefiniteness

Unlike Horn & Johnson and Zhang, we never adopt a complex domain of test with real matrices. So motivated is our consideration of proper statements of positive semidefiniteness under real domain of test. This restriction, ironically, complicates the facts when compared to the corresponding statements for the complex case (found elsewhere, [180] [352]).

We state several fundamental facts regarding positive semidefiniteness of real matrix A and the product AB and sum $A + B$ of real matrices under fundamental real test (1344); a few require proof as they depart from the standard texts, while those remaining are well established or obvious.

^{A.6}It is a little more difficult to find a counter-example in $\mathbb{R}^{2 \times 2}$ or $\mathbb{R}^{3 \times 3}$; which may have served to advance any confusion.

A.3.0.0.1 Theorem. *Positive (semi)definite matrix.*

$A \in \mathbb{S}^M$ is positive semidefinite if and only if for each and every vector $x \in \mathbb{R}^M$ of unit norm, $\|x\| = 1$, ^{A.7} we have $x^T A x \geq 0$ (1344);

$$A \succeq 0 \Leftrightarrow \operatorname{tr}(xx^T A) = x^T A x \geq 0 \quad \forall xx^T \quad (1353)$$

Matrix $A \in \mathbb{S}^M$ is positive definite if and only if for each and every $\|x\| = 1$ we have $x^T A x > 0$;

$$A \succ 0 \Leftrightarrow \operatorname{tr}(xx^T A) = x^T A x > 0 \quad \forall xx^T, xx^T \neq \mathbf{0} \quad (1354)$$

◇

Proof. Statements (1353) and (1354) are each a particular instance of dual generalized inequalities (§2.13.2) with respect to the positive semidefinite cone; *videlicet*, [320]

$$\begin{aligned} A \succeq 0 &\Leftrightarrow \langle xx^T, A \rangle \geq 0 \quad \forall xx^T (\succeq 0) \\ A \succ 0 &\Leftrightarrow \langle xx^T, A \rangle > 0 \quad \forall xx^T (\succeq 0), xx^T \neq \mathbf{0} \end{aligned} \quad (1355)$$

This says: positive semidefinite matrix A must belong to the normal side of every hyperplane whose normal is an extreme direction of the positive semidefinite cone. Relations (1353) and (1354) remain true when xx^T is replaced with “for each and every” positive semidefinite matrix $X \in \mathbb{S}_+^M$ (§2.13.5) of unit norm, $\|X\| = 1$, as in

$$\begin{aligned} A \succeq 0 &\Leftrightarrow \operatorname{tr}(XA) \geq 0 \quad \forall X \in \mathbb{S}_+^M \\ A \succ 0 &\Leftrightarrow \operatorname{tr}(XA) > 0 \quad \forall X \in \mathbb{S}_+^M, X \neq \mathbf{0} \end{aligned} \quad (1356)$$

But that condition is more than what is necessary. By the *discretized membership theorem* in §2.13.4.2.1, the extreme directions xx^T of the positive semidefinite cone constitute a minimal set of generators necessary and sufficient for discretization of dual generalized inequalities (1356) certifying membership to that cone. ◆

^{A.7}The traditional condition requiring all $x \in \mathbb{R}^M$ for defining positive (semi)definiteness is actually more than what is necessary. The set of norm-1 vectors is necessary and sufficient to establish positive semidefiniteness; actually, any particular norm and any nonzero norm-constant will work.

A.3.1 Semidefiniteness, eigenvalues, nonsymmetric

When $A \in \mathbb{R}^{n \times n}$, let $\lambda(\frac{1}{2}(A + A^T)) \in \mathbb{R}^n$ denote eigenvalues of the symmetrized matrix^{A.8} arranged in nonincreasing order.

- By positive semidefiniteness of $A \in \mathbb{R}^{n \times n}$ we mean,^{A.9} [241, §1.3.1] (confer §A.3.1.0.1)

$$x^T A x \geq 0 \quad \forall x \in \mathbb{R}^n \Leftrightarrow A + A^T \succeq 0 \Leftrightarrow \lambda(A + A^T) \succeq 0 \quad (1357)$$

- (§2.9.0.1)

$$A \succeq 0 \Rightarrow A^T = A \quad (1358)$$

$$A \succeq B \Leftrightarrow A - B \succeq 0 \not\Rightarrow A \succeq 0 \text{ or } B \succeq 0 \quad (1359)$$

$$x^T A x \geq 0 \quad \forall x \not\Rightarrow A^T = A \quad (1360)$$

- Matrix symmetry is not intrinsic to positive semidefiniteness;

$$A^T = A, \quad \lambda(A) \succeq 0 \Rightarrow x^T A x \geq 0 \quad \forall x \quad (1361)$$

$$\lambda(A) \succeq 0 \Leftarrow A^T = A, \quad x^T A x \geq 0 \quad \forall x \quad (1362)$$

- If $A^T = A$ then

$$\lambda(A) \succeq 0 \Leftrightarrow A \succeq 0 \quad (1363)$$

meaning, matrix A belongs to the positive semidefinite cone in the subspace of symmetric matrices if and only if its eigenvalues belong to the nonnegative orthant.

$$\langle A, A \rangle = \langle \lambda(A), \lambda(A) \rangle \quad (39)$$

- For $\mu \in \mathbb{R}$, $A \in \mathbb{R}^{n \times n}$, and vector $\lambda(A) \in \mathbb{C}^n$ holding the ordered eigenvalues of A

$$\lambda(\mu I + A) = \mu \mathbf{1} + \lambda(A) \quad (1364)$$

Proof: $A = M J M^{-1}$ and $\mu I + A = M(\mu I + J)M^{-1}$ where J is the *Jordan form* for A ; [294, §5.6, App.B] *id est*, $\delta(J) = \lambda(A)$, so $\lambda(\mu I + A) = \delta(\mu I + J)$ because $\mu I + J$ is also a Jordan form. ♦

^{A.8}The symmetrization of A is $(A + A^T)/2$. $\lambda(\frac{1}{2}(A + A^T)) = \lambda(A + A^T)/2$.

^{A.9}Strang agrees [294, p.334] it is not $\lambda(A)$ that requires observation. Yet he is mistaken by proposing the Hermitian part alone $x^H(A + A^H)x$ be tested, because the anti-Hermitian part does not vanish under complex test unless A is Hermitian. (1348)

By similar reasoning,

$$\lambda(I + \mu A) = \mathbf{1} + \lambda(\mu A) \quad (1365)$$

For vector $\sigma(A)$ holding the singular values of any matrix A

$$\sigma(I + \mu A^T A) = \pi(|\mathbf{1} + \mu \sigma(A^T A)|) \quad (1366)$$

$$\sigma(\mu I + A^T A) = \pi(|\mu \mathbf{1} + \sigma(A^T A)|) \quad (1367)$$

where π is the nonlinear permutation-operator sorting its vector argument into nonincreasing order.

- For $A \in \mathbb{S}^M$ and each and every $\|w\| = 1$ [180, §7.7, prob.9]

$$w^T A w \leq \mu \Leftrightarrow A \preceq \mu I \Leftrightarrow \lambda(A) \preceq \mu \mathbf{1} \quad (1368)$$

- [180, §2.5.4] (confer (38))

$$A \text{ is normal} \Leftrightarrow \|A\|_F^2 = \lambda(A)^T \lambda(A) \quad (1369)$$

- For $A \in \mathbb{R}^{m \times n}$

$$A^T A \succeq 0, \quad A A^T \succeq 0 \quad (1370)$$

because, for dimensionally compatible vector x , $x^T A^T A x = \|Ax\|_2^2$, $x^T A A^T x = \|A^T x\|_2^2$.

- For $A \in \mathbb{R}^{n \times n}$ and $c \in \mathbb{R}$

$$\text{tr}(cA) = c \text{tr}(A) = c \mathbf{1}^T \lambda(A) \quad (\S A.1.1 \text{ no.4})$$

For m a nonnegative integer, (1755)

$$\det(A^m) = \prod_{i=1}^n \lambda(A)_i^m \quad (1371)$$

$$\text{tr}(A^m) = \sum_{i=1}^n \lambda(A)_i^m \quad (1372)$$

- For A diagonalizable (§A.5), $A = S\Lambda S^{-1}$, (confer [294, p.255])

$$\text{rank } A = \text{rank } \delta(\lambda(A)) = \text{rank } \Lambda \quad (1373)$$

meaning, rank is equal to the number of nonzero eigenvalues in vector

$$\lambda(A) \triangleq \delta(\Lambda) \quad (1374)$$

by the 0 *eigenvalues theorem* (§A.7.3.0.1).

- (Ky Fan) For $A, B \in \mathbb{S}^n$ [48, §1.2] (confer (1632))

$$\text{tr}(AB) \leq \lambda(A)^T \lambda(B) \quad (1618)$$

with equality (Theobald) when A and B are simultaneously diagonalizable [180] with the same ordering of eigenvalues.

- For $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times m}$

$$\text{tr}(AB) = \text{tr}(BA) \quad (1375)$$

and η eigenvalues of the product and commuted product are identical, including their multiplicity; [180, §1.3.20] *id est*,

$$\lambda(AB)_{1:\eta} = \lambda(BA)_{1:\eta}, \quad \eta \triangleq \min\{m, n\} \quad (1376)$$

Any eigenvalues remaining are zero. By the 0 *eigenvalues theorem* (§A.7.3.0.1),

$$\text{rank}(AB) = \text{rank}(BA), \quad AB \text{ and } BA \text{ diagonalizable} \quad (1377)$$

- For any compatible matrices A, B [180, §0.4]

$$\min\{\text{rank } A, \text{rank } B\} \geq \text{rank}(AB) \quad (1378)$$

- For $A, B \in \mathbb{S}_+^n$

$$\text{rank } A + \text{rank } B \geq \text{rank}(A + B) \geq \min\{\text{rank } A, \text{rank } B\} \geq \text{rank}(AB) \quad (1379)$$

- For $A, B \in \mathbb{S}_+^n$ linearly independent (§B.1.1),

$$\text{rank } A + \text{rank } B = \text{rank}(A + B) > \min\{\text{rank } A, \text{rank } B\} \geq \text{rank}(AB) \quad (1380)$$

- Because $\mathcal{R}(A^T A) = \mathcal{R}(A^T)$ and $\mathcal{R}(A A^T) = \mathcal{R}(A)$ (p.582), for any $A \in \mathbb{R}^{m \times n}$

$$\text{rank}(A A^T) = \text{rank}(A^T A) = \text{rank } A = \text{rank } A^T \quad (1381)$$

- For $A \in \mathbb{R}^{m \times n}$ having no nullspace, and for any $B \in \mathbb{R}^{n \times k}$

$$\text{rank}(AB) = \text{rank}(B) \quad (1382)$$

Proof. For any compatible matrix C , $\mathcal{N}(CAB) \supseteq \mathcal{N}(AB) \supseteq \mathcal{N}(B)$ is obvious. By assumption $\exists A^\dagger \ni A^\dagger A = I$. Let $C = A^\dagger$, then $\mathcal{N}(AB) = \mathcal{N}(B)$ and the stated result follows by conservation of dimension (1481). \blacklozenge

- For $A \in \mathbb{S}^n$ and any nonsingular matrix Y

$$\text{inertia}(A) = \text{inertia}(YAY^T) \quad (1383)$$

a.k.a, *Sylvester's law of inertia.* (1425) [96, §2.4.3]

- For $A, B \in \mathbb{R}^{n \times n}$ square, [180, §0.3.5]

$$\det(AB) = \det(BA) \quad (1384)$$

$$\det(AB) = \det A \det B \quad (1385)$$

Yet for $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times m}$ [68, p.72]

$$\det(I + AB) = \det(I + BA) \quad (1386)$$

- For $A, B \in \mathbb{S}^n$, product AB is symmetric if and only if AB is commutative;

$$(AB)^T = AB \Leftrightarrow AB = BA \quad (1387)$$

Proof. (\Rightarrow) Suppose $AB = (AB)^T$. $(AB)^T = B^T A^T = BA$.
 $AB = (AB)^T \Rightarrow AB = BA$.

(\Leftarrow) Suppose $AB = BA$. $BA = B^T A^T = (AB)^T$. $AB = BA \Rightarrow AB = (AB)^T$. \blacklozenge

Commutativity alone is insufficient for symmetry of the product.
 [294, p.26]

- Diagonalizable matrices $A, B \in \mathbb{R}^{n \times n}$ commute if and only if they are simultaneously diagonalizable. [180, §1.3.12] A product of diagonal matrices is always commutative.
- For $A, B \in \mathbb{R}^{n \times n}$ and $AB = BA$

$$x^T A x \geq 0, x^T B x \geq 0 \quad \forall x \Rightarrow \lambda(A + A^T)_i \lambda(B + B^T)_i \geq 0 \quad \forall i \Leftrightarrow x^T A B x \geq 0 \quad \forall x \quad (1388)$$

the negative result arising because of the schism between the product of eigenvalues $\lambda(A + A^T)_i \lambda(B + B^T)_i$ and the eigenvalues of the symmetrized matrix product $\lambda(AB + (AB)^T)_i$. For example, X^2 is generally not positive semidefinite unless matrix X is symmetric; then (1370) applies. Simply substituting symmetric matrices changes the outcome:

- For $A, B \in \mathbb{S}^n$ and $AB = BA$

$$A \succeq 0, B \succeq 0 \Rightarrow \lambda(AB)_i = \lambda(A)_i \lambda(B)_i \geq 0 \quad \forall i \Leftrightarrow AB \succeq 0 \quad (1389)$$

Positive semidefiniteness of A and B is sufficient but not a necessary condition for positive semidefiniteness of the product AB .

Proof. Because all symmetric matrices are diagonalizable, (§A.5.2) [294, §5.6] we have $A = S\Lambda S^T$ and $B = T\Delta T^T$, where Λ and Δ are real diagonal matrices while S and T are orthogonal matrices. Because $(AB)^T = AB$, then T must equal S , [180, §1.3] and the eigenvalues of A are ordered in the same way as those of B ; *id est*, $\lambda(A)_i = \delta(\Lambda)_i$ and $\lambda(B)_i = \delta(\Delta)_i$ correspond to the same eigenvector.

(\Rightarrow) Assume $\lambda(A)_i \lambda(B)_i \geq 0$ for $i = 1 \dots n$. $AB = S\Lambda\Delta S^T$ is symmetric and has nonnegative eigenvalues contained in diagonal matrix $\Lambda\Delta$ by assumption; hence positive semidefinite by (1357). Now assume $A, B \succeq 0$. That, of course, implies $\lambda(A)_i \lambda(B)_i \geq 0$ for all i because all the individual eigenvalues are nonnegative.

(\Leftarrow) Suppose $AB = S\Lambda\Delta S^T \succeq 0$. Then $\Lambda\Delta \succeq 0$ by (1357), and so all products $\lambda(A)_i \lambda(B)_i$ must be nonnegative; meaning, $\text{sgn}(\lambda(A)) = \text{sgn}(\lambda(B))$. We may, therefore, conclude nothing about the semidefiniteness of A and B . \blacklozenge

- For $A, B \in \mathbb{S}^n$ and $A \succeq 0, B \succeq 0$ (Example [A.2.1.0.1](#))

$$AB = BA \Rightarrow \lambda(AB)_i = \lambda(A)_i \lambda(B)_i \geq 0 \quad \forall i \Rightarrow AB \succeq 0 \quad (1390)$$

$$AB = BA \Rightarrow \lambda(AB)_i \geq 0, \lambda(A)_i \lambda(B)_i \geq 0 \quad \forall i \Leftrightarrow AB \succeq 0 \quad (1391)$$

- For $A, B \in \mathbb{S}^n$ [[181](#), §4.2.13]

$$A \succeq 0, B \succeq 0 \Rightarrow A \otimes B \succeq 0 \quad (1392)$$

$$A \succ 0, B \succ 0 \Rightarrow A \otimes B \succ 0 \quad (1393)$$

and the Kronecker product is symmetric.

- For $A, B \in \mathbb{S}^n$ [[352](#), §6.2]

$$A \succeq 0 \Rightarrow \operatorname{tr} A \geq 0 \quad (1394)$$

$$A \succeq 0, B \succeq 0 \Rightarrow \operatorname{tr} A \operatorname{tr} B \geq \operatorname{tr}(AB) \geq 0 \quad (1395)$$

Because $A \succeq 0, B \succeq 0 \Rightarrow \lambda(AB) = \lambda(\sqrt{A}B\sqrt{A}) \succeq 0$ by [\(1376\)](#) and Corollary [A.3.1.0.5](#), then we have $\operatorname{tr}(AB) \geq 0$.

$$A \succeq 0 \Leftrightarrow \operatorname{tr}(AB) \geq 0 \quad \forall B \succeq 0 \quad (342)$$

- For $A, B, C \in \mathbb{S}^n$ (Löwner)

$$\begin{aligned} A \preceq B, B \preceq C &\Rightarrow A \preceq C && \text{(transitivity)} \\ A \preceq B &\Leftrightarrow A + C \preceq B + C && \text{(additivity)} \\ A \preceq B, A \succeq B &\Rightarrow A = B && \text{(antisymmetry)} \\ A \preceq A &&& \text{(reflexivity)} \end{aligned} \quad (1396)$$

- For $A, B \in \mathbb{R}^{n \times n}$

$$x^T A x \geq x^T B x \quad \forall x \Rightarrow \operatorname{tr} A \geq \operatorname{tr} B \quad (1397)$$

Proof. $x^T A x \geq x^T B x \quad \forall x \Leftrightarrow \lambda((A-B) + (A-B)^T)/2 \succeq 0 \Rightarrow \operatorname{tr}(A+A^T - (B+B^T))/2 = \operatorname{tr}(A-B) \geq 0$. There is no converse. \blacklozenge

- For $A, B \in \mathbb{S}^n$ [352, §6.2] (Theorem A.3.1.0.4)

$$A \succeq B \Rightarrow \operatorname{tr} A \geq \operatorname{tr} B \quad (1398)$$

$$A \succeq B \Rightarrow \delta(A) \succeq \delta(B) \quad (1399)$$

There is no converse, and restriction to the positive semidefinite cone does not improve the situation. All-strict versions hold.

$$A \succeq B \succeq 0 \Rightarrow \operatorname{rank} A \geq \operatorname{rank} B \quad (1400)$$

$$A \succ B \succeq 0 \Rightarrow \det A \geq \det B \geq 0 \quad (1401)$$

$$A \succ B \succeq 0 \Rightarrow \det A > \det B \geq 0 \quad (1402)$$

- For $A, B \in \operatorname{int} \mathbb{S}_+^n$ [31, §4.2] [180, §7.7.4]

$$A \succeq B \Leftrightarrow A^{-1} \preceq B^{-1}, \quad A \succ 0 \Leftrightarrow A^{-1} \succ 0 \quad (1403)$$

- For $A, B \in \mathbb{S}^n$ [352, §6.2]

$$A \succeq B \succeq 0 \Rightarrow \sqrt{A} \succeq \sqrt{B} \quad (1404)$$

- For $A, B \in \mathbb{S}^n$ and $AB = BA$ [352, §6.2, prob.3]

$$A \succeq B \succeq 0 \Rightarrow A^k \succeq B^k, \quad k=1, 2, \dots \quad (1405)$$

A.3.1.0.1 Theorem. *Positive semidefinite ordering of eigenvalues.*

For $A, B \in \mathbb{R}^{M \times M}$, place the eigenvalues of each symmetrized matrix into the respective vectors $\lambda(\frac{1}{2}(A + A^T)), \lambda(\frac{1}{2}(B + B^T)) \in \mathbb{R}^M$. Then, [294, §6]

$$x^T A x \geq 0 \quad \forall x \Leftrightarrow \lambda(A + A^T) \succeq 0 \quad (1406)$$

$$x^T A x > 0 \quad \forall x \neq \mathbf{0} \Leftrightarrow \lambda(A + A^T) \succ 0 \quad (1407)$$

because $x^T(A - A^T)x = 0$. (1345) Now arrange the entries of $\lambda(\frac{1}{2}(A + A^T))$ and $\lambda(\frac{1}{2}(B + B^T))$ in nonincreasing order so $\lambda(\frac{1}{2}(A + A^T))_1$ holds the largest eigenvalue of symmetrized A while $\lambda(\frac{1}{2}(B + B^T))_1$ holds the largest eigenvalue of symmetrized B , and so on. Then [180, §7.7, prob.1, prob.9] for $\kappa \in \mathbb{R}$

$$\begin{aligned} x^T A x \geq x^T B x \quad \forall x &\Rightarrow \lambda(A + A^T) \succeq \lambda(B + B^T) \\ x^T A x \geq x^T I x \kappa \quad \forall x &\Leftrightarrow \lambda(\frac{1}{2}(A + A^T)) \succeq \kappa \mathbf{1} \end{aligned} \quad (1408)$$

Now let $A, B \in \mathbb{S}^M$ have diagonalizations $A = Q\Lambda Q^T$ and $B = U\Upsilon U^T$ with $\lambda(A) = \delta(\Lambda)$ and $\lambda(B) = \delta(\Upsilon)$ arranged in nonincreasing order. Then

$$A \succeq B \Leftrightarrow \lambda(A - B) \succeq 0 \quad (1409)$$

$$A \succeq B \Rightarrow \lambda(A) \succeq \lambda(B) \quad (1410)$$

$$A \succeq B \not\Leftrightarrow \lambda(A) \succeq \lambda(B) \quad (1411)$$

$$S^T A S \succeq B \Leftrightarrow \lambda(A) \succeq \lambda(B) \quad (1412)$$

where $S = QU^T$. [352, §7.5] \diamond

A.3.1.0.2 Theorem. (Weyl) *Eigenvalues of sum.* [180, §4.3.1]

For $A, B \in \mathbb{R}^{M \times M}$, place the eigenvalues of each symmetrized matrix into the respective vectors $\lambda(\frac{1}{2}(A + A^T))$, $\lambda(\frac{1}{2}(B + B^T)) \in \mathbb{R}^M$ in nonincreasing order so $\lambda(\frac{1}{2}(A + A^T))_1$ holds the largest eigenvalue of symmetrized A while $\lambda(\frac{1}{2}(B + B^T))_1$ holds the largest eigenvalue of symmetrized B , and so on. Then, for any $k \in \{1 \dots M\}$

$$\lambda(A + A^T)_k + \lambda(B + B^T)_M \leq \lambda((A + A^T) + (B + B^T))_k \leq \lambda(A + A^T)_k + \lambda(B + B^T)_1 \quad (1413)$$

\diamond

Weyl's theorem establishes positive semidefiniteness of a sum of positive semidefinite matrices; for $A, B \in \mathbb{S}_+^M$

$$\lambda(A)_k + \lambda(B)_M \leq \lambda(A + B)_k \leq \lambda(A)_k + \lambda(B)_1 \quad (1414)$$

Because \mathbb{S}_+^M is a convex cone (§2.9.0.0.1), then by (158)

$$A, B \succeq 0 \Rightarrow \zeta A + \xi B \succeq 0 \text{ for all } \zeta, \xi \geq 0 \quad (1415)$$

A.3.1.0.3 Corollary. *Eigenvalues of sum and difference.* [180, §4.3]

For $A \in \mathbb{S}^M$ and $B \in \mathbb{S}_+^M$, place the eigenvalues of each matrix into the respective vectors $\lambda(A), \lambda(B) \in \mathbb{R}^M$ in nonincreasing order so $\lambda(A)_1$ holds the largest eigenvalue of A while $\lambda(B)_1$ holds the largest eigenvalue of B , and so on. Then, for any $k \in \{1 \dots M\}$

$$\lambda(A - B)_k \leq \lambda(A)_k \leq \lambda(A + B)_k \quad (1416)$$

\diamond

When B is rank-one positive semidefinite, the eigenvalues interlace; *id est*, for $B = qq^T$

$$\lambda(A)_{k-1} \leq \lambda(A - qq^T)_k \leq \lambda(A)_k \leq \lambda(A + qq^T)_k \leq \lambda(A)_{k+1} \quad (1417)$$

A.3.1.0.4 Theorem. *Positive (semi)definite principal submatrices.* [A.10](#)

- $A \in \mathbb{S}^M$ is positive semidefinite if and only if all M principal submatrices of dimension $M-1$ are positive semidefinite and $\det A$ is nonnegative.
- $A \in \mathbb{S}^M$ is positive definite if and only if any one principal submatrix of dimension $M-1$ is positive definite and $\det A$ is positive. \diamond

If any one principal submatrix of dimension $M-1$ is not positive definite, conversely, then A can neither be. Regardless of symmetry, if $A \in \mathbb{R}^{M \times M}$ is positive (semi)definite, then the determinant of each and every principal submatrix is (nonnegative) positive. [241, §1.3.1]

A.3.1.0.5 Corollary. *Positive (semi)definite symmetric products.*

[180, p.399]

- If $A \in \mathbb{S}^M$ is positive definite and any particular dimensionally compatible matrix Z has no nullspace, then $Z^T A Z$ is positive definite.
- If matrix $A \in \mathbb{S}^M$ is positive (semi)definite then, for any matrix Z of compatible dimension, $Z^T A Z$ is positive semidefinite.
- $A \in \mathbb{S}^M$ is positive (semi)definite if and only if there exists a nonsingular Z such that $Z^T A Z$ is positive (semi)definite.
- If $A \in \mathbb{S}^M$ is positive semidefinite and singular it remains possible, for some skinny $Z \in \mathbb{R}^{M \times N}$ with $N < M$, that $Z^T A Z$ becomes positive definite. [A.11](#) \diamond

A.10 A recursive condition for positive (semi)definiteness, this theorem is a synthesis of facts from [180, §7.2] [294, §6.3] (*confer* [241, §1.3.1]). A *principal submatrix* is formed by discarding any subset of rows and columns having the same indices. There are $M!/(1!(M-1)!)$ principal 1×1 submatrices, $M!/(2!(M-2)!)$ principal 2×2 submatrices, and so on, totaling $2^M - 1$ principal submatrices including A itself. By loading y in $y^T A y$ with various patterns of ones and zeros, it follows that any principal submatrix must be positive (semi)definite whenever A is.

A.11 Using the interpretation in §E.6.4.3, this means coefficients of orthogonal projection of vectorized A on a subset of extreme directions from \mathbb{S}_+^M determined by Z can be positive.

We can deduce from these, given nonsingular matrix Z and any particular dimensionally compatible Y : matrix $A \in \mathbb{S}^M$ is positive semidefinite if and only if $\begin{bmatrix} Z^T \\ Y^T \end{bmatrix} A [Z \ Y]$ is positive semidefinite. In other words, from the Corollary it follows: for dimensionally compatible Z

- $A \succeq 0 \Leftrightarrow Z^T A Z \succeq 0$ **and** Z^T has a left inverse

Products such as $Z^\dagger Z$ and $Z Z^\dagger$ are symmetric and positive semidefinite although, given $A \succeq 0$, $Z^\dagger A Z$ and $Z A Z^\dagger$ are neither necessarily symmetric or positive semidefinite.

A.3.1.0.6 Theorem. *Symmetric projector semidefinite.* [19, §III] [20, §6] [197, p.55] For symmetric idempotent matrices P and R

$$P, R \succeq 0 \tag{1418}$$

$$P \succeq R \Leftrightarrow \mathcal{R}(P) \supseteq \mathcal{R}(R) \Leftrightarrow \mathcal{N}(P) \subseteq \mathcal{N}(R)$$

Projector P is never positive definite [296, §6.5, prob.20] unless it is the identity matrix. \diamond

A.3.1.0.7 Theorem. *Symmetric positive semidefinite.*
Given real matrix Ψ with $\text{rank } \Psi = 1$

$$\Psi \succeq 0 \Leftrightarrow \Psi = uu^T \tag{1419}$$

where u is some real vector; *id est*, symmetry is necessary and sufficient for positive semidefiniteness of a rank-1 matrix. \diamond

Proof. Any rank-one matrix must have the form $\Psi = uv^T$. (§B.1) Suppose Ψ is symmetric; *id est*, $v = u$. For all $y \in \mathbb{R}^M$, $y^T u u^T y \geq 0$. Conversely, suppose uv^T is positive semidefinite. We know that can hold if and only if $uv^T + vu^T \succeq 0 \Leftrightarrow$ for all normalized $y \in \mathbb{R}^M$, $2y^T u v^T y \geq 0$; but that is possible only if $v = u$. \blacklozenge

The same does not hold true for matrices of higher rank, as Example A.2.1.0.1 shows.

A.4 Schur complement

Consider *Schur-form* partitioned matrix G : Given $A^T = A$ and $C^T = C$, then [51]

$$\begin{aligned} G &= \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \succeq 0 \\ \Leftrightarrow A \succeq 0, \quad B^T(I - AA^\dagger) &= \mathbf{0}, \quad C - B^T A^\dagger B \succeq 0 \\ \Leftrightarrow C \succeq 0, \quad B(I - CC^\dagger) &= \mathbf{0}, \quad A - BC^\dagger B^T \succeq 0 \end{aligned} \tag{1420}$$

where A^\dagger denotes the Moore-Penrose (pseudo)inverse (§E). In the first instance, $I - AA^\dagger$ is a symmetric projection matrix orthogonally projecting on $\mathcal{N}(A^T)$. (1797) It is apparently required

$$\mathcal{R}(B) \perp \mathcal{N}(A^T) \tag{1421}$$

which precludes $A = \mathbf{0}$ when B is any nonzero matrix. Note that $A \succ 0 \Rightarrow A^\dagger = A^{-1}$; thereby, the projection matrix vanishes. Likewise, in the second instance, $I - CC^\dagger$ projects orthogonally on $\mathcal{N}(C^T)$. It is required

$$\mathcal{R}(B^T) \perp \mathcal{N}(C^T) \tag{1422}$$

which precludes $C = \mathbf{0}$ for B nonzero. Again, $C \succ 0 \Rightarrow C^\dagger = C^{-1}$. So we get, for A or C nonsingular,

$$\begin{aligned} G &= \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \succeq 0 \\ \Leftrightarrow \\ A \succ 0, \quad C - B^T A^{-1} B &\succeq 0 \\ \text{or} \\ C \succ 0, \quad A - BC^{-1} B^T &\succeq 0 \end{aligned} \tag{1423}$$

When A is full-rank then, for all B of compatible dimension, $\mathcal{R}(B)$ is in $\mathcal{R}(A)$. Likewise, when C is full-rank, $\mathcal{R}(B^T)$ is in $\mathcal{R}(C)$. Thus the flavor, for A and C nonsingular,

$$\begin{aligned} G &= \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \succ 0 \\ \Leftrightarrow A \succ 0, \quad C - B^T A^{-1} B &\succ 0 \\ \Leftrightarrow C \succ 0, \quad A - BC^{-1} B^T &\succ 0 \end{aligned} \tag{1424}$$

where $C - B^T A^{-1} B$ is called the *Schur complement of A in G* , while the *Schur complement of C in G* is $A - BC^{-1} B^T$. [132, §4.8]

Origin of the term *Schur complement* is from complementary *inertia*: [96, §2.4.4] Define

$$\text{inertia}(G \in \mathbb{S}^M) \triangleq \{p, z, n\} \quad (1425)$$

where p, z, n respectively represent number of positive, zero, and negative eigenvalues of G ; *id est*,

$$M = p + z + n \quad (1426)$$

Then, when A is invertible,

$$\text{inertia}(G) = \text{inertia}(A) + \text{inertia}(C - B^T A^{-1} B) \quad (1427)$$

and when C is invertible,

$$\text{inertia}(G) = \text{inertia}(C) + \text{inertia}(A - B C^{-1} B^T) \quad (1428)$$

When $A = C = \mathbf{0}$, denoting by $\sigma(B) \in \mathbb{R}_+^m$ the nonincreasingly ordered singular values of matrix $B \in \mathbb{R}^{m \times m}$, then we have the eigenvalues [48, §1.2, prob.17]

$$\lambda(G) = \lambda\left(\begin{bmatrix} \mathbf{0} & B \\ B^T & \mathbf{0} \end{bmatrix}\right) = \begin{bmatrix} \sigma(B) \\ -\Xi \sigma(B) \end{bmatrix} \quad (1429)$$

and

$$\text{inertia}(G) = \text{inertia}(B^T B) + \text{inertia}(-B^T B) \quad (1430)$$

where Ξ is the order-reversing permutation matrix defined in (1619).

A.4.0.0.1 Example. *Nonnegative polynomial.* [31, p.163] Schur-form positive semidefiniteness is sufficient for quadratic polynomial convexity in x , but it is necessary and sufficient for nonnegativity; *videlicet*, for all compatible x

$$\begin{bmatrix} x^T & 1 \end{bmatrix} \begin{bmatrix} A & b \\ b^T & c \end{bmatrix} \begin{bmatrix} x \\ 1 \end{bmatrix} \geq 0 \quad \Leftrightarrow \quad x^T A x + 2b^T x + c \geq 0 \quad (1431)$$

Sublevel set $\{x \mid x^T A x + 2b^T x + c \leq 0\}$ is convex if $A \succeq 0$, but the quadratic polynomial is a convex function if and only if $A \succeq 0$. \square

A.4.0.0.2 Example. *Schur conditions.*

From (1394),

$$\begin{aligned}
\begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \succeq 0 &\quad \Rightarrow \quad \text{tr}(A + C) \geq 0 \\
\Downarrow & \\
\begin{bmatrix} A & \mathbf{0} \\ \mathbf{0}^T & C - B^T A^{-1} B \end{bmatrix} \succeq 0 &\quad \Rightarrow \quad \text{tr}(C - B^T A^{-1} B) \geq 0 \quad (1432) \\
\Downarrow & \\
\begin{bmatrix} A - B C^{-1} B^T & \mathbf{0} \\ \mathbf{0}^T & C \end{bmatrix} \succeq 0 &\quad \Rightarrow \quad \text{tr}(A - B C^{-1} B^T) \geq 0
\end{aligned}$$

Since $\text{tr}(C - B^T A^{-1} B) \geq 0 \Leftrightarrow \text{tr} C \geq \text{tr}(B^T A^{-1} B) \geq 0$ for example, then minimization of $\text{tr} C$ is necessary and sufficient for minimization of $\text{tr}(C - B^T A^{-1} B)$ when both are under constraint $\begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \succeq 0$. \square

A.4.0.0.3 Example. *Sparse Schur conditions.*

Setting matrix A to the identity simplifies the Schur conditions. One consequence relates the definiteness of three quantities:

$$\begin{bmatrix} I & B \\ B^T & C \end{bmatrix} \succeq 0 \Leftrightarrow C - B^T B \succeq 0 \Leftrightarrow \begin{bmatrix} I & \mathbf{0} \\ \mathbf{0}^T & C - B^T B \end{bmatrix} \succeq 0 \quad (1433)$$

\square

A.4.0.0.4 Exercise. *Eigenvalues λ of sparse Schur-form.*

Prove: given $C - B^T B = \mathbf{0}$, for $B \in \mathbb{R}^{m \times n}$ and $C \in \mathbb{S}^n$

$$\lambda \left(\begin{bmatrix} I & B \\ B^T & C \end{bmatrix} \right)_i = \begin{cases} 1 + \lambda(C)_i, & 1 \leq i \leq n \\ 1, & n < i \leq m \\ 0, & \text{otherwise} \end{cases} \quad (1434)$$

\blacktriangledown

A.4.0.0.5 Theorem. *Rank of partitioned matrices.*

When symmetric matrix A is invertible and C is symmetric,

$$\begin{aligned} \operatorname{rank} \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} &= \operatorname{rank} \begin{bmatrix} A & \mathbf{0} \\ \mathbf{0}^T & C - B^T A^{-1} B \end{bmatrix} \\ &= \operatorname{rank} A + \operatorname{rank}(C - B^T A^{-1} B) \end{aligned} \quad (1435)$$

equals rank of a block on the main diagonal plus rank of its Schur complement [352, §2.2, prob.7]. Similarly, when symmetric matrix C is invertible and A is symmetric,

$$\begin{aligned} \operatorname{rank} \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} &= \operatorname{rank} \begin{bmatrix} A - BC^{-1}B^T & \mathbf{0} \\ \mathbf{0}^T & C \end{bmatrix} \\ &= \operatorname{rank}(A - BC^{-1}B^T) + \operatorname{rank} C \end{aligned} \quad (1436)$$

◇

Proof. The first assertion (1435) holds if and only if [180, §0.4.6(c)]

$$\exists \text{ nonsingular } X, Y \ni X \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} Y = \begin{bmatrix} A & \mathbf{0} \\ \mathbf{0}^T & C - B^T A^{-1} B \end{bmatrix} \quad (1437)$$

Let [180, §7.7.6]

$$Y = X^T = \begin{bmatrix} I & -A^{-1}B \\ \mathbf{0}^T & I \end{bmatrix} \quad (1438)$$

◆

A.4.0.0.6 Lemma. *Rank of Schur-form block.* [119] [117]

Matrix $B \in \mathbb{R}^{m \times n}$ has $\operatorname{rank} B \leq \rho$ if and only if there exist matrices $A \in \mathbb{S}^m$ and $C \in \mathbb{S}^n$ such that

$$\operatorname{rank} \begin{bmatrix} A & \mathbf{0} \\ \mathbf{0}^T & C \end{bmatrix} \leq 2\rho \quad \text{and} \quad G = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \succeq 0 \quad (1439)$$

◇

Schur-form positive semidefiniteness alone implies $\operatorname{rank} A \geq \operatorname{rank} B$ and $\operatorname{rank} C \geq \operatorname{rank} B$. But, even in absence of semidefiniteness, we must always have $\operatorname{rank} G \geq \operatorname{rank} A, \operatorname{rank} B, \operatorname{rank} C$ by fundamental linear algebra.

A.4.1 Determinant

$$G = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \quad (1440)$$

We consider again a matrix G partitioned like (1420), but not necessarily positive (semi)definite, where A and C are symmetric.

- When A is invertible,

$$\det G = \det A \det(C - B^T A^{-1} B) \quad (1441)$$

When C is invertible,

$$\det G = \det C \det(A - B C^{-1} B^T) \quad (1442)$$

- When B is full-rank and skinny, $C = \mathbf{0}$, and $A \succeq 0$, then [53, §10.1.1]

$$\det G \neq 0 \Leftrightarrow A + B B^T \succ 0 \quad (1443)$$

When B is a (column) vector, then for all $C \in \mathbb{R}$ and all A of dimension compatible with G

$$\det G = \det(A) C - B^T A_{\text{cof}}^T B \quad (1444)$$

while for $C \neq 0$

$$\det G = C \det\left(A - \frac{1}{C} B B^T\right) \quad (1445)$$

where A_{cof} is the matrix of cofactors [294, §4] corresponding to A .

- When B is full-rank and fat, $A = \mathbf{0}$, and $C \succeq 0$, then

$$\det G \neq 0 \Leftrightarrow C + B^T B \succ 0 \quad (1446)$$

When B is a row-vector, then for $A \neq 0$ and all C of dimension compatible with G

$$\det G = A \det\left(C - \frac{1}{A} B^T B\right) \quad (1447)$$

while for all $A \in \mathbb{R}$

$$\det G = \det(C) A - B C_{\text{cof}}^T B^T \quad (1448)$$

where C_{cof} is the matrix of cofactors corresponding to C .

A.5 Eigen decomposition

When a square matrix $X \in \mathbb{R}^{m \times m}$ is *diagonalizable*, [294, §5.6] then

$$X = S\Lambda S^{-1} = [s_1 \cdots s_m] \Lambda \begin{bmatrix} w_1^T \\ \vdots \\ w_m^T \end{bmatrix} = \sum_{i=1}^m \lambda_i s_i w_i^T \quad (1449)$$

where $\{s_i \in \mathcal{N}(X - \lambda_i I) \subseteq \mathbb{C}^m\}$ are l.i. (right-)eigenvectors^{A.12} constituting the columns of $S \in \mathbb{C}^{m \times m}$ defined by

$$XS = S\Lambda \quad (1450)$$

$\{w_i \in \mathcal{N}(X^T - \lambda_i I) \subseteq \mathbb{C}^m\}$ are linearly independent *left-eigenvectors* of X (eigenvectors of X^T) constituting the rows of S^{-1} defined by [180]

$$S^{-1}X = \Lambda S^{-1} \quad (1451)$$

and where $\{\lambda_i \in \mathbb{C}\}$ are eigenvalues (populating diagonal matrix $\Lambda \in \mathbb{C}^{m \times m}$) corresponding to both left and right eigenvectors; *id est*, $\lambda(X) = \lambda(X^T)$.

There is no connection between diagonalizability and invertibility of X . [294, §5.2] Diagonalizability is guaranteed by a full set of linearly independent eigenvectors, whereas invertibility is guaranteed by all nonzero eigenvalues.

$$\begin{aligned} \text{distinct eigenvalues} &\Rightarrow \text{l.i. eigenvectors} \Leftrightarrow \text{diagonalizable} \\ \text{not diagonalizable} &\Rightarrow \text{repeated eigenvalue} \end{aligned} \quad (1452)$$

A.5.0.0.1 Theorem. *Real eigenvector.* Eigenvectors of a real matrix corresponding to real eigenvalues must be real. \diamond

Proof. $Ax = \lambda x$. Given $\lambda = \lambda^*$, $x^H Ax = \lambda x^H x = \lambda \|x\|^2 = x^T A x^* \Rightarrow x = x^*$, where $x^H = x^{*T}$. The converse is equally simple. \blacklozenge

A.5.0.1 Uniqueness

From the *fundamental theorem of algebra*, [194] which guarantees existence of zeros for a given polynomial, it follows: eigenvalues, including their

^{A.12}Eigenvectors must, of course, be nonzero. The prefix *eigen* is from the German; in this context meaning, something akin to “characteristic”. [291, p.14]

multiplicity, for a given square matrix are unique; meaning, there is no other set of eigenvalues for that matrix. (Conversely, many different matrices may share the same unique set of eigenvalues; *e.g.*, for any X , $\lambda(X) = \lambda(X^T)$.)

Uniqueness of eigenvectors, in contrast, disallows multiplicity of the same direction:

A.5.0.1.1 Definition. *Unique eigenvectors.*

When eigenvectors are *unique*, we mean: unique to within a real nonzero scaling, and their directions are distinct. \triangle

If S is a matrix of eigenvectors of X as in (1449), for example, then $-S$ is certainly another matrix of eigenvectors decomposing X with the same eigenvalues.

For any square matrix, the eigenvector corresponding to a distinct eigenvalue is unique; [291, p.220]

$$\text{distinct eigenvalues} \Rightarrow \text{eigenvectors unique} \quad (1453)$$

Eigenvectors corresponding to a repeated eigenvalue are not unique for a diagonalizable matrix;

$$\text{repeated eigenvalue} \Rightarrow \text{eigenvectors not unique} \quad (1454)$$

Proof follows from the observation: any linear combination of distinct eigenvectors of diagonalizable X , corresponding to a particular eigenvalue, produces another eigenvector. For eigenvalue λ whose multiplicity^{A.13} $\dim \mathcal{N}(X - \lambda I)$ exceeds 1, in other words, any choice of independent vectors from $\mathcal{N}(X - \lambda I)$ (of the same multiplicity) constitutes eigenvectors corresponding to λ . \blacklozenge

Caveat diagonalizability insures linear independence which implies existence of distinct eigenvectors. We may conclude, for diagonalizable matrices,

$$\text{distinct eigenvalues} \Leftrightarrow \text{eigenvectors unique} \quad (1455)$$

^{A.13}A matrix is diagonalizable iff *algebraic multiplicity* (number of occurrences of same eigenvalue) equals *geometric multiplicity* $\dim \mathcal{N}(X - \lambda I) = m - \text{rank}(X - \lambda I)$ [291, p.15] (number of *Jordan blocks* w.r.t λ or number of corresponding l.i. eigenvectors).

A.5.0.2 Invertible matrix

When diagonalizable matrix $X \in \mathbb{R}^{m \times m}$ is nonsingular (no zero eigenvalues), then it has an inverse obtained simply by inverting eigenvalues in (1449):

$$X^{-1} = S\Lambda^{-1}S^{-1} \quad (1456)$$

A.5.1 eigenmatrix

The (right-)eigenvectors $\{s_i\}$ (1449) are naturally orthogonal $w_i^T s_j = 0$ to left-eigenvectors $\{w_i\}$ except, for $i=1 \dots m$, $w_i^T s_i = 1$; called a *biorthogonality condition* [325, §2.2.4] [180] because neither set of left or right eigenvectors is necessarily an orthogonal set. Consequently, each dyad from a diagonalization is an independent (§B.1.1) nonorthogonal projector because

$$s_i w_i^T s_i w_i^T = s_i w_i^T \quad (1457)$$

(whereas the dyads of singular value decomposition are not inherently projectors (*confer* (1462))).

The dyads of eigen decomposition can be termed *eigenmatrices* because

$$X s_i w_i^T = \lambda_i s_i w_i^T \quad (1458)$$

Sum of the eigenmatrices is the identity;

$$\sum_{i=1}^m s_i w_i^T = I \quad (1459)$$

A.5.2 Symmetric matrix diagonalization

The set of *normal matrices* is, precisely, that set of all real matrices having a complete orthonormal set of eigenvectors; [352, §8.1] [296, prob.10.2.31] *id est*, any matrix X for which $XX^T = X^T X$; [139, §7.1.3] [291, p.3] *e.g.*, orthogonal and circulant matrices [147]. All normal matrices are diagonalizable. A symmetric matrix is a special normal matrix whose eigenvalues must be real^{A.14} and whose eigenvectors can be chosen to make

A.14Proof. Suppose λ_i is an eigenvalue, of real $A = A^T$, corresponding to eigenvector s_i . Then $s_i^H A s_i = s_i^T A s_i^*$ (by transposition) $\Rightarrow s_i^{*T} \lambda_i s_i = s_i^T \lambda_i^* s_i^*$ because $(A s_i)^* = (\lambda_i s_i)^*$ by assumption. So we have $\lambda_i \|s_i\|^2 = \lambda_i^* \|s_i\|^2$. There is no converse. \blacklozenge

a real orthonormal set; [296, §6.4] [294, p.315] *id est*, for $X \in \mathbb{S}^m$

$$X = S\Lambda S^T = [s_1 \cdots s_m] \Lambda \begin{bmatrix} s_1^T \\ \vdots \\ s_m^T \end{bmatrix} = \sum_{i=1}^m \lambda_i s_i s_i^T \quad (1460)$$

where $\delta^2(\Lambda) = \Lambda \in \mathbb{S}^m$ (§A.1) and $S^{-1} = S^T \in \mathbb{R}^{m \times m}$ (orthogonal matrix, §B.5) because of symmetry: $S\Lambda S^{-1} = S^{-T}\Lambda S^T$.

Because the arrangement of eigenvectors and their corresponding eigenvalues is arbitrary, we almost always arrange eigenvalues in nonincreasing order as is the convention for singular value decomposition. Then to diagonalize a symmetric matrix that is already a diagonal matrix, orthogonal matrix S becomes a permutation matrix.

A.5.2.1 Positive semidefinite matrix square root

When $X \in \mathbb{S}_+^m$, its unique positive semidefinite matrix square root is defined

$$\sqrt{X} \triangleq S\sqrt{\Lambda}S^T \in \mathbb{S}_+^m \quad (1461)$$

where the square root of nonnegative diagonal matrix $\sqrt{\Lambda}$ is taken entrywise and positive. Then $X = \sqrt{X}\sqrt{X}$.

A.6 Singular value decomposition, SVD

A.6.1 Compact SVD

[139, §2.5.4] For any $A \in \mathbb{R}^{m \times n}$

$$A = U\Sigma Q^T = [u_1 \cdots u_\eta] \Sigma \begin{bmatrix} q_1^T \\ \vdots \\ q_\eta^T \end{bmatrix} = \sum_{i=1}^{\eta} \sigma_i u_i q_i^T \quad (1462)$$

$$U \in \mathbb{R}^{m \times \eta}, \quad \Sigma \in \mathbb{R}^{\eta \times \eta}, \quad Q \in \mathbb{R}^{n \times \eta}$$

where U and Q are always skinny-or-square each having orthonormal columns, and where

$$\eta \triangleq \min\{m, n\} \quad (1463)$$

Square matrix Σ is diagonal (§A.1.1)

$$\delta^2(\Sigma) = \Sigma \in \mathbb{R}^{\eta \times \eta} \quad (1464)$$

holding the singular values $\{\sigma_i \in \mathbb{R}\}$ of A which are always arranged in nonincreasing order by convention and are related to eigenvalues λ by [A.15](#)

$$\sigma(A)_i = \sigma(A^T)_i = \begin{cases} \sqrt{\lambda(A^T A)_i} = \sqrt{\lambda(A A^T)_i} = \lambda(\sqrt{A^T A})_i = \lambda(\sqrt{A A^T})_i > 0, & 1 \leq i \leq \rho \\ 0, & \rho < i \leq \eta \end{cases} \quad (1465)$$

of which the last $\eta - \rho$ are 0, [A.16](#) where

$$\rho \triangleq \text{rank } A = \text{rank } \Sigma \quad (1466)$$

A point sometimes lost: Any real matrix may be decomposed in terms of its real singular values $\sigma(A) \in \mathbb{R}^\eta$ and real matrices U and Q as in [\(1462\)](#), where [\[139, §2.5.3\]](#)

$$\begin{aligned} \mathcal{R}\{u_i \mid \sigma_i \neq 0\} &= \mathcal{R}(A) \\ \mathcal{R}\{u_i \mid \sigma_i = 0\} &\subseteq \mathcal{N}(A^T) \\ \mathcal{R}\{q_i \mid \sigma_i \neq 0\} &= \mathcal{R}(A^T) \\ \mathcal{R}\{q_i \mid \sigma_i = 0\} &\subseteq \mathcal{N}(A) \end{aligned} \quad (1467)$$

A.6.2 Subcompact SVD

Some authors allow only nonzero singular values. In that case the compact decomposition can be made smaller; it can be redimensioned in terms of rank ρ because, for any $A \in \mathbb{R}^{m \times n}$

$$\rho = \text{rank } A = \text{rank } \Sigma = \max\{i \in \{1 \dots \eta\} \mid \sigma_i \neq 0\} \leq \eta \quad (1468)$$

- There are η singular values. For any flavor SVD, rank is equivalent to the number of nonzero singular values on the main diagonal of Σ .

Now

$$A = U \Sigma Q^T = [u_1 \cdots u_\rho] \Sigma \begin{bmatrix} q_1^T \\ \vdots \\ q_\rho^T \end{bmatrix} = \sum_{i=1}^{\rho} \sigma_i u_i q_i^T \quad (1469)$$

$$U \in \mathbb{R}^{m \times \rho}, \quad \Sigma \in \mathbb{R}^{\rho \times \rho}, \quad Q \in \mathbb{R}^{n \times \rho}$$

[A.15](#) When matrix A is normal, $\sigma(A) = |\lambda(A)|$. [\[352, §8.1\]](#)

[A.16](#) For $\eta = n$, $\sigma(A) = \sqrt{\lambda(A^T A)} = \lambda(\sqrt{A^T A})$.

For $\eta = m$, $\sigma(A) = \sqrt{\lambda(A A^T)} = \lambda(\sqrt{A A^T})$.

where the main diagonal of diagonal matrix Σ has no 0 entries, and

$$\begin{aligned}\mathcal{R}\{u_i\} &= \mathcal{R}(A) \\ \mathcal{R}\{q_i\} &= \mathcal{R}(A^T)\end{aligned}\tag{1470}$$

A.6.3 Full SVD

Another common and useful expression of the SVD makes U and Q square; making the decomposition larger than compact SVD. Completing the nullspace bases in U and Q from (1467) provides what is called the *full singular value decomposition* of $A \in \mathbb{R}^{m \times n}$ [294, App.A]. Orthonormal matrices U and Q become orthogonal matrices (§B.5):

$$\begin{aligned}\mathcal{R}\{u_i \mid \sigma_i \neq 0\} &= \mathcal{R}(A) \\ \mathcal{R}\{u_i \mid \sigma_i = 0\} &= \mathcal{N}(A^T) \\ \mathcal{R}\{q_i \mid \sigma_i \neq 0\} &= \mathcal{R}(A^T) \\ \mathcal{R}\{q_i \mid \sigma_i = 0\} &= \mathcal{N}(A)\end{aligned}\tag{1471}$$

For any matrix A having rank ρ ($= \text{rank } \Sigma$)

$$\begin{aligned}A &= U\Sigma Q^T = [u_1 \cdots u_m] \Sigma \begin{bmatrix} q_1^T \\ \vdots \\ q_n^T \end{bmatrix} = \sum_{i=1}^{\eta} \sigma_i u_i q_i^T \\ &= \begin{bmatrix} m \times \rho \text{ basis } \mathcal{R}(A) & m \times m - \rho \text{ basis } \mathcal{N}(A^T) \end{bmatrix} \begin{bmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \end{bmatrix} \begin{bmatrix} (n \times \rho \text{ basis } \mathcal{R}(A^T))^T \\ (n \times n - \rho \text{ basis } \mathcal{N}(A))^T \end{bmatrix} \\ &U \in \mathbb{R}^{m \times m}, \quad \Sigma \in \mathbb{R}^{m \times n}, \quad Q \in \mathbb{R}^{n \times n}\end{aligned}\tag{1472}$$

where upper limit of summation η is defined in (1463). Matrix Σ is no longer necessarily square, now padded with respect to (1464) by $m - \eta$ zero rows or $n - \eta$ zero columns; the nonincreasingly ordered (possibly 0) singular values appear along its main diagonal as for compact SVD (1465).




Figure 134: Geometrical interpretation of full SVD [240]: Image of circle $\{x \in \mathbb{R}^2 \mid \|x\|_2 = 1\}$ under matrix multiplication Ax is, in general, an ellipse. For the example illustrated, $U \triangleq [u_1 \ u_2] \in \mathbb{R}^{2 \times 2}$, $Q \triangleq [q_1 \ q_2] \in \mathbb{R}^{2 \times 2}$.

An important geometrical interpretation of SVD is given in Figure 134 for $m = n = 2$: The image of the unit sphere under any $m \times n$ matrix multiplication is an ellipse. Considering the three factors of the SVD separately, note that Q^T is a pure rotation of the circle. Figure 134 shows how the axes q_1 and q_2 are first rotated by Q^T to coincide with the coordinate axes. Second, the circle is stretched by Σ in the directions of the coordinate axes to form an ellipse. The third step rotates the ellipse by U into its final position. Note how q_1 and q_2 are rotated to end up as u_1 and u_2 , the principal axes of the final ellipse. A direct calculation shows that $Aq_j = \sigma_j u_j$. Thus q_j is first rotated to coincide with the j^{th} coordinate axis, stretched by a factor σ_j , and then rotated to point in the direction of u_j . All of this is beautifully illustrated for 2×2 matrices by the MATLAB code `eigshow.m` (see [297]).

A direct consequence of the geometric interpretation is that the largest singular value σ_1 measures the “magnitude” of A (its 2-norm):

$$\|A\|_2 = \sup_{\|x\|_2=1} \|Ax\|_2 = \sigma_1 \quad (1473)$$

This means that $\|A\|_2$ is the length of the longest principal semiaxis of the ellipse.

Expressions for U , Q , and Σ follow readily from (1472),

$$AA^T U = U \Sigma \Sigma^T \quad \text{and} \quad A^T A Q = Q \Sigma^T \Sigma \quad (1474)$$

demonstrating that the columns of U are the eigenvectors of AA^T and the columns of Q are the eigenvectors of $A^T A$. —Neil Muller et alii [240]

A.6.4 Pseudoinverse by SVD

Matrix pseudoinverse (§E) is nearly synonymous with singular value decomposition because of the elegant expression, given $A = U \Sigma Q^T$

$$A^\dagger = Q \Sigma^\dagger U^T \in \mathbb{R}^{n \times m} \quad (1475)$$

that applies to all three flavors of SVD, where Σ^\dagger simply inverts nonzero entries of matrix Σ .

Given symmetric matrix $A \in \mathbb{S}^n$ and its diagonalization $A = S \Lambda S^T$ (§A.5.2), its pseudoinverse simply inverts all nonzero eigenvalues;

$$A^\dagger = S \Lambda^\dagger S^T \quad (1476)$$

A.6.5 SVD of symmetric matrices

A.6.5.0.1 Definition. *Step function.* (confer §4.3.2.0.1)

Define the signum-like quasilinear function $\psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ that takes value 1 corresponding to a 0-valued entry in its argument:

$$\psi(a) \triangleq \left[\lim_{x_i \rightarrow a_i} \frac{x_i}{|x_i|} = \begin{cases} 1, & a_i \geq 0 \\ -1, & a_i < 0 \end{cases}, \quad i=1 \dots n \right] \in \mathbb{R}^n \quad (1477)$$

△

Eigenvalue signs of a symmetric matrix having diagonalization $A = S\Lambda S^T$ (1460) can be absorbed either into real U or real Q from the full SVD; [310, p.34] (confer §C.4.2.1)

$$A = S\Lambda S^T = S\delta(\psi(\delta(\Lambda)))|\Lambda|S^T \triangleq U\Sigma Q^T \in \mathbb{S}^n \quad (1478)$$

or

$$A = S\Lambda S^T = S|\Lambda|\delta(\psi(\delta(\Lambda)))S^T \triangleq U\Sigma Q^T \in \mathbb{S}^n \quad (1479)$$

where matrix of singular values $\Sigma = |\Lambda|$ denotes entrywise absolute value of diagonal eigenvalue matrix Λ .

A.7 Zeros

A.7.1 norm zero

For any given norm, by definition,

$$\|x\|_\ell = 0 \quad \Leftrightarrow \quad x = \mathbf{0} \quad (1480)$$

Consequently, a generally nonconvex constraint in x like $\|Ax - b\| = \kappa$ becomes convex when $\kappa = 0$.

A.7.2 0 entry

If a positive semidefinite matrix $A = [A_{ij}] \in \mathbb{R}^{n \times n}$ has a 0 entry A_{ii} on its main diagonal, then $A_{ij} + A_{ji} = 0 \quad \forall j$. [241, §1.3.1]

Any symmetric positive semidefinite matrix having a 0 entry on its main diagonal must be $\mathbf{0}$ along the entire row and column to which that 0 entry belongs. [139, §4.2.8] [180, §7.1, prob.2]

A.7.3 0 eigenvalues theorem

This theorem is simple, powerful, and widely applicable:

A.7.3.0.1 Theorem. *Number of 0 eigenvalues.*

For any matrix $A \in \mathbb{R}^{m \times n}$

$$\text{rank}(A) + \dim \mathcal{N}(A) = n \quad (1481)$$

by conservation of dimension. [180, §0.4.4]

For any square matrix $A \in \mathbb{R}^{m \times m}$, the number of 0 eigenvalues is at least equal to $\dim \mathcal{N}(A)$

$$\dim \mathcal{N}(A) \leq \text{number of 0 eigenvalues} \leq m \quad (1482)$$

while the eigenvectors corresponding to those 0 eigenvalues belong to $\mathcal{N}(A)$. [294, §5.1] ^{A.17}

For diagonalizable matrix A (§A.5), the number of 0 eigenvalues is precisely $\dim \mathcal{N}(A)$ while the corresponding eigenvectors span $\mathcal{N}(A)$. The real and imaginary parts of the eigenvectors remaining span $\mathcal{R}(A)$.

(TRANSPOSE.)

Likewise, for any matrix $A \in \mathbb{R}^{m \times n}$

$$\text{rank}(A^T) + \dim \mathcal{N}(A^T) = m \quad (1483)$$

For any square $A \in \mathbb{R}^{m \times m}$, the number of 0 eigenvalues is at least equal to $\dim \mathcal{N}(A^T) = \dim \mathcal{N}(A)$ while the left-eigenvectors (eigenvectors of A^T) corresponding to those 0 eigenvalues belong to $\mathcal{N}(A^T)$.

For diagonalizable A , the number of 0 eigenvalues is precisely $\dim \mathcal{N}(A^T)$ while the corresponding left-eigenvectors span $\mathcal{N}(A^T)$. The real and imaginary parts of the left-eigenvectors remaining span $\mathcal{R}(A^T)$. \diamond

^{A.17}We take as given the well-known fact that the number of 0 eigenvalues cannot be less than the dimension of the nullspace. We offer an example of the converse:

$$A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

$\dim \mathcal{N}(A) = 2$, $\lambda(A) = [0 \ 0 \ 0 \ 1]^T$; three eigenvectors in the nullspace but only two are independent. The right-hand side of (1482) is tight for nonzero matrices; e.g., (§B.1) dyad $uv^T \in \mathbb{R}^{m \times m}$ has m 0-eigenvalues when $u \in v^\perp$.

Proof. First we show, for a diagonalizable matrix, the number of 0 eigenvalues is precisely the dimension of its nullspace while the eigenvectors corresponding to those 0 eigenvalues span the nullspace:

Any diagonalizable matrix $A \in \mathbb{R}^{m \times m}$ must possess a complete set of linearly independent eigenvectors. If A is full-rank (invertible), then all $m = \text{rank}(A)$ eigenvalues are nonzero. [294, §5.1]

Suppose $\text{rank}(A) < m$. Then $\dim \mathcal{N}(A) = m - \text{rank}(A)$. Thus there is a set of $m - \text{rank}(A)$ linearly independent vectors spanning $\mathcal{N}(A)$. Each of those can be an eigenvector associated with a 0 eigenvalue because A is diagonalizable $\Leftrightarrow \exists m$ linearly independent eigenvectors. [294, §5.2] Eigenvectors of a real matrix corresponding to 0 eigenvalues must be real. **A.18** Thus A has at least $m - \text{rank}(A)$ eigenvalues equal to 0.

Now suppose A has more than $m - \text{rank}(A)$ eigenvalues equal to 0. Then there are more than $m - \text{rank}(A)$ linearly independent eigenvectors associated with 0 eigenvalues, and each of those eigenvectors must be in $\mathcal{N}(A)$. Thus there are more than $m - \text{rank}(A)$ linearly independent vectors in $\mathcal{N}(A)$; a contradiction.

Therefore diagonalizable A has $\text{rank}(A)$ nonzero eigenvalues and exactly $m - \text{rank}(A)$ eigenvalues equal to 0 whose corresponding eigenvectors span $\mathcal{N}(A)$.

By similar argument, the left-eigenvectors corresponding to 0 eigenvalues span $\mathcal{N}(A^T)$.

Next we show when A is diagonalizable, the real and imaginary parts of its eigenvectors (corresponding to nonzero eigenvalues) span $\mathcal{R}(A)$:

The right-eigenvectors of a diagonalizable matrix $A \in \mathbb{R}^{m \times m}$ are linearly independent if and only if the left-eigenvectors are. So, matrix A has a representation in terms of its right- and left-eigenvectors; from the diagonalization (1449), assuming 0 eigenvalues are ordered last,

$$A = \sum_{i=1}^m \lambda_i s_i w_i^T = \sum_{\substack{i=1 \\ \lambda_i \neq 0}}^{k \leq m} \lambda_i s_i w_i^T \quad (1484)$$

From the *linearly independent dyads theorem* (§B.1.1.0.2), the dyads $\{s_i w_i^T\}$ must be independent because each set of eigenvectors are; hence $\text{rank } A = k$, the number of nonzero eigenvalues. Complex eigenvectors and eigenvalues

A.18 Proof. Let $*$ denote complex conjugation. Suppose $A = A^*$ and $As_i = \mathbf{0}$. Then $s_i = s_i^* \Rightarrow As_i = As_i^* \Rightarrow As_i^* = \mathbf{0}$. Conversely, $As_i^* = \mathbf{0} \Rightarrow As_i = As_i^* \Rightarrow s_i = s_i^*$. \blacklozenge

are common for real matrices, and must come in complex conjugate pairs for the summation to remain real. Assume that conjugate pairs of eigenvalues appear in sequence. Given any particular conjugate pair from (1484), we get the partial summation

$$\begin{aligned}\lambda_i s_i w_i^T + \lambda_i^* s_i^* w_i^{*\text{T}} &= 2 \operatorname{Re}(\lambda_i s_i w_i^T) \\ &= 2(\operatorname{Re} s_i \operatorname{Re}(\lambda_i w_i^T) - \operatorname{Im} s_i \operatorname{Im}(\lambda_i w_i^T))\end{aligned}\quad (1485)$$

where^{A.19} $\lambda_i^* \triangleq \lambda_{i+1}$, $s_i^* \triangleq s_{i+1}$, and $w_i^* \triangleq w_{i+1}$. Then (1484) is equivalently written

$$A = 2 \sum_{\substack{i \\ \lambda_i \in \mathbb{C} \\ \lambda_i \neq 0}} \operatorname{Re} s_{2i} \operatorname{Re}(\lambda_{2i} w_{2i}^T) - \operatorname{Im} s_{2i} \operatorname{Im}(\lambda_{2i} w_{2i}^T) + \sum_{\substack{j \\ \lambda_j \in \mathbb{R} \\ \lambda_j \neq 0}} \lambda_j s_j w_j^T \quad (1486)$$

The summation (1486) shows: A is a linear combination of real and imaginary parts of its right-eigenvectors corresponding to nonzero eigenvalues. The k vectors $\{\operatorname{Re} s_i \in \mathbb{R}^m, \operatorname{Im} s_i \in \mathbb{R}^m \mid \lambda_i \neq 0, i \in \{1 \dots m\}\}$ must therefore span the range of diagonalizable matrix A .

The argument is similar regarding the span of the left-eigenvectors. \blacklozenge

A.7.4 0 trace and matrix product

For $X, A \in \mathbb{R}_+^{M \times N}$ (33)

$$\operatorname{tr}(X^T A) = 0 \Leftrightarrow X \circ A = A \circ X = \mathbf{0} \quad (1487)$$

For $X, A \in \mathbb{S}_+^M$ [31, §2.6.1, exer.2.8] [319, §3.1]

$$\operatorname{tr}(XA) = 0 \Leftrightarrow XA = AX = \mathbf{0} \quad (1488)$$

Proof. (\Leftarrow) Suppose $XA = AX = \mathbf{0}$. Then $\operatorname{tr}(XA) = 0$ is obvious.
 (\Rightarrow) Suppose $\operatorname{tr}(XA) = 0$. $\operatorname{tr}(XA) = \operatorname{tr}(\sqrt{A} X \sqrt{A})$ whose argument is positive semidefinite by Corollary A.3.1.0.5. Trace of any square matrix is equivalent to the sum of its eigenvalues. Eigenvalues of a positive semidefinite matrix can total 0 if and only if each and every nonnegative eigenvalue

^{A.19}Complex conjugate of w is denoted w^* . Conjugate transpose is denoted $w^H = w^{*\text{T}}$.

is 0. The only feasible positive semidefinite matrix, having all 0 eigenvalues, resides at the origin; (*confer* (1512)) *id est*,

$$\sqrt{A} X \sqrt{A} = \left(\sqrt{X} \sqrt{A} \right)^T \sqrt{X} \sqrt{A} = \mathbf{0} \quad (1489)$$

implying $\sqrt{X} \sqrt{A} = \mathbf{0}$ which in turn implies $\sqrt{X} (\sqrt{X} \sqrt{A}) \sqrt{A} = X A = \mathbf{0}$. Arguing similarly yields $A X = \mathbf{0}$. \blacklozenge

Diagonalizable matrices A and X are *simultaneously diagonalizable* if and only if they are commutative under multiplication; [180, §1.3.12] *id est*, iff they share a complete set of eigenvectors.

A.7.4.0.1 Example. *An equivalence in nonisomorphic spaces.*

Identity (1488) leads to an unusual equivalence relating convex geometry to traditional linear algebra: The convex sets, given $A \succeq 0$

$$\{X \mid \langle X, A \rangle = 0\} \cap \{X \succeq 0\} \equiv \{X \mid \mathcal{N}(X) \supseteq \mathcal{R}(A)\} \cap \{X \succeq 0\} \quad (1490)$$

(one expressed in terms of a hyperplane, the other in terms of nullspace and range) are equivalent only when symmetric matrix A is positive semidefinite.

We might apply this equivalence to the geometric center subspace, for example,

$$\begin{aligned} \mathbb{S}_c^M &= \{Y \in \mathbb{S}^M \mid Y \mathbf{1} = \mathbf{0}\} \\ &= \{Y \in \mathbb{S}^M \mid \mathcal{N}(Y) \supseteq \mathbf{1}\} = \{Y \in \mathbb{S}^M \mid \mathcal{R}(Y) \subseteq \mathcal{N}(\mathbf{1}^T)\} \end{aligned} \quad (1885)$$

from which we derive (*confer* (909))

$$\mathbb{S}_c^M \cap \mathbb{S}_+^M \equiv \{X \succeq 0 \mid \langle X, \mathbf{1} \mathbf{1}^T \rangle = 0\} \quad (1491)$$

□

A.7.5 Zero definite

The domain over which an arbitrary real matrix A is zero definite can exceed its left and right nullspaces. For any positive semidefinite matrix $A \in \mathbb{R}^{M \times M}$ (for $A + A^T \succeq 0$)

$$\{x \mid x^T A x = 0\} = \mathcal{N}(A + A^T) \quad (1492)$$

because $\exists R \ni A+A^T=R^TR$, $\|Rx\|=0 \Leftrightarrow Rx=\mathbf{0}$, and $\mathcal{N}(A+A^T)=\mathcal{N}(R)$. Then given any particular vector x_p , $x_p^T A x_p = 0 \Leftrightarrow x_p \in \mathcal{N}(A+A^T)$. For any positive definite matrix A (for $A+A^T \succ 0$)

$$\{x \mid x^T A x = 0\} = \mathbf{0} \quad (1493)$$

Further, [352, §3.2, prob.5]

$$\{x \mid x^T A x = 0\} = \mathbb{R}^M \Leftrightarrow A^T = -A \quad (1494)$$

while

$$\{x \mid x^H A x = 0\} = \mathbb{C}^M \Leftrightarrow A = \mathbf{0} \quad (1495)$$

The positive semidefinite matrix

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \quad (1496)$$

for example, has no nullspace. Yet

$$\{x \mid x^T A x = 0\} = \{x \mid \mathbf{1}^T x = 0\} \subset \mathbb{R}^2 \quad (1497)$$

which is the nullspace of the symmetrized matrix. Symmetric matrices are not spared from the excess; *videlicet*,

$$B = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \quad (1498)$$

has eigenvalues $\{-1, 3\}$, no nullspace, but is zero definite on ^{A.20}

$$\mathcal{X} \triangleq \{x \in \mathbb{R}^2 \mid x_2 = (-2 \pm \sqrt{3})x_1\} \quad (1499)$$

^{A.20}These two lines represent the limit in the union of two generally distinct hyperbolae; *id est*, for matrix B and set \mathcal{X} as defined

$$\lim_{\varepsilon \rightarrow 0^+} \{x \in \mathbb{R}^2 \mid x^T B x = \varepsilon\} = \mathcal{X}$$

A.7.5.0.1 Proposition. (Sturm) *Dyad-decompositions.* [299, §5.2]

Let positive semidefinite matrix $X \in \mathbb{S}_+^M$ have rank ρ . Then given symmetric matrix $A \in \mathbb{S}^M$, $\langle A, X \rangle = 0$ if and only if there exists a dyad-decomposition

$$X = \sum_{j=1}^{\rho} x_j x_j^T \quad (1500)$$

satisfying

$$\langle A, x_j x_j^T \rangle = 0 \text{ for each and every } j \in \{1 \dots \rho\} \quad (1501)$$

◇

The dyad-decomposition of X proposed is generally not that obtained from a standard diagonalization by eigen decomposition, unless $\rho = 1$ or the given matrix A is diagonalizable simultaneously (§A.7.4) with X . That means, elemental dyads in decomposition (1500) constitute a generally nonorthogonal set. Sturm & Zhang give a simple procedure for constructing the dyad-decomposition (*Wikimization*); matrix A may be regarded as a parameter.

A.7.5.0.2 Example. *Dyad.*

The dyad $uv^T \in \mathbb{R}^{M \times M}$ (§B.1) is zero definite on all x for which either $x^T u = 0$ or $x^T v = 0$;

$$\{x \mid x^T uv^T x = 0\} = \{x \mid x^T u = 0\} \cup \{x \mid v^T x = 0\} \quad (1502)$$

id est, on $u^\perp \cup v^\perp$. Symmetrizing the dyad does not change the outcome:

$$\{x \mid x^T (uv^T + vu^T)x/2 = 0\} = \{x \mid x^T u = 0\} \cup \{x \mid v^T x = 0\} \quad (1503)$$

□

Appendix B

Simple matrices

*Mathematicians also attempted to develop algebra of vectors but there was no natural definition of the product of two vectors that held in arbitrary dimensions. The first vector algebra that involved a noncommutative vector product (that is, $v \times w$ need not equal $w \times v$) was proposed by Hermann Grassmann in his book *Ausdehnungslehre* (1844). Grassmann's text also introduced the product of a column matrix and a row matrix, which resulted in what is now called a simple or a rank-one matrix. In the late 19th century the American mathematical physicist Willard Gibbs published his famous treatise on vector analysis. In that treatise Gibbs represented general matrices, which he called dyadics, as sums of simple matrices, which Gibbs called dyads. Later the physicist P. A. M. Dirac introduced the term "bra-ket" for what we now call the scalar product of a "bra" (row) vector times a "ket" (column) vector and the term "ket-bra" for the product of a ket times a bra, resulting in what we now call a simple matrix, as above. Our convention of identifying column matrices and vectors was introduced by physicists in the 20th century.*

–Marie A. Vitulli [328]

B.1 Rank-one matrix (dyad)

Any matrix formed from the unsigned outer product of two vectors,

$$\Psi = uv^T \in \mathbb{R}^{M \times N} \quad (1504)$$

where $u \in \mathbb{R}^M$ and $v \in \mathbb{R}^N$, is rank-one and called a *dyad*. Conversely, any rank-one matrix must have the form Ψ . [180, prob.1.4.1] Product $-uv^T$ is a *negative dyad*. For matrix products AB^T , in general, we have

$$\mathcal{R}(AB^T) \subseteq \mathcal{R}(A), \quad \mathcal{N}(AB^T) \supseteq \mathcal{N}(B^T) \quad (1505)$$

with equality when $B=A$ [294, §3.3, §3.6]^{B.1} or respectively when B is invertible and $\mathcal{N}(A)=\mathbf{0}$. Yet for all nonzero dyads we have

$$\mathcal{R}(uv^T) = \mathcal{R}(u), \quad \mathcal{N}(uv^T) = \mathcal{N}(v^T) \equiv v^\perp \quad (1506)$$

where $\dim v^\perp = N - 1$.

It is obvious a dyad can be $\mathbf{0}$ only when u or v is $\mathbf{0}$;

$$\Psi = uv^T = \mathbf{0} \Leftrightarrow u = \mathbf{0} \text{ or } v = \mathbf{0} \quad (1507)$$

The matrix 2-norm for Ψ is equivalent to the Frobenius norm;

$$\|\Psi\|_2 = \sigma_1 = \|uv^T\|_F = \|uv^T\|_2 = \|u\| \|v\| \quad (1508)$$

When u and v are normalized, the pseudoinverse is the transposed dyad. Otherwise,

$$\Psi^\dagger = (uv^T)^\dagger = \frac{vu^T}{\|u\|^2 \|v\|^2} \quad (1509)$$

^{B.1}**Proof.** $\mathcal{R}(AA^T) \subseteq \mathcal{R}(A)$ is obvious.

$$\begin{aligned} \mathcal{R}(AA^T) &= \{AA^T y \mid y \in \mathbb{R}^m\} \\ &\supseteq \{AA^T y \mid A^T y \in \mathcal{R}(A^T)\} = \mathcal{R}(A) \text{ by (133)} \quad \blacklozenge \end{aligned}$$




Figure 135: The four fundamental subspaces [296, §3.6] of any dyad $\Psi = uv^T \in \mathbb{R}^{M \times N}$. $\Psi(x) \triangleq uv^T x$ is a linear mapping from \mathbb{R}^N to \mathbb{R}^M . The map from $\mathcal{R}(v)$ to $\mathcal{R}(u)$ is bijective. [294, §3.1]

When dyad $uv^T \in \mathbb{R}^{N \times N}$ is square, uv^T has at least $N - 1$ 0-eigenvalues and corresponding eigenvectors spanning v^\perp . The remaining eigenvector u spans the range of uv^T with corresponding eigenvalue

$$\lambda = v^T u = \text{tr}(uv^T) \in \mathbb{R} \tag{1510}$$

Determinant is a product of the eigenvalues; so, it is always true that

$$\det \Psi = \det(uv^T) = 0 \tag{1511}$$

When $\lambda = 1$, the square dyad is a nonorthogonal projector projecting on its range ($\Psi^2 = \Psi$, §E.6); a *projector dyad*. It is quite possible that $u \in v^\perp$ making the remaining eigenvalue instead 0; **B.2** $\lambda = 0$ together with the first $N - 1$ 0-eigenvalues; *id est*, it is possible uv^T were nonzero while all its eigenvalues are 0. The matrix

$$\begin{bmatrix} 1 \\ -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix} \tag{1512}$$

for example, has two 0-eigenvalues. In other words, eigenvector u may simultaneously be a member of the nullspace and range of the dyad. The explanation is, simply, because u and v share the same dimension, $\dim u = M = \dim v = N$:

B.2A dyad is not always diagonalizable (§A.5) because its eigenvectors are not necessarily independent.

Proof. Figure 135 shows the four fundamental subspaces for the dyad. Linear operator $\Psi : \mathbb{R}^N \rightarrow \mathbb{R}^M$ provides a map between vector spaces that remain distinct when $M = N$;

$$\begin{aligned} u &\in \mathcal{R}(uv^T) \\ u &\in \mathcal{N}(uv^T) \Leftrightarrow v^T u = 0 \\ \mathcal{R}(uv^T) \cap \mathcal{N}(uv^T) &= \emptyset \end{aligned} \quad (1513)$$

◆

B.1.0.1 rank-one modification

If $A \in \mathbb{R}^{N \times N}$ is any nonsingular matrix and $1 + v^T A^{-1} u \neq 0$, then [196, App.6] [352, §2.3, prob.16] [132, §4.11.2] (Sherman-Morrison)

$$(A + uv^T)^{-1} = A^{-1} - \frac{A^{-1} uv^T A^{-1}}{1 + v^T A^{-1} u} \quad (1514)$$

B.1.0.2 dyad symmetry

In the specific circumstance that $v = u$, then $uu^T \in \mathbb{R}^{N \times N}$ is symmetric, rank-one, and positive semidefinite having exactly $N - 1$ 0-eigenvalues. In fact, (Theorem A.3.1.0.7)

$$uv^T \succeq 0 \Leftrightarrow v = u \quad (1515)$$

and the remaining eigenvalue is almost always positive;

$$\lambda = u^T u = \text{tr}(uu^T) > 0 \text{ unless } u = \mathbf{0} \quad (1516)$$

The matrix

$$\begin{bmatrix} \Psi & u \\ u^T & 1 \end{bmatrix} \quad (1517)$$

for example, is rank-1 positive semidefinite if and only if $\Psi = uu^T$.

B.1.1 Dyad independence

Now we consider a sum of dyads like (1504) as encountered in diagonalization and singular value decomposition:

$$\mathcal{R}\left(\sum_{i=1}^k s_i w_i^T\right) = \sum_{i=1}^k \mathcal{R}(s_i w_i^T) = \sum_{i=1}^k \mathcal{R}(s_i) \Leftrightarrow w_i \forall i \text{ are l.i.} \quad (1518)$$

range of summation is the vector sum of ranges.^{B.3} (Theorem B.1.1.1.1) Under the assumption the dyads are linearly independent (l.i.), then the vector sums are unique (p.714): for $\{w_i\}$ l.i. and $\{s_i\}$ l.i.

$$\mathcal{R}\left(\sum_{i=1}^k s_i w_i^T\right) = \mathcal{R}(s_1 w_1^T) \oplus \dots \oplus \mathcal{R}(s_k w_k^T) = \mathcal{R}(s_1) \oplus \dots \oplus \mathcal{R}(s_k) \quad (1519)$$

B.1.1.0.1 Definition. *Linearly independent dyads.* [186, p.29, thm.11] [301, p.2] The set of k dyads

$$\{s_i w_i^T \mid i=1 \dots k\} \quad (1520)$$

where $s_i \in \mathbb{C}^M$ and $w_i \in \mathbb{C}^N$, is said to be linearly independent iff

$$\text{rank}\left(SW^T \triangleq \sum_{i=1}^k s_i w_i^T\right) = k \quad (1521)$$

where $S \triangleq [s_1 \dots s_k] \in \mathbb{C}^{M \times k}$ and $W \triangleq [w_1 \dots w_k] \in \mathbb{C}^{N \times k}$. \triangle

As defined, dyad independence does not preclude existence of a nullspace $\mathcal{N}(SW^T)$, nor does it imply SW^T is full-rank. In absence of an assumption of independence, generally, $\text{rank } SW^T \leq k$. Conversely, any rank- k matrix can be written in the form SW^T by singular value decomposition. (§A.6)

B.1.1.0.2 Theorem. *Linearly independent (l.i.) dyads.*

Vectors $\{s_i \in \mathbb{C}^M, i=1 \dots k\}$ are l.i. and vectors $\{w_i \in \mathbb{C}^N, i=1 \dots k\}$ are l.i. if and only if dyads $\{s_i w_i^T \in \mathbb{C}^{M \times N}, i=1 \dots k\}$ are l.i. \diamond

Proof. Linear independence of k dyads is identical to definition (1521). (\Rightarrow) Suppose $\{s_i\}$ and $\{w_i\}$ are each linearly independent sets. Invoking Sylvester's rank inequality, [180, §0.4] [352, §2.4]

$$\text{rank } S + \text{rank } W - k \leq \text{rank}(SW^T) \leq \min\{\text{rank } S, \text{rank } W\} (\leq k) \quad (1522)$$

Then $k \leq \text{rank}(SW^T) \leq k$ which implies the dyads are independent.

(\Leftarrow) Conversely, suppose $\text{rank}(SW^T) = k$. Then

$$k \leq \min\{\text{rank } S, \text{rank } W\} \leq k \quad (1523)$$

implying the vector sets are each independent. \blacklozenge

^{B.3}Move of range \mathcal{R} to inside the summation depends on linear independence of $\{w_i\}$.

B.1.1.1 Biorthogonality condition, Range and Nullspace of Sum

Dyads characterized by a biorthogonality condition $W^T S = I$ are independent; *id est*, for $S \in \mathbb{C}^{M \times k}$ and $W \in \mathbb{C}^{N \times k}$, if $W^T S = I$ then $\text{rank}(SW^T) = k$ by the *linearly independent dyads theorem* because (confer §E.1.1)

$$W^T S = I \Rightarrow \text{rank } S = \text{rank } W = k \leq M = N \quad (1524)$$

To see that, we need only show: $\mathcal{N}(S) = \mathbf{0} \Leftrightarrow \exists B \ni BS = I$. **B.4**
 (\Leftarrow) Assume $BS = I$. Then $\mathcal{N}(BS) = \mathbf{0} = \{x \mid BSx = \mathbf{0}\} \supseteq \mathcal{N}(S)$. (1505)
 (\Rightarrow) If $\mathcal{N}(S) = \mathbf{0}$ then S must be full-rank skinny-or-square.

$$\therefore \exists A, B, C \ni \begin{bmatrix} B \\ C \end{bmatrix} [S \ A] = I \text{ (id est, } [S \ A] \text{ is invertible)} \Rightarrow BS = I.$$

Left inverse B is given as W^T here. Because of reciprocity with S , it immediately follows: $\mathcal{N}(W) = \mathbf{0} \Leftrightarrow \exists S \ni S^T W = I$. \blacklozenge

Dyads produced by diagonalization, for example, are independent because of their inherent biorthogonality. (§A.5.1) The converse is generally false; *id est*, linearly independent dyads are not necessarily biorthogonal.

B.1.1.1.1 Theorem. Nullspace and range of dyad sum.

Given a sum of dyads represented by SW^T where $S \in \mathbb{C}^{M \times k}$ and $W \in \mathbb{C}^{N \times k}$

$$\begin{aligned} \mathcal{N}(SW^T) = \mathcal{N}(W^T) &\Leftarrow \exists B \ni BS = I \\ \mathcal{R}(SW^T) = \mathcal{R}(S) &\Leftarrow \exists Z \ni W^T Z = I \end{aligned} \quad (1525)$$

\diamond

Proof. (\Rightarrow) $\mathcal{N}(SW^T) \supseteq \mathcal{N}(W^T)$ and $\mathcal{R}(SW^T) \subseteq \mathcal{R}(S)$ are obvious.
 (\Leftarrow) Assume the existence of a left inverse $B \in \mathbb{R}^{k \times N}$ and a right inverse $Z \in \mathbb{R}^{N \times k}$. **B.5**

$$\mathcal{N}(SW^T) = \{x \mid SW^T x = \mathbf{0}\} \subseteq \{x \mid BSW^T x = \mathbf{0}\} = \mathcal{N}(W^T) \quad (1526)$$

$$\mathcal{R}(SW^T) = \{SW^T x \mid x \in \mathbb{R}^N\} \supseteq \{SW^T Z y \mid Z y \in \mathbb{R}^N\} = \mathcal{R}(S) \quad (1527)$$

\blacklozenge

B.4Left inverse is not unique, in general.

B.5By counter-example, the theorem's converse cannot be true; *e.g.*, $S = W = [\mathbf{1} \ \mathbf{0}]$.




Figure 136: Four fundamental subspaces [296, §3.6] of a doublet $\Pi = uv^T + vu^T \in \mathbb{S}^N$. $\Pi(x) = (uv^T + vu^T)x$ is a linear bijective mapping from $\mathcal{R}([u \ v])$ to $\mathcal{R}([u \ v])$.

B.2 Doublet

Consider a sum of two linearly independent square dyads, one a transposition of the other:

$$\Pi = uv^T + vu^T = [u \ v] \begin{bmatrix} v^T \\ u^T \end{bmatrix} = SW^T \in \mathbb{S}^N \quad (1528)$$

where $u, v \in \mathbb{R}^N$. Like the dyad, a doublet can be $\mathbf{0}$ only when u or v is $\mathbf{0}$;

$$\Pi = uv^T + vu^T = \mathbf{0} \Leftrightarrow u = \mathbf{0} \text{ or } v = \mathbf{0} \quad (1529)$$

By assumption of independence, a nonzero doublet has two nonzero eigenvalues

$$\lambda_1 \triangleq u^T v + \|uv^T\|, \quad \lambda_2 \triangleq u^T v - \|uv^T\| \quad (1530)$$

where $\lambda_1 > 0 > \lambda_2$, with corresponding eigenvectors

$$x_1 \triangleq \frac{u}{\|u\|} + \frac{v}{\|v\|}, \quad x_2 \triangleq \frac{u}{\|u\|} - \frac{v}{\|v\|} \quad (1531)$$

spanning the doublet range. Eigenvalue λ_1 cannot be 0 unless u and v have opposing directions, but that is antithetical since then the dyads would no longer be independent. Eigenvalue λ_2 is 0 if and only if u and v share the same direction, again antithetical. Generally we have $\lambda_1 > 0$ and $\lambda_2 < 0$, so Π is indefinite.




Figure 137: $v^T u = 1/\zeta$. The four fundamental subspaces [296, §3.6] of elementary matrix E as a linear mapping $E(x) = \left(I - \frac{uv^T}{v^T u}\right)x$.

By the *nullspace and range of dyad sum theorem*, doublet Π has $N-2$ zero-eigenvalues remaining and corresponding eigenvectors spanning $\mathcal{N}\left(\begin{bmatrix} v^T \\ u^T \end{bmatrix}\right)$. We therefore have

$$\mathcal{R}(\Pi) = \mathcal{R}([u \ v]), \quad \mathcal{N}(\Pi) = v^\perp \cap u^\perp \quad (1532)$$

of respective dimension 2 and $N-2$.

B.3 Elementary matrix

A matrix of the form

$$E = I - \zeta uv^T \in \mathbb{R}^{N \times N} \quad (1533)$$

where $\zeta \in \mathbb{R}$ is finite and $u, v \in \mathbb{R}^N$, is called an *elementary matrix* or a *rank-one modification of the identity*. [182] Any elementary matrix in $\mathbb{R}^{N \times N}$ has $N-1$ eigenvalues equal to 1 corresponding to real eigenvectors that span v^\perp . The remaining eigenvalue

$$\lambda = 1 - \zeta v^T u \quad (1534)$$

corresponds to eigenvector u . ^{B.6} From [196, App.7.A.26] the determinant:

$$\det E = 1 - \text{tr}(\zeta uv^T) = \lambda \quad (1535)$$

^{B.6}Elementary matrix E is not always diagonalizable because eigenvector u need not be independent of the others; *id est*, $u \in v^\perp$ is possible.

If $\lambda \neq 0$ then E is invertible; [132]

$$E^{-1} = I + \frac{\zeta}{\lambda} uv^T \tag{1536}$$

Eigenvectors corresponding to 0 eigenvalues belong to $\mathcal{N}(E)$, and the number of 0 eigenvalues must be at least $\dim \mathcal{N}(E)$ which, here, can be at most one. (§A.7.3.0.1) The nullspace exists, therefore, when $\lambda=0$; *id est*, when $v^T u = 1/\zeta$, rather, whenever u belongs to the hyperplane $\{z \in \mathbb{R}^N \mid v^T z = 1/\zeta\}$. Then (when $\lambda=0$) elementary matrix E is a nonorthogonal projector projecting on its range ($E^2 = E$, §E.1) and $\mathcal{N}(E) = \mathcal{R}(u)$; eigenvector u spans the nullspace when it exists. By conservation of dimension, $\dim \mathcal{R}(E) = N - \dim \mathcal{N}(E)$. It is apparent from (1533) that $v^\perp \subseteq \mathcal{R}(E)$, but $\dim v^\perp = N - 1$. Hence $\mathcal{R}(E) \equiv v^\perp$ when the nullspace exists, and the remaining eigenvectors span it.

In summary, when a nontrivial nullspace of E exists,

$$\mathcal{R}(E) = \mathcal{N}(v^T), \quad \mathcal{N}(E) = \mathcal{R}(u), \quad v^T u = 1/\zeta \tag{1537}$$

illustrated in Figure 137, which is opposite to the assignment of subspaces for a dyad (Figure 135). Otherwise, $\mathcal{R}(E) = \mathbb{R}^N$.

When $E = E^T$, the spectral norm is

$$\|E\|_2 = \max\{1, |\lambda|\} \tag{1538}$$

B.3.1 Householder matrix

An elementary matrix is called a Householder matrix when it has the defining form, for nonzero vector u [139, §5.1.2] [132, §4.10.1] [294, §7.3] [180, §2.2]

$$H = I - 2 \frac{uu^T}{u^T u} \in \mathbb{S}^N \tag{1539}$$

which is a symmetric orthogonal (reflection) matrix ($H^{-1} = H^T = H$ (§B.5.2)). Vector u is normal to an $N - 1$ -dimensional subspace u^\perp through which this particular H effects pointwise reflection; *e.g.*, $Hu^\perp = u^\perp$ while $Hu = -u$.

Matrix H has $N - 1$ orthonormal eigenvectors spanning that reflecting subspace u^\perp with corresponding eigenvalues equal to 1. The remaining eigenvector u has corresponding eigenvalue -1 ; so

$$\det H = -1 \tag{1540}$$

Due to symmetry of H , the matrix 2-norm (the spectral norm) is equal to the largest eigenvalue-magnitude. A Householder matrix is thus characterized,

$$H^T = H, \quad H^{-1} = H^T, \quad \|H\|_2 = 1, \quad H \neq 0 \quad (1541)$$

For example, the permutation matrix

$$\Xi = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad (1542)$$

is a Householder matrix having $u = [0 \ 1 \ -1]^T / \sqrt{2}$. Not all permutation matrices are Householder matrices, although all permutation matrices are orthogonal matrices (§B.5.1, $\Xi^T \Xi = I$) [294, §3.4] because they are made by permuting rows and columns of the identity matrix. Neither are all symmetric permutation matrices Householder matrices; *e.g.*,

$$\Xi = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \quad (1619) \text{ is not a Householder matrix.}$$

B.4 Auxiliary V -matrices

B.4.1 Auxiliary projector matrix V

It is convenient to define a matrix V that arises naturally as a consequence of translating the geometric center α_c (§5.5.1.0.1) of some list X to the origin. In place of $X - \alpha_c \mathbf{1}^T$ we may write XV as in (889) where

$$V = I - \frac{1}{N} \mathbf{1} \mathbf{1}^T \in \mathbb{S}^N \quad (831)$$

is an elementary matrix called the *geometric centering matrix*.

Any elementary matrix in $\mathbb{R}^{N \times N}$ has $N - 1$ eigenvalues equal to 1. For the particular elementary matrix V , the N^{th} eigenvalue equals 0. The number of 0 eigenvalues must equal $\dim \mathcal{N}(V) = 1$, by the 0 *eigenvalues theorem* (§A.7.3.0.1), because $V = V^T$ is diagonalizable. Because

$$V \mathbf{1} = \mathbf{0} \quad (1543)$$

the nullspace $\mathcal{N}(V) = \mathcal{R}(\mathbf{1})$ is spanned by the eigenvector $\mathbf{1}$. The remaining eigenvectors span $\mathcal{R}(V) \equiv \mathbf{1}^\perp = \mathcal{N}(\mathbf{1}^\top)$ that has dimension $N - 1$.

Because

$$V^2 = V \tag{1544}$$

and $V^\top = V$, elementary matrix V is also a projection matrix (§E.3) projecting orthogonally on its range $\mathcal{N}(\mathbf{1}^\top)$ which is a hyperplane containing the origin in \mathbb{R}^N

$$V = I - \mathbf{1}(\mathbf{1}^\top \mathbf{1})^{-1} \mathbf{1}^\top \tag{1545}$$

The $\{0, 1\}$ eigenvalues also indicate diagonalizable V is a projection matrix. [352, §4.1, thm.4.1] Symmetry of V denotes orthogonal projection; from (1802),

$$V^\top = V, \quad V^\dagger = V, \quad \|V\|_2 = 1, \quad V \succeq 0 \tag{1546}$$

Matrix V is also circulant [147].

B.4.1.0.1 Example. *Relationship of auxiliary to Householder matrix.*

Let $H \in \mathbb{S}^N$ be a Householder matrix (1539) defined by

$$u = \begin{bmatrix} 1 \\ \vdots \\ 1 \\ 1 + \sqrt{N} \end{bmatrix} \in \mathbb{R}^N \tag{1547}$$

Then we have [135, §2]

$$V = H \begin{bmatrix} I & \mathbf{0} \\ \mathbf{0}^\top & 0 \end{bmatrix} H \tag{1548}$$

Let $D \in \mathbb{S}_h^N$ and define

$$-HDH \triangleq - \begin{bmatrix} A & b \\ b^\top & c \end{bmatrix} \tag{1549}$$

where b is a vector. Then because H is nonsingular (§A.3.1.0.5) [163, §3]

$$-VDV = -H \begin{bmatrix} A & \mathbf{0} \\ \mathbf{0}^\top & 0 \end{bmatrix} H \succeq 0 \Leftrightarrow -A \succeq 0 \tag{1550}$$

and affine dimension is $r = \text{rank } A$ when D is a Euclidean distance matrix.

□

B.4.2 Schoenberg auxiliary matrix $V_{\mathcal{N}}$

1. $V_{\mathcal{N}} = \frac{1}{\sqrt{2}} \begin{bmatrix} -\mathbf{1}^T \\ I \end{bmatrix} \in \mathbb{R}^{N \times N-1}$
2. $V_{\mathcal{N}}^T \mathbf{1} = \mathbf{0}$
3. $I - e_1 \mathbf{1}^T = [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]$
4. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] V_{\mathcal{N}} = V_{\mathcal{N}}$
5. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] V = V$
6. $V [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] = [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]$
7. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] = [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]$
8. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]^\dagger = \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} V$
9. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]^\dagger V = [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]^\dagger$
10. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]^\dagger = V$
11. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]^\dagger [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] = \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix}$
12. $[\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} = [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}]$
13. $\begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} [\mathbf{0} \quad \sqrt{2} V_{\mathcal{N}}] = \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix}$
14. $[V_{\mathcal{N}} \quad \frac{1}{\sqrt{2}} \mathbf{1}]^{-1} = \begin{bmatrix} V_{\mathcal{N}}^\dagger \\ \frac{\sqrt{2}}{N} \mathbf{1}^T \end{bmatrix}$
15. $V_{\mathcal{N}}^\dagger = \sqrt{2} [-\frac{1}{N} \mathbf{1} \quad I - \frac{1}{N} \mathbf{1} \mathbf{1}^T] \in \mathbb{R}^{N-1 \times N}, \quad (I - \frac{1}{N} \mathbf{1} \mathbf{1}^T \in \mathbb{S}^{N-1})$
16. $V_{\mathcal{N}}^\dagger \mathbf{1} = \mathbf{0}$
17. $V_{\mathcal{N}}^\dagger V_{\mathcal{N}} = I$

18. $V^T = V = V_{\mathcal{N}}V_{\mathcal{N}}^\dagger = I - \frac{1}{N}\mathbf{1}\mathbf{1}^T \in \mathbb{S}^N$

19. $-V_{\mathcal{N}}^\dagger(\mathbf{1}\mathbf{1}^T - I)V_{\mathcal{N}} = I$, $(\mathbf{1}\mathbf{1}^T - I \in \mathbb{EDM}^N)$

20. $D = [d_{ij}] \in \mathbb{S}_h^N$ (833)

$$\text{tr}(-VDV) = \text{tr}(-VD) = \text{tr}(-V_{\mathcal{N}}^\dagger DV_{\mathcal{N}}) = \frac{1}{N}\mathbf{1}^T D \mathbf{1} = \frac{1}{N} \text{tr}(\mathbf{1}\mathbf{1}^T D) = \frac{1}{N} \sum_{i,j} d_{ij}$$

Any elementary matrix $E \in \mathbb{S}^N$ of the particular form

$$E = k_1 I - k_2 \mathbf{1}\mathbf{1}^T \tag{1551}$$

where $k_1, k_2 \in \mathbb{R}$, **B.7** will make $\text{tr}(-ED)$ proportional to $\sum d_{ij}$.

21. $D = [d_{ij}] \in \mathbb{S}^N$

$$\text{tr}(-VDV) = \frac{1}{N} \sum_{\substack{i,j \\ i \neq j}} d_{ij} - \frac{N-1}{N} \sum_i d_{ii} = \mathbf{1}^T D \mathbf{1} \frac{1}{N} - \text{tr} D$$

22. $D = [d_{ij}] \in \mathbb{S}_h^N$

$$\text{tr}(-V_{\mathcal{N}}^T D V_{\mathcal{N}}) = \sum_j d_{1j}$$

23. For $Y \in \mathbb{S}^N$

$$V(Y - \delta(Y\mathbf{1}))V = Y - \delta(Y\mathbf{1})$$

B.4.3 Orthonormal auxiliary matrix $V_{\mathcal{W}}$

The skinny matrix

$$V_{\mathcal{W}} \triangleq \begin{bmatrix} \frac{-1}{\sqrt{N}} & \frac{-1}{\sqrt{N}} & \cdots & \frac{-1}{\sqrt{N}} \\ 1 + \frac{-1}{N+\sqrt{N}} & \frac{-1}{N+\sqrt{N}} & \cdots & \frac{-1}{N+\sqrt{N}} \\ \frac{-1}{N+\sqrt{N}} & \ddots & \ddots & \frac{-1}{N+\sqrt{N}} \\ \vdots & \ddots & \ddots & \vdots \\ \frac{-1}{N+\sqrt{N}} & \frac{-1}{N+\sqrt{N}} & \cdots & 1 + \frac{-1}{N+\sqrt{N}} \end{bmatrix} \in \mathbb{R}^{N \times N-1} \tag{1552}$$

B.7 If k_1 is $1-\rho$ while k_2 equals $-\rho \in \mathbb{R}$, then all eigenvalues of E for $-1/(N-1) < \rho < 1$ are guaranteed positive and therefore E is guaranteed positive definite. [266]

has $\mathcal{R}(V_{\mathcal{W}}) = \mathcal{N}(\mathbf{1}^T)$ and orthonormal columns. [5] We defined three auxiliary V -matrices: V , $V_{\mathcal{N}}$ (814), and $V_{\mathcal{W}}$ sharing some attributes listed in Table B.4.4. For example, V can be expressed

$$V = V_{\mathcal{W}}V_{\mathcal{W}}^T = V_{\mathcal{N}}V_{\mathcal{N}}^\dagger \quad (1553)$$

but $V_{\mathcal{W}}^T V_{\mathcal{W}} = I$ means V is an orthogonal projector (1799) and

$$V_{\mathcal{W}}^\dagger = V_{\mathcal{W}}^T, \quad \|V_{\mathcal{W}}\|_2 = 1, \quad V_{\mathcal{W}}^T \mathbf{1} = \mathbf{0} \quad (1554)$$

B.4.4 Auxiliary V -matrix Table

	$\dim V$	$\text{rank } V$	$\mathcal{R}(V)$	$\mathcal{N}(V^T)$	$V^T V$	$V V^T$	$V V^\dagger$
V	$N \times N$	$N - 1$	$\mathcal{N}(\mathbf{1}^T)$	$\mathcal{R}(\mathbf{1})$	V	V	V
$V_{\mathcal{N}}$	$N \times (N - 1)$	$N - 1$	$\mathcal{N}(\mathbf{1}^T)$	$\mathcal{R}(\mathbf{1})$	$\frac{1}{2}(I + \mathbf{1}\mathbf{1}^T)$	$\frac{1}{2} \begin{bmatrix} N-1 & -\mathbf{1}^T \\ -\mathbf{1} & I \end{bmatrix}$	V
$V_{\mathcal{W}}$	$N \times (N - 1)$	$N - 1$	$\mathcal{N}(\mathbf{1}^T)$	$\mathcal{R}(\mathbf{1})$	I	V	V

B.4.5 More auxiliary matrices

Mathar shows [229, §2] that any elementary matrix (§B.3) of the form

$$V_{\mathcal{M}} = I - b\mathbf{1}^T \in \mathbb{R}^{N \times N} \quad (1555)$$

such that $b^T \mathbf{1} = 1$ (confer [141, §2]), is an auxiliary V -matrix having

$$\begin{aligned} \mathcal{R}(V_{\mathcal{M}}^T) &= \mathcal{N}(b^T), & \mathcal{R}(V_{\mathcal{M}}) &= \mathcal{N}(\mathbf{1}^T) \\ \mathcal{N}(V_{\mathcal{M}}) &= \mathcal{R}(b), & \mathcal{N}(V_{\mathcal{M}}^T) &= \mathcal{R}(\mathbf{1}) \end{aligned} \quad (1556)$$

Given $X \in \mathbb{R}^{n \times N}$, the choice $b = \frac{1}{N} \mathbf{1}$ ($V_{\mathcal{M}} = V$) minimizes $\|X(I - b\mathbf{1}^T)\|_F$. [143, §3.2.1]

B.5 Orthogonal matrix

B.5.1 Vector rotation

The property $Q^{-1} = Q^T$ completely defines an orthogonal matrix $Q \in \mathbb{R}^{n \times n}$ employed to effect vector rotation; [294, §2.6, §3.4] [296, §6.5] [180, §2.1] for any $x \in \mathbb{R}^n$

$$\|Qx\| = \|x\| \quad (1557)$$

A *unitary matrix* is a complex generalization of the orthogonal matrix. The conjugate transpose defines it: $U^{-1} = U^H$. An orthogonal matrix is simply a real unitary matrix.

Orthogonal matrix Q is a normal matrix further characterized:

$$Q^{-1} = Q^T, \quad \|Q\|_2 = 1 \quad (1558)$$

Applying characterization (1558) to Q^T we see it too is an orthogonal matrix. Hence the rows and columns of Q respectively form an orthonormal set. Normalcy guarantees diagonalization (§A.5.2) so, for $Q \triangleq S\Lambda S^H$

$$S\Lambda^{-1}S^H = S^*\Lambda S^T, \quad \|\delta(\Lambda)\|_\infty = 1 \quad (1559)$$

characterizes an orthogonal matrix in terms of eigenvalues and eigenvectors.

All permutation matrices Ξ , for example, are orthogonal matrices. Any product of permutation matrices remains a permutator. Any product of a permutation matrix with an orthogonal matrix remains orthogonal. In fact, any product of orthogonal matrices AQ remains orthogonal by definition. Given any other dimensionally compatible orthogonal matrix U , the mapping $g(A) = U^T A Q$ is a linear bijection on the domain of orthogonal matrices (a nonconvex manifold of dimension $\frac{1}{2}n(n-1)$ [45]). [211, §2.1] [212]

The largest magnitude entry of an orthogonal matrix is 1; for each and every $j \in 1 \dots n$

$$\begin{aligned} \|Q(j, :)\|_\infty &\leq 1 \\ \|Q(:, j)\|_\infty &\leq 1 \end{aligned} \quad (1560)$$

Each and every eigenvalue of a (real) orthogonal matrix has magnitude 1

$$\lambda(Q) \in \mathbb{C}^n, \quad |\lambda(Q)| = 1 \quad (1561)$$

while only the identity matrix can be simultaneously positive definite and orthogonal.

B.5.2 Reflection

A matrix for pointwise reflection is defined by imposing symmetry upon the orthogonal matrix; *id est*, a *reflection matrix* is completely defined by $Q^{-1} = Q^T = Q$. The reflection matrix is an orthogonal matrix, characterized:

$$Q^T = Q, \quad Q^{-1} = Q^T, \quad \|Q\|_2 = 1 \quad (1562)$$

The Householder matrix (§B.3.1) is an example of a symmetric orthogonal (reflection) matrix.

Reflection matrices have eigenvalues equal to ± 1 and so $\det Q = \pm 1$. It is natural to expect a relationship between reflection and projection matrices because all projection matrices have eigenvalues belonging to $\{0, 1\}$. In fact, any reflection matrix Q is related to some orthogonal projector P by [182, §1, prob.44]

$$Q = I - 2P \quad (1563)$$

Yet P is, generally, neither orthogonal or invertible. (§E.3.2)

$$\lambda(Q) \in \mathbb{R}^n, \quad |\lambda(Q)| = 1 \quad (1564)$$

Reflection is with respect to $\mathcal{R}(P)^\perp$. Matrix $2P - I$ represents antireflection.

Every orthogonal matrix can be expressed as the product of a rotation and a reflection. The collection of all orthogonal matrices of particular dimension does not form a convex set.

B.5.3 Rotation of range and rowspace

Given orthogonal matrix Q , column vectors of a matrix X are simultaneously rotated about the origin via product QX . In three dimensions ($X \in \mathbb{R}^{3 \times N}$), the precise meaning of rotation is best illustrated in Figure 138 where the gimbal aids visualization of achievable rotation.

B.5.3.0.1 Example. *One axis of revolution.*

Partition an $n + 1$ -dimensional Euclidean space $\mathbb{R}^{n+1} \triangleq \begin{bmatrix} \mathbb{R}^n \\ \mathbb{R} \end{bmatrix}$ and define an n -dimensional subspace

$$\mathcal{R} \triangleq \{\lambda \in \mathbb{R}^{n+1} \mid \mathbf{1}^T \lambda = 0\} \quad (1565)$$




Figure 138: *Gimbal*: a mechanism imparting three degrees of dimensional freedom to a Euclidean body suspended at the device's center. Each ring is free to rotate about one axis. (Courtesy of The MathWorks Inc.)

(a hyperplane through the origin). We want an orthogonal matrix that rotates a list in the columns of matrix $X \in \mathbb{R}^{n+1 \times N}$ through the dihedral angle between \mathbb{R}^n and \mathcal{R} (§2.4.3)

$$\sphericalangle(\mathbb{R}^n, \mathcal{R}) = \arccos\left(\frac{\langle e_{n+1}, \mathbf{1} \rangle}{\|e_{n+1}\| \|\mathbf{1}\|}\right) = \arccos\left(\frac{1}{\sqrt{n+1}}\right) \text{ radians} \quad (1566)$$

The vertex-description of the nonnegative orthant in \mathbb{R}^{n+1} is

$$\{[e_1 \ e_2 \ \cdots \ e_{n+1}]a \mid a \geq 0\} = \{a \geq 0\} = \mathbb{R}_+^{n+1} \subset \mathbb{R}^{n+1} \quad (1567)$$

Consider rotation of these vertices via orthogonal matrix

$$Q \triangleq [\mathbf{1}_{\sqrt{n+1}} \ \Xi V_{\mathcal{W}}] \Xi \in \mathbb{R}^{n+1 \times n+1} \quad (1568)$$

where permutation matrix $\Xi \in \mathbb{S}^{n+1}$ is defined in (1619), and $V_{\mathcal{W}} \in \mathbb{R}^{n+1 \times n}$ is the orthonormal auxiliary matrix defined in §B.4.3. This particular orthogonal matrix is selected because it rotates any point in subspace \mathbb{R}^n about one axis of revolution onto \mathcal{R} ; *e.g.*, rotation Qe_{n+1} aligns the last standard basis vector with subspace normal $\mathcal{R}^\perp = \mathbf{1}$. The rotated standard basis vectors remaining are orthonormal spanning \mathcal{R} . \square

Another interpretation of product QX is rotation/reflection of $\mathcal{R}(X)$. Rotation of X as in QXQ^T is the simultaneous rotation/reflection of range and rowspace. **B.8**

Proof. Any matrix can be expressed as a singular value decomposition $X = U\Sigma W^T$ (1462) where $\delta^2(\Sigma) = \Sigma$, $\mathcal{R}(U) \supseteq \mathcal{R}(X)$, and $\mathcal{R}(W) \supseteq \mathcal{R}(X^T)$. \blacklozenge

B.5.4 Matrix rotation

Orthogonal matrices are also employed to rotate/reflect other matrices like vectors: [139, §12.4.1] Given orthogonal matrix Q , the product $Q^T A$ will rotate $A \in \mathbb{R}^{n \times n}$ in the Euclidean sense in \mathbb{R}^{n^2} because the Frobenius norm is orthogonally invariant (§2.2.1);

$$\|Q^T A\|_F = \sqrt{\text{tr}(A^T Q Q^T A)} = \|A\|_F \quad (1569)$$

(likewise for AQ). Were A symmetric, such a rotation would depart from \mathbb{S}^n . One remedy is to instead form the product $Q^T A Q$ because

$$\|Q^T A Q\|_F = \sqrt{\text{tr}(Q^T A^T Q Q^T A Q)} = \|A\|_F \quad (1570)$$

Matrix A is *orthogonally equivalent* to B if $B = S^T A S$ for some orthogonal matrix S . Every square matrix, for example, is orthogonally equivalent to a matrix having equal entries along the main diagonal. [180, §2.2, prob.3]

B.8The product $Q^T A Q$ can be regarded as a coordinate transformation; *e.g.*, given linear map $y = Ax : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and orthogonal Q , the transformation $Qy = A Qx$ is a rotation/reflection of the range and rowspace (132) of matrix A where $Qy \in \mathcal{R}(A)$ and $Qx \in \mathcal{R}(A^T)$ (133).

Appendix C

Some analytical optimal results

C.1 Properties of infima

•

$$\begin{aligned}\inf \emptyset &\triangleq \infty \\ \sup \emptyset &\triangleq -\infty\end{aligned}\tag{1571}$$

- Given $f(x) : \mathcal{X} \rightarrow \mathbb{R}$ defined on arbitrary set \mathcal{X} [177, §0.1.2]

$$\begin{aligned}\inf_{x \in \mathcal{X}} f(x) &= -\sup_{x \in \mathcal{X}} -f(x) \\ \sup_{x \in \mathcal{X}} f(x) &= -\inf_{x \in \mathcal{X}} -f(x)\end{aligned}\tag{1572}$$

$$\begin{aligned}\arg \inf_{x \in \mathcal{X}} f(x) &= \arg \sup_{x \in \mathcal{X}} -f(x) \\ \arg \sup_{x \in \mathcal{X}} f(x) &= \arg \inf_{x \in \mathcal{X}} -f(x)\end{aligned}\tag{1573}$$

- Given $f(x) : \mathcal{X} \rightarrow \mathbb{R}$ and $g(x) : \mathcal{X} \rightarrow \mathbb{R}$ defined on arbitrary set \mathcal{X} [177, §0.1.2]

$$\inf_{x \in \mathcal{X}} (f(x) + g(x)) \geq \inf_{x \in \mathcal{X}} f(x) + \inf_{x \in \mathcal{X}} g(x)\tag{1574}$$

- Given $f(x) : \mathcal{X} \cup \mathcal{Y} \rightarrow \mathbb{R}$ and arbitrary sets \mathcal{X} and \mathcal{Y} [177, §0.1.2]

$$\mathcal{X} \subset \mathcal{Y} \Rightarrow \inf_{x \in \mathcal{X}} f(x) \geq \inf_{x \in \mathcal{Y}} f(x)\tag{1575}$$

$$\inf_{x \in \mathcal{X} \cup \mathcal{Y}} f(x) = \min\{\inf_{x \in \mathcal{X}} f(x), \inf_{x \in \mathcal{Y}} f(x)\}\tag{1576}$$

$$\inf_{x \in \mathcal{X} \cap \mathcal{Y}} f(x) \geq \max\{\inf_{x \in \mathcal{X}} f(x), \inf_{x \in \mathcal{Y}} f(x)\}\tag{1577}$$

C.2 Trace, singular and eigen values

- For $A \in \mathbb{R}^{m \times n}$ and $\sigma(A)$ denoting its singular values, the *nuclear norm* (Ky Fan norm) of matrix A (confer (38), §A.1.1 no.15, [181, p.200]) is

$$\begin{aligned} \sum_i \sigma(A)_i &= \operatorname{tr} \sqrt{A^T A} = \sup_{\|X\|_2 \leq 1} \operatorname{tr}(X^T A) = \underset{X \in \mathbb{R}^{m \times n}}{\text{maximize}} \operatorname{tr}(X^T A) \\ &\text{subject to } \begin{bmatrix} I & X \\ X^T & I \end{bmatrix} \succeq 0 \\ &= \frac{1}{2} \underset{X \in \mathbb{S}^m, Y \in \mathbb{S}^n}{\text{minimize}} \operatorname{tr} X + \operatorname{tr} Y \\ &\text{subject to } \begin{bmatrix} X & A \\ A^T & Y \end{bmatrix} \succeq 0 \end{aligned} \quad (1578)$$

This nuclear norm is dual to the spectral norm. [181, p.214] [53, §A.1.6] Given singular value decomposition $A = S \Sigma Q^T \in \mathbb{R}^{m \times n}$ (A.6), then $X^* = S Q^T \in \mathbb{R}^{m \times n}$ is an optimal solution to the maximization (§2.3.2.0.5) while $X^* = S \Sigma S^T \in \mathbb{S}^m$ and $Y^* = Q \Sigma Q^T \in \mathbb{S}^n$ is an optimal solution to the minimization [118]. Srebro [286] asserts

$$\begin{aligned} \sum_i \sigma(A)_i &= \frac{1}{2} \underset{U, V}{\text{minimize}} \|U\|_F^2 + \|V\|_F^2 \\ &\text{subject to } A = UV^T \\ &= \underset{U, V}{\text{minimize}} \|U\|_F \|V\|_F \\ &\text{subject to } A = UV^T \end{aligned} \quad (1579)$$

C.2.0.0.1 Exercise. *Optimal matrix factorization.*
Prove (1579). **C.1** ▼

C.1 Hint: Write $A = S \Sigma Q^T \in \mathbb{R}^{m \times n}$ and

$$\begin{bmatrix} X & A \\ A^T & Y \end{bmatrix} = \begin{bmatrix} U \\ V \end{bmatrix} \begin{bmatrix} U^T & V^T \end{bmatrix} \succeq 0$$

Show $U^* = S \sqrt{\Sigma} \in \mathbb{R}^{m \times \min\{m, n\}}$ and $V^* = Q \sqrt{\Sigma} \in \mathbb{R}^{n \times \min\{m, n\}}$, hence $\|U^*\|_F^2 = \|V^*\|_F^2$.

- For $X \in \mathbb{S}^m$, $Y \in \mathbb{S}^n$, $A \in \mathcal{C} \subseteq \mathbb{R}^{m \times n}$ for set \mathcal{C} convex, and $\sigma(A)$ denoting the singular values of A [118, §3]

$$\begin{aligned} \underset{A}{\text{minimize}} \quad & \sum_i \sigma(A)_i \\ \text{subject to} \quad & A \in \mathcal{C} \end{aligned} \quad \equiv \quad \begin{aligned} & \frac{1}{2} \underset{A, X, Y}{\text{minimize}} \quad \text{tr } X + \text{tr } Y \\ & \text{subject to} \quad \begin{bmatrix} X & A \\ A^T & Y \end{bmatrix} \succeq 0 \\ & \quad \quad \quad A \in \mathcal{C} \end{aligned} \quad (1580)$$

- For $A \in \mathbb{S}_+^N$ and $\beta \in \mathbb{R}$

$$\begin{aligned} \beta \text{tr } A &= \underset{X \in \mathbb{S}^N}{\text{maximize}} \quad \text{tr}(XA) \\ &\text{subject to} \quad X \preceq \beta I \end{aligned} \quad (1581)$$

But the following statement is numerically stable, preventing an unbounded solution in direction of a 0 eigenvalue:

$$\begin{aligned} & \underset{X \in \mathbb{S}^N}{\text{maximize}} \quad \text{sgn}(\beta) \text{tr}(XA) \\ & \text{subject to} \quad X \preceq |\beta| I \\ & \quad \quad \quad X \succeq -|\beta| I \end{aligned} \quad (1582)$$

where $\beta \text{tr } A = \text{tr}(X^*A)$. If $\beta \geq 0$, then $X \succeq -|\beta|I \leftarrow X \succeq 0$.

- For $A \in \mathbb{S}^N$ having eigenvalues $\lambda(A) \in \mathbb{R}^N$, its smallest and largest eigenvalue is respectively [10, §4.1] [38, §I.6.15] [180, §4.2] [211, §2.1] [212]

$$\begin{aligned} \min_i \{\lambda(A)_i\} &= \inf_{\|x\|=1} x^T A x = \underset{X \in \mathbb{S}_+^N}{\text{minimize}} \quad \text{tr}(XA) = \underset{t \in \mathbb{R}}{\text{maximize}} \quad t \\ & \quad \quad \quad \text{subject to} \quad \text{tr } X = 1 \quad \text{subject to} \quad A \succeq t I \end{aligned} \quad (1583)$$

$$\begin{aligned} \max_i \{\lambda(A)_i\} &= \sup_{\|x\|=1} x^T A x = \underset{X \in \mathbb{S}_+^N}{\text{maximize}} \quad \text{tr}(XA) = \underset{t \in \mathbb{R}}{\text{minimize}} \quad t \\ & \quad \quad \quad \text{subject to} \quad \text{tr } X = 1 \quad \text{subject to} \quad A \preceq t I \end{aligned} \quad (1584)$$

The smallest eigenvalue of any symmetric matrix is always a concave function of its entries, while the largest eigenvalue is always convex. [53, exmp.3.10] For v_1 a normalized eigenvector of A corresponding to

the largest eigenvalue, and v_N a normalized eigenvector corresponding to the smallest eigenvalue,

$$v_N = \arg \inf_{\|x\|=1} x^T A x \quad (1585)$$

$$v_1 = \arg \sup_{\|x\|=1} x^T A x \quad (1586)$$

- For $A \in \mathbb{S}^N$ having eigenvalues $\lambda(A) \in \mathbb{R}^N$, consider the unconstrained nonconvex optimization that is a projection on the rank-1 subset (§2.9.2.1, §3.1.8.0.1) of the boundary of positive semidefinite cone \mathbb{S}_+^N : Defining $\lambda_1 \triangleq \max_i \{\lambda(A)_i\}$ and corresponding eigenvector v_1

$$\begin{aligned} \underset{x}{\text{minimize}} \quad \|xx^T - A\|_F^2 &= \underset{x}{\text{minimize}} \quad \text{tr}(xx^T(x^T x) - 2Axx^T + A^T A) \\ &= \begin{cases} \|\lambda(A)\|^2, & \lambda_1 \leq 0 \\ \|\lambda(A)\|^2 - \lambda_1^2, & \lambda_1 > 0 \end{cases} \end{aligned} \quad (1587)$$

$$\arg \underset{x}{\text{minimize}} \quad \|xx^T - A\|_F^2 = \begin{cases} \mathbf{0}, & \lambda_1 \leq 0 \\ v_1 \sqrt{\lambda_1}, & \lambda_1 > 0 \end{cases} \quad (1588)$$

Proof. This is simply the Eckart & Young solution from §7.1.2:

$$x^* x^{*\top} = \begin{cases} \mathbf{0}, & \lambda_1 \leq 0 \\ \lambda_1 v_1 v_1^T, & \lambda_1 > 0 \end{cases} \quad (1589)$$

Given nonincreasingly ordered diagonalization $A = Q\Lambda Q^T$ where $\Lambda = \delta(\lambda(A))$ (§A.5), then (1587) has minimum value

$$\underset{x}{\text{minimize}} \quad \|xx^T - A\|_F^2 = \begin{cases} \|Q\Lambda Q^T\|_F^2 = \|\delta(\Lambda)\|^2, & \lambda_1 \leq 0 \\ \left\| Q \left(\begin{bmatrix} \lambda_1 & & & \\ & 0 & & \\ & & \ddots & \\ & & & 0 \end{bmatrix} - \Lambda \right) Q^T \right\|_F^2 = \left\| \begin{bmatrix} \lambda_1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} - \delta(\Lambda) \right\|^2, & \lambda_1 > 0 \end{cases} \quad (1590)$$

◆

C.2.0.0.2 Exercise. *Rank-1 approximation.*

Given symmetric matrix $A \in \mathbb{S}^N$, prove:

$$\begin{aligned} v_1 = \arg \underset{x}{\text{minimize}} \quad & \|xx^T - A\|_F^2 \\ \text{subject to} \quad & \|x\| = 1 \end{aligned} \quad (1591)$$

where v_1 is a normalized eigenvector of A corresponding to its largest eigenvalue. What is the objective's optimal value? ▼

- (Ky Fan, 1949) For $B \in \mathbb{S}^N$ whose eigenvalues $\lambda(B) \in \mathbb{R}^N$ are arranged in nonincreasing order, and for $1 \leq k \leq N$ [10, §4.1] [190] [180, §4.3.18] [319, §2] [211, §2.1] [212]

$$\begin{aligned} \sum_{i=N-k+1}^N \lambda(B)_i = \inf_{\substack{U \in \mathbb{R}^{N \times k} \\ U^T U = I}} \text{tr}(UU^T B) = \underset{X \in \mathbb{S}_+^N}{\text{minimize}} \quad & \text{tr}(XB) \\ \text{subject to} \quad & X \preceq I \\ & \text{tr } X = k \end{aligned} \quad (a)$$

$$\begin{aligned} = \underset{\mu \in \mathbb{R}, Z \in \mathbb{S}_+^N}{\text{maximize}} \quad & (k - N)\mu + \text{tr}(B - Z) \\ \text{subject to} \quad & \mu I + Z \succeq B \end{aligned} \quad (b)$$

$$\begin{aligned} \sum_{i=1}^k \lambda(B)_i = \sup_{\substack{U \in \mathbb{R}^{N \times k} \\ U^T U = I}} \text{tr}(UU^T B) = \underset{X \in \mathbb{S}_+^N}{\text{maximize}} \quad & \text{tr}(XB) \\ \text{subject to} \quad & X \preceq I \\ & \text{tr } X = k \end{aligned} \quad (c)$$

$$\begin{aligned} = \underset{\mu \in \mathbb{R}, Z \in \mathbb{S}_+^N}{\text{minimize}} \quad & k\mu + \text{tr } Z \\ \text{subject to} \quad & \mu I + Z \succeq B \end{aligned} \quad (d) \quad (1592)$$

Given ordered diagonalization $B = Q\Lambda Q^T$, (§A.5.2) then an optimal U for the infimum is $U^* = Q(:, N-k+1:N) \in \mathbb{R}^{N \times k}$ whereas $U^* = Q(:, 1:k) \in \mathbb{R}^{N \times k}$ for the supremum. In both cases, $X^* = U^*U^{*\top}$. Optimization (a) searches the convex hull of the outer product UU^T of all $N \times k$ orthonormal matrices. (§2.3.2.0.1)

- For $B \in \mathbb{S}^N$ whose eigenvalues $\lambda(B) \in \mathbb{R}^N$ are arranged in nonincreasing order, and for diagonal matrix $\Upsilon \in \mathbb{S}^k$ whose diagonal entries are arranged in nonincreasing order where $1 \leq k \leq N$, we utilize the main-diagonal δ operator's self-adjointness property (1331): [11, §4.2]

$$\begin{aligned}
\sum_{i=1}^k \Upsilon_{ii} \lambda(B)_{N-i+1} &= \inf_{\substack{U \in \mathbb{R}^{N \times k} \\ U^T U = I}} \text{tr}(\Upsilon U^T B U) = \inf_{\substack{U \in \mathbb{R}^{N \times k} \\ U^T U = I}} \delta(\Upsilon)^T \delta(U^T B U) \\
&= \underset{V_i \in \mathbb{S}^N}{\text{minimize}} \text{tr} \left(B \sum_{i=1}^k (\Upsilon_{ii} - \Upsilon_{i+1, i+1}) V_i \right) \\
&\quad \text{subject to} \quad \text{tr} V_i = i, \quad i = 1 \dots k \\
&\quad \quad \quad I \succeq V_i \succeq 0, \quad i = 1 \dots k
\end{aligned} \tag{1593}$$

where $\Upsilon_{k+1, k+1} \triangleq 0$. We speculate,

$$\sum_{i=1}^k \Upsilon_{ii} \lambda(B)_i = \sup_{\substack{U \in \mathbb{R}^{N \times k} \\ U^T U = I}} \text{tr}(\Upsilon U^T B U) = \sup_{\substack{U \in \mathbb{R}^{N \times k} \\ U^T U = I}} \delta(\Upsilon)^T \delta(U^T B U) \tag{1594}$$

Alizadeh shows: [10, §4.2]

$$\begin{aligned}
\sum_{i=1}^k \Upsilon_{ii} \lambda(B)_i &= \underset{\substack{\mu \in \mathbb{R}^k, Z_i \in \mathbb{S}^N}}{\text{minimize}} \sum_{i=1}^k i \mu_i + \text{tr} Z_i \\
&\quad \text{subject to} \quad \mu_i I + Z_i - (\Upsilon_{ii} - \Upsilon_{i+1, i+1}) B \succeq 0, \quad i = 1 \dots k \\
&\quad \quad \quad Z_i \succeq 0, \quad i = 1 \dots k \\
&= \underset{V_i \in \mathbb{S}^N}{\text{maximize}} \text{tr} \left(B \sum_{i=1}^k (\Upsilon_{ii} - \Upsilon_{i+1, i+1}) V_i \right) \\
&\quad \text{subject to} \quad \text{tr} V_i = i, \quad i = 1 \dots k \\
&\quad \quad \quad I \succeq V_i \succeq 0, \quad i = 1 \dots k
\end{aligned} \tag{1595}$$

where $\Upsilon_{k+1, k+1} \triangleq 0$.

- The largest eigenvalue magnitude μ of $A \in \mathbb{S}^N$

$$\begin{aligned}
\max_i \{ |\lambda(A)_i| \} &= \underset{\mu \in \mathbb{R}}{\text{minimize}} \quad \mu \\
&\quad \text{subject to} \quad -\mu I \preceq A \preceq \mu I
\end{aligned} \tag{1596}$$

is minimized over convex set \mathcal{C} by semidefinite program: (confer §7.1.5)

$$\begin{aligned} \underset{A}{\text{minimize}} \quad \|A\|_2 &\equiv \underset{\mu, A}{\text{minimize}} \quad \mu \\ \text{subject to} \quad A \in \mathcal{C} &\equiv \text{subject to} \quad -\mu I \preceq A \preceq \mu I \\ &\quad A \in \mathcal{C} \end{aligned} \quad (1597)$$

id est,

$$\mu^* \triangleq \max_i \{ |\lambda(A^*)_i|, i = 1 \dots N \} \in \mathbb{R}_+ \quad (1598)$$

- For $B \in \mathbb{S}^N$ whose eigenvalues $\lambda(B) \in \mathbb{R}^N$ are arranged in nonincreasing order, let $\Pi \lambda(B)$ be a permutation of eigenvalues $\lambda(B)$ such that their absolute value becomes arranged in nonincreasing order: $|\Pi \lambda(B)|_1 \geq |\Pi \lambda(B)|_2 \geq \dots \geq |\Pi \lambda(B)|_N$. Then, for $1 \leq k \leq N$ [10, §4.3] **C.2**

$$\begin{aligned} \sum_{i=1}^k |\Pi \lambda(B)|_i &= \underset{\mu \in \mathbb{R}, Z \in \mathbb{S}_+^N}{\text{minimize}} \quad k\mu + \text{tr} Z &= \underset{V, W \in \mathbb{S}_+^N}{\text{maximize}} \quad \langle B, V - W \rangle \\ \text{subject to} \quad &\mu I + Z + B \succeq 0 &\text{subject to} \quad I \succeq V, W \\ &\mu I + Z - B \succeq 0 &\text{tr}(V + W) = k \end{aligned} \quad (1599)$$

For diagonal matrix $\Upsilon \in \mathbb{S}^k$ whose diagonal entries are arranged in nonincreasing order where $1 \leq k \leq N$

$$\begin{aligned} \sum_{i=1}^k \Upsilon_{ii} |\Pi \lambda(B)|_i &= \underset{\mu \in \mathbb{R}^k, Z_i \in \mathbb{S}^N}{\text{minimize}} \quad \sum_{i=1}^k i\mu_i + \text{tr} Z_i \\ \text{subject to} \quad &\mu_i I + Z_i + (\Upsilon_{ii} - \Upsilon_{i+1, i+1})B \succeq 0, \quad i = 1 \dots k \\ &\mu_i I + Z_i - (\Upsilon_{ii} - \Upsilon_{i+1, i+1})B \succeq 0, \quad i = 1 \dots k \\ &Z_i \succeq 0, \quad i = 1 \dots k \\ &= \underset{V_i, W_i \in \mathbb{S}^N}{\text{maximize}} \quad \text{tr} \left(B \sum_{i=1}^k (\Upsilon_{ii} - \Upsilon_{i+1, i+1})(V_i - W_i) \right) \\ \text{subject to} \quad &\text{tr}(V_i + W_i) = i, \quad i = 1 \dots k \\ &I \succeq V_i \succeq 0, \quad i = 1 \dots k \\ &I \succeq W_i \succeq 0, \quad i = 1 \dots k \end{aligned} \quad (1600)$$

where $\Upsilon_{k+1, k+1} \triangleq 0$.

C.2We eliminate a redundant positive semidefinite variable from Alizadeh's minimization. There exist typographical errors in [258, (6.49) (6.55)] for this minimization.

- For $A, B \in \mathbb{S}^N$ whose eigenvalues $\lambda(A), \lambda(B) \in \mathbb{R}^N$ are respectively arranged in nonincreasing order, and for nonincreasingly ordered diagonalizations $A = W_A \Upsilon W_A^T$ and $B = W_B \Lambda W_B^T$ [178] [211, §2.1] [212]

$$\lambda(A)^T \lambda(B) = \sup_{\substack{U \in \mathbb{R}^{N \times N} \\ U^T U = I}} \text{tr}(A^T U^T B U) \geq \text{tr}(A^T B) \quad (1618)$$

(confer (1623)) where optimal U is

$$U^* = W_B W_A^T \in \mathbb{R}^{N \times N} \quad (1615)$$

We can push that upper bound higher using a result in §C.4.2.1:

$$|\lambda(A)|^T |\lambda(B)| = \sup_{\substack{U \in \mathbb{C}^{N \times N} \\ U^H U = I}} \text{Re tr}(A^T U^H B U) \quad (1601)$$

For step function ψ as defined in (1477), optimal U becomes

$$U^* = W_B \sqrt{\delta(\psi(\delta(\Lambda)))}^H \sqrt{\delta(\psi(\delta(\Upsilon)))} W_A^T \in \mathbb{C}^{N \times N} \quad (1602)$$

C.3 Orthogonal Procrustes problem

Given matrices $A, B \in \mathbb{R}^{n \times N}$, their product having full singular value decomposition (§A.6.3)

$$AB^T \triangleq U \Sigma Q^T \in \mathbb{R}^{n \times n} \quad (1603)$$

then an optimal solution R^* to the orthogonal Procrustes problem

$$\begin{aligned} & \underset{R}{\text{minimize}} \quad \|A - R^T B\|_F \\ & \text{subject to} \quad R^T = R^{-1} \end{aligned} \quad (1604)$$

maximizes $\text{tr}(A^T R^T B)$ over the nonconvex manifold of orthogonal matrices: [180, §7.4.8]

$$R^* = Q U^T \in \mathbb{R}^{n \times n} \quad (1605)$$

A necessary and sufficient condition for optimality

$$AB^T R^* \succeq 0 \quad (1606)$$

holds whenever R^* is an orthogonal matrix. [143, §4]

Optimal solution R^* can reveal rotation/reflection (§5.5.2, §B.5) of one list in the columns of matrix A with respect to another list in B . Solution is unique if $\text{rank } BV_{\mathcal{N}} = n$. [96, §2.4.1] In the case that A is a vector and permutation of B , solution R^* is not necessarily a permutation matrix (§4.6.0.0.3) although the optimal objective will be zero. More generally, the optimal value for objective of minimization is

$$\begin{aligned} \text{tr}(A^T A + B^T B - 2AB^T R^*) &= \text{tr}(A^T A) + \text{tr}(B^T B) - 2\text{tr}(U\Sigma U^T) \\ &= \|A\|_F^2 + \|B\|_F^2 - 2\delta(\Sigma)^T \mathbf{1} \end{aligned} \quad (1607)$$

while the optimal value for corresponding trace maximization is

$$\sup_{R^T=R^{-1}} \text{tr}(A^T R^T B) = \text{tr}(A^T R^{*T} B) = \delta(\Sigma)^T \mathbf{1} \geq \text{tr}(A^T B) \quad (1608)$$

The same optimal solution R^* solves

$$\begin{aligned} &\underset{R}{\text{maximize}} \quad \|A + R^T B\|_F \\ &\text{subject to} \quad R^T = R^{-1} \end{aligned} \quad (1609)$$

C.3.1 Effect of translation

Consider the impact on problem (1604) of dc offset in known lists $A, B \in \mathbb{R}^{n \times N}$. Rotation of B there is with respect to the origin, so better results may be obtained if offset is first accounted. Because the geometric centers of the lists AV and BV are the origin, instead we solve

$$\begin{aligned} &\underset{R}{\text{minimize}} \quad \|AV - R^T B V\|_F \\ &\text{subject to} \quad R^T = R^{-1} \end{aligned} \quad (1610)$$

where $V \in \mathbb{S}^N$ is the geometric centering matrix (§B.4.1). Now we define the full singular value decomposition

$$AVB^T \triangleq U\Sigma Q^T \in \mathbb{R}^{n \times n} \quad (1611)$$

and an optimal rotation matrix

$$R^* = QU^T \in \mathbb{R}^{n \times n} \quad (1605)$$

The desired result is an optimally rotated offset list

$$R^{*T} B V + A(I - V) \approx A \quad (1612)$$

which most closely matches the list in A . Equality is attained when the lists are precisely related by a rotation/reflection and an offset. When $R^{*T} B = A$ or $B\mathbf{1} = A\mathbf{1} = \mathbf{0}$, this result (1612) reduces to $R^{*T} B \approx A$.

C.3.1.1 Translation of extended list

Suppose an optimal rotation matrix $R^* \in \mathbb{R}^{n \times n}$ were derived as before from matrix $B \in \mathbb{R}^{n \times N}$, but B is part of a larger list in the columns of $[C \ B] \in \mathbb{R}^{n \times M+N}$ where $C \in \mathbb{R}^{n \times M}$. In that event, we wish to apply the rotation/reflection and translation to the larger list. The expression supplanting the approximation in (1612) makes $\mathbf{1}^T$ of compatible dimension;

$$R^{*\text{T}}[C - B\mathbf{1}\mathbf{1}^{\text{T}}\frac{1}{N} \quad BV] + A\mathbf{1}\mathbf{1}^{\text{T}}\frac{1}{N} \quad (1613)$$

id est, $C - B\mathbf{1}\mathbf{1}^{\text{T}}\frac{1}{N} \in \mathbb{R}^{n \times M}$ and $A\mathbf{1}\mathbf{1}^{\text{T}}\frac{1}{N} \in \mathbb{R}^{n \times M+N}$.

C.4 Two-sided orthogonal Procrustes

C.4.0.1 Minimization

Given symmetric $A, B \in \mathbb{S}^N$, each having diagonalization (§A.5.2)

$$A \triangleq Q_A \Lambda_A Q_A^{\text{T}}, \quad B \triangleq Q_B \Lambda_B Q_B^{\text{T}} \quad (1614)$$

where eigenvalues are arranged in their respective diagonal matrix Λ in nonincreasing order, then an optimal solution [113]

$$R^* = Q_B Q_A^{\text{T}} \in \mathbb{R}^{N \times N} \quad (1615)$$

to the two-sided orthogonal Procrustes problem

$$\begin{aligned} \underset{R}{\text{minimize}} \quad & \|A - R^{\text{T}}BR\|_{\text{F}} &= \underset{R}{\text{minimize}} \quad & \text{tr}(A^{\text{T}}A - 2A^{\text{T}}R^{\text{T}}BR + B^{\text{T}}B) \\ \text{subject to} \quad & R^{\text{T}} = R^{-1} & \text{subject to} \quad & R^{\text{T}} = R^{-1} \end{aligned} \quad (1616)$$

maximizes $\text{tr}(A^{\text{T}}R^{\text{T}}BR)$ over the nonconvex manifold of orthogonal matrices. Optimal product $R^{*\text{T}}BR^*$ has the eigenvectors of A but the eigenvalues of B . [143, §7.5.1] The optimal value for the objective of minimization is, by (42)

$$\|Q_A \Lambda_A Q_A^{\text{T}} - R^{*\text{T}}Q_B \Lambda_B Q_B^{\text{T}}R^*\|_{\text{F}} = \|Q_A(\Lambda_A - \Lambda_B)Q_A^{\text{T}}\|_{\text{F}} = \|\Lambda_A - \Lambda_B\|_{\text{F}} \quad (1617)$$

while the corresponding trace maximization has optimal value

$$\sup_{R^{\text{T}}=R^{-1}} \text{tr}(A^{\text{T}}R^{\text{T}}BR) = \text{tr}(A^{\text{T}}R^{*\text{T}}BR^*) = \text{tr}(\Lambda_A \Lambda_B) \geq \text{tr}(A^{\text{T}}B) \quad (1618)$$

C.4.0.2 Maximization

Any permutation matrix is an orthogonal matrix. Defining a row- and column-swapping permutation matrix (a reflection matrix, §B.5.2)

$$\Xi = \Xi^T = \begin{bmatrix} \mathbf{0} & & & 1 \\ & & \cdot & \\ & & \cdot & \\ & 1 & & \\ 1 & & & \mathbf{0} \end{bmatrix} \quad (1619)$$

then an optimal solution R^* to the maximization problem [sic]

$$\begin{aligned} & \underset{R}{\text{maximize}} && \|A - R^T B R\|_F \\ & \text{subject to} && R^T = R^{-1} \end{aligned} \quad (1620)$$

minimizes $\text{tr}(A^T R^T B R)$: [178] [211, §2.1] [212]

$$R^* = Q_B \Xi Q_A^T \in \mathbb{R}^{N \times N} \quad (1621)$$

The optimal value for the objective of maximization is

$$\begin{aligned} \|Q_A \Lambda_A Q_A^T - R^{*T} Q_B \Lambda_B Q_B^T R^*\|_F &= \|Q_A \Lambda_A Q_A^T - Q_A \Xi^T \Lambda_B \Xi Q_A^T\|_F \\ &= \|\Lambda_A - \Xi \Lambda_B \Xi\|_F \end{aligned} \quad (1622)$$

while the corresponding trace minimization has optimal value

$$\inf_{R^T=R^{-1}} \text{tr}(A^T R^T B R) = \text{tr}(A^T R^{*T} B R^*) = \text{tr}(\Lambda_A \Xi \Lambda_B \Xi) \quad (1623)$$

C.4.1 Procrustes' relation to linear programming

Although these two-sided Procrustes problems are nonconvex, a connection with linear programming [82] was discovered by Anstreicher & Wolkowicz [11, §3] [211, §2.1] [212]: Given $A, B \in \mathbb{S}^N$, this semidefinite program in S and T

$$\begin{aligned} & \underset{R}{\text{minimize}} && \text{tr}(A^T R^T B R) = \underset{S, T \in \mathbb{S}^N}{\text{maximize}} && \text{tr}(S + T) \\ & \text{subject to} && R^T = R^{-1} && \text{subject to} && A^T \otimes B - I \otimes S - T \otimes I \succeq 0 \end{aligned} \quad (1624)$$

(where \otimes signifies Kronecker product (§D.1.2.1)) has optimal objective value (1623). These two problems are strong duals (§2.13.1.0.3). Given ordered diagonalizations (1614), make the observation:

$$\inf_R \operatorname{tr}(A^T R^T B R) = \inf_{\hat{R}} \operatorname{tr}(\Lambda_A \hat{R}^T \Lambda_B \hat{R}) \quad (1625)$$

because $\hat{R} \triangleq Q_B^T R Q_A$ on the set of orthogonal matrices (which includes the permutation matrices) is a bijection. This means, basically, diagonal matrices of eigenvalues Λ_A and Λ_B may be substituted for A and B , so only the main diagonals of S and T come into play;

$$\begin{aligned} & \underset{S, T \in \mathbb{S}^N}{\text{maximize}} && \mathbf{1}^T \delta(S + T) \\ & \text{subject to} && \delta(\Lambda_A \otimes (\Xi \Lambda_B \Xi) - I \otimes S - T \otimes I) \succeq 0 \end{aligned} \quad (1626)$$

a linear program in $\delta(S)$ and $\delta(T)$ having the same optimal objective value as the semidefinite program (1624).

We relate their results to Procrustes problem (1616) by manipulating signs (1572) and permuting eigenvalues:

$$\begin{aligned} \underset{R}{\text{maximize}} \quad \operatorname{tr}(A^T R^T B R) &= \underset{S, T \in \mathbb{S}^N}{\text{minimize}} \quad \mathbf{1}^T \delta(S + T) \\ \text{subject to} \quad R^T &= R^{-1} && \text{subject to} \quad \delta(I \otimes S + T \otimes I - \Lambda_A \otimes \Lambda_B) \succeq 0 \\ &&& = \underset{S, T \in \mathbb{S}^N}{\text{minimize}} \quad \operatorname{tr}(S + T) \\ &&& \text{subject to} \quad I \otimes S + T \otimes I - A^T \otimes B \succeq 0 \end{aligned} \quad (1627)$$

This formulation has optimal objective value identical to that in (1618).

C.4.2 Two-sided orthogonal Procrustes via SVD

By making left- and right-side orthogonal matrices independent, we can push the upper bound on trace (1618) a little further: Given real matrices A, B each having full singular value decomposition (§A.6.3)

$$A \triangleq U_A \Sigma_A Q_A^T \in \mathbb{R}^{m \times n}, \quad B \triangleq U_B \Sigma_B Q_B^T \in \mathbb{R}^{m \times n} \quad (1628)$$

then a well-known optimal solution R^*, S^* to the problem

$$\begin{aligned} & \underset{R, S}{\text{minimize}} && \|A - SBR\|_F \\ & \text{subject to} && R^H = R^{-1} \\ & && S^H = S^{-1} \end{aligned} \quad (1629)$$

maximizes $\operatorname{Re} \operatorname{tr}(A^T S B R)$: [278] [255] [45] [173] optimal orthogonal matrices

$$S^* = U_A U_B^H \in \mathbb{R}^{m \times m}, \quad R^* = Q_B Q_A^H \in \mathbb{R}^{n \times n} \quad (1630)$$

[*sic*] are not necessarily unique [180, §7.4.13] because the feasible set is not convex. The optimal value for the objective of minimization is, by (42)

$$\|U_A \Sigma_A Q_A^H - S^* U_B \Sigma_B Q_B^H R^*\|_F = \|U_A (\Sigma_A - \Sigma_B) Q_A^H\|_F = \|\Sigma_A - \Sigma_B\|_F \quad (1631)$$

while the corresponding trace maximization has optimal value [38, §III.6.12]

$$\sup_{\substack{R^H=R^{-1} \\ S^H=S^{-1}}} |\operatorname{tr}(A^T S B R)| = \sup_{\substack{R^H=R^{-1} \\ S^H=S^{-1}}} \operatorname{Re} \operatorname{tr}(A^T S B R) = \operatorname{Re} \operatorname{tr}(A^T S^* B R^*) = \operatorname{tr}(\Sigma_A^T \Sigma_B) \geq \operatorname{tr}(A^T B) \quad (1632)$$

for which it is necessary

$$A^T S^* B R^* \succeq 0, \quad B R^* A^T S^* \succeq 0 \quad (1633)$$

The lower bound on inner product of singular values in (1632) is due to von Neumann. Equality is attained if $U_A^H U_B = I$ and $Q_B^H Q_A = I$.

C.4.2.1 Symmetric matrices

Now optimizing over the complex manifold of unitary matrices (§B.5.1), the upper bound on trace (1618) is thereby raised: Suppose we are given diagonalizations for (real) symmetric A, B (§A.5)

$$A = W_A \Upsilon W_A^T \in \mathbb{S}^n, \quad \delta(\Upsilon) \in \mathcal{K}_{\mathcal{M}} \quad (1634)$$

$$B = W_B \Lambda W_B^T \in \mathbb{S}^n, \quad \delta(\Lambda) \in \mathcal{K}_{\mathcal{M}} \quad (1635)$$

having their respective eigenvalues in diagonal matrices $\Upsilon, \Lambda \in \mathbb{S}^n$ arranged in nonincreasing order (membership to the monotone cone $\mathcal{K}_{\mathcal{M}}$ (396)). Then by splitting eigenvalue signs, we invent a symmetric SVD-like decomposition

$$A \triangleq U_A \Sigma_A Q_A^H \in \mathbb{S}^n, \quad B \triangleq U_B \Sigma_B Q_B^H \in \mathbb{S}^n \quad (1636)$$

where $U_A, U_B, Q_A, Q_B \in \mathbb{C}^{n \times n}$ are unitary matrices defined by (*confer* §A.6.5)

$$U_A \triangleq W_A \sqrt{\delta(\psi(\delta(\Upsilon)))}, \quad Q_A \triangleq W_A \sqrt{\delta(\psi(\delta(\Upsilon)))}^H, \quad \Sigma_A = |\Upsilon| \quad (1637)$$

$$U_B \triangleq W_B \sqrt{\delta(\psi(\delta(\Lambda)))}, \quad Q_B \triangleq W_B \sqrt{\delta(\psi(\delta(\Lambda)))}^H, \quad \Sigma_B = |\Lambda| \quad (1638)$$

where step function ψ is defined in (1477). In this circumstance,

$$S^* = U_A U_B^H = R^{*\text{T}} \in \mathbb{C}^{n \times n} \quad (1639)$$

optimal matrices (1630) now unitary are related by transposition. The optimal value of objective (1631) is

$$\|U_A \Sigma_A Q_A^H - S^* U_B \Sigma_B Q_B^H R^*\|_F = \||\Upsilon| - |\Lambda|\|_F \quad (1640)$$

while the corresponding optimal value of trace maximization (1632) is

$$\sup_{\substack{R^H = R^{-1} \\ S^H = S^{-1}}} \text{Re tr}(A^T S B R) = \text{tr}(|\Upsilon| |\Lambda|) \quad (1641)$$

C.4.2.2 Diagonal matrices

Now suppose A and B are diagonal matrices

$$A = \Upsilon = \delta^2(\Upsilon) \in \mathbb{S}^n, \quad \delta(\Upsilon) \in \mathcal{K}_{\mathcal{M}} \quad (1642)$$

$$B = \Lambda = \delta^2(\Lambda) \in \mathbb{S}^n, \quad \delta(\Lambda) \in \mathcal{K}_{\mathcal{M}} \quad (1643)$$

both having their respective main diagonal entries arranged in nonincreasing order:

$$\begin{aligned} & \underset{R, S}{\text{minimize}} && \|\Upsilon - S \Lambda R\|_F \\ & \text{subject to} && R^H = R^{-1} \\ & && S^H = S^{-1} \end{aligned} \quad (1644)$$

Then we have a symmetric decomposition from unitary matrices as in (1636) where

$$U_A \triangleq \sqrt{\delta(\psi(\delta(\Upsilon)))}, \quad Q_A \triangleq \sqrt{\delta(\psi(\delta(\Upsilon)))}^H, \quad \Sigma_A = |\Upsilon| \quad (1645)$$

$$U_B \triangleq \sqrt{\delta(\psi(\delta(\Lambda)))}, \quad Q_B \triangleq \sqrt{\delta(\psi(\delta(\Lambda)))}^H, \quad \Sigma_B = |\Lambda| \quad (1646)$$

Procrustes solution (1630) again sees the transposition relationship

$$S^* = U_A U_B^H = R^{*\text{T}} \in \mathbb{C}^{n \times n} \quad (1639)$$

but both optimal unitary matrices are now themselves diagonal. So,

$$S^* \Lambda R^* = \delta(\psi(\delta(\Upsilon))) \Lambda \delta(\psi(\delta(\Lambda))) = \delta(\psi(\delta(\Upsilon))) |\Lambda| \quad (1647)$$

Appendix D

Matrix calculus

From too much study, and from extreme passion, cometh madness.

– Isaac Newton [134, §5]

D.1 Directional derivative, Taylor series

D.1.1 Gradients

Gradient of a differentiable real function $f(x) : \mathbb{R}^K \rightarrow \mathbb{R}$ with respect to its vector argument is defined

$$\nabla f(x) = \begin{bmatrix} \frac{\partial f(x)}{\partial x_1} \\ \frac{\partial f(x)}{\partial x_2} \\ \vdots \\ \frac{\partial f(x)}{\partial x_K} \end{bmatrix} \in \mathbb{R}^K \quad (1648)$$

while the second-order gradient of the twice differentiable real function with respect to its vector argument is traditionally called the *Hessian*;

$$\nabla^2 f(x) = \begin{bmatrix} \frac{\partial^2 f(x)}{\partial^2 x_1} & \frac{\partial^2 f(x)}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f(x)}{\partial x_1 \partial x_K} \\ \frac{\partial^2 f(x)}{\partial x_2 \partial x_1} & \frac{\partial^2 f(x)}{\partial^2 x_2} & \cdots & \frac{\partial^2 f(x)}{\partial x_2 \partial x_K} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f(x)}{\partial x_K \partial x_1} & \frac{\partial^2 f(x)}{\partial x_K \partial x_2} & \cdots & \frac{\partial^2 f(x)}{\partial^2 x_K} \end{bmatrix} \in \mathbb{S}^K \quad (1649)$$

The gradient of vector-valued function $v(x) : \mathbb{R} \rightarrow \mathbb{R}^N$ on real domain is a row-vector

$$\nabla v(x) \triangleq \left[\frac{\partial v_1(x)}{\partial x} \quad \frac{\partial v_2(x)}{\partial x} \quad \dots \quad \frac{\partial v_N(x)}{\partial x} \right] \in \mathbb{R}^N \quad (1650)$$

while the second-order gradient is

$$\nabla^2 v(x) \triangleq \left[\frac{\partial^2 v_1(x)}{\partial x^2} \quad \frac{\partial^2 v_2(x)}{\partial x^2} \quad \dots \quad \frac{\partial^2 v_N(x)}{\partial x^2} \right] \in \mathbb{R}^N \quad (1651)$$

Gradient of vector-valued function $h(x) : \mathbb{R}^K \rightarrow \mathbb{R}^N$ on vector domain is

$$\begin{aligned} \nabla h(x) &\triangleq \begin{bmatrix} \frac{\partial h_1(x)}{\partial x_1} & \frac{\partial h_2(x)}{\partial x_1} & \dots & \frac{\partial h_N(x)}{\partial x_1} \\ \frac{\partial h_1(x)}{\partial x_2} & \frac{\partial h_2(x)}{\partial x_2} & \dots & \frac{\partial h_N(x)}{\partial x_2} \\ \vdots & \vdots & & \vdots \\ \frac{\partial h_1(x)}{\partial x_K} & \frac{\partial h_2(x)}{\partial x_K} & \dots & \frac{\partial h_N(x)}{\partial x_K} \end{bmatrix} \\ &= [\nabla h_1(x) \quad \nabla h_2(x) \quad \dots \quad \nabla h_N(x)] \in \mathbb{R}^{K \times N} \end{aligned} \quad (1652)$$

while the second-order gradient has a three-dimensional representation dubbed *cubix*; ^{D.1}

$$\begin{aligned} \nabla^2 h(x) &\triangleq \begin{bmatrix} \nabla \frac{\partial h_1(x)}{\partial x_1} & \nabla \frac{\partial h_2(x)}{\partial x_1} & \dots & \nabla \frac{\partial h_N(x)}{\partial x_1} \\ \nabla \frac{\partial h_1(x)}{\partial x_2} & \nabla \frac{\partial h_2(x)}{\partial x_2} & \dots & \nabla \frac{\partial h_N(x)}{\partial x_2} \\ \vdots & \vdots & & \vdots \\ \nabla \frac{\partial h_1(x)}{\partial x_K} & \nabla \frac{\partial h_2(x)}{\partial x_K} & \dots & \nabla \frac{\partial h_N(x)}{\partial x_K} \end{bmatrix} \\ &= [\nabla^2 h_1(x) \quad \nabla^2 h_2(x) \quad \dots \quad \nabla^2 h_N(x)] \in \mathbb{R}^{K \times N \times K} \end{aligned} \quad (1653)$$

where the gradient of each real entry is with respect to vector x as in (1648).

^{D.1}The word *matrix* comes from the Latin for *womb*; related to the prefix *matri-* derived from *mater* meaning *mother*.

The gradient of real function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}$ on matrix domain is

$$\begin{aligned} \nabla g(X) &\triangleq \begin{bmatrix} \frac{\partial g(X)}{\partial X_{11}} & \frac{\partial g(X)}{\partial X_{12}} & \dots & \frac{\partial g(X)}{\partial X_{1L}} \\ \frac{\partial g(X)}{\partial X_{21}} & \frac{\partial g(X)}{\partial X_{22}} & \dots & \frac{\partial g(X)}{\partial X_{2L}} \\ \vdots & \vdots & & \vdots \\ \frac{\partial g(X)}{\partial X_{K1}} & \frac{\partial g(X)}{\partial X_{K2}} & \dots & \frac{\partial g(X)}{\partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L} \\ &= \begin{bmatrix} \nabla_{X(:,1)} g(X) \\ \nabla_{X(:,2)} g(X) \\ \dots \\ \nabla_{X(:,L)} g(X) \end{bmatrix} \in \mathbb{R}^{K \times 1 \times L} \end{aligned} \tag{1654}$$

where the gradient $\nabla_{X(:,i)}$ is with respect to the i^{th} column of X . The strange appearance of (1654) in $\mathbb{R}^{K \times 1 \times L}$ is meant to suggest a third dimension perpendicular to the page (not a diagonal matrix). The second-order gradient has representation

$$\begin{aligned} \nabla^2 g(X) &\triangleq \begin{bmatrix} \nabla \frac{\partial g(X)}{\partial X_{11}} & \nabla \frac{\partial g(X)}{\partial X_{12}} & \dots & \nabla \frac{\partial g(X)}{\partial X_{1L}} \\ \nabla \frac{\partial g(X)}{\partial X_{21}} & \nabla \frac{\partial g(X)}{\partial X_{22}} & \dots & \nabla \frac{\partial g(X)}{\partial X_{2L}} \\ \vdots & \vdots & & \vdots \\ \nabla \frac{\partial g(X)}{\partial X_{K1}} & \nabla \frac{\partial g(X)}{\partial X_{K2}} & \dots & \nabla \frac{\partial g(X)}{\partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L \times K \times L} \\ &= \begin{bmatrix} \nabla \nabla_{X(:,1)} g(X) \\ \nabla \nabla_{X(:,2)} g(X) \\ \dots \\ \nabla \nabla_{X(:,L)} g(X) \end{bmatrix} \in \mathbb{R}^{K \times 1 \times L \times K \times L} \end{aligned} \tag{1655}$$

where the gradient ∇ is with respect to matrix X .

Gradient of vector-valued function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^N$ on matrix domain is a cubix

$$\begin{aligned} \nabla g(X) &\triangleq \begin{bmatrix} \nabla_{X(:,1)} g_1(X) & \nabla_{X(:,1)} g_2(X) & \cdots & \nabla_{X(:,1)} g_N(X) \\ \nabla_{X(:,2)} g_1(X) & \nabla_{X(:,2)} g_2(X) & \cdots & \nabla_{X(:,2)} g_N(X) \\ \vdots & \vdots & \ddots & \vdots \\ \nabla_{X(:,L)} g_1(X) & \nabla_{X(:,L)} g_2(X) & \cdots & \nabla_{X(:,L)} g_N(X) \end{bmatrix} \\ &= [\nabla g_1(X) \quad \nabla g_2(X) \quad \cdots \quad \nabla g_N(X)] \in \mathbb{R}^{K \times N \times L} \end{aligned} \quad (1656)$$

while the second-order gradient has a five-dimensional representation;

$$\begin{aligned} \nabla^2 g(X) &\triangleq \begin{bmatrix} \nabla \nabla_{X(:,1)} g_1(X) & \nabla \nabla_{X(:,1)} g_2(X) & \cdots & \nabla \nabla_{X(:,1)} g_N(X) \\ \nabla \nabla_{X(:,2)} g_1(X) & \nabla \nabla_{X(:,2)} g_2(X) & \cdots & \nabla \nabla_{X(:,2)} g_N(X) \\ \vdots & \vdots & \ddots & \vdots \\ \nabla \nabla_{X(:,L)} g_1(X) & \nabla \nabla_{X(:,L)} g_2(X) & \cdots & \nabla \nabla_{X(:,L)} g_N(X) \end{bmatrix} \\ &= [\nabla^2 g_1(X) \quad \nabla^2 g_2(X) \quad \cdots \quad \nabla^2 g_N(X)] \in \mathbb{R}^{K \times N \times L \times K \times L} \end{aligned} \quad (1657)$$

The gradient of matrix-valued function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^{M \times N}$ on matrix domain has a four-dimensional representation called *quartix*

$$\nabla g(X) \triangleq \begin{bmatrix} \nabla g_{11}(X) & \nabla g_{12}(X) & \cdots & \nabla g_{1N}(X) \\ \nabla g_{21}(X) & \nabla g_{22}(X) & \cdots & \nabla g_{2N}(X) \\ \vdots & \vdots & \ddots & \vdots \\ \nabla g_{M1}(X) & \nabla g_{M2}(X) & \cdots & \nabla g_{MN}(X) \end{bmatrix} \in \mathbb{R}^{M \times N \times K \times L} \quad (1658)$$

while the second-order gradient has six-dimensional representation

$$\nabla^2 g(X) \triangleq \begin{bmatrix} \nabla^2 g_{11}(X) & \nabla^2 g_{12}(X) & \cdots & \nabla^2 g_{1N}(X) \\ \nabla^2 g_{21}(X) & \nabla^2 g_{22}(X) & \cdots & \nabla^2 g_{2N}(X) \\ \vdots & \vdots & \ddots & \vdots \\ \nabla^2 g_{M1}(X) & \nabla^2 g_{M2}(X) & \cdots & \nabla^2 g_{MN}(X) \end{bmatrix} \in \mathbb{R}^{M \times N \times K \times L \times K \times L} \quad (1659)$$

and so on.

D.1.2 Product rules for matrix-functions

Given dimensionally compatible matrix-valued functions of matrix variable $f(X)$ and $g(X)$

$$\nabla_X (f(X)^T g(X)) = \nabla_X (f) g + \nabla_X (g) f \tag{1660}$$

while [46, §8.3] [279]

$$\nabla_X \operatorname{tr}(f(X)^T g(X)) = \nabla_X \left(\operatorname{tr}(f(X)^T g(Z)) + \operatorname{tr}(g(X) f(Z)^T) \right) \Big|_{Z=X} \tag{1661}$$

These expressions implicitly apply as well to scalar-, vector-, or matrix-valued functions of scalar, vector, or matrix arguments.

D.1.2.0.1 Example. Cubix.

Suppose $f(X) : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}^2 = X^T a$ and $g(X) : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}^2 = X b$. We wish to find

$$\nabla_X (f(X)^T g(X)) = \nabla_X a^T X^2 b \tag{1662}$$

using the product rule. Formula (1660) calls for

$$\nabla_X a^T X^2 b = \nabla_X (X^T a) X b + \nabla_X (X b) X^T a \tag{1663}$$

Consider the first of the two terms:

$$\begin{aligned} \nabla_X (f) g &= \nabla_X (X^T a) X b \\ &= [\nabla (X^T a)_1 \quad \nabla (X^T a)_2] X b \end{aligned} \tag{1664}$$

The gradient of $X^T a$ forms a cubix in $\mathbb{R}^{2 \times 2 \times 2}$.

$$\nabla_X (X^T a) X b = \left[\begin{array}{cc} \frac{\partial (X^T a)_1}{\partial X_{11}} & \dots & \frac{\partial (X^T a)_2}{\partial X_{11}} \\ \vdots & \frac{\partial (X^T a)_1}{\partial X_{12}} & \dots & \frac{\partial (X^T a)_2}{\partial X_{12}} \\ \frac{\partial (X^T a)_1}{\partial X_{21}} & \dots & \frac{\partial (X^T a)_2}{\partial X_{21}} & \vdots \\ \vdots & \frac{\partial (X^T a)_1}{\partial X_{22}} & \dots & \frac{\partial (X^T a)_2}{\partial X_{22}} \end{array} \right] \left[\begin{array}{c} (X b)_1 \\ (X b)_2 \end{array} \right] \in \mathbb{R}^{2 \times 1 \times 2} \tag{1665}$$

Because gradient of the product (1662) requires total change with respect to change in each entry of matrix X , the Xb vector must make an inner product with each vector in the second dimension of the cubix (indicated by dotted line segments);

$$\begin{aligned} \nabla_X(X^T a) X b &= \begin{bmatrix} a_1 & 0 & & \\ & 0 & a_1 & \\ a_2 & 0 & & \\ & 0 & a_2 & \end{bmatrix} \begin{bmatrix} b_1 X_{11} + b_2 X_{12} \\ b_1 X_{21} + b_2 X_{22} \end{bmatrix} \in \mathbb{R}^{2 \times 1 \times 2} \\ &= \begin{bmatrix} a_1(b_1 X_{11} + b_2 X_{12}) & a_1(b_1 X_{21} + b_2 X_{22}) \\ a_2(b_1 X_{11} + b_2 X_{12}) & a_2(b_1 X_{21} + b_2 X_{22}) \end{bmatrix} \in \mathbb{R}^{2 \times 2} \\ &= ab^T X^T \end{aligned} \quad (1666)$$

where the cubix appears as a complete $2 \times 2 \times 2$ matrix. In like manner for the second term $\nabla_X(g) f$

$$\begin{aligned} \nabla_X(Xb) X^T a &= \begin{bmatrix} b_1 & 0 & & \\ & b_2 & 0 & \\ 0 & & b_1 & \\ & 0 & & b_2 \end{bmatrix} \begin{bmatrix} X_{11} a_1 + X_{21} a_2 \\ X_{12} a_1 + X_{22} a_2 \end{bmatrix} \in \mathbb{R}^{2 \times 1 \times 2} \\ &= X^T ab^T \in \mathbb{R}^{2 \times 2} \end{aligned} \quad (1667)$$

The solution

$$\nabla_X a^T X^2 b = ab^T X^T + X^T ab^T \quad (1668)$$

can be found from Table D.2.1 or verified using (1661). \square

D.1.2.1 Kronecker product

A partial remedy for venturing into *hyperdimensional* representations, such as the cubix or quartix, is to first vectorize matrices as in (31). This device gives rise to the Kronecker product of matrices \otimes ; **a.k.a.**, *direct product* or *tensor product*. Although it sees reversal in the literature, [290, §2.1] we adopt the definition: for $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{p \times q}$

$$B \otimes A \triangleq \begin{bmatrix} B_{11}A & B_{12}A & \cdots & B_{1q}A \\ B_{21}A & B_{22}A & \cdots & B_{2q}A \\ \vdots & \vdots & \ddots & \vdots \\ B_{p1}A & B_{p2}A & \cdots & B_{pq}A \end{bmatrix} \in \mathbb{R}^{pm \times qn} \quad (1669)$$

for which $A \otimes 1 = 1 \otimes A = A$ (real unity acts like identity).

One advantage to vectorization is existence of a traditional two-dimensional matrix representation for the second-order gradient of a real function with respect to a vectorized matrix. For example, from §A.1.1 no.33 (§D.2.1) for square $A, B \in \mathbb{R}^{n \times n}$ [145, §5.2] [11, §3]

$$\nabla_{\text{vec } X}^2 \text{tr}(AXBX^T) = \nabla_{\text{vec } X}^2 \text{vec}(X)^T (B^T \otimes A) \text{vec } X = B \otimes A^T + B^T \otimes A \in \mathbb{R}^{n^2 \times n^2} \quad (1670)$$

To disadvantage is a large new but known set of algebraic rules (§A.1.1) and the fact that its mere use does not generally guarantee two-dimensional matrix representation of gradients.

Another application of the Kronecker product is to reverse order of appearance in a matrix product: Suppose we wish to weight the columns of a matrix $S \in \mathbb{R}^{M \times N}$, for example, by respective entries w_i from the main-diagonal in

$$W \triangleq \begin{bmatrix} w_1 & & \mathbf{0} \\ & \ddots & \\ \mathbf{0}^T & & w_N \end{bmatrix} \in \mathbb{S}^N \quad (1671)$$

The conventional way of accomplishing that is to multiply S by diagonal matrix W on the right-hand side:^{D.2}

$$SW = S \begin{bmatrix} w_1 & & \mathbf{0} \\ & \ddots & \\ \mathbf{0}^T & & w_N \end{bmatrix} = [S(:, 1)w_1 \quad \cdots \quad S(:, N)w_N] \in \mathbb{R}^{M \times N} \quad (1672)$$

To reverse product order such that diagonal matrix W instead appears to the left of S : (Sze Wan)

$$SW = (\delta(W)^T \otimes I) \begin{bmatrix} S(:, 1) & \mathbf{0} & & \mathbf{0} \\ \mathbf{0} & S(:, 2) & \ddots & \\ & \ddots & \ddots & \mathbf{0} \\ \mathbf{0} & & \mathbf{0} & S(:, N) \end{bmatrix} \in \mathbb{R}^{M \times N} \quad (1673)$$

^{D.2}Multiplying on the left by $W \in \mathbb{S}^M$ would instead weight the rows of S .

where $I \in \mathbb{S}^M$. For any matrices of like size, $S, Y \in \mathbb{R}^{M \times N}$

$$S \circ Y = [\delta(Y(:, 1)) \cdots \delta(Y(:, N))] \begin{bmatrix} S(:, 1) & \mathbf{0} & & \mathbf{0} \\ \mathbf{0} & S(:, 2) & \cdots & \\ & \ddots & \ddots & \mathbf{0} \\ \mathbf{0} & & \mathbf{0} & S(:, N) \end{bmatrix} \in \mathbb{R}^{M \times N} \quad (1674)$$

which converts a Hadamard product into a standard matrix product. In the special case that $S = s$ and $Y = y$ are vectors in \mathbb{R}^M

$$s \circ y = \delta(s)y \quad (1675)$$

$$\begin{aligned} s^T \otimes y &= ys^T \\ s \otimes y^T &= sy^T \end{aligned} \quad (1676)$$

D.1.3 Chain rules for composite matrix-functions

Given dimensionally compatible matrix-valued functions of matrix variable $f(X)$ and $g(X)$ [194, §15.7]

$$\nabla_X g(f(X)^T) = \nabla_X f^T \nabla_f g \quad (1677)$$

$$\nabla_X^2 g(f(X)^T) = \nabla_X (\nabla_X f^T \nabla_f g) = \nabla_X^2 f \nabla_f g + \nabla_X f^T \nabla_f^2 g \nabla_X f \quad (1678)$$

D.1.3.1 Two arguments

$$\nabla_X g(f(X)^T, h(X)^T) = \nabla_X f^T \nabla_f g + \nabla_X h^T \nabla_h g \quad (1679)$$

D.1.3.1.1 Example. *Chain rule for two arguments.* [37, §1.1]

$$g(f(x)^T, h(x)^T) = (f(x) + h(x))^T A (f(x) + h(x)) \quad (1680)$$

$$f(x) = \begin{bmatrix} x_1 \\ \varepsilon x_2 \end{bmatrix}, \quad h(x) = \begin{bmatrix} \varepsilon x_1 \\ x_2 \end{bmatrix} \quad (1681)$$

$$\nabla_x g(f(x)^T, h(x)^T) = \begin{bmatrix} 1 & 0 \\ 0 & \varepsilon \end{bmatrix} (A + A^T)(f + h) + \begin{bmatrix} \varepsilon & 0 \\ 0 & 1 \end{bmatrix} (A + A^T)(f + h) \quad (1682)$$

$$\nabla_x g(f(x)^T, h(x)^T) = \begin{bmatrix} 1 + \varepsilon & 0 \\ 0 & 1 + \varepsilon \end{bmatrix} (A + A^T) \left(\begin{bmatrix} x_1 \\ \varepsilon x_2 \end{bmatrix} + \begin{bmatrix} \varepsilon x_1 \\ x_2 \end{bmatrix} \right) \quad (1683)$$

$$\lim_{\varepsilon \rightarrow 0} \nabla_x g(f(x)^T, h(x)^T) = (A + A^T)x \quad (1684)$$

from Table [D.2.1](#). \square

These foregoing formulae remain correct when gradient produces hyperdimensional representation:

D.1.4 First directional derivative

Assume that a differentiable function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^{M \times N}$ has continuous first- and second-order gradients ∇g and $\nabla^2 g$ over $\text{dom } g$ which is an open set. We seek simple expressions for the first and second directional derivatives in direction $Y \in \mathbb{R}^{K \times L}$: respectively, $\overset{\rightarrow Y}{dg} \in \mathbb{R}^{M \times N}$ and $\overset{\rightarrow Y}{dg^2} \in \mathbb{R}^{M \times N}$.

Assuming that the limit exists, we may state the partial derivative of the mn^{th} entry of g with respect to the kl^{th} entry of X ;

$$\frac{\partial g_{mn}(X)}{\partial X_{kl}} = \lim_{\Delta t \rightarrow 0} \frac{g_{mn}(X + \Delta t e_k e_l^T) - g_{mn}(X)}{\Delta t} \in \mathbb{R} \quad (1685)$$

where e_k is the k^{th} standard basis vector in \mathbb{R}^K while e_l is the l^{th} standard basis vector in \mathbb{R}^L . The total number of partial derivatives equals $KLMN$ while the gradient is defined in their terms; the mn^{th} entry of the gradient is

$$\nabla g_{mn}(X) = \begin{bmatrix} \frac{\partial g_{mn}(X)}{\partial X_{11}} & \frac{\partial g_{mn}(X)}{\partial X_{12}} & \cdots & \frac{\partial g_{mn}(X)}{\partial X_{1L}} \\ \frac{\partial g_{mn}(X)}{\partial X_{21}} & \frac{\partial g_{mn}(X)}{\partial X_{22}} & \cdots & \frac{\partial g_{mn}(X)}{\partial X_{2L}} \\ \vdots & \vdots & & \vdots \\ \frac{\partial g_{mn}(X)}{\partial X_{K1}} & \frac{\partial g_{mn}(X)}{\partial X_{K2}} & \cdots & \frac{\partial g_{mn}(X)}{\partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L} \quad (1686)$$

while the gradient is a quartix

$$\nabla g(X) = \begin{bmatrix} \nabla g_{11}(X) & \nabla g_{12}(X) & \cdots & \nabla g_{1N}(X) \\ \nabla g_{21}(X) & \nabla g_{22}(X) & \cdots & \nabla g_{2N}(X) \\ \vdots & \vdots & & \vdots \\ \nabla g_{M1}(X) & \nabla g_{M2}(X) & \cdots & \nabla g_{MN}(X) \end{bmatrix} \in \mathbb{R}^{M \times N \times K \times L} \quad (1687)$$

By simply rotating our perspective of the four-dimensional representation of gradient matrix, we find one of three useful transpositions of this quartix (connoted T_1):

$$\nabla g(X)^{T_1} = \begin{bmatrix} \frac{\partial g(X)}{\partial X_{11}} & \frac{\partial g(X)}{\partial X_{12}} & \dots & \frac{\partial g(X)}{\partial X_{1L}} \\ \frac{\partial g(X)}{\partial X_{21}} & \frac{\partial g(X)}{\partial X_{22}} & \dots & \frac{\partial g(X)}{\partial X_{2L}} \\ \vdots & \vdots & & \vdots \\ \frac{\partial g(X)}{\partial X_{K1}} & \frac{\partial g(X)}{\partial X_{K2}} & \dots & \frac{\partial g(X)}{\partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L \times M \times N} \quad (1688)$$

When the limit for $\Delta t \in \mathbb{R}$ exists, it is easy to show by substitution of variables in (1685)

$$\frac{\partial g_{mn}(X)}{\partial X_{kl}} Y_{kl} = \lim_{\Delta t \rightarrow 0} \frac{g_{mn}(X + \Delta t Y_{kl} e_k e_l^T) - g_{mn}(X)}{\Delta t} \in \mathbb{R} \quad (1689)$$

which may be interpreted as the change in g_{mn} at X when the change in X_{kl} is equal to Y_{kl} , the kl^{th} entry of any $Y \in \mathbb{R}^{K \times L}$. Because the total change in $g_{mn}(X)$ due to Y is the sum of change with respect to each and every X_{kl} , the mn^{th} entry of the directional derivative is the corresponding total differential [194, §15.8]

$$dg_{mn}(X)|_{dX \rightarrow Y} = \sum_{k,l} \frac{\partial g_{mn}(X)}{\partial X_{kl}} Y_{kl} = \text{tr}(\nabla g_{mn}(X)^T Y) \quad (1690)$$

$$= \sum_{k,l} \lim_{\Delta t \rightarrow 0} \frac{g_{mn}(X + \Delta t Y_{kl} e_k e_l^T) - g_{mn}(X)}{\Delta t} \quad (1691)$$

$$= \lim_{\Delta t \rightarrow 0} \frac{g_{mn}(X + \Delta t Y) - g_{mn}(X)}{\Delta t} \quad (1692)$$

$$= \left. \frac{d}{dt} \right|_{t=0} g_{mn}(X + tY) \quad (1693)$$

where $t \in \mathbb{R}$. Assuming finite Y , equation (1692) is called the *Gâteaux differential* [36, App.A.5] [177, §D.2.1] [318, §5.28] whose existence is implied by existence of the *Fréchet differential* (the sum in (1690)). [220, §7.2] Each may be understood as the change in g_{mn} at X when the change in X is equal

in magnitude and direction to Y .^{D.3} Hence the directional derivative,

$$\begin{aligned}
\overset{\rightarrow Y}{dg}(X) &\triangleq \left[\begin{array}{cccc} dg_{11}(X) & dg_{12}(X) & \cdots & dg_{1N}(X) \\ dg_{21}(X) & dg_{22}(X) & \cdots & dg_{2N}(X) \\ \vdots & \vdots & & \vdots \\ dg_{M1}(X) & dg_{M2}(X) & \cdots & dg_{MN}(X) \end{array} \right] \Bigg|_{dX \rightarrow Y} \in \mathbb{R}^{M \times N} \\
&= \left[\begin{array}{cccc} \text{tr}(\nabla g_{11}(X)^T Y) & \text{tr}(\nabla g_{12}(X)^T Y) & \cdots & \text{tr}(\nabla g_{1N}(X)^T Y) \\ \text{tr}(\nabla g_{21}(X)^T Y) & \text{tr}(\nabla g_{22}(X)^T Y) & \cdots & \text{tr}(\nabla g_{2N}(X)^T Y) \\ \vdots & \vdots & & \vdots \\ \text{tr}(\nabla g_{M1}(X)^T Y) & \text{tr}(\nabla g_{M2}(X)^T Y) & \cdots & \text{tr}(\nabla g_{MN}(X)^T Y) \end{array} \right] \\
&= \left[\begin{array}{cccc} \sum_{k,l} \frac{\partial g_{11}(X)}{\partial X_{kl}} Y_{kl} & \sum_{k,l} \frac{\partial g_{12}(X)}{\partial X_{kl}} Y_{kl} & \cdots & \sum_{k,l} \frac{\partial g_{1N}(X)}{\partial X_{kl}} Y_{kl} \\ \sum_{k,l} \frac{\partial g_{21}(X)}{\partial X_{kl}} Y_{kl} & \sum_{k,l} \frac{\partial g_{22}(X)}{\partial X_{kl}} Y_{kl} & \cdots & \sum_{k,l} \frac{\partial g_{2N}(X)}{\partial X_{kl}} Y_{kl} \\ \vdots & \vdots & & \vdots \\ \sum_{k,l} \frac{\partial g_{M1}(X)}{\partial X_{kl}} Y_{kl} & \sum_{k,l} \frac{\partial g_{M2}(X)}{\partial X_{kl}} Y_{kl} & \cdots & \sum_{k,l} \frac{\partial g_{MN}(X)}{\partial X_{kl}} Y_{kl} \end{array} \right] \quad (1694)
\end{aligned}$$

from which it follows

$$\overset{\rightarrow Y}{dg}(X) = \sum_{k,l} \frac{\partial g(X)}{\partial X_{kl}} Y_{kl} \quad (1695)$$

Yet for all $X \in \text{dom } g$, any $Y \in \mathbb{R}^{K \times L}$, and some open interval of $t \in \mathbb{R}$

$$g(X + tY) = g(X) + t \overset{\rightarrow Y}{dg}(X) + o(t^2) \quad (1696)$$

which is the first-order Taylor series expansion about X . [194, §18.4] [133, §2.3.4] Differentiation with respect to t and subsequent t -zeroing isolates the second term of expansion. Thus differentiating and zeroing $g(X + tY)$ in t is an operation equivalent to individually differentiating and zeroing every entry $g_{mn}(X + tY)$ as in (1693). So the directional derivative of $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^{M \times N}$ in any direction $Y \in \mathbb{R}^{K \times L}$ evaluated at $X \in \text{dom } g$ becomes

$$\overset{\rightarrow Y}{dg}(X) = \frac{d}{dt} \Bigg|_{t=0} g(X + tY) \in \mathbb{R}^{M \times N} \quad (1697)$$

^{D.3} Although Y is a matrix, we may regard it as a vector in \mathbb{R}^{KL} .




Figure 139: Strictly convex quadratic bowl in $\mathbb{R}^2 \times \mathbb{R}$; $f(x) = x^T x : \mathbb{R}^2 \rightarrow \mathbb{R}$ versus x on some open disc in \mathbb{R}^2 . Plane slice $\partial\mathcal{H}$ is perpendicular to function domain. Slice intersection with domain connotes bidirectional vector y . Slope of tangent line \mathcal{T} at point $(\alpha, f(\alpha))$ is value of $\nabla_x f(\alpha)^T y$ directional derivative (1722) at α in slice direction y . Negative gradient $-\nabla_x f(x) \in \mathbb{R}^2$ is direction of *steepest descent*. [334] [194, §15.6] [133] When vector $v \in \mathbb{R}^3$ entry v_3 is half directional derivative in gradient direction at α and when $\begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \nabla_x f(\alpha)$, then $-v$ points directly toward bowl bottom.

[244, §2.1, §5.4.5] [31, §6.3.1] which is simplest. In case of a real function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}$

$$\overrightarrow{dg}(X) = \text{tr}(\nabla g(X)^T Y) \quad (1719)$$

In case $g(X) : \mathbb{R}^K \rightarrow \mathbb{R}$

$$\overrightarrow{dg}(X) = \nabla g(X)^T Y \quad (1722)$$

Unlike gradient, directional derivative does not expand dimension; directional derivative (1697) retains the dimensions of g . The derivative with respect to t makes the directional derivative resemble ordinary calculus (§D.2); e.g., when $g(X)$ is linear, $\overrightarrow{dg}(X) = g(Y)$. [220, §7.2]

D.1.4.1 Interpretation of directional derivative

In the case of any differentiable real function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}$, the directional derivative of $g(X)$ at X in any direction Y yields the slope of g along the line $\{X + tY \mid t \in \mathbb{R}\}$ through its domain evaluated at $t=0$. For higher-dimensional functions, by (1694), this slope interpretation can be applied to each entry of the directional derivative.

Figure 139, for example, shows a plane slice of a real convex bowl-shaped function $f(x)$ along a line $\{\alpha + ty \mid t \in \mathbb{R}\}$ through its domain. The slice reveals a one-dimensional real function of t ; $f(\alpha + ty)$. The directional derivative at $x = \alpha$ in direction y is the slope of $f(\alpha + ty)$ with respect to t at $t=0$. In the case of a real function having vector argument $h(X) : \mathbb{R}^K \rightarrow \mathbb{R}$, its directional derivative in the normalized direction of its gradient is the gradient magnitude. (1722) For a real function of real variable, the directional derivative evaluated at any point in the function domain is just the slope of that function there scaled by the real direction. (*confer* §3.1.8)

Directional derivative generalizes our one-dimensional notion of derivative to a multidimensional domain. When direction Y coincides with a member of the standard unit Cartesian basis $e_k e_l^T$ (54), then a single partial derivative $\partial g(X) / \partial X_{kl}$ is obtained from directional derivative (1695); such is each entry of gradient $\nabla g(X)$ in equalities (1719) and (1722), for example.

D.1.4.1.1 Theorem. Directional derivative optimality condition.

[220, §7.4] Suppose $f(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}$ is minimized on convex set $\mathcal{C} \subseteq \mathbb{R}^{p \times k}$ by X^* , and the directional derivative of f exists there. Then for all $X \in \mathcal{C}$

$$df(X) \stackrel{\rightarrow{X-X^*}}{\geq} 0 \quad (1698)$$

◇

D.1.4.1.2 Example. Simple bowl.

Bowl function (Figure 139)

$$f(x) : \mathbb{R}^K \rightarrow \mathbb{R} \stackrel{\Delta}{=} (x - a)^T(x - a) - b \quad (1699)$$

has function offset $-b \in \mathbb{R}$, axis of revolution at $x = a$, and positive definite Hessian (1649) everywhere in its domain (an open *hyperdisc* in \mathbb{R}^K); *id est*, strictly convex quadratic $f(x)$ has unique global minimum equal to $-b$ at

$x = a$. A vector $-v$ based anywhere in $\text{dom } f \times \mathbb{R}$ pointing toward the unique bowl-bottom is specified:

$$v \propto \begin{bmatrix} x - a \\ f(x) + b \end{bmatrix} \in \mathbb{R}^K \times \mathbb{R} \quad (1700)$$

Such a vector is

$$v = \begin{bmatrix} \nabla_x f(x) \\ -\nabla_x f(x) \\ \frac{1}{2} df(x) \end{bmatrix} \quad (1701)$$

since the gradient is

$$\nabla_x f(x) = 2(x - a) \quad (1702)$$

and the directional derivative in the direction of the gradient is (1722)

$$\begin{aligned} \begin{matrix} \rightarrow \\ \nabla_x f(x) \end{matrix} df(x) &= \nabla_x f(x)^\top \nabla_x f(x) = 4(x - a)^\top (x - a) = 4(f(x) + b) \quad (1703) \\ &\quad \square \end{aligned}$$

D.1.5 Second directional derivative

By similar argument, it so happens: the second directional derivative is equally simple. Given $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^{M \times N}$ on open domain,

$$\nabla \frac{\partial g_{mn}(X)}{\partial X_{kl}} = \frac{\partial \nabla g_{mn}(X)}{\partial X_{kl}} = \begin{bmatrix} \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{11}} & \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{12}} & \cdots & \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{1L}} \\ \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{21}} & \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{22}} & \cdots & \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{2L}} \\ \vdots & \vdots & & \vdots \\ \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{K1}} & \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{K2}} & \cdots & \frac{\partial^2 g_{mn}(X)}{\partial X_{ki} \partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L} \quad (1704)$$

$$\begin{aligned} \nabla^2 g_{mn}(X) &= \begin{bmatrix} \nabla \frac{\partial g_{mn}(X)}{\partial X_{11}} & \nabla \frac{\partial g_{mn}(X)}{\partial X_{12}} & \cdots & \nabla \frac{\partial g_{mn}(X)}{\partial X_{1L}} \\ \nabla \frac{\partial g_{mn}(X)}{\partial X_{21}} & \nabla \frac{\partial g_{mn}(X)}{\partial X_{22}} & \cdots & \nabla \frac{\partial g_{mn}(X)}{\partial X_{2L}} \\ \vdots & \vdots & & \vdots \\ \nabla \frac{\partial g_{mn}(X)}{\partial X_{K1}} & \nabla \frac{\partial g_{mn}(X)}{\partial X_{K2}} & \cdots & \nabla \frac{\partial g_{mn}(X)}{\partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L \times K \times L} \\ &\quad (1705) \end{aligned}$$

$$= \begin{bmatrix} \frac{\partial \nabla g_{mn}(X)}{\partial X_{11}} & \frac{\partial \nabla g_{mn}(X)}{\partial X_{12}} & \cdots & \frac{\partial \nabla g_{mn}(X)}{\partial X_{1L}} \\ \frac{\partial \nabla g_{mn}(X)}{\partial X_{21}} & \frac{\partial \nabla g_{mn}(X)}{\partial X_{22}} & \cdots & \frac{\partial \nabla g_{mn}(X)}{\partial X_{2L}} \\ \vdots & \vdots & & \vdots \\ \frac{\partial \nabla g_{mn}(X)}{\partial X_{K1}} & \frac{\partial \nabla g_{mn}(X)}{\partial X_{K2}} & \cdots & \frac{\partial \nabla g_{mn}(X)}{\partial X_{KL}} \end{bmatrix}$$

Rotating our perspective, we get several views of the second-order gradient:

$$\nabla^2 g(X) = \begin{bmatrix} \nabla^2 g_{11}(X) & \nabla^2 g_{12}(X) & \cdots & \nabla^2 g_{1N}(X) \\ \nabla^2 g_{21}(X) & \nabla^2 g_{22}(X) & \cdots & \nabla^2 g_{2N}(X) \\ \vdots & \vdots & \ddots & \vdots \\ \nabla^2 g_{M1}(X) & \nabla^2 g_{M2}(X) & \cdots & \nabla^2 g_{MN}(X) \end{bmatrix} \in \mathbb{R}^{M \times N \times K \times L \times K \times L} \quad (1706)$$

$$\nabla^2 g(X)^{T_1} = \begin{bmatrix} \nabla \frac{\partial g(X)}{\partial X_{11}} & \nabla \frac{\partial g(X)}{\partial X_{12}} & \cdots & \nabla \frac{\partial g(X)}{\partial X_{1L}} \\ \nabla \frac{\partial g(X)}{\partial X_{21}} & \nabla \frac{\partial g(X)}{\partial X_{22}} & \cdots & \nabla \frac{\partial g(X)}{\partial X_{2L}} \\ \vdots & \vdots & \ddots & \vdots \\ \nabla \frac{\partial g(X)}{\partial X_{K1}} & \nabla \frac{\partial g(X)}{\partial X_{K2}} & \cdots & \nabla \frac{\partial g(X)}{\partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L \times M \times N \times K \times L} \quad (1707)$$

$$\nabla^2 g(X)^{T_2} = \begin{bmatrix} \frac{\partial \nabla g(X)}{\partial X_{11}} & \frac{\partial \nabla g(X)}{\partial X_{12}} & \cdots & \frac{\partial \nabla g(X)}{\partial X_{1L}} \\ \frac{\partial \nabla g(X)}{\partial X_{21}} & \frac{\partial \nabla g(X)}{\partial X_{22}} & \cdots & \frac{\partial \nabla g(X)}{\partial X_{2L}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial \nabla g(X)}{\partial X_{K1}} & \frac{\partial \nabla g(X)}{\partial X_{K2}} & \cdots & \frac{\partial \nabla g(X)}{\partial X_{KL}} \end{bmatrix} \in \mathbb{R}^{K \times L \times K \times L \times M \times N} \quad (1708)$$

Assuming the limits exist, we may state the partial derivative of the mn^{th} entry of g with respect to the kl^{th} and ij^{th} entries of X ;

$$\frac{\partial^2 g_{mn}(X)}{\partial X_{kl} \partial X_{ij}} = \lim_{\Delta\tau, \Delta t \rightarrow 0} \frac{g_{mn}(X + \Delta t e_k e_l^T + \Delta\tau e_i e_j^T) - g_{mn}(X + \Delta t e_k e_l^T) - (g_{mn}(X + \Delta\tau e_i e_j^T) - g_{mn}(X))}{\Delta\tau \Delta t} \quad (1709)$$

Differentiating (1689) and then scaling by Y_{ij}

$$\begin{aligned} \frac{\partial^2 g_{mn}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} &= \lim_{\Delta t \rightarrow 0} \frac{\partial g_{mn}(X + \Delta t Y_{kl} e_k e_l^T) - \partial g_{mn}(X)}{\partial X_{ij} \Delta t} Y_{ij} \\ &= \lim_{\Delta\tau, \Delta t \rightarrow 0} \frac{g_{mn}(X + \Delta t Y_{kl} e_k e_l^T + \Delta\tau Y_{ij} e_i e_j^T) - g_{mn}(X + \Delta t Y_{kl} e_k e_l^T) - (g_{mn}(X + \Delta\tau Y_{ij} e_i e_j^T) - g_{mn}(X))}{\Delta\tau \Delta t} \end{aligned} \quad (1710)$$

which can be proved by substitution of variables in (1709). The mn^{th} second-order total differential due to any $Y \in \mathbb{R}^{K \times L}$ is

$$d^2g_{mn}(X)|_{dX \rightarrow Y} = \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{mn}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} = \text{tr} \left(\nabla_X \text{tr}(\nabla g_{mn}(X)^T Y)^T Y \right) \quad (1711)$$

$$= \sum_{i,j} \lim_{\Delta t \rightarrow 0} \frac{\partial g_{mn}(X + \Delta t Y) - \partial g_{mn}(X)}{\partial X_{ij} \Delta t} Y_{ij} \quad (1712)$$

$$= \lim_{\Delta t \rightarrow 0} \frac{g_{mn}(X + 2\Delta t Y) - 2g_{mn}(X + \Delta t Y) + g_{mn}(X)}{\Delta t^2} \quad (1713)$$

$$= \left. \frac{d^2}{dt^2} \right|_{t=0} g_{mn}(X + t Y) \quad (1714)$$

Hence the second directional derivative,

$$\begin{aligned} \overset{-Y}{dg^2}(X) &\triangleq \left[\begin{array}{cccc} d^2g_{11}(X) & d^2g_{12}(X) & \cdots & d^2g_{1N}(X) \\ d^2g_{21}(X) & d^2g_{22}(X) & \cdots & d^2g_{2N}(X) \\ \vdots & \vdots & & \vdots \\ d^2g_{M1}(X) & d^2g_{M2}(X) & \cdots & d^2g_{MN}(X) \end{array} \right] \Bigg|_{dX \rightarrow Y} \in \mathbb{R}^{M \times N} \\ &= \left[\begin{array}{cccc} \text{tr}(\nabla \text{tr}(\nabla g_{11}(X)^T Y)^T Y) & \text{tr}(\nabla \text{tr}(\nabla g_{12}(X)^T Y)^T Y) & \cdots & \text{tr}(\nabla \text{tr}(\nabla g_{1N}(X)^T Y)^T Y) \\ \text{tr}(\nabla \text{tr}(\nabla g_{21}(X)^T Y)^T Y) & \text{tr}(\nabla \text{tr}(\nabla g_{22}(X)^T Y)^T Y) & \cdots & \text{tr}(\nabla \text{tr}(\nabla g_{2N}(X)^T Y)^T Y) \\ \vdots & \vdots & & \vdots \\ \text{tr}(\nabla \text{tr}(\nabla g_{M1}(X)^T Y)^T Y) & \text{tr}(\nabla \text{tr}(\nabla g_{M2}(X)^T Y)^T Y) & \cdots & \text{tr}(\nabla \text{tr}(\nabla g_{MN}(X)^T Y)^T Y) \end{array} \right] \\ &= \left[\begin{array}{cccc} \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{11}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} & \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{12}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} & \cdots & \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{1N}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} \\ \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{21}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} & \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{22}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} & \cdots & \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{2N}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} \\ \vdots & \vdots & & \vdots \\ \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{M1}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} & \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{M2}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} & \cdots & \sum_{i,j} \sum_{k,l} \frac{\partial^2 g_{MN}(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} \end{array} \right] \quad (1715) \end{aligned}$$

from which it follows

$$\overset{\rightarrow Y}{dg^2}(X) = \sum_{i,j} \sum_{k,l} \frac{\partial^2 g(X)}{\partial X_{kl} \partial X_{ij}} Y_{kl} Y_{ij} = \sum_{i,j} \frac{\partial}{\partial X_{ij}} \overset{\rightarrow Y}{dg}(X) Y_{ij} \quad (1716)$$

Yet for all $X \in \text{dom } g$, any $Y \in \mathbb{R}^{K \times L}$, and some open interval of $t \in \mathbb{R}$

$$g(X + tY) = g(X) + t \overset{\rightarrow Y}{dg}(X) + \frac{1}{2!} t^2 \overset{\rightarrow Y}{dg^2}(X) + o(t^3) \quad (1717)$$

which is the second-order Taylor series expansion about X . [194, §18.4] [133, §2.3.4] Differentiating twice with respect to t and subsequent t -zeroing isolates the third term of the expansion. Thus differentiating and zeroing $g(X + tY)$ in t is an operation equivalent to individually differentiating and zeroing every entry $g_{mn}(X + tY)$ as in (1714). So the second directional derivative of $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^{M \times N}$ becomes [244, §2.1, §5.4.5] [31, §6.3.1]

$$\overset{\rightarrow Y}{dg^2}(X) = \left. \frac{d^2}{dt^2} \right|_{t=0} g(X + tY) \in \mathbb{R}^{M \times N} \quad (1718)$$

which is again simplest. (*confer*(1697)) Directional derivative retains the dimensions of g .

D.1.6 directional derivative expressions

In the case of a real function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}$, all its directional derivatives are in \mathbb{R} :

$$\overset{\rightarrow Y}{dg}(X) = \text{tr}(\nabla g(X)^T Y) \quad (1719)$$

$$\overset{\rightarrow Y}{dg^2}(X) = \text{tr}(\nabla_X \text{tr}(\nabla g(X)^T Y)^T Y) = \text{tr}(\nabla_X \overset{\rightarrow Y}{dg}(X)^T Y) \quad (1720)$$

$$\overset{\rightarrow Y}{dg^3}(X) = \text{tr}(\nabla_X \text{tr}(\nabla_X \text{tr}(\nabla g(X)^T Y)^T Y)^T Y) = \text{tr}(\nabla_X \overset{\rightarrow Y}{dg^2}(X)^T Y) \quad (1721)$$

In the case $g(X) : \mathbb{R}^K \rightarrow \mathbb{R}$ has vector argument, they further simplify:

$$\overset{\rightarrow Y}{dg}(X) = \nabla g(X)^T Y \quad (1722)$$

$$\overset{\rightarrow Y}{dg^2}(X) = Y^T \nabla^2 g(X) Y \quad (1723)$$

$$\overset{\rightarrow Y}{dg^3}(X) = \nabla_X (Y^T \nabla^2 g(X) Y)^T Y \quad (1724)$$

and so on.

D.1.7 Taylor series

Series expansions of the differentiable matrix-valued function $g(X)$, of matrix argument, were given earlier in (1696) and (1717). Assuming $g(X)$ has continuous first-, second-, and third-order gradients over the open set $\text{dom } g$, then for $X \in \text{dom } g$ and any $Y \in \mathbb{R}^{K \times L}$ the complete Taylor series on some open interval of $\mu \in \mathbb{R}$ is expressed

$$g(X + \mu Y) = g(X) + \mu \overset{\rightarrow Y}{dg}(X) + \frac{1}{2!} \mu^2 \overset{\rightarrow Y}{dg^2}(X) + \frac{1}{3!} \mu^3 \overset{\rightarrow Y}{dg^3}(X) + o(\mu^4) \quad (1725)$$

or on some open interval of $\|Y\|_2$

$$g(Y) = g(X) + \overset{\rightarrow Y-X}{dg}(X) + \frac{1}{2!} \overset{\rightarrow Y-X}{dg^2}(X) + \frac{1}{3!} \overset{\rightarrow Y-X}{dg^3}(X) + o(\|Y\|^4) \quad (1726)$$

which are third-order expansions about X . The *mean value theorem* from calculus is what insures finite order of the series. [194] [37, §1.1] [36, App.A.5] [177, §0.4]

D.1.7.0.1 Example. Inverse matrix function.

Say $g(Y) = Y^{-1}$. Let's find its Taylor series expansion about $X = I$. Continuing the table on page 636,

$$\overset{\rightarrow Y}{dg}(X) = \left. \frac{d}{dt} \right|_{t=0} g(X + tY) = -X^{-1} Y X^{-1} \quad (1727)$$

$$\overset{\rightarrow Y}{dg^2}(X) = \left. \frac{d^2}{dt^2} \right|_{t=0} g(X + tY) = 2X^{-1} Y X^{-1} Y X^{-1} \quad (1728)$$

$$\overset{\rightarrow Y}{dg^3}(X) = \left. \frac{d^3}{dt^3} \right|_{t=0} g(X + tY) = -6X^{-1} Y X^{-1} Y X^{-1} Y X^{-1} \quad (1729)$$

Since $g(I) = I$, ($\mu = 1$) for $\|Y\|_2 < 1$

$$g(I + Y) = (I + Y)^{-1} = I - Y + Y^2 - Y^3 + \dots \quad (1730)$$

If Y is small, $(I + Y)^{-1} \approx I - Y$. (Had we instead set $g(Y) = (I + Y)^{-1}$, then the equivalent expansion would have been about $X = \mathbf{0}$.) \square

D.1.7.0.2 Exercise. *log det.* (confer [53, p.644])
Find the first three terms of a Taylor series expansion for $\log \det Y$. Specify an open interval over which the expansion holds in vicinity of X . \blacktriangledown

D.1.8 Correspondence of gradient to derivative

From the foregoing expressions for directional derivative, we derive a relationship between the gradient with respect to matrix X and the derivative with respect to real variable t :

D.1.8.1 first-order

Removing from (1697) the evaluation at $t = 0$, ^{D.4} we find an expression for the directional derivative of $g(X)$ in direction Y evaluated anywhere along a line $\{X + tY \mid t \in \mathbb{R}\}$ intersecting $\text{dom } g$

$$\overset{\rightarrow Y}{dg}(X + tY) = \frac{d}{dt}g(X + tY) \quad (1731)$$

In the general case $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^{M \times N}$, from (1690) and (1693) we find

$$\text{tr}(\nabla_X g_{mn}(X + tY)^T Y) = \frac{d}{dt}g_{mn}(X + tY) \quad (1732)$$

which is valid at $t = 0$, of course, when $X \in \text{dom } g$. In the important case of a real function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}$, from (1719) we have simply

$$\text{tr}(\nabla_X g(X + tY)^T Y) = \frac{d}{dt}g(X + tY) \quad (1733)$$

^{D.4}Justified by replacing X with $X + tY$ in (1690)-(1692); beginning,

$$dg_{mn}(X + tY)|_{dX \rightarrow Y} = \sum_{k,l} \frac{\partial g_{mn}(X + tY)}{\partial X_{kl}} Y_{kl}$$

When, additionally, $g(X) : \mathbb{R}^K \rightarrow \mathbb{R}$ has vector argument,

$$\nabla_X g(X + tY)^T Y = \frac{d}{dt} g(X + tY) \quad (1734)$$

D.1.8.1.1 Example. *Gradient.*

$g(X) = w^T X^T X w$, $X \in \mathbb{R}^{K \times L}$, $w \in \mathbb{R}^L$. Using the tables in §D.2,

$$\text{tr}(\nabla_X g(X + tY)^T Y) = \text{tr}(2ww^T(X^T + tY^T)Y) \quad (1735)$$

$$= 2w^T(X^T Y + tY^T Y)w \quad (1736)$$

Applying the equivalence (1733),

$$\frac{d}{dt} g(X + tY) = \frac{d}{dt} w^T (X + tY)^T (X + tY) w \quad (1737)$$

$$= w^T (X^T Y + Y^T X + 2tY^T Y) w \quad (1738)$$

$$= 2w^T (X^T Y + tY^T Y) w \quad (1739)$$

which is the same as (1736); hence, equivalence is demonstrated.

It is easy to extract $\nabla g(X)$ from (1739) knowing only (1733):

$$\begin{aligned} \text{tr}(\nabla_X g(X + tY)^T Y) &= 2w^T (X^T Y + tY^T Y) w \\ &= 2 \text{tr}(ww^T (X^T + tY^T) Y) \end{aligned} \quad (1740)$$

$$\text{tr}(\nabla_X g(X)^T Y) = 2 \text{tr}(ww^T X^T Y) \quad (1740)$$

\Leftrightarrow

$$\nabla_X g(X) = 2Xww^T$$

□

D.1.8.2 second-order

Likewise removing the evaluation at $t = 0$ from (1718),

$$\overset{\rightarrow Y}{dg^2}(X + tY) = \frac{d^2}{dt^2} g(X + tY) \quad (1741)$$

we can find a similar relationship between the second-order gradient and the second derivative: In the general case $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}^{M \times N}$ from (1711) and (1714),

$$\text{tr}\left(\nabla_X \text{tr}(\nabla_X g_{mn}(X + tY)^T Y)^T Y\right) = \frac{d^2}{dt^2} g_{mn}(X + tY) \quad (1742)$$

In the case of a real function $g(X) : \mathbb{R}^{K \times L} \rightarrow \mathbb{R}$ we have, of course,

$$\operatorname{tr}\left(\nabla_X \operatorname{tr}\left(\nabla_X g(X+tY)^T Y\right)^T Y\right) = \frac{d^2}{dt^2} g(X+tY) \quad (1743)$$

From (1723), the simpler case, where the real function $g(X) : \mathbb{R}^K \rightarrow \mathbb{R}$ has vector argument,

$$Y^T \nabla_X^2 g(X+tY) Y = \frac{d^2}{dt^2} g(X+tY) \quad (1744)$$

D.1.8.2.1 Example. *Second-order gradient.*

Given real function $g(X) = \log \det X$ having domain $\operatorname{int} \mathbb{S}_+^K$, we want to find $\nabla^2 g(X) \in \mathbb{R}^{K \times K \times K \times K}$. From the tables in §D.2,

$$h(X) \triangleq \nabla g(X) = X^{-1} \in \operatorname{int} \mathbb{S}_+^K \quad (1745)$$

so $\nabla^2 g(X) = \nabla h(X)$. By (1732) and (1696), for $Y \in \mathbb{S}^K$

$$\operatorname{tr}(\nabla h_{mn}(X)^T Y) = \left. \frac{d}{dt} \right|_{t=0} h_{mn}(X+tY) \quad (1746)$$

$$= \left(\left. \frac{d}{dt} \right|_{t=0} h(X+tY) \right)_{mn} \quad (1747)$$

$$= \left(\left. \frac{d}{dt} \right|_{t=0} (X+tY)^{-1} \right)_{mn} \quad (1748)$$

$$= -(X^{-1} Y X^{-1})_{mn} \quad (1749)$$

Setting Y to a member of $\{e_k e_l^T \in \mathbb{R}^{K \times K} \mid k, l = 1 \dots K\}$, and employing a property (33) of the trace function we find

$$\nabla^2 g(X)_{mnkl} = \operatorname{tr}(\nabla h_{mn}(X)^T e_k e_l^T) = \nabla h_{mn}(X)_{kl} = -(X^{-1} e_k e_l^T X^{-1})_{mn} \quad (1750)$$

$$\nabla^2 g(X)_{kl} = \nabla h(X)_{kl} = -(X^{-1} e_k e_l^T X^{-1}) \in \mathbb{R}^{K \times K} \quad (1751)$$

□

From all these first- and second-order expressions, we may generate new ones by evaluating both sides at arbitrary t (in some open interval) but only after the differentiation.

D.2 Tables of gradients and derivatives

[145] [57]

- When proving results for symmetric matrices algebraically, it is critical to take gradients ignoring symmetry and to then substitute symmetric entries afterward.
- $a, b \in \mathbb{R}^n$, $x, y \in \mathbb{R}^k$, $A, B \in \mathbb{R}^{m \times n}$, $X, Y \in \mathbb{R}^{K \times L}$, $t, \mu \in \mathbb{R}$, $i, j, k, \ell, K, L, m, n, M, N$ are integers, unless otherwise noted.
- x^μ means $\delta(\delta(x)^\mu)$ for $\mu \in \mathbb{R}$; *id est*, entrywise vector exponentiation. δ is the main-diagonal linear operator (1328). $x^0 \triangleq \mathbf{1}$, $X^0 \triangleq I$ if square.
- $\frac{d}{dx} \triangleq \begin{bmatrix} \frac{d}{dx_1} \\ \vdots \\ \frac{d}{dx_k} \end{bmatrix}$, $\overset{\rightarrow y}{dg}(x)$, $\overset{\rightarrow y}{dg^2}(x)$ (directional derivatives §D.1), $\log x$, e^x , $|x|$, $\operatorname{sgn} x$, x/y (Hadamard quotient), \sqrt{x} (entrywise square root), *etcetera*, are maps $f: \mathbb{R}^k \rightarrow \mathbb{R}^k$ that maintain dimension; *e.g.*, (§A.1.1)

$$\frac{d}{dx} x^{-1} \triangleq \nabla_x \mathbf{1}^T \delta(x)^{-1} \mathbf{1} \quad (1752)$$

- For A a scalar or square matrix, we have the Taylor series [68, §3.6]

$$e^A \triangleq \sum_{k=0}^{\infty} \frac{1}{k!} A^k \quad (1753)$$

Further, [294, §5.4]

$$e^A \succ 0 \quad \forall A \in \mathbb{S}^m \quad (1754)$$

- For all square A and integer k

$$\det^k A = \det A^k \quad (1755)$$

D.2.1 algebraic

$\nabla_x x = \nabla_x x^T = I \in \mathbb{R}^{k \times k}$	$\nabla_X X = \nabla_X X^T \triangleq I \in \mathbb{R}^{K \times L \times K \times L}$ (identity)
$\nabla_x (Ax - b) = A^T$	
$\nabla_x (x^T A - b^T) = A$	
$\nabla_x (Ax - b)^T (Ax - b) = 2A^T (Ax - b)$	
$\nabla_x^2 (Ax - b)^T (Ax - b) = 2A^T A$	
$\nabla_x z^T Ax - b = A^T \delta(z) \operatorname{sgn}(Ax - b)$, $z_i \neq 0 \Rightarrow (Ax - b)_i \neq 0$	
$\nabla_x \mathbf{1}^T f(Ax - b) = A^T \delta\left(\frac{df(y)}{dy}\bigg _{y= Ax-b }\right) \operatorname{sgn}(Ax - b)$	
$\nabla_x (x^T Ax + 2x^T B y + y^T C y) = (A + A^T)x + 2B y$	
$\nabla_x (x + y)^T A (x + y) = (A + A^T)(x + y)$	
$\nabla_x^2 (x^T Ax + 2x^T B y + y^T C y) = A + A^T$	
	$\nabla_X a^T X b = \nabla_X b^T X^T a = ab^T$
	$\nabla_X a^T X^2 b = X^T ab^T + ab^T X^T$
	$\nabla_X a^T X^{-1} b = -X^{-T} ab^T X^{-T}$
	$\nabla_X (X^{-1})_{kl} = \frac{\partial X^{-1}}{\partial X_{kl}} = -X^{-1} e_k e_l^T X^{-1}$, <i>confer</i> (1688)
	(1751)
$\nabla_x a^T x^T x b = 2x a^T b$	$\nabla_X a^T X^T X b = X(ab^T + ba^T)$
$\nabla_x a^T x x^T b = (ab^T + ba^T)x$	$\nabla_X a^T X X^T b = (ab^T + ba^T)X$
$\nabla_x a^T x^T x a = 2x a^T a$	$\nabla_X a^T X^T X a = 2X a a^T$
$\nabla_x a^T x x^T a = 2a a^T x$	$\nabla_X a^T X X^T a = 2a a^T X$
$\nabla_x a^T y x^T b = b a^T y$	$\nabla_X a^T Y X^T b = b a^T Y$
$\nabla_x a^T y^T x b = y b^T a$	$\nabla_X a^T Y^T X b = Y a b^T$
$\nabla_x a^T x y^T b = a b^T y$	$\nabla_X a^T X Y^T b = a b^T Y$
$\nabla_x a^T x^T y b = y a^T b$	$\nabla_X a^T X^T Y b = Y b a^T$

algebraic continued

$$\frac{d}{dt}(X + tY) = Y$$

$$\frac{d}{dt}B^T(X + tY)^{-1}A = -B^T(X + tY)^{-1}Y(X + tY)^{-1}A$$

$$\frac{d}{dt}B^T(X + tY)^{-T}A = -B^T(X + tY)^{-T}Y^T(X + tY)^{-T}A$$

$$\frac{d}{dt}B^T(X + tY)^\mu A = \dots, \quad -1 \leq \mu \leq 1, \quad X, Y \in \mathbb{S}_+^M$$

$$\frac{d^2}{dt^2}B^T(X + tY)^{-1}A = 2B^T(X + tY)^{-1}Y(X + tY)^{-1}Y(X + tY)^{-1}A$$

$$\frac{d}{dt}((X + tY)^T A (X + tY)) = Y^T A X + X^T A Y + 2tY^T A Y$$

$$\frac{d^2}{dt^2}((X + tY)^T A (X + tY)) = 2Y^T A Y$$

$$\frac{d}{dt}((X + tY)A(X + tY)) = YAX + XAY + 2tYAY$$

$$\frac{d^2}{dt^2}((X + tY)A(X + tY)) = 2YAY$$

D.2.1.0.1 Exercise. *Expand these tables.*

Provide unfinished table entries indicated by ... throughout §D.2. ▼

D.2.2 trace Kronecker

$$\nabla_{\text{vec } X} \text{tr}(AXBX^T) = \nabla_{\text{vec } X} \text{vec}(X)^T (B^T \otimes A) \text{vec } X = (B \otimes A^T + B^T \otimes A) \text{vec } X$$

$$\nabla_{\text{vec } X}^2 \text{tr}(AXBX^T) = \nabla_{\text{vec } X}^2 \text{vec}(X)^T (B^T \otimes A) \text{vec } X = B \otimes A^T + B^T \otimes A$$

D.2.3 trace

$\nabla_x \mu x = \mu I$	$\nabla_X \operatorname{tr} \mu X = \nabla_X \mu \operatorname{tr} X = \mu I$
$\nabla_x \mathbf{1}^T \delta(x)^{-1} \mathbf{1} = \frac{d}{dx} x^{-1} = -x^{-2}$	$\nabla_X \operatorname{tr} X^{-1} = -X^{-2T}$
$\nabla_x \mathbf{1}^T \delta(x)^{-1} y = -\delta(x)^{-2} y$	$\nabla_X \operatorname{tr}(X^{-1} Y) = \nabla_X \operatorname{tr}(Y X^{-1}) = -X^{-T} Y^T X^{-T}$
$\frac{d}{dx} x^\mu = \mu x^{\mu-1}$	$\nabla_X \operatorname{tr} X^\mu = \mu X^{\mu-1}, \quad X \in \mathbb{S}^M$
	$\nabla_X \operatorname{tr} X^j = j X^{(j-1)T}$
$\nabla_x (b - a^T x)^{-1} = (b - a^T x)^{-2} a$	$\nabla_X \operatorname{tr}((B - AX)^{-1}) = ((B - AX)^{-2} A)^T$
$\nabla_x (b - a^T x)^\mu = -\mu (b - a^T x)^{\mu-1} a$	
$\nabla_x x^T y = \nabla_x y^T x = y$	$\nabla_X \operatorname{tr}(X^T Y) = \nabla_X \operatorname{tr}(Y X^T) = \nabla_X \operatorname{tr}(Y^T X) = \nabla_X \operatorname{tr}(X Y^T) = Y$
	$\nabla_X \operatorname{tr}(A X B X^T) = \nabla_X \operatorname{tr}(X B X^T A) = A^T X B^T + A X B$
	$\nabla_X \operatorname{tr}(A X B X) = \nabla_X \operatorname{tr}(X B X A) = A^T X^T B^T + B^T X^T A^T$
	$\nabla_X \operatorname{tr}(A X A X A X) = \nabla_X \operatorname{tr}(X A X A X A) = 3(A X A X A)^T$
	$\nabla_X \operatorname{tr}(Y X^k) = \nabla_X \operatorname{tr}(X^k Y) = \sum_{i=0}^{k-1} (X^i Y X^{k-1-i})^T$
	$\nabla_X \operatorname{tr}(Y^T X X^T Y) = \nabla_X \operatorname{tr}(X^T Y Y^T X) = 2 Y Y^T X$
	$\nabla_X \operatorname{tr}(Y^T X^T X Y) = \nabla_X \operatorname{tr}(X Y Y^T X^T) = 2 X Y Y^T$
	$\nabla_X \operatorname{tr}((X + Y)^T (X + Y)) = 2(X + Y) = \nabla_X \ X + Y\ _F^2$
	$\nabla_X \operatorname{tr}((X + Y)(X + Y)) = 2(X + Y)^T$
	$\nabla_X \operatorname{tr}(A^T X B) = \nabla_X \operatorname{tr}(X^T A B^T) = AB^T$
	$\nabla_X \operatorname{tr}(A^T X^{-1} B) = \nabla_X \operatorname{tr}(X^{-T} A B^T) = -X^{-T} A B^T X^{-T}$
	$\nabla_X a^T X b = \nabla_X \operatorname{tr}(b a^T X) = \nabla_X \operatorname{tr}(X b a^T) = a b^T$
	$\nabla_X b^T X^T a = \nabla_X \operatorname{tr}(X^T a b^T) = \nabla_X \operatorname{tr}(a b^T X^T) = a b^T$
	$\nabla_X a^T X^{-1} b = \nabla_X \operatorname{tr}(X^{-T} a b^T) = -X^{-T} a b^T X^{-T}$
	$\nabla_X a^T X^\mu b = \dots$

trace continued

$$\frac{d}{dt} \operatorname{tr} g(X + tY) = \operatorname{tr} \frac{d}{dt} g(X + tY) \quad [181, \text{p.491}]$$

$$\frac{d}{dt} \operatorname{tr}(X + tY) = \operatorname{tr} Y$$

$$\frac{d}{dt} \operatorname{tr}^j(X + tY) = j \operatorname{tr}^{j-1}(X + tY) \operatorname{tr} Y$$

$$\frac{d}{dt} \operatorname{tr}(X + tY)^j = j \operatorname{tr}((X + tY)^{j-1} Y) \quad (\forall j)$$

$$\frac{d}{dt} \operatorname{tr}((X + tY)Y) = \operatorname{tr} Y^2$$

$$\frac{d}{dt} \operatorname{tr}((X + tY)^k Y) = \frac{d}{dt} \operatorname{tr}(Y(X + tY)^k) = k \operatorname{tr}((X + tY)^{k-1} Y^2), \quad k \in \{0, 1, 2\}$$

$$\frac{d}{dt} \operatorname{tr}((X + tY)^k Y) = \frac{d}{dt} \operatorname{tr}(Y(X + tY)^k) = \operatorname{tr} \sum_{i=0}^{k-1} (X + tY)^i Y (X + tY)^{k-1-i} Y$$

$$\frac{d}{dt} \operatorname{tr}((X + tY)^{-1} Y) = - \operatorname{tr}((X + tY)^{-1} Y (X + tY)^{-1} Y)$$

$$\frac{d}{dt} \operatorname{tr}(B^T (X + tY)^{-1} A) = - \operatorname{tr}(B^T (X + tY)^{-1} Y (X + tY)^{-1} A)$$

$$\frac{d}{dt} \operatorname{tr}(B^T (X + tY)^{-T} A) = - \operatorname{tr}(B^T (X + tY)^{-T} Y^T (X + tY)^{-T} A)$$

$$\frac{d}{dt} \operatorname{tr}(B^T (X + tY)^{-k} A) = \dots, \quad k > 0$$

$$\frac{d}{dt} \operatorname{tr}(B^T (X + tY)^\mu A) = \dots, \quad -1 \leq \mu \leq 1, \quad X, Y \in \mathbb{S}_+^M$$

$$\frac{d^2}{dt^2} \operatorname{tr}(B^T (X + tY)^{-1} A) = 2 \operatorname{tr}(B^T (X + tY)^{-1} Y (X + tY)^{-1} Y (X + tY)^{-1} A)$$

$$\frac{d}{dt} \operatorname{tr}((X + tY)^T A (X + tY)) = \operatorname{tr}(Y^T A X + X^T A Y + 2t Y^T A Y)$$

$$\frac{d^2}{dt^2} \operatorname{tr}((X + tY)^T A (X + tY)) = 2 \operatorname{tr}(Y^T A Y)$$

$$\frac{d}{dt} \operatorname{tr}((X + tY) A (X + tY)) = \operatorname{tr}(Y A X + X A Y + 2t Y A Y)$$

$$\frac{d^2}{dt^2} \operatorname{tr}((X + tY) A (X + tY)) = 2 \operatorname{tr}(Y A Y)$$

D.2.4 logarithmic determinant

$x > 0$, $\det X > 0$ on some neighborhood of X , and $\det(X + tY) > 0$ on some open interval of t ; otherwise, $\log(\)$ would be discontinuous.

$\frac{d}{dx} \log x = x^{-1}$	$\nabla_X \log \det X = X^{-T}$
	$\nabla_X^2 \log \det(X)_{kl} = \frac{\partial X^{-T}}{\partial X_{kl}} = -(X^{-1} e_k e_l^T X^{-1})^T$, confer (1705)(1751)
$\frac{d}{dx} \log x^{-1} = -x^{-1}$	$\nabla_X \log \det X^{-1} = -X^{-T}$
$\frac{d}{dx} \log x^\mu = \mu x^{-1}$	$\nabla_X \log \det^\mu X = \mu X^{-T}$
	$\nabla_X \log \det X^\mu = \mu X^{-T}$
	$\nabla_X \log \det X^k = \nabla_X \log \det^k X = kX^{-T}$
	$\nabla_X \log \det^\mu(X + tY) = \mu(X + tY)^{-T}$
$\nabla_x \log(a^T x + b) = a \frac{1}{a^T x + b}$	$\nabla_X \log \det(AX + B) = A^T(AX + B)^{-T}$
	$\nabla_X \log \det(I \pm A^T X A) = \pm A(I \pm A^T X A)^{-T} A^T$
	$\nabla_X \log \det(X + tY)^k = \nabla_X \log \det^k(X + tY) = k(X + tY)^{-T}$
	$\frac{d}{dt} \log \det(X + tY) = \text{tr}((X + tY)^{-1} Y)$
	$\frac{d^2}{dt^2} \log \det(X + tY) = -\text{tr}((X + tY)^{-1} Y(X + tY)^{-1} Y)$
	$\frac{d}{dt} \log \det(X + tY)^{-1} = -\text{tr}((X + tY)^{-1} Y)$
	$\frac{d^2}{dt^2} \log \det(X + tY)^{-1} = \text{tr}((X + tY)^{-1} Y(X + tY)^{-1} Y)$
	$\frac{d}{dt} \log \det(\delta(A(x + ty) + a)^2 + \mu I)$ $= \text{tr}((\delta(A(x + ty) + a)^2 + \mu I)^{-1} 2\delta(A(x + ty) + a)\delta(Ay))$

D.2.5 determinant

$$\nabla_X \det X = \nabla_X \det X^T = \det(X)X^{-T}$$

$$\nabla_X \det X^{-1} = -\det(X^{-1})X^{-T} = -\det(X)^{-1}X^{-T}$$

$$\nabla_X \det^\mu X = \mu \det^\mu(X)X^{-T}$$

$$\nabla_X \det X^\mu = \mu \det(X^\mu)X^{-T}$$

$$\nabla_X \det X^k = k \det^{k-1}(X)(\text{tr}(X)I - X^T), \quad X \in \mathbb{R}^{2 \times 2}$$

$$\nabla_X \det X^k = \nabla_X \det^k X = k \det(X^k)X^{-T} = k \det^k(X)X^{-T}$$

$$\nabla_X \det^\mu(X + tY) = \mu \det^\mu(X + tY)(X + tY)^{-T}$$

$$\nabla_X \det(X + tY)^k = \nabla_X \det^k(X + tY) = k \det^k(X + tY)(X + tY)^{-T}$$

$$\frac{d}{dt} \det(X + tY) = \det(X + tY) \text{tr}((X + tY)^{-1}Y)$$

$$\frac{d^2}{dt^2} \det(X + tY) = \det(X + tY)(\text{tr}^2((X + tY)^{-1}Y) - \text{tr}((X + tY)^{-1}Y(X + tY)^{-1}Y))$$

$$\frac{d}{dt} \det(X + tY)^{-1} = -\det(X + tY)^{-1} \text{tr}((X + tY)^{-1}Y)$$

$$\frac{d^2}{dt^2} \det(X + tY)^{-1} = \det(X + tY)^{-1}(\text{tr}^2((X + tY)^{-1}Y) + \text{tr}((X + tY)^{-1}Y(X + tY)^{-1}Y))$$

$$\frac{d}{dt} \det^\mu(X + tY) = \mu \det^\mu(X + tY) \text{tr}((X + tY)^{-1}Y)$$

D.2.6 logarithmic

Matrix logarithm.

$$\frac{d}{dt} \log(X + tY)^\mu = \mu Y(X + tY)^{-1} = \mu(X + tY)^{-1}Y, \quad XY = YX$$

$$\frac{d}{dt} \log(I - tY)^\mu = -\mu Y(I - tY)^{-1} = -\mu(I - tY)^{-1}Y \quad [181, \text{p.493}]$$

D.2.7 exponential

Matrix exponential. [68, §3.6, §4.5] [294, §5.4]

$$\nabla_X e^{\text{tr}(Y^T X)} = \nabla_X \det e^{Y^T X} = e^{\text{tr}(Y^T X)} Y \quad (\forall X, Y)$$

$$\nabla_X \text{tr} e^{Y X} = e^{Y^T X^T} Y^T = Y^T e^{X^T Y^T}$$

$$\nabla_x \mathbf{1}^T e^{Ax} = A^T e^{Ax}$$

$$\nabla_x \mathbf{1}^T e^{|Ax|} = A^T \delta(\text{sgn}(Ax)) e^{|Ax|} \quad (Ax)_i \neq 0$$

$$\nabla_x \log(\mathbf{1}^T e^x) = \frac{1}{\mathbf{1}^T e^x} e^x$$

$$\nabla_x^2 \log(\mathbf{1}^T e^x) = \frac{1}{\mathbf{1}^T e^x} \left(\delta(e^x) - \frac{1}{\mathbf{1}^T e^x} e^x e^{x^T} \right)$$

$$\nabla_x \prod_{i=1}^k x_i^{\frac{1}{k}} = \frac{1}{k} \left(\prod_{i=1}^k x_i^{\frac{1}{k}} \right) \mathbf{1}/x$$

$$\nabla_x^2 \prod_{i=1}^k x_i^{\frac{1}{k}} = -\frac{1}{k} \left(\prod_{i=1}^k x_i^{\frac{1}{k}} \right) \left(\delta(x)^{-2} - \frac{1}{k} (\mathbf{1}/x)(\mathbf{1}/x)^T \right)$$

$$\frac{d}{dt} e^{tY} = e^{tY} Y = Y e^{tY}$$

$$\frac{d}{dt} e^{X+tY} = e^{X+tY} Y = Y e^{X+tY}, \quad XY = YX$$

$$\frac{d^2}{dt^2} e^{X+tY} = e^{X+tY} Y^2 = Y e^{X+tY} Y = Y^2 e^{X+tY}, \quad XY = YX$$

$$\frac{d^j}{dt^j} e^{\text{tr}(X+tY)} = e^{\text{tr}(X+tY)} \text{tr}^j(Y)$$

Appendix E

Projection

For any $A \in \mathbb{R}^{m \times n}$, the *pseudoinverse* [180, §7.3, prob.9] [220, §6.12, prob.19] [139, §5.5.4] [294, App.A]

$$A^\dagger \triangleq \lim_{t \rightarrow 0^+} (A^T A + tI)^{-1} A^T = \lim_{t \rightarrow 0^+} A^T (A A^T + tI)^{-1} \in \mathbb{R}^{n \times m} \quad (1756)$$

is a unique matrix from the optimal solution set to $\min_X \|AX - I\|_F^2$ (§3.1.8.0.2). For any $t > 0$

$$I - A(A^T A + tI)^{-1} A^T = t(A A^T + tI)^{-1} \quad (1757)$$

Equivalently, pseudoinverse A^\dagger is that unique matrix satisfying the *Moore-Penrose conditions*: [182, §1.3] [333]

1. $AA^\dagger A = A$
2. $A^\dagger AA^\dagger = A^\dagger$
3. $(AA^\dagger)^T = AA^\dagger$
4. $(A^\dagger A)^T = A^\dagger A$

which are necessary and sufficient to establish the pseudoinverse whose principal action is to injectively map $\mathcal{R}(A)$ onto $\mathcal{R}(A^T)$ (Figure 140). Conditions 1 and 3 are necessary and sufficient for AA^\dagger to be the orthogonal projector on $\mathcal{R}(A)$, while conditions 2 and 4 hold iff $A^\dagger A$ is the orthogonal projector on $\mathcal{R}(A^T)$.

Range and nullspace of the pseudoinverse [236] [291, §III.1, exer.1]

$$\mathcal{R}(A^\dagger) = \mathcal{R}(A^T), \quad \mathcal{R}(A^{\dagger T}) = \mathcal{R}(A) \quad (1758)$$

$$\mathcal{N}(A^\dagger) = \mathcal{N}(A^T), \quad \mathcal{N}(A^{\dagger T}) = \mathcal{N}(A) \quad (1759)$$

can be derived by singular value decomposition (§A.6).




Figure 140: (*confer* Figure 15) Pseudoinverse $A^\dagger \in \mathbb{R}^{n \times m}$ action: [294, p.449] Component of vector b in $\mathcal{N}(A^\top)$ maps to origin, while component of b in $\mathcal{R}(A)$ maps to rowspace $\mathcal{R}(A^\top)$. For any $A \in \mathbb{R}^{m \times n}$, inversion is bijective $\forall p \in \mathcal{R}(A)$.

The following relations reliably hold without qualification:

- $A^{\top\dagger} = A^{\dagger\top}$
- $A^{\dagger\dagger} = A$
- $(AA^\top)^\dagger = A^{\dagger\top}A^\dagger$
- $(A^\top A)^\dagger = A^\dagger A^{\dagger\top}$
- $(AA^\dagger)^\dagger = AA^\dagger$
- $(A^\dagger A)^\dagger = A^\dagger A$

Yet for arbitrary A, B it is generally true that $(AB)^\dagger \neq B^\dagger A^\dagger$:

E.0.0.0.1 Theorem. *Pseudoinverse of product.* [148] [49] [213, exer.7.23]
For $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times k}$

$$(AB)^\dagger = B^\dagger A^\dagger \quad (1760)$$

if and only if

$$\mathcal{R}(A^\top AB) \subseteq \mathcal{R}(B) \quad \text{and} \quad \mathcal{R}(BB^\top A^\top) \subseteq \mathcal{R}(A^\top) \quad (1761)$$

◇

$U^T = U^\dagger$ for orthonormal (including the orthogonal) matrices U . So, for orthonormal matrices U, Q and arbitrary A

$$(UAQ^T)^\dagger = QA^\dagger U^T \quad (1762)$$

E.0.0.0.2 Exercise. *Kronecker inverse.*

Prove:

$$(A \otimes B)^\dagger = A^\dagger \otimes B^\dagger \quad (1763)$$

▼

E.0.1 Logical deductions

When A is invertible, $A^\dagger = A^{-1}$; so $A^\dagger A = AA^\dagger = I$. Otherwise, for $A \in \mathbb{R}^{m \times n}$ [132, §5.3.3.1] [213, §7] [261]

- g. $A^\dagger A = I$, $A^\dagger = (A^T A)^{-1} A^T$, $\text{rank } A = n$
- h. $AA^\dagger = I$, $A^\dagger = A^T (AA^T)^{-1}$, $\text{rank } A = m$
- i. $A^\dagger A \omega = \omega$, $\omega \in \mathcal{R}(A^T)$
- j. $AA^\dagger v = v$, $v \in \mathcal{R}(A)$
- k. $A^\dagger A = AA^\dagger$, A normal
- l. $A^{k\dagger} = A^{\dagger k}$, A normal, k an integer

Equivalent to the corresponding Moore-Penrose condition:

- 1. $A^T = A^T A A^\dagger$ **or** $A^T = A^\dagger A A^T$
- 2. $A^{\dagger T} = A^{\dagger T} A^\dagger A$ **or** $A^{\dagger T} = A A^\dagger A^{\dagger T}$

When A is symmetric, A^\dagger is symmetric and (§A.6)

$$A \succeq 0 \Leftrightarrow A^\dagger \succeq 0 \quad (1764)$$

E.0.1.0.1 Example. *Solution to classical linear equation $Ax = b$.*

In §2.5.1.1, the solution set to matrix equation $Ax = b$ was represented as an intersection of hyperplanes. Regardless of rank of A or its shape (fat or skinny), interpretation as a hyperplane intersection describing a possibly empty affine set generally holds. If matrix A is rank deficient or fat, there is an infinity of solutions x when $b \in \mathcal{R}(A)$. A unique solution occurs when the hyperplanes intersect at a single point.

For any shape of matrix A of any rank, and given any vector b that may or may not be in $\mathcal{R}(A)$, we wish to find a best Euclidean solution x^* to

$$Ax = b \quad (1765)$$

(more generally, $Ax \approx b$ given arbitrary matrices) by solving

$$\underset{x}{\text{minimize}} \quad \|Ax - b\|^2 \quad (1766)$$

Necessary and sufficient condition for optimal solution to this unconstrained optimization is the so-called *normal equation* that results from zeroing the convex objective's gradient: (§D.2.1)

$$A^T A x = A^T b \quad (1767)$$

normal because error vector $b - Ax$ is perpendicular to $\mathcal{R}(A)$; *id est*, $A^T(b - Ax) = \mathbf{0}$. Given any matrix A and any vector b , the normal equation is solvable exactly; always so, because $\mathcal{R}(A^T A) = \mathcal{R}(A^T)$ and $A^T b \in \mathcal{R}(A^T)$.

When A is skinny-or-square full-rank, normal equation (1767) can be solved exactly by inversion:

$$x^* = (A^T A)^{-1} A^T b \equiv A^\dagger b \quad (1768)$$

For matrix A of arbitrary rank and shape, on the other hand, $A^T A$ might not be invertible. Yet the normal equation can always be solved exactly by: (1756)

$$x^* = \lim_{t \rightarrow 0^+} (A^T A + tI)^{-1} A^T b = A^\dagger b \quad (1769)$$

invertible for any positive value of t by (1364). The exact inversion (1768) and this pseudoinverse solution (1769) each solve

$$\lim_{t \rightarrow 0^+} \underset{x}{\text{minimize}} \quad \|Ax - b\|^2 + t \|x\|^2 \quad (1770)$$

simultaneously providing least squares solution to (1766) and the classical *least norm* solution^{E.1} [294, App.A.4] (*confer* §3.1.3.0.1)

$$\begin{aligned} & \arg \underset{x}{\text{minimize}} \quad \|x\|^2 \\ & \text{subject to} \quad Ax = AA^\dagger b \end{aligned} \quad (1771)$$

^{E.1}This means: optimal solutions of lesser norm than the so-called *least norm* solution (1771) can be obtained (at expense of approximation $Ax \approx b$ hence, of perpendicularity) by ignoring the limiting operation and introducing finite positive values of t into (1770).

where $AA^\dagger b$ is the orthogonal projection of vector b on $\mathcal{R}(A)$. Least norm solution can be interpreted as orthogonal projection of the origin $\mathbf{0}$ on affine subset $\mathcal{A} = \{x \mid Ax = AA^\dagger b\}$; (§E.5.0.0.5, §E.5.0.0.6)

$$\begin{aligned} \arg \underset{x}{\text{minimize}} \quad & \|x - \mathbf{0}\|^2 \\ \text{subject to} \quad & x \in \mathcal{A} \end{aligned} \tag{1772}$$

equivalently, maximization of the Euclidean ball until it kisses \mathcal{A} ; rather, $\text{dist}(\mathbf{0}, \mathcal{A})$. \square

E.1 Idempotent matrices

Projection matrices are square and defined by *idempotence*, $P^2 = P$; [294, §2.6] [182, §1.3] equivalent to the condition, P be diagonalizable [180, §3.3, prob.3] with eigenvalues $\phi_i \in \{0, 1\}$. [352, §4.1, thm.4.1] Idempotent matrices are not necessarily symmetric. The transpose of an idempotent matrix remains idempotent; $P^T P^T = P^T$. Solely excepting $P = I$, all projection matrices are neither orthogonal (§B.5) or invertible. [294, §3.4] The collection of all projection matrices of particular dimension does not form a convex set.

Suppose we wish to project nonorthogonally (*obliquely*) on the range of any particular matrix $A \in \mathbb{R}^{m \times n}$. All idempotent matrices projecting nonorthogonally on $\mathcal{R}(A)$ may be expressed:

$$P = A(A^\dagger + BZ^T) \in \mathbb{R}^{m \times m} \tag{1773}$$

where $\mathcal{R}(P) = \mathcal{R}(A)$, ^{E.2} $B \in \mathbb{R}^{n \times k}$ for $k \in \{1 \dots m\}$ is otherwise arbitrary, and $Z \in \mathbb{R}^{m \times k}$ is any matrix whose range is in $\mathcal{N}(A^T)$; *id est*,

$$A^T Z = A^\dagger Z = \mathbf{0} \tag{1774}$$

Evidently, the collection of nonorthogonal projectors projecting on $\mathcal{R}(A)$ is an affine subset

$$\mathcal{P}_k = \{A(A^\dagger + BZ^T) \mid B \in \mathbb{R}^{n \times k}\} \tag{1775}$$

^{E.2}**Proof.** $\mathcal{R}(P) \subseteq \mathcal{R}(A)$ is obvious [294, §3.6]. By (132) and (133),

$$\begin{aligned} \mathcal{R}(A^\dagger + BZ^T) &= \{(A^\dagger + BZ^T)y \mid y \in \mathbb{R}^m\} \\ &\supseteq \{(A^\dagger + BZ^T)y \mid y \in \mathcal{R}(A)\} = \mathcal{R}(A^T) \\ \mathcal{R}(P) &= \{A(A^\dagger + BZ^T)y \mid y \in \mathbb{R}^m\} \\ &\supseteq \{A(A^\dagger + BZ^T)y \mid (A^\dagger + BZ^T)y \in \mathcal{R}(A^T)\} = \mathcal{R}(A) \quad \blacklozenge \end{aligned}$$

When matrix A in (1773) is skinny full-rank ($A^\dagger A = I$) or has orthonormal columns ($A^T A = I$), either property leads to a biorthogonal characterization of nonorthogonal projection:

E.1.1 Biorthogonal characterization

Any nonorthogonal projector $P^2 = P \in \mathbb{R}^{m \times m}$ projecting on nontrivial $\mathcal{R}(U)$ can be defined by a biorthogonality condition $Q^T U = I$; the *biorthogonal decomposition* of P being (*confer* (1773))

$$P = UQ^T, \quad Q^T U = I \quad (1776)$$

where^{E.3} (§B.1.1.1)

$$\mathcal{R}(P) = \mathcal{R}(U), \quad \mathcal{N}(P) = \mathcal{N}(Q^T) \quad (1777)$$

and where generally (*confer* (1802))^{E.4}

$$P^T \neq P, \quad P^\dagger \neq P, \quad \|P\|_2 \neq 1, \quad P \not\leq 0 \quad (1778)$$

and P is not nonexpansive (1803).

(\Leftarrow) To verify assertion (1776) we observe: because idempotent matrices are diagonalizable (§A.5), [180, §3.3, prob.3] they must have the form (1449)

$$P = S\Phi S^{-1} = \sum_{i=1}^m \phi_i s_i w_i^T = \sum_{i=1}^{k \leq m} s_i w_i^T \quad (1779)$$

that is a sum of $k = \text{rank } P$ independent projector dyads (idempotent dyads, §B.1.1, §E.6.2.1) where $\phi_i \in \{0, 1\}$ are the eigenvalues of P [352, §4.1, thm.4.1] in diagonal matrix $\Phi \in \mathbb{R}^{m \times m}$ arranged in nonincreasing

^{E.3}**Proof.** Obviously, $\mathcal{R}(P) \subseteq \mathcal{R}(U)$. Because $Q^T U = I$

$$\begin{aligned} \mathcal{R}(P) &= \{UQ^T x \mid x \in \mathbb{R}^m\} \\ &\supseteq \{UQ^T U y \mid y \in \mathbb{R}^k\} = \mathcal{R}(U) \end{aligned} \quad \blacklozenge$$

^{E.4}Orthonormal decomposition (1799) (*confer* §E.3.4) is a special case of biorthogonal decomposition (1776) characterized by (1802). So, these characteristics (1778) are not necessary conditions for biorthogonality.

order, and where $s_i, w_i \in \mathbb{R}^m$ are the right- and left-eigenvectors of P , respectively, which are independent and real.^{E.5} Therefore

$$U \triangleq S(:, 1:k) = [s_1 \cdots s_k] \in \mathbb{R}^{m \times k} \quad (1780)$$

is the full-rank matrix $S \in \mathbb{R}^{m \times m}$ having $m - k$ columns truncated (corresponding to 0 eigenvalues), while

$$Q^T \triangleq S^{-1}(1:k, :) = \begin{bmatrix} w_1^T \\ \vdots \\ w_k^T \end{bmatrix} \in \mathbb{R}^{k \times m} \quad (1781)$$

is matrix S^{-1} having the corresponding $m - k$ rows truncated. By the 0 eigenvalues theorem (§A.7.3.0.1), $\mathcal{R}(U) = \mathcal{R}(P)$, $\mathcal{R}(Q) = \mathcal{R}(P^T)$, and

$$\begin{aligned} \mathcal{R}(P) &= \text{span} \{s_i \mid \phi_i = 1 \ \forall i\} \\ \mathcal{N}(P) &= \text{span} \{s_i \mid \phi_i = 0 \ \forall i\} \\ \mathcal{R}(P^T) &= \text{span} \{w_i \mid \phi_i = 1 \ \forall i\} \\ \mathcal{N}(P^T) &= \text{span} \{w_i \mid \phi_i = 0 \ \forall i\} \end{aligned} \quad (1782)$$

Thus biorthogonality $Q^T U = I$ is a necessary condition for idempotence, and so the collection of nonorthogonal projectors projecting on $\mathcal{R}(U)$ is the affine subset $\mathcal{P}_k = U \mathcal{Q}_k^T$ where $\mathcal{Q}_k = \{Q \mid Q^T U = I, Q \in \mathbb{R}^{m \times k}\}$.

(\Rightarrow) Biorthogonality is a sufficient condition for idempotence;

$$P^2 = \sum_{i=1}^k s_i w_i^T \sum_{j=1}^k s_j w_j^T = P \quad (1783)$$

id est, if the cross-products are annihilated, then $P^2 = P$. ◆

Nonorthogonal projection of x on $\mathcal{R}(P)$ has expression like a biorthogonal expansion,

$$Px = U Q^T x = \sum_{i=1}^k w_i^T x s_i \quad (1784)$$

^{E.5}Eigenvectors of a real matrix corresponding to real eigenvalues must be real. (§A.5.0.0.1)




Figure 141: Nonorthogonal projection of $x \in \mathbb{R}^3$ on $\mathcal{R}(U) = \mathbb{R}^2$ under biorthogonality condition; *id est*, $Px = UQ^T x$ such that $Q^T U = I$. Any point along imaginary line \mathcal{T} connecting x to Px will be projected nonorthogonally on Px with respect to horizontal plane constituting $\mathbb{R}^2 = \text{aff cone}(U)$ in this example. Extreme directions of $\text{cone}(U)$ correspond to two columns of U ; likewise for $\text{cone}(Q)$. For purpose of illustration, we truncate each conic hull by truncating coefficients of conic combination at unity. Conic hull $\text{cone}(Q)$ is headed upward at an angle, out of plane of page. Nonorthogonal projection would fail were $\mathcal{N}(Q^T)$ in $\mathcal{R}(U)$ (were \mathcal{T} parallel to a line in $\mathcal{R}(U)$).

When the domain is restricted to range of P , say $x = U\xi$ for $\xi \in \mathbb{R}^k$, then $x = Px = UQ^T U\xi = U\xi$ and expansion is unique because the eigenvectors are linearly independent. Otherwise, any component of x in $\mathcal{N}(P) = \mathcal{N}(Q^T)$ will be annihilated. Direction of nonorthogonal projection is orthogonal to $\mathcal{R}(Q) \Leftrightarrow Q^T U = I$; *id est*, for $Px \in \mathcal{R}(U)$

$$Px - x \perp \mathcal{R}(Q) \text{ in } \mathbb{R}^m \quad (1785)$$

E.1.1.0.1 Example. *Illustration of nonorthogonal projector.*

Figure 141 shows $\text{cone}(U)$, the conic hull of the columns of

$$U = \begin{bmatrix} 1 & 1 \\ -0.5 & 0.3 \\ 0 & 0 \end{bmatrix} \quad (1786)$$

from nonorthogonal projector $P = UQ^T$. Matrix U has a limitless number of left inverses because $\mathcal{N}(U^T)$ is nontrivial. Similarly depicted is left inverse Q^T from (1773)

$$\begin{aligned} Q = U^{\dagger T} + ZB^T &= \begin{bmatrix} 0.3750 & 0.6250 \\ -1.2500 & 1.2500 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} [0.5 \ 0.5] \\ &= \begin{bmatrix} 0.3750 & 0.6250 \\ -1.2500 & 1.2500 \\ 0.5000 & 0.5000 \end{bmatrix} \end{aligned} \quad (1787)$$

where $Z \in \mathcal{N}(U^T)$ and matrix B is selected arbitrarily; *id est*, $Q^T U = I$ because U is full-rank.

Direction of projection on $\mathcal{R}(U)$ is orthogonal to $\mathcal{R}(Q)$. Any point along line \mathcal{T} in the figure, for example, will have the same projection. Were matrix Z instead equal to $\mathbf{0}$, then $\text{cone}(Q)$ would become the relative dual to $\text{cone}(U)$ (sharing the same affine hull; §2.13.8, *confer* Figure 52(a)). In that case, projection $Px = UU^\dagger x$ of x on $\mathcal{R}(U)$ becomes orthogonal projection (and unique minimum-distance). \square

E.1.2 Idempotence summary

Nonorthogonal subspace-projector P is a linear operator defined by idempotence or biorthogonal decomposition (1776), but characterized not by symmetry nor positive semidefiniteness nor nonexpansivity (1803).

E.2 $I - P$, Projection on algebraic complement

It follows from the diagonalizability of idempotent matrices that $I - P$ must also be a projection matrix because it too is idempotent, and because it may be expressed

$$I - P = S(I - \Phi)S^{-1} = \sum_{i=1}^m (1 - \phi_i) s_i w_i^T \quad (1788)$$

where $(1 - \phi_i) \in \{1, 0\}$ are the eigenvalues of $I - P$ (1365) whose eigenvectors s_i, w_i are identical to those of P in (1779). A consequence of that complementary relationship of eigenvalues is the fact, [305, §2] [300, §2] for subspace projector $P = P^2 \in \mathbb{R}^{m \times m}$

$$\begin{aligned} \mathcal{R}(P) &= \text{span} \{s_i \mid \phi_i = 1 \ \forall i\} = \text{span} \{s_i \mid (1 - \phi_i) = 0 \ \forall i\} = \mathcal{N}(I - P) \\ \mathcal{N}(P) &= \text{span} \{s_i \mid \phi_i = 0 \ \forall i\} = \text{span} \{s_i \mid (1 - \phi_i) = 1 \ \forall i\} = \mathcal{R}(I - P) \\ \mathcal{R}(P^T) &= \text{span} \{w_i \mid \phi_i = 1 \ \forall i\} = \text{span} \{w_i \mid (1 - \phi_i) = 0 \ \forall i\} = \mathcal{N}(I - P^T) \\ \mathcal{N}(P^T) &= \text{span} \{w_i \mid \phi_i = 0 \ \forall i\} = \text{span} \{w_i \mid (1 - \phi_i) = 1 \ \forall i\} = \mathcal{R}(I - P^T) \end{aligned} \quad (1789)$$

that is easy to see from (1779) and (1788). Idempotent $I - P$ therefore projects vectors on its range, $\mathcal{N}(P)$. Because all eigenvectors of a real idempotent matrix are real and independent, the algebraic complement of $\mathcal{R}(P)$ [201, §3.3] is equivalent to $\mathcal{N}(P)$; **E.6** *id est*,

$$\mathcal{R}(P) \oplus \mathcal{N}(P) = \mathcal{R}(P^T) \oplus \mathcal{N}(P^T) = \mathcal{R}(P^T) \oplus \mathcal{N}(P) = \mathcal{R}(P) \oplus \mathcal{N}(P^T) = \mathbb{R}^m \quad (1790)$$

because $\mathcal{R}(P) \oplus \mathcal{R}(I - P) = \mathbb{R}^m$. For idempotent $P \in \mathbb{R}^{m \times m}$, consequently,

$$\text{rank } P + \text{rank}(I - P) = m \quad (1791)$$

E.2.0.0.1 Theorem. *Rank/Trace.* [352, §4.1, prob.9] (*confer* (1807))

$$\begin{aligned} P^2 &= P \\ &\Leftrightarrow \\ \text{rank } P &= \text{tr } P \quad \mathbf{and} \quad \text{rank}(I - P) = \text{tr}(I - P) \end{aligned} \quad (1792)$$

◇

E.6 The same phenomenon occurs with symmetric (nonidempotent) matrices, for example. When the summands in $A \oplus B = \mathbb{R}^m$ are orthogonal vector spaces, the algebraic complement is the orthogonal complement.

E.2.1 Universal projector characteristic

Although projection is not necessarily orthogonal and $\mathcal{R}(P) \not\perp \mathcal{R}(I - P)$ in general, still for any projector P and any $x \in \mathbb{R}^m$

$$Px + (I - P)x = x \quad (1793)$$

must hold where $\mathcal{R}(I - P) = \mathcal{N}(P)$ is the algebraic complement of $\mathcal{R}(P)$. The algebraic complement of closed convex cone \mathcal{K} , for example, is the negative dual cone $-\mathcal{K}^*$. (1911)

E.3 Symmetric idempotent matrices

When idempotent matrix P is symmetric, P is an orthogonal projector. In other words, the direction of projection of point $x \in \mathbb{R}^m$ on subspace $\mathcal{R}(P)$ is orthogonal to $\mathcal{R}(P)$; *id est*, for $P^2 = P \in \mathbb{S}^m$ and projection $Px \in \mathcal{R}(P)$

$$Px - x \perp \mathcal{R}(P) \text{ in } \mathbb{R}^m \quad (1794)$$

Perpendicularity is a necessary and sufficient condition for orthogonal projection on a subspace. [92, §4.9]

A condition equivalent to (1794) is: Norm of direction $x - Px$ is the infimum over all nonorthogonal projections of x on $\mathcal{R}(P)$; [220, §3.3] for $P^2 = P \in \mathbb{S}^m$, $\mathcal{R}(P) = \mathcal{R}(A)$, matrices A, B, Z and positive integer k as defined for (1773), and given $x \in \mathbb{R}^m$

$$\|x - Px\|_2 = \|x - AA^\dagger x\|_2 = \inf_{B \in \mathbb{R}^{n \times k}} \|x - A(A^\dagger + BZ^\top)x\|_2 \quad (1795)$$

The infimum is attained for $\mathcal{R}(B) \subseteq \mathcal{N}(A)$ over any affine subset of nonorthogonal projectors (1775) indexed by k .

Proof is straightforward: The vector 2-norm is a convex function. Setting gradient of the norm-square to $\mathbf{0}$, applying §D.2,

$$\begin{aligned} (A^\top ABZ^\top - A^\top(I - AA^\dagger))xx^\top A &= \mathbf{0} \\ \Leftrightarrow \\ A^\top ABZ^\top xx^\top A &= \mathbf{0} \end{aligned} \quad (1796)$$

because $A^\top = A^\top AA^\dagger$. Projector $P = AA^\dagger$ is therefore unique; the minimum-distance projector is the orthogonal projector, and *vice versa*.

◆

We get $P = AA^\dagger$ so this projection matrix must be symmetric. Then for any matrix $A \in \mathbb{R}^{m \times n}$, symmetric idempotent P projects a given vector x in \mathbb{R}^m orthogonally on $\mathcal{R}(A)$. Under either condition (1794) or (1795), the projection Px is unique minimum-distance; for subspaces, perpendicularity and minimum-distance conditions are equivalent.

E.3.1 Four subspaces

We summarize the orthogonal projectors projecting on the four fundamental subspaces: for $A \in \mathbb{R}^{m \times n}$

$$\begin{aligned} A^\dagger A &: \mathbb{R}^n \text{ on } \mathcal{R}(A^\dagger A) &= \mathcal{R}(A^T) \\ AA^\dagger &: \mathbb{R}^m \text{ on } \mathcal{R}(AA^\dagger) &= \mathcal{R}(A) \\ I - A^\dagger A &: \mathbb{R}^n \text{ on } \mathcal{R}(I - A^\dagger A) &= \mathcal{N}(A) \\ I - AA^\dagger &: \mathbb{R}^m \text{ on } \mathcal{R}(I - AA^\dagger) &= \mathcal{N}(A^T) \end{aligned} \quad (1797)$$

For completeness:^{E.7} (1789)

$$\begin{aligned} \mathcal{N}(A^\dagger A) &= \mathcal{N}(A) \\ \mathcal{N}(AA^\dagger) &= \mathcal{N}(A^T) \\ \mathcal{N}(I - A^\dagger A) &= \mathcal{R}(A^T) \\ \mathcal{N}(I - AA^\dagger) &= \mathcal{R}(A) \end{aligned} \quad (1798)$$

E.3.2 Orthogonal characterization

Any symmetric projector $P^2 = P \in \mathbb{S}^m$ projecting on nontrivial $\mathcal{R}(Q)$ can be defined by the orthonormality condition $Q^T Q = I$. When skinny matrix $Q \in \mathbb{R}^{m \times k}$ has orthonormal columns, then $Q^\dagger = Q^T$ by the Moore-Penrose conditions. Hence, any P having an *orthonormal decomposition* (§E.3.4)

$$P = QQ^T, \quad Q^T Q = I \quad (1799)$$

where [294, §3.3] (1505)

$$\mathcal{R}(P) = \mathcal{R}(Q), \quad \mathcal{N}(P) = \mathcal{N}(Q^T) \quad (1800)$$

^{E.7}**Proof** is by singular value decomposition (§A.6.2): $\mathcal{N}(A^\dagger A) \subseteq \mathcal{N}(A)$ is obvious. Conversely, suppose $A^\dagger Ax = \mathbf{0}$. Then $x^T A^\dagger Ax = x^T Q Q^T x = \|Q^T x\|^2 = 0$ where $A = U \Sigma Q^T$ is the subcompact singular value decomposition. Because $\mathcal{R}(Q) = \mathcal{R}(A^T)$, then $x \in \mathcal{N}(A)$ which implies $\mathcal{N}(A^\dagger A) \supseteq \mathcal{N}(A)$. $\therefore \mathcal{N}(A^\dagger A) = \mathcal{N}(A)$. \blacklozenge

is an orthogonal projector projecting on $\mathcal{R}(Q)$ having, for $Px \in \mathcal{R}(Q)$ (confer (1785))

$$Px - x \perp \mathcal{R}(Q) \text{ in } \mathbb{R}^m \tag{1801}$$

From (1799), orthogonal projector P is obviously positive semidefinite (§A.3.1.0.6); necessarily,

$$P^T = P, \quad P^\dagger = P, \quad \|P\|_2 = 1, \quad P \succeq 0 \tag{1802}$$

and $\|Px\| = \|QQ^T x\| = \|Q^T x\|$ because $\|Qy\| = \|y\| \forall y \in \mathbb{R}^k$. All orthogonal projectors are therefore *nonexpansive* because

$$\sqrt{\langle Px, x \rangle} = \|Px\| = \|Q^T x\| \leq \|x\| \quad \forall x \in \mathbb{R}^m \tag{1803}$$

the Bessel inequality, [92] [201] with equality when $x \in \mathcal{R}(Q)$.

From the diagonalization of idempotent matrices (1779) on page 648

$$P = S\Phi S^T = \sum_{i=1}^m \phi_i s_i s_i^T = \sum_{i=1}^{k \leq m} s_i s_i^T \tag{1804}$$

orthogonal projection of point x on $\mathcal{R}(P)$ has expression like an orthogonal expansion [92, §4.10]

$$Px = QQ^T x = \sum_{i=1}^k s_i^T x s_i \tag{1805}$$

where

$$Q = S(:, 1:k) = [s_1 \cdots s_k] \in \mathbb{R}^{m \times k} \tag{1806}$$

and where the s_i [sic] are orthonormal eigenvectors of symmetric idempotent P . When the domain is restricted to range of P , say $x = Q\xi$ for $\xi \in \mathbb{R}^k$, then $x = Px = QQ^T Q\xi = Q\xi$ and expansion is unique. Otherwise, any component of x in $\mathcal{N}(Q^T)$ will be annihilated.

E.3.2.0.1 Theorem. *Symmetric rank/trace.* (confer (1792) (1369))

$$\begin{aligned} P^T = P, \quad P^2 = P \\ \Leftrightarrow \\ \text{rank } P = \text{tr } P = \|P\|_F^2 \quad \text{and} \quad \text{rank}(I - P) = \text{tr}(I - P) = \|I - P\|_F^2 \end{aligned} \tag{1807}$$

◇

Proof. We take as given Theorem E.2.0.0.1 establishing idempotence. We have left only to show $\text{tr } P = \|P\|_F^2 \Rightarrow P^T = P$, established in [352, §7.1].

◆

E.3.3 Summary, symmetric idempotent

In summary, orthogonal projector P is a linear operator defined [177, §A.3.1] by idempotence and symmetry, and characterized by positive semidefiniteness and nonexpansivity. The algebraic complement (§E.2) to $\mathcal{R}(P)$ becomes the orthogonal complement $\mathcal{R}(I - P)$; *id est*, $\mathcal{R}(P) \perp \mathcal{R}(I - P)$.

E.3.4 Orthonormal decomposition

When $Z = \mathbf{0}$ in the general nonorthogonal projector $A(A^\dagger + BZ^T)$ (1773), an orthogonal projector results (for any matrix A) characterized principally by idempotence and symmetry. Any real orthogonal projector may, in fact, be represented by an orthonormal decomposition such as (1799). [182, §1, prob.42]

To verify that assertion for the four fundamental subspaces (1797), we need only to express A by subcompact singular value decomposition (§A.6.2): From pseudoinverse (1475) of $A = U\Sigma Q^T \in \mathbb{R}^{m \times n}$

$$\begin{aligned} AA^\dagger &= U\Sigma\Sigma^\dagger U^T = UU^T, & A^\dagger A &= Q\Sigma^\dagger\Sigma Q^T = QQ^T \\ I - AA^\dagger &= I - UU^T = U^\perp U^{\perp T}, & I - A^\dagger A &= I - QQ^T = Q^\perp Q^{\perp T} \end{aligned} \quad (1808)$$

where $U^\perp \in \mathbb{R}^{m \times m - \text{rank } A}$ holds columnar an orthonormal basis for the orthogonal complement of $\mathcal{R}(U)$, and likewise for $Q^\perp \in \mathbb{R}^{n \times n - \text{rank } A}$. Existence of an orthonormal decomposition is sufficient to establish idempotence and symmetry of an orthogonal projector (1799). ◆

E.3.5 Unifying trait of all projectors: direction

Relation (1808) shows: orthogonal projectors simultaneously possess a biorthogonal decomposition (*confer* §E.1.1) (for example, AA^\dagger for skinny-or-square A full-rank) and an orthonormal decomposition (UU^T whence $Px = UU^T x$).

E.3.5.1 orthogonal projector, orthonormal decomposition

Consider orthogonal expansion of $x \in \mathcal{R}(U)$:

$$x = UU^T x = \sum_{i=1}^n u_i u_i^T x \quad (1809)$$

a sum of one-dimensional orthogonal projections (§E.6.3), where

$$U \triangleq [u_1 \cdots u_n] \quad \text{and} \quad U^T U = I \quad (1810)$$

and where the subspace projector has two expressions, (1808)

$$AA^\dagger \triangleq UU^T \quad (1811)$$

where $A \in \mathbb{R}^{m \times n}$ has rank n . The direction of projection of x on u_j for some $j \in \{1 \dots n\}$, for example, is orthogonal to u_j but parallel to a vector in the span of all the remaining vectors constituting the columns of U ;

$$\begin{aligned} u_j^T (u_j u_j^T x - x) &= 0 \\ u_j u_j^T x - x &= u_j u_j^T x - UU^T x \in \mathcal{R}(\{u_i \mid i=1 \dots n, i \neq j\}) \end{aligned} \quad (1812)$$

E.3.5.2 orthogonal projector, biorthogonal decomposition

We get a similar result for the biorthogonal expansion of $x \in \mathcal{R}(A)$. Define

$$A \triangleq [a_1 \ a_2 \ \cdots \ a_n] \in \mathbb{R}^{m \times n} \quad (1813)$$

and the rows of the pseudoinverse^{E.8}

$$A^\dagger \triangleq \begin{bmatrix} a_1^{*T} \\ a_2^{*T} \\ \vdots \\ a_n^{*T} \end{bmatrix} \in \mathbb{R}^{n \times m} \quad (1814)$$

under biorthogonality condition $A^\dagger A = I$. In biorthogonal expansion (§2.13.8)

$$x = AA^\dagger x = \sum_{i=1}^n a_i a_i^{*T} x \quad (1815)$$

^{E.8}Notation * in this context connotes extreme direction of a dual cone; e.g., (367) or Example E.5.0.0.3.

the direction of projection of x on a_j for some particular $j \in \{1 \dots n\}$, for example, is orthogonal to a_j^* and parallel to a vector in the span of all the remaining vectors constituting the columns of A ;

$$\begin{aligned} a_j^{*\text{T}}(a_j a_j^{*\text{T}} x - x) &= 0 \\ a_j a_j^{*\text{T}} x - x &= a_j a_j^{*\text{T}} x - AA^\dagger x \in \mathcal{R}(\{a_i \mid i=1 \dots n, i \neq j\}) \end{aligned} \quad (1816)$$

E.3.5.3 nonorthogonal projector, biorthogonal decomposition

Because the result in §E.3.5.2 is independent of matrix symmetry $AA^\dagger = (AA^\dagger)^\text{T}$, we must get the same result for any nonorthogonal projector characterized by a biorthogonality condition; namely, for nonorthogonal projector $P = UQ^\text{T}$ (1776) under biorthogonality condition $Q^\text{T}U = I$, in the biorthogonal expansion of $x \in \mathcal{R}(U)$

$$x = UQ^\text{T}x = \sum_{i=1}^k u_i q_i^\text{T} x \quad (1817)$$

where

$$\begin{aligned} U &\triangleq [u_1 \dots u_k] \in \mathbb{R}^{m \times k} \\ Q^\text{T} &\triangleq \begin{bmatrix} q_1^\text{T} \\ \vdots \\ q_k^\text{T} \end{bmatrix} \in \mathbb{R}^{k \times m} \end{aligned} \quad (1818)$$

the direction of projection of x on u_j is orthogonal to q_j and parallel to a vector in the span of the remaining u_i :

$$\begin{aligned} q_j^\text{T}(u_j q_j^\text{T} x - x) &= 0 \\ u_j q_j^\text{T} x - x &= u_j q_j^\text{T} x - UQ^\text{T}x \in \mathcal{R}(\{u_i \mid i=1 \dots k, i \neq j\}) \end{aligned} \quad (1819)$$

E.4 Algebra of projection on affine subsets

Let $P_{\mathcal{A}}x$ denote projection of x on affine subset $\mathcal{A} \triangleq \mathcal{R} + \alpha$ where \mathcal{R} is a subspace and $\alpha \in \mathcal{A}$. Then, because \mathcal{R} is parallel to \mathcal{A} , it holds:

$$\begin{aligned} P_{\mathcal{A}}x &= P_{\mathcal{R}+\alpha}x = (I - P_{\mathcal{R}})(\alpha) + P_{\mathcal{R}}x \\ &= P_{\mathcal{R}}(x - \alpha) + \alpha \end{aligned} \quad (1820)$$

Subspace projector $P_{\mathcal{R}}$ is a linear operator ($P_{\mathcal{A}}$ is not), and $P_{\mathcal{R}}(x + y) = P_{\mathcal{R}}x$ whenever $y \perp \mathcal{R}$ and $P_{\mathcal{R}}$ is an orthogonal projector.

E.4.0.0.1 Theorem. *Orthogonal projection on affine subset.* [92, §9.26] Let $\mathcal{A} = \mathcal{R} + \alpha$ be an affine subset where $\alpha \in \mathcal{A}$, and let \mathcal{R}^\perp be the orthogonal complement of subspace \mathcal{R} . Then $P_{\mathcal{A}}x$ is the orthogonal projection of $x \in \mathbb{R}^n$ on \mathcal{A} if and only if

$$P_{\mathcal{A}}x \in \mathcal{A}, \quad \langle P_{\mathcal{A}}x - x, a - \alpha \rangle = 0 \quad \forall a \in \mathcal{A} \quad (1821)$$

or if and only if

$$P_{\mathcal{A}}x \in \mathcal{A}, \quad P_{\mathcal{A}}x - x \in \mathcal{R}^\perp \quad (1822)$$

◇

E.5 Projection examples

E.5.0.0.1 Example. *Orthogonal projection on orthogonal basis.*

Orthogonal projection on a subspace can instead be accomplished by orthogonally projecting on the individual members of an orthogonal basis for that subspace. Suppose, for example, matrix $A \in \mathbb{R}^{m \times n}$ holds an orthonormal basis for $\mathcal{R}(A)$ in its columns; $A \triangleq [a_1 \ a_2 \ \cdots \ a_n]$. Then orthogonal projection of vector $x \in \mathbb{R}^n$ on $\mathcal{R}(A)$ is a sum of one-dimensional orthogonal projections

$$Px = AA^\dagger x = A(A^\top A)^{-1}A^\top x = AA^\top x = \sum_{i=1}^n a_i a_i^\top x \quad (1823)$$

where each symmetric dyad $a_i a_i^\top$ is an orthogonal projector projecting on $\mathcal{R}(a_i)$. (§E.6.3) Because $\|x - Px\|$ is minimized by orthogonal projection, Px is considered to be the best approximation (in the Euclidean sense) to x from the set $\mathcal{R}(A)$. [92, §4.9] □

E.5.0.0.2 Example. *Orthogonal projection on span of nonorthogonal basis.*

Orthogonal projection on a subspace can also be accomplished by projecting nonorthogonally on the individual members of any nonorthogonal basis for that subspace. This interpretation is in fact the principal application of the pseudoinverse we discussed. Now suppose matrix A holds a nonorthogonal basis for $\mathcal{R}(A)$ in its columns,

$$A = [a_1 \ a_2 \ \cdots \ a_n] \in \mathbb{R}^{m \times n} \quad (1813)$$

and define the rows $a_i^{*\text{T}}$ of its pseudoinverse A^\dagger as in (1814). Then orthogonal projection of vector $x \in \mathbb{R}^n$ on $\mathcal{R}(A)$ is a sum of one-dimensional nonorthogonal projections

$$Px = AA^\dagger x = \sum_{i=1}^n a_i a_i^{*\text{T}} x \quad (1824)$$

where each nonsymmetric dyad $a_i a_i^{*\text{T}}$ is a nonorthogonal projector projecting on $\mathcal{R}(a_i)$, (§E.6.1) idempotent because of biorthogonality condition $A^\dagger A = I$.

The projection Px is regarded as the best approximation to x from the set $\mathcal{R}(A)$, as it was in Example E.5.0.0.1. \square

E.5.0.0.3 Example. *Biorthogonal expansion as nonorthogonal projection.* Biorthogonal expansion can be viewed as a sum of components, each a nonorthogonal projection on the range of an extreme direction of a pointed polyhedral cone \mathcal{K} ; e.g., Figure 142.

Suppose matrix $A \in \mathbb{R}^{m \times n}$ holds a nonorthogonal basis for $\mathcal{R}(A)$ in its columns as in (1813), and the rows of pseudoinverse A^\dagger are defined as in (1814). Assuming the most general biorthogonality condition $(A^\dagger + BZ^\text{T})A = I$ with BZ^T defined as for (1773), then biorthogonal expansion of vector x is a sum of one-dimensional nonorthogonal projections; for $x \in \mathcal{R}(A)$

$$x = A(A^\dagger + BZ^\text{T})x = AA^\dagger x = \sum_{i=1}^n a_i a_i^{*\text{T}} x \quad (1825)$$

where each dyad $a_i a_i^{*\text{T}}$ is a nonorthogonal projector projecting on $\mathcal{R}(a_i)$. (§E.6.1) The extreme directions of $\mathcal{K} = \text{cone}(A)$ are $\{a_1, \dots, a_n\}$ the linearly independent columns of A , while the extreme directions $\{a_1^*, \dots, a_n^*\}$ of relative dual cone $\mathcal{K}^* \cap \text{aff } \mathcal{K} = \text{cone}(A^{\dagger\text{T}})$ (§2.13.9.4) correspond to the linearly independent (§B.1.1.1) rows of A^\dagger . Directions of nonorthogonal projection are determined by the pseudoinverse; *id est*, direction of projection $a_i a_i^{*\text{T}} x - x$ on $\mathcal{R}(a_i)$ is orthogonal to a_i^* . **E.9**

E.9 This remains true in high dimension although only a little more difficult to visualize in \mathbb{R}^3 ; confer, Figure 59.




Figure 142: (*confer* Figure 58) Biorthogonal expansion of point $x \in \text{aff } \mathcal{K}$ is found by projecting x nonorthogonally on extreme directions of polyhedral cone $\mathcal{K} \subset \mathbb{R}^2$. (Dotted lines of projection bound this translated negated cone.) Direction of projection on extreme direction a_1 is orthogonal to extreme direction a_1^* of dual cone \mathcal{K}^* and parallel to a_2 (§E.3.5); similarly, direction of projection on a_2 is orthogonal to a_2^* and parallel to a_1 . Point x is sum of nonorthogonal projections: x on $\mathcal{R}(a_1)$ and x on $\mathcal{R}(a_2)$. Expansion is unique because extreme directions of \mathcal{K} are linearly independent. Were a_1 orthogonal to a_2 , then \mathcal{K} would be identical to \mathcal{K}^* and nonorthogonal projections would become orthogonal.

Because the extreme directions of this cone \mathcal{K} are linearly independent, the component projections are unique in the sense:

- there is only one linear combination of extreme directions of \mathcal{K} that yields a particular point $x \in \mathcal{R}(A)$ whenever

$$\mathcal{R}(A) = \text{aff } \mathcal{K} = \mathcal{R}(a_1) \oplus \mathcal{R}(a_2) \oplus \dots \oplus \mathcal{R}(a_n) \quad (1826)$$

□

E.5.0.0.4 Example. *Nonorthogonal projection on elementary matrix.* Suppose $P_{\mathcal{Y}}$ is a linear nonorthogonal projector projecting on subspace \mathcal{Y} , and suppose the range of a vector u is linearly independent of \mathcal{Y} ; *id est*, for some other subspace \mathcal{M} containing \mathcal{Y} suppose

$$\mathcal{M} = \mathcal{R}(u) \oplus \mathcal{Y} \quad (1827)$$

Assuming $P_{\mathcal{M}}x = P_u x + P_{\mathcal{Y}}x$ holds, then it follows for vector $x \in \mathcal{M}$

$$P_u x = x - P_{\mathcal{Y}}x, \quad P_{\mathcal{Y}}x = x - P_u x \quad (1828)$$

nonorthogonal projection of x on $\mathcal{R}(u)$ can be determined from nonorthogonal projection of x on \mathcal{Y} , and *vice versa*.

Such a scenario is realizable were there some arbitrary basis for \mathcal{Y} populating a full-rank skinny-or-square matrix A

$$A \triangleq [\text{basis } \mathcal{Y} \quad u] \in \mathbb{R}^{n+1} \quad (1829)$$

Then $P_{\mathcal{M}} = AA^\dagger$ fulfills the requirements, with $P_u = A(:, n+1)A^\dagger(n+1, :)$ and $P_{\mathcal{Y}} = A(:, 1:n)A^\dagger(1:n, :)$. Observe, $P_{\mathcal{M}}$ is an orthogonal projector whereas $P_{\mathcal{Y}}$ and P_u are nonorthogonal projectors.

Now suppose, for example, $P_{\mathcal{Y}}$ is an elementary matrix (§B.3); in particular,

$$P_{\mathcal{Y}} = I - e_1 \mathbf{1}^T = \begin{bmatrix} \mathbf{0} & \sqrt{2}V_{\mathcal{N}} \end{bmatrix} \in \mathbb{R}^{N \times N} \quad (1830)$$

where $\mathcal{Y} = \mathcal{N}(\mathbf{1}^T)$. We have $\mathcal{M} = \mathbb{R}^N$, $A = [\sqrt{2}V_{\mathcal{N}} \quad e_1]$, and $u = e_1$. Thus $P_u = e_1 \mathbf{1}^T$ is a nonorthogonal projector projecting on $\mathcal{R}(u)$ in a direction parallel to a vector in \mathcal{Y} (§E.3.5), and $P_{\mathcal{Y}}x = x - e_1 \mathbf{1}^T x$ is a nonorthogonal projection of x on \mathcal{Y} in a direction parallel to u . □

E.5.0.0.5 Example. *Projecting the origin on a hyperplane.*

(confer §2.4.2.0.2) Given the hyperplane representation having $b \in \mathbb{R}$ and nonzero normal $a \in \mathbb{R}^m$

$$\partial\mathcal{H} = \{y \mid a^T y = b\} \subset \mathbb{R}^m \quad (106)$$

orthogonal projection of the origin $P\mathbf{0}$ on that hyperplane is the unique optimal solution to a minimization problem: (1795)

$$\begin{aligned} \|P\mathbf{0} - \mathbf{0}\|_2 &= \inf_{y \in \partial\mathcal{H}} \|y - \mathbf{0}\|_2 \\ &= \inf_{\xi \in \mathbb{R}^{m-1}} \|Z\xi + x\|_2 \end{aligned} \quad (1831)$$

where x is any solution to $a^T y = b$, and where the columns of $Z \in \mathbb{R}^{m \times m-1}$ constitute a basis for $\mathcal{N}(a^T)$ so that $y = Z\xi + x \in \partial\mathcal{H}$ for all $\xi \in \mathbb{R}^{m-1}$.

The infimum can be found by setting the gradient (with respect to ξ) of the strictly convex norm-square to $\mathbf{0}$. We find the minimizing argument

$$\xi^* = -(Z^T Z)^{-1} Z^T x \quad (1832)$$

so

$$y^* = (I - Z(Z^T Z)^{-1} Z^T)x \quad (1833)$$

and from (1797)

$$P\mathbf{0} = y^* = a(a^T a)^{-1} a^T x = \frac{a}{\|a\|} \frac{a^T}{\|a\|} x = \frac{a}{\|a\|^2} a^T x \triangleq A^\dagger A x = a \frac{b}{\|a\|^2} \quad (1834)$$

In words, any point x in the hyperplane $\partial\mathcal{H}$ projected on its normal a (confer (1859)) yields that point y^* in the hyperplane closest to the origin.

□

E.5.0.0.6 Example. *Projection on affine subset.*

The technique of Example E.5.0.0.5 is extensible. Given an intersection of hyperplanes

$$\mathcal{A} = \{y \mid Ay = b\} \subset \mathbb{R}^m \quad (1835)$$

where each row of $A \in \mathbb{R}^{m \times n}$ is nonzero and $b \in \mathcal{R}(A)$, then the orthogonal projection Px of any point $x \in \mathbb{R}^n$ on \mathcal{A} is the solution to a minimization problem:

$$\begin{aligned} \|Px - x\|_2 &= \inf_{y \in \mathcal{A}} \|y - x\|_2 \\ &= \inf_{\xi \in \mathbb{R}^{n - \text{rank } A}} \|Z\xi + y_p - x\|_2 \end{aligned} \quad (1836)$$

where y_p is any solution to $Ay = b$, and where the columns of $Z \in \mathbb{R}^{n \times n - \text{rank } A}$ constitute a basis for $\mathcal{N}(A)$ so that $y = Z\xi + y_p \in \mathcal{A}$ for all $\xi \in \mathbb{R}^{n - \text{rank } A}$.

The infimum is found by setting the gradient of the strictly convex norm-square to $\mathbf{0}$. The minimizing argument is

$$\xi^* = -(Z^T Z)^{-1} Z^T (y_p - x) \quad (1837)$$

so

$$y^* = (I - Z(Z^T Z)^{-1} Z^T)(y_p - x) + x \quad (1838)$$

and from (1797),

$$\begin{aligned} Px &= y^* = x - A^\dagger(Ax - b) \\ &= (I - A^\dagger A)x + A^\dagger A y_p \end{aligned} \quad (1839)$$

which is a projection of x on $\mathcal{N}(A)$ then translated perpendicularly with respect to the nullspace until it meets the affine subset \mathcal{A} . \square

E.5.0.0.7 Example. Projection on affine subset, vertex-description.

Suppose now we instead describe the affine subset \mathcal{A} in terms of some given minimal set of generators arranged columnar in $X \in \mathbb{R}^{n \times N}$ (69); *id est*,

$$\mathcal{A} \triangleq \text{aff } X = \{Xa \mid a^T \mathbf{1} = 1\} \subseteq \mathbb{R}^n \quad (1840)$$

Here *minimal set* means $XV_N = [x_2 - x_1 \quad x_3 - x_1 \quad \cdots \quad x_N - x_1] / \sqrt{2}$ (875) is full-rank (§2.4.2.2) where $V_N \in \mathbb{R}^{N \times N-1}$ is the Schoenberg auxiliary matrix (§B.4.2). Then the orthogonal projection Px of any point $x \in \mathbb{R}^n$ on \mathcal{A} is the solution to a minimization problem:

$$\begin{aligned} \|Px - x\|_2 &= \inf_{a^T \mathbf{1} = 1} \|Xa - x\|_2 \\ &= \inf_{\xi \in \mathbb{R}^{N-1}} \|X(V_N \xi + a_p) - x\|_2 \end{aligned} \quad (1841)$$

where a_p is any solution to $a^T \mathbf{1} = 1$. We find the minimizing argument

$$\xi^* = -(V_N^T X^T X V_N)^{-1} V_N^T X^T (X a_p - x) \quad (1842)$$

and so the orthogonal projection is [183, §3]

$$Px = X a^* = (I - X V_N (X V_N)^\dagger) X a_p + X V_N (X V_N)^\dagger x \quad (1843)$$

a projection of point x on $\mathcal{R}(X V_N)$ then translated perpendicularly with respect to that range until it meets the affine subset \mathcal{A} . \square

E.5.0.0.8 Example. *Projecting on hyperplane, halfspace, slab.*

Given the hyperplane representation having $b \in \mathbb{R}$ and nonzero normal $a \in \mathbb{R}^m$

$$\partial\mathcal{H} = \{y \mid a^T y = b\} \subset \mathbb{R}^m \quad (106)$$

the orthogonal projection of any point $x \in \mathbb{R}^m$ on that hyperplane is

$$Px = x - a(a^T a)^{-1}(a^T x - b) \quad (1844)$$

Orthogonal projection of x on the halfspace parametrized by $b \in \mathbb{R}$ and nonzero normal $a \in \mathbb{R}^m$

$$\mathcal{H}_- = \{y \mid a^T y \leq b\} \subset \mathbb{R}^m \quad (98)$$

is the point

$$Px = x - a(a^T a)^{-1} \max\{0, a^T x - b\} \quad (1845)$$

Orthogonal projection of x on the convex slab (Figure 11), for $c < b$

$$\mathcal{B} \triangleq \{y \mid c \leq a^T y \leq b\} \subset \mathbb{R}^m \quad (1846)$$

is the point [128, §5.1]

$$Px = x - a(a^T a)^{-1} (\max\{0, a^T x - b\} - \max\{0, c - a^T x\}) \quad (1847)$$

□

E.6 Vectorization interpretation, projection on a matrix

E.6.1 Nonorthogonal projection on a vector

Nonorthogonal projection of vector x on the range of vector y is accomplished using a normalized dyad P_0 (§B.1); *videlicet*,

$$\frac{\langle z, x \rangle}{\langle z, y \rangle} y = \frac{z^T x}{z^T y} y = \frac{y z^T}{z^T y} x \triangleq P_0 x \quad (1848)$$

where $\langle z, x \rangle / \langle z, y \rangle$ is the coefficient of projection on y . Because $P_0^2 = P_0$ and $\mathcal{R}(P_0) = \mathcal{R}(y)$, rank-one matrix P_0 is a nonorthogonal *projector dyad* projecting on $\mathcal{R}(y)$. Direction of nonorthogonal projection is orthogonal to z ; *id est*,

$$P_0 x - x \perp \mathcal{R}(P_0^T) \quad (1849)$$

E.6.2 Nonorthogonal projection on vectorized matrix

Formula (1848) is extensible. Given $X, Y, Z \in \mathbb{R}^{m \times n}$, we have the one-dimensional nonorthogonal projection of X in isomorphic \mathbb{R}^{mn} on the range of vectorized Y : (§2.2)

$$\frac{\langle Z, X \rangle}{\langle Z, Y \rangle} Y, \quad \langle Z, Y \rangle \neq 0 \quad (1850)$$

where $\langle Z, X \rangle / \langle Z, Y \rangle$ is the coefficient of projection. The inequality accounts for the fact: projection on $\mathcal{R}(\text{vec } Y)$ is in a direction orthogonal to $\text{vec } Z$.

E.6.2.1 Nonorthogonal projection on dyad

Now suppose we have nonorthogonal projector dyad

$$P_0 = \frac{yz^T}{z^T y} \in \mathbb{R}^{m \times m} \quad (1851)$$

Analogous to (1848), for $X \in \mathbb{R}^{m \times m}$

$$P_0 X P_0 = \frac{yz^T}{z^T y} X \frac{yz^T}{z^T y} = \frac{z^T X y}{(z^T y)^2} yz^T = \frac{\langle zy^T, X \rangle}{\langle zy^T, yz^T \rangle} yz^T \quad (1852)$$

is a nonorthogonal projection of matrix X on the range of vectorized dyad P_0 ; from which it follows:

$$P_0 X P_0 = \frac{z^T X y}{z^T y} \frac{yz^T}{z^T y} = \left\langle \frac{zy^T}{z^T y}, X \right\rangle \frac{yz^T}{z^T y} = \langle P_0^T, X \rangle P_0 = \frac{\langle P_0^T, X \rangle}{\langle P_0^T, P_0 \rangle} P_0 \quad (1853)$$

Yet this relationship between matrix product and vector inner-product only holds for a dyad projector. When nonsymmetric projector P_0 is rank-one as in (1851), therefore,

$$\mathcal{R}(\text{vec } P_0 X P_0) = \mathcal{R}(\text{vec } P_0) \text{ in } \mathbb{R}^{m^2} \quad (1854)$$

and

$$P_0 X P_0 - X \perp P_0^T \text{ in } \mathbb{R}^{m^2} \quad (1855)$$

E.6.2.1.1 Example. λ as coefficients of nonorthogonal projection.
Any diagonalization (§A.5)

$$X = S\Lambda S^{-1} = \sum_{i=1}^m \lambda_i s_i w_i^T \in \mathbb{R}^{m \times m} \quad (1449)$$

may be expressed as a sum of one-dimensional nonorthogonal projections of X , each on the range of a vectorized eigenmatrix $P_j \triangleq s_j w_j^T$;

$$\begin{aligned} X &= \sum_{i,j=1}^m \langle (S e_i e_j^T S^{-1})^T, X \rangle S e_i e_j^T S^{-1} \\ &= \sum_{j=1}^m \langle (s_j w_j^T)^T, X \rangle s_j w_j^T + \sum_{\substack{i,j=1 \\ j \neq i}}^m \langle (S e_i e_j^T S^{-1})^T, S\Lambda S^{-1} \rangle S e_i e_j^T S^{-1} \\ &= \sum_{j=1}^m \langle (s_j w_j^T)^T, X \rangle s_j w_j^T \quad (1856) \\ &\triangleq \sum_{j=1}^m \langle P_j^T, X \rangle P_j = \sum_{j=1}^m s_j w_j^T X s_j w_j^T = \sum_{j=1}^m P_j X P_j \\ &= \sum_{j=1}^m \lambda_j s_j w_j^T \end{aligned}$$

This biorthogonal expansion of matrix X is a sum of nonorthogonal projections because the term outside the projection coefficient $\langle \rangle$ is not identical to the inside-term. (§E.6.4) The eigenvalues λ_j are coefficients of nonorthogonal projection of X , while the remaining $M(M-1)/2$ coefficients (for $i \neq j$) are zeroed by projection. When P_j is rank-one as in (1856),

$$\mathcal{R}(\text{vec } P_j X P_j) = \mathcal{R}(\text{vec } s_j w_j^T) = \mathcal{R}(\text{vec } P_j) \text{ in } \mathbb{R}^{m^2} \quad (1857)$$

and

$$P_j X P_j - X \perp P_j^T \text{ in } \mathbb{R}^{m^2} \quad (1858)$$

Were matrix X symmetric, then its eigenmatrices would also be. So the one-dimensional projections would become orthogonal. (§E.6.4.1.1) \square

E.6.3 Orthogonal projection on a vector

The formula for orthogonal projection of vector x on the range of vector y (*one-dimensional projection*) is basic analytic geometry; [12, §3.3] [294, §3.2] [325, §2.2] [339, §1-8]

$$\frac{\langle y, x \rangle}{\langle y, y \rangle} y = \frac{y^T x}{y^T y} y = \frac{y y^T}{y^T y} x \triangleq P_1 x \quad (1859)$$

where $\langle y, x \rangle / \langle y, y \rangle$ is the coefficient of projection on $\mathcal{R}(y)$. An equivalent description is: Vector $P_1 x$ is the orthogonal projection of vector x on $\mathcal{R}(P_1) = \mathcal{R}(y)$. Rank-one matrix P_1 is a projection matrix because $P_1^2 = P_1$. The direction of projection is orthogonal

$$P_1 x - x \perp \mathcal{R}(P_1) \quad (1860)$$

because $P_1^T = P_1$.

E.6.4 Orthogonal projection on a vectorized matrix

From (1859), given instead $X, Y \in \mathbb{R}^{m \times n}$, we have the one-dimensional orthogonal projection of matrix X in isomorphic \mathbb{R}^{mn} on the range of vectorized Y : (§2.2)

$$\frac{\langle Y, X \rangle}{\langle Y, Y \rangle} Y \quad (1861)$$

where $\langle Y, X \rangle / \langle Y, Y \rangle$ is the coefficient of projection.

For orthogonal projection, the term outside the vector inner-products $\langle \rangle$ must be identical to the terms inside in three places.

E.6.4.1 Orthogonal projection on dyad

There is opportunity for insight when Y is a dyad yz^T (§B.1): Instead given $X \in \mathbb{R}^{m \times n}$, $y \in \mathbb{R}^m$, and $z \in \mathbb{R}^n$

$$\frac{\langle yz^T, X \rangle}{\langle yz^T, yz^T \rangle} yz^T = \frac{y^T X z}{y^T y z^T z} yz^T \quad (1862)$$

is the one-dimensional orthogonal projection of X in isomorphic \mathbb{R}^{mn} on the range of vectorized yz^T . To reveal the obscured symmetric projection

matrices P_1 and P_2 we rewrite (1862):

$$\frac{y^T X z}{y^T y z^T z} y z^T = \frac{y y^T}{y^T y} X \frac{z z^T}{z^T z} \triangleq P_1 X P_2 \quad (1863)$$

So for projector dyads, projection (1863) is the orthogonal projection in \mathbb{R}^{mn} if and only if projectors P_1 and P_2 are symmetric; ^{E.10} in other words,

- for orthogonal projection on the range of a vectorized dyad yz^T , the term outside the vector inner-products $\langle \rangle$ in (1862) must be identical to the terms inside in three places.

When P_1 and P_2 are rank-one symmetric projectors as in (1863), (31)

$$\mathcal{R}(\text{vec } P_1 X P_2) = \mathcal{R}(\text{vec } y z^T) \text{ in } \mathbb{R}^{mn} \quad (1864)$$

and

$$P_1 X P_2 - X \perp y z^T \text{ in } \mathbb{R}^{mn} \quad (1865)$$

When $y = z$ then $P_1 = P_2 = P_2^T$ and

$$P_1 X P_1 = \langle P_1, X \rangle P_1 = \frac{\langle P_1, X \rangle}{\langle P_1, P_1 \rangle} P_1 \quad (1866)$$

meaning, $P_1 X P_1$ is equivalent to orthogonal projection of matrix X on the range of vectorized projector dyad P_1 . Yet this relationship between matrix product and vector inner-product does not hold for general symmetric projector matrices.

^{E.10}For diagonalizable $X \in \mathbb{R}^{m \times m}$ (§A.5), its orthogonal projection in isomorphic \mathbb{R}^{m^2} on the range of vectorized $yz^T \in \mathbb{R}^{m \times m}$ becomes:

$$P_1 X P_2 = \sum_{i=1}^m \lambda_i P_1 s_i w_i^T P_2$$

When $\mathcal{R}(P_1) = \mathcal{R}(w_j)$ and $\mathcal{R}(P_2) = \mathcal{R}(s_j)$, the j^{th} dyad term from the diagonalization is isolated but only, in general, to within a scale factor because neither set of left or right eigenvectors is necessarily orthonormal unless X is normal [352, §3.2]. Yet when $\mathcal{R}(P_2) = \mathcal{R}(s_k)$, $k \neq j \in \{1 \dots m\}$, then $P_1 X P_2 = \mathbf{0}$.

E.6.4.1.1 Example. *Eigenvalues λ as coefficients of orthogonal projection.* Let \mathcal{C} represent any convex subset of subspace \mathbb{S}^M , and let \mathcal{C}_1 be any element of \mathcal{C} . Then \mathcal{C}_1 can be expressed as the orthogonal expansion

$$\mathcal{C}_1 = \sum_{i=1}^M \sum_{\substack{j=1 \\ j \geq i}}^M \langle E_{ij}, \mathcal{C}_1 \rangle E_{ij} \in \mathcal{C} \subset \mathbb{S}^M \quad (1867)$$

where $E_{ij} \in \mathbb{S}^M$ is a member of the standard orthonormal basis for \mathbb{S}^M (53). This expansion is a sum of one-dimensional orthogonal projections of \mathcal{C}_1 ; each projection on the range of a vectorized standard basis matrix. Vector inner-product $\langle E_{ij}, \mathcal{C}_1 \rangle$ is the coefficient of projection of $\text{svec } \mathcal{C}_1$ on $\mathcal{R}(\text{svec } E_{ij})$.

When \mathcal{C}_1 is any member of a convex set \mathcal{C} whose dimension is L , *Carathéodory's theorem* [96] [271] [177] [36] [37] guarantees that no more than $L + 1$ affinely independent members from \mathcal{C} are required to faithfully represent \mathcal{C}_1 by their linear combination. **E.11**

Dimension of \mathbb{S}^M is $L = M(M+1)/2$ in isometrically isomorphic $\mathbb{R}^{M(M+1)/2}$. Yet because any symmetric matrix can be diagonalized, (§A.5.2) $\mathcal{C}_1 \in \mathbb{S}^M$ is a linear combination of its M eigenmatrices $q_i q_i^T$ (§A.5.1) weighted by its eigenvalues λ_i ;

$$\mathcal{C}_1 = Q \Lambda Q^T = \sum_{i=1}^M \lambda_i q_i q_i^T \quad (1868)$$

where $\Lambda \in \mathbb{S}^M$ is a diagonal matrix having $\delta(\Lambda)_i = \lambda_i$, and $Q = [q_1 \cdots q_M]$ is an orthogonal matrix in $\mathbb{R}^{M \times M}$ containing corresponding eigenvectors.

To derive eigen decomposition (1868) from expansion (1867), M standard basis matrices E_{ij} are rotated (§B.5) into alignment with the M eigenmatrices $q_i q_i^T$ of \mathcal{C}_1 by applying a *similarity transformation*; [294, §5.6]

$$\{Q E_{ij} Q^T\} = \left\{ \begin{array}{ll} q_i q_i^T, & i = j = 1 \dots M \\ \frac{1}{\sqrt{2}}(q_i q_j^T + q_j q_i^T), & 1 \leq i < j \leq M \end{array} \right\} \quad (1869)$$

E.11 Carathéodory's theorem guarantees existence of a biorthogonal expansion for any element in $\text{aff } \mathcal{C}$ when \mathcal{C} is any pointed closed convex cone.

which remains an orthonormal basis for \mathbb{S}^M . Then remarkably

$$\begin{aligned}
\mathcal{C}_1 &= \sum_{\substack{i,j=1 \\ j \geq i}}^M \langle QE_{ij}Q^T, \mathcal{C}_1 \rangle QE_{ij}Q^T \\
&= \sum_{i=1}^M \langle q_i q_i^T, \mathcal{C}_1 \rangle q_i q_i^T + \sum_{\substack{i,j=1 \\ j > i}}^M \langle QE_{ij}Q^T, Q\Lambda Q^T \rangle QE_{ij}Q^T \\
&= \sum_{i=1}^M \langle q_i q_i^T, \mathcal{C}_1 \rangle q_i q_i^T \\
&\triangleq \sum_{i=1}^M \langle P_i, \mathcal{C}_1 \rangle P_i = \sum_{i=1}^M q_i q_i^T \mathcal{C}_1 q_i q_i^T = \sum_{i=1}^M P_i \mathcal{C}_1 P_i \\
&= \sum_{i=1}^M \lambda_i q_i q_i^T
\end{aligned} \tag{1870}$$

this orthogonal expansion becomes the diagonalization; still a sum of one-dimensional orthogonal projections. The eigenvalues

$$\lambda_i = \langle q_i q_i^T, \mathcal{C}_1 \rangle \tag{1871}$$

are clearly coefficients of projection of \mathcal{C}_1 on the range of each vectorized eigenmatrix. (confer §E.6.2.1.1) The remaining $M(M-1)/2$ coefficients ($i \neq j$) are zeroed by projection. When P_i is rank-one symmetric as in (1870),

$$\mathcal{R}(\text{svec } P_i \mathcal{C}_1 P_i) = \mathcal{R}(\text{svec } q_i q_i^T) = \mathcal{R}(\text{svec } P_i) \text{ in } \mathbb{R}^{M(M+1)/2} \tag{1872}$$

and

$$P_i \mathcal{C}_1 P_i - \mathcal{C}_1 \perp P_i \text{ in } \mathbb{R}^{M(M+1)/2} \tag{1873}$$

□

E.6.4.2 Positive semidefiniteness test as orthogonal projection

For any given $X \in \mathbb{R}^{m \times m}$ the familiar quadratic construct $y^T X y \geq 0$, over broad domain, is a fundamental test for positive semidefiniteness. (§A.2) It is a fact that $y^T X y$ is always proportional to a coefficient of orthogonal projection; letting z in formula (1862) become $y \in \mathbb{R}^m$, then $P_2 = P_1 = yy^T / y^T y = yy^T / \|yy^T\|_2$ (confer (1508)) and formula (1863) becomes

$$\frac{\langle yy^T, X \rangle}{\langle yy^T, yy^T \rangle} yy^T = \frac{y^T X y}{y^T y} \frac{yy^T}{y^T y} = \frac{yy^T}{y^T y} X \frac{yy^T}{y^T y} \triangleq P_1 X P_1 \tag{1874}$$

By (1861), product $P_1 X P_1$ is the one-dimensional orthogonal projection of X in isomorphic \mathbb{R}^{m^2} on the range of vectorized P_1 because, for $\text{rank } P_1 = 1$ and $P_1^2 = P_1 \in \mathbb{S}^m$ (confer (1853))

$$P_1 X P_1 = \frac{y^T X y}{y^T y} \frac{y y^T}{y^T y} = \left\langle \frac{y y^T}{y^T y}, X \right\rangle \frac{y y^T}{y^T y} = \langle P_1, X \rangle P_1 = \frac{\langle P_1, X \rangle}{\langle P_1, P_1 \rangle} P_1 \quad (1875)$$

The coefficient of orthogonal projection $\langle P_1, X \rangle = y^T X y / (y^T y)$ is also known as *Rayleigh's quotient*.^{E.12} When P_1 is rank-one symmetric as in (1874),

$$\mathcal{R}(\text{vec } P_1 X P_1) = \mathcal{R}(\text{vec } P_1) \text{ in } \mathbb{R}^{m^2} \quad (1876)$$

and

$$P_1 X P_1 - X \perp P_1 \text{ in } \mathbb{R}^{m^2} \quad (1877)$$

The test for positive semidefiniteness, then, is a test for nonnegativity of the coefficient of orthogonal projection of X on the range of each and every vectorized extreme direction yy^T (§2.8.1) from the positive semidefinite cone in the ambient space of symmetric matrices.

^{E.12}When y becomes the j^{th} eigenvector s_j of diagonalizable X , for example, $\langle P_1, X \rangle$ becomes the j^{th} eigenvalue: [174, §III]

$$\langle P_1, X \rangle|_{y=s_j} = \frac{s_j^T \left(\sum_{i=1}^m \lambda_i s_i w_i^T \right) s_j}{s_j^T s_j} = \lambda_j$$

Similarly for $y = w_j$, the j^{th} left-eigenvector,

$$\langle P_1, X \rangle|_{y=w_j} = \frac{w_j^T \left(\sum_{i=1}^m \lambda_i s_i w_i^T \right) w_j}{w_j^T w_j} = \lambda_j$$

A quandary may arise regarding the potential annihilation of the antisymmetric part of X when $s_j^T X s_j$ is formed. Were annihilation to occur, it would imply the eigenvalue thus found came instead from the symmetric part of X . The quandary is resolved recognizing that diagonalization of real X admits complex eigenvectors; hence, annihilation could only come about by forming $\text{Re}(s_j^H X s_j) = s_j^H (X + X^T) s_j / 2$ [180, §7.1] where $(X + X^T)/2$ is the symmetric part of X , and s_j^H denotes conjugate transpose.

E.6.4.3 $PXP \succeq 0$

In some circumstances, it may be desirable to limit the domain of test $y^T X y \geq 0$ for positive semidefiniteness; *e.g.*, $\{\|y\|=1\}$. Another example of limiting domain-of-test is central to Euclidean distance geometry: For $\mathcal{R}(V) = \mathcal{N}(\mathbf{1}^T)$, the test $-VDV \succeq 0$ determines whether $D \in \mathbb{S}_h^N$ is a Euclidean distance matrix. The same test may be stated: For $D \in \mathbb{S}_h^N$ (and optionally $\|y\|=1$)

$$D \in \text{EDM}^N \Leftrightarrow -y^T D y = \langle yy^T, -D \rangle \geq 0 \quad \forall y \in \mathcal{R}(V) \quad (1878)$$

The test $-VDV \succeq 0$ is therefore equivalent to a test for nonnegativity of the coefficient of orthogonal projection of $-D$ on the range of each and every vectorized extreme direction yy^T from the positive semidefinite cone \mathbb{S}_+^N such that $\mathcal{R}(yy^T) = \mathcal{R}(y) \subseteq \mathcal{R}(V)$. (The validity of this result is independent of whether V is itself a projection matrix.)

E.7 Projection on matrix subspaces

E.7.1 PXP misinterpretation for higher-rank P

For a projection matrix P of rank greater than 1, PXP is generally not commensurate with $\frac{\langle P, X \rangle}{\langle P, P \rangle} P$ as is the case for projector dyads (1875). Yet for a symmetric idempotent matrix P of any rank we are tempted to say “ PXP is the orthogonal projection of $X \in \mathbb{S}^m$ on $\mathcal{R}(\text{vec } P)$ ”. The fallacy is: $\text{vec } PXP$ does not necessarily belong to the range of vectorized P ; the most basic requirement for projection on $\mathcal{R}(\text{vec } P)$.

E.7.2 Orthogonal projection on matrix subspaces

With $A_1 \in \mathbb{R}^{m \times n}$, $B_1 \in \mathbb{R}^{n \times k}$, $Z_1 \in \mathbb{R}^{m \times k}$, $A_2 \in \mathbb{R}^{p \times n}$, $B_2 \in \mathbb{R}^{n \times k}$, $Z_2 \in \mathbb{R}^{p \times k}$ as defined for nonorthogonal projector (1773), and defining

$$P_1 \triangleq A_1 A_1^\dagger \in \mathbb{S}^m, \quad P_2 \triangleq A_2 A_2^\dagger \in \mathbb{S}^p \quad (1879)$$

then, given compatible X

$$\|X - P_1 X P_2\|_F = \inf_{B_1, B_2 \in \mathbb{R}^{n \times k}} \|X - A_1 (A_1^\dagger + B_1 Z_1^T) X (A_2^{\dagger T} + Z_2 B_2^T) A_2^T\|_F \quad (1880)$$

As for all subspace projectors, range of the projector is the subspace on which projection is made: $\{P_1 Y P_2 \mid Y \in \mathbb{R}^{m \times p}\}$. Altogether, for projectors P_1 and P_2 of any rank, this means projection $P_1 X P_2$ is unique minimum-distance, orthogonal

$$P_1 X P_2 - X \perp \{P_1 Y P_2 \mid Y \in \mathbb{R}^{m \times p}\} \text{ in } \mathbb{R}^{mp} \quad (1881)$$

and P_1 and P_2 must each be symmetric (*confer* (1863)) to attain the infimum.

E.7.2.0.1 Proof. Minimum Frobenius norm (1880).

Defining $P \triangleq A_1(A_1^\dagger + B_1 Z_1^\top)$,

$$\begin{aligned} & \inf_{B_1, B_2} \|X - A_1(A_1^\dagger + B_1 Z_1^\top)X(A_2^{\dagger\top} + Z_2 B_2^\top)A_2^\top\|_F^2 \\ &= \inf_{B_1, B_2} \|X - PX(A_2^{\dagger\top} + Z_2 B_2^\top)A_2^\top\|_F^2 \\ &= \inf_{B_1, B_2} \text{tr}\left((X^\top - A_2(A_2^\dagger + B_2 Z_2^\top)X^\top P^\top)(X - PX(A_2^{\dagger\top} + Z_2 B_2^\top)A_2^\top)\right) \quad (1882) \\ &= \inf_{B_1, B_2} \text{tr}\left(X^\top X - X^\top P X(A_2^{\dagger\top} + Z_2 B_2^\top)A_2^\top - A_2(A_2^\dagger + B_2 Z_2^\top)X^\top P^\top X \right. \\ & \quad \left. + A_2(A_2^\dagger + B_2 Z_2^\top)X^\top P^\top P X(A_2^{\dagger\top} + Z_2 B_2^\top)A_2^\top\right) \end{aligned}$$

Necessary conditions for a global minimum are $\nabla_{B_1} = \mathbf{0}$ and $\nabla_{B_2} = \mathbf{0}$. Terms not containing B_2 in (1882) will vanish from gradient ∇_{B_2} ; (§D.2.3)

$$\begin{aligned} & \nabla_{B_2} \text{tr}\left(-X^\top P X Z_2 B_2^\top A_2^\top - A_2 B_2 Z_2^\top X^\top P^\top X + A_2 A_2^\dagger X^\top P^\top P X Z_2 B_2^\top A_2^\top \right. \\ & \quad \left. + A_2 B_2 Z_2^\top X^\top P^\top P X A_2^{\dagger\top} A_2^\top + A_2 B_2 Z_2^\top X^\top P^\top P X Z_2 B_2^\top A_2^\top\right) \\ &= -2A_2^\top X^\top P X Z_2 + 2A_2^\top A_2 A_2^\dagger X^\top P^\top P X Z_2 + 2A_2^\top A_2 B_2 Z_2^\top X^\top P^\top P X Z_2 \\ &= A_2^\top \left(-X^\top + A_2 A_2^\dagger X^\top P^\top + A_2 B_2 Z_2^\top X^\top P^\top\right) P X Z_2 \quad (1883) \\ &= \mathbf{0} \quad \Leftrightarrow \\ & \quad \mathcal{R}(B_1) \subseteq \mathcal{N}(A_1) \quad \text{and} \quad \mathcal{R}(B_2) \subseteq \mathcal{N}(A_2) \end{aligned}$$

(or $Z_2 = \mathbf{0}$) because $A^\top = A^\top A A^\dagger$. Symmetry requirement (1879) is implicit. Were instead $P^\top \triangleq (A_2^{\dagger\top} + Z_2 B_2^\top)A_2^\top$ and the gradient with respect to B_1 observed, then similar results are obtained. The projector is unique. Perpendicularity (1881) establishes uniqueness [92, §4.9] of projection $P_1 X P_2$ on a matrix subspace. The minimum-distance projector is the orthogonal projector, and *vice versa*. \blacklozenge

E.7.2.0.2 Example. PXP redux & $\mathcal{N}(\mathbf{V})$.

Suppose we define a subspace of $m \times n$ matrices, each elemental matrix having columns constituting a list whose geometric center (§5.5.1.0.1) is the origin in \mathbb{R}^m :

$$\begin{aligned}\mathbb{R}_c^{m \times n} &\triangleq \{Y \in \mathbb{R}^{m \times n} \mid Y\mathbf{1} = \mathbf{0}\} \\ &= \{Y \in \mathbb{R}^{m \times n} \mid \mathcal{N}(Y) \supseteq \mathbf{1}\} = \{Y \in \mathbb{R}^{m \times n} \mid \mathcal{R}(Y^T) \subseteq \mathcal{N}(\mathbf{1}^T)\} \\ &= \{XV \mid X \in \mathbb{R}^{m \times n}\} \subset \mathbb{R}^{m \times n}\end{aligned}\tag{1884}$$

the *nonsymmetric geometric center subspace*. Further suppose $V \in \mathbb{S}^n$ is a projection matrix having $\mathcal{N}(V) = \mathcal{R}(\mathbf{1})$ and $\mathcal{R}(V) = \mathcal{N}(\mathbf{1}^T)$. Then linear mapping $T(X) = XV$ is the orthogonal projection of any $X \in \mathbb{R}^{m \times n}$ on $\mathbb{R}_c^{m \times n}$ in the Euclidean (vectorization) sense because V is symmetric, $\mathcal{N}(XV) \supseteq \mathbf{1}$, and $\mathcal{R}(VX^T) \subseteq \mathcal{N}(\mathbf{1}^T)$.

Now suppose we define a subspace of symmetric $n \times n$ matrices each of whose columns constitute a list having the origin in \mathbb{R}^n as geometric center,

$$\begin{aligned}\mathbb{S}_c^n &\triangleq \{Y \in \mathbb{S}^n \mid Y\mathbf{1} = \mathbf{0}\} \\ &= \{Y \in \mathbb{S}^n \mid \mathcal{N}(Y) \supseteq \mathbf{1}\} = \{Y \in \mathbb{S}^n \mid \mathcal{R}(Y) \subseteq \mathcal{N}(\mathbf{1}^T)\}\end{aligned}\tag{1885}$$

the *geometric center subspace*. Further suppose $V \in \mathbb{S}^n$ is a projection matrix, the same as before. Then VXV is the orthogonal projection of any $X \in \mathbb{S}^n$ on \mathbb{S}_c^n in the Euclidean sense (1881) because V is symmetric, $VXV\mathbf{1} = \mathbf{0}$, and $\mathcal{R}(VXV) \subseteq \mathcal{N}(\mathbf{1}^T)$. Two-sided projection is necessary only to remain in the ambient symmetric matrix subspace. Then

$$\mathbb{S}_c^n = \{VXV \mid X \in \mathbb{S}^n\} \subset \mathbb{S}^n\tag{1886}$$

has $\dim \mathbb{S}_c^n = n(n-1)/2$ in isomorphic $\mathbb{R}^{n(n+1)/2}$. We find its orthogonal complement as the aggregate of all negative directions of orthogonal projection on \mathbb{S}_c^n : the *translation-invariant subspace* (§5.5.1.1)

$$\begin{aligned}\mathbb{S}_c^{\perp} &\triangleq \{X - VXV \mid X \in \mathbb{S}^n\} \subset \mathbb{S}^n \\ &= \{u\mathbf{1}^T + \mathbf{1}u^T \mid u \in \mathbb{R}^n\}\end{aligned}\tag{1887}$$

characterized by the doublet $u\mathbf{1}^T + \mathbf{1}u^T$ (§B.2).^{E.13} Defining the

E.13 Proof.

$$\begin{aligned}\{X - VXV \mid X \in \mathbb{S}^n\} &= \{X - (I - \frac{1}{n}\mathbf{1}\mathbf{1}^T)X(I - \mathbf{1}\mathbf{1}^T\frac{1}{n}) \mid X \in \mathbb{S}^n\} \\ &= \{\frac{1}{n}\mathbf{1}\mathbf{1}^T X + X\mathbf{1}\mathbf{1}^T\frac{1}{n} - \frac{1}{n}\mathbf{1}\mathbf{1}^T X \mathbf{1}\mathbf{1}^T\frac{1}{n} \mid X \in \mathbb{S}^n\}\end{aligned}$$

geometric center mapping $\mathbf{V}(X) = -VXV\frac{1}{2}$ consistently with (911), then $\mathcal{N}(\mathbf{V}) = \mathcal{R}(I - \mathbf{V})$ on domain \mathbb{S}^n analogously to vector projectors (§E.2); *id est*,

$$\mathcal{N}(\mathbf{V}) = \mathbb{S}_c^{n\perp} \quad (1888)$$

a subspace of \mathbb{S}^n whose dimension is $\dim \mathbb{S}_c^{n\perp} = n$ in isomorphic $\mathbb{R}^{n(n+1)/2}$. Intuitively, operator \mathbf{V} is an orthogonal projector; any argument duplicitously in its range is a fixed point. So, this symmetric operator's nullspace must be orthogonal to its range.

Now compare the subspace of symmetric matrices having all zeros in the first row and column

$$\begin{aligned} \mathbb{S}_1^n &\triangleq \{Y \in \mathbb{S}^n \mid Ye_1 = \mathbf{0}\} \\ &= \left\{ \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} X \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} \mid X \in \mathbb{S}^n \right\} \\ &= \left\{ [\mathbf{0} \quad \sqrt{2}V_N]^T Z [\mathbf{0} \quad \sqrt{2}V_N] \mid Z \in \mathbb{S}^N \right\} \end{aligned} \quad (1889)$$

where $P = \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix}$ is an orthogonal projector. Then, similarly, PXP is the orthogonal projection of any $X \in \mathbb{S}^n$ on \mathbb{S}_1^n in the Euclidean sense (1881), and

$$\begin{aligned} \mathbb{S}_1^{n\perp} &\triangleq \left\{ \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} X \begin{bmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & I \end{bmatrix} - X \mid X \in \mathbb{S}^n \right\} \subset \mathbb{S}^n \\ &= \{ue_1^T + e_1u^T \mid u \in \mathbb{R}^n\} \end{aligned} \quad (1890)$$

Obviously, $\mathbb{S}_1^n \oplus \mathbb{S}_1^{n\perp} = \mathbb{S}^n$. □

Because $\{X\mathbf{1} \mid X \in \mathbb{S}^n\} = \mathbb{R}^n$,

$$\begin{aligned} \{X - VXV \mid X \in \mathbb{S}^n\} &= \{\mathbf{1}\zeta^T + \zeta\mathbf{1}^T - \mathbf{1}\mathbf{1}^T(\mathbf{1}^T\zeta\frac{1}{n}) \mid \zeta \in \mathbb{R}^n\} \\ &= \{\mathbf{1}\zeta^T(I - \mathbf{1}\mathbf{1}^T\frac{1}{2n}) + (I - \frac{1}{2n}\mathbf{1}\mathbf{1}^T)\zeta\mathbf{1}^T \mid \zeta \in \mathbb{R}^n\} \end{aligned}$$

where $I - \frac{1}{2n}\mathbf{1}\mathbf{1}^T$ is invertible. ◆

E.8 Range/Rowspace interpretation

For idempotent matrices P_1 and P_2 of any rank, $P_1XP_2^T$ is a projection of $\mathcal{R}(X)$ on $\mathcal{R}(P_1)$ and a projection of $\mathcal{R}(X^T)$ on $\mathcal{R}(P_2)$: For any given $X = U\Sigma Q^T \in \mathbb{R}^{m \times p}$, as in compact singular value decomposition (1462),

$$P_1XP_2^T = \sum_{i=1}^{\eta} \sigma_i P_1 u_i q_i^T P_2^T = \sum_{i=1}^{\eta} \sigma_i P_1 u_i (P_2 q_i)^T \quad (1891)$$

where $\eta \triangleq \min\{m, p\}$. Recall $u_i \in \mathcal{R}(X)$ and $q_i \in \mathcal{R}(X^T)$ when the corresponding singular value σ_i is nonzero. (§A.6.1) So P_1 projects u_i on $\mathcal{R}(P_1)$ while P_2 projects q_i on $\mathcal{R}(P_2)$; *id est*, the range and rowspace of any X are respectively projected on the ranges of P_1 and P_2 . ^{E.14}

E.9 Projection on convex set

Thus far we have discussed only projection on subspaces. Now we generalize, considering projection on arbitrary convex sets in Euclidean space; convex because projection is, then, unique minimum-distance and a convex optimization problem:

For projection P_Cx of point x on any closed set $\mathcal{C} \subseteq \mathbb{R}^n$ it is obvious:

$$\mathcal{C} \equiv \{P_Cx \mid x \in \mathbb{R}^n\} \quad (1892)$$

If $\mathcal{C} \subseteq \mathbb{R}^n$ is a closed convex set, then for each and every $x \in \mathbb{R}^n$ there exists a unique point Px belonging to \mathcal{C} that is closest to x in the Euclidean sense. Like (1795), unique projection Px (or P_Cx) of a point x on convex set \mathcal{C} is that point in \mathcal{C} closest to x ; [220, §3.12]

$$\|x - Px\|_2 = \inf_{y \in \mathcal{C}} \|x - y\|_2 \quad (1893)$$

There exists a converse (in finite-dimensional Euclidean space):

^{E.14}When P_1 and P_2 are symmetric and $\mathcal{R}(P_1) = \mathcal{R}(u_j)$ and $\mathcal{R}(P_2) = \mathcal{R}(q_j)$, then the j^{th} dyad term from the singular value decomposition of X is isolated by the projection. Yet if $\mathcal{R}(P_2) = \mathcal{R}(q_\ell)$, $\ell \neq j \in \{1 \dots \eta\}$, then $P_1XP_2 = \mathbf{0}$.

E.9.0.0.1 Theorem. (Bunt-Motzkin) *Convex set if projections unique.* [331, §7.5] [175] If $\mathcal{C} \subseteq \mathbb{R}^n$ is a nonempty closed set and if for each and every x in \mathbb{R}^n there is a unique Euclidean projection Px of x on \mathcal{C} belonging to \mathcal{C} , then \mathcal{C} is convex. \diamond

Borwein & Lewis propose, for closed convex set \mathcal{C} [48, §3.3, exer.12(d)]

$$\nabla \|x - Px\|_2^2 = 2(x - Px) \quad (1894)$$

for any point x whereas, for $x \notin \mathcal{C}$

$$\nabla \|x - Px\|_2 = (x - Px) \|x - Px\|_2^{-1} \quad (1895)$$

E.9.0.0.2 Exercise. *Norm gradient.*
Prove (1894) and (1895). (Not proved in [48].) \blacktriangledown

A well-known equivalent characterization of projection on a convex set is a generalization of the perpendicularity condition (1794) for projection on a subspace:

E.9.1 Dual interpretation of projection on convex set

E.9.1.0.1 Definition. *Normal vector.* [271, p.15]

Vector z is *normal* to convex set \mathcal{C} at point $Px \in \mathcal{C}$ if

$$\langle z, y - Px \rangle \leq 0 \quad \forall y \in \mathcal{C} \quad (1896)$$

\triangle

A convex set has at least one nonzero normal at each of its boundary points. [271, p.100] (Figure 62) Hence, the *normal* or *dual* interpretation of projection:

E.9.1.0.2 Theorem. *Unique minimum-distance projection.* [177, §A.3.1] [220, §3.12] [92, §4.1] [69] (Figure 147(b), p.695) Given a closed convex set $\mathcal{C} \subseteq \mathbb{R}^n$, point Px is the unique projection of a given point $x \in \mathbb{R}^n$ on \mathcal{C} (Px is that point in \mathcal{C} nearest x) if and only if

$$Px \in \mathcal{C}, \quad \langle x - Px, y - Px \rangle \leq 0 \quad \forall y \in \mathcal{C} \quad (1897)$$

\diamond

As for subspace projection, operator P is idempotent in the sense: for each and every $x \in \mathbb{R}^n$, $P(Px) = Px$. Yet operator P is not linear;

- projector P is a linear operator if and only if convex set \mathcal{C} (on which projection is made) is a subspace. (§E.4)

E.9.1.0.3 Theorem. *Unique projection via normal cone.*^{E.15} [92, §4.3] Given closed convex set $\mathcal{C} \subseteq \mathbb{R}^n$, point Px is the unique projection of a given point $x \in \mathbb{R}^n$ on \mathcal{C} if and only if

$$Px \in \mathcal{C}, \quad Px - x \in (\mathcal{C} - Px)^* \quad (1898)$$

In other words, Px is that point in \mathcal{C} nearest x if and only if $Px - x$ belongs to that cone dual to translate $\mathcal{C} - Px$. \diamond

E.9.1.1 Dual interpretation as optimization

Deutsch [94, thm.2.3] [95, §2] and Luenberger [220, p.134] carry forward Nirenberg's dual interpretation of projection [245] as solution to a maximization problem: Minimum distance from a point $x \in \mathbb{R}^n$ to a convex set $\mathcal{C} \subset \mathbb{R}^n$ can be found by maximizing distance from x to hyperplane $\partial\mathcal{H}$ over the set of all hyperplanes separating x from \mathcal{C} . Existence of a separating hyperplane (§2.4.2.7) presumes point x lies on the boundary or exterior to set \mathcal{C} .

The optimal separating hyperplane is characterized by the fact it also supports \mathcal{C} . Any hyperplane supporting \mathcal{C} (Figure 27(a)) has form

$$\underline{\partial\mathcal{H}}_- = \{y \in \mathbb{R}^n \mid a^T y = \sigma_{\mathcal{C}}(a)\} \quad (121)$$

where the support function is convex, defined

$$\sigma_{\mathcal{C}}(a) \triangleq \sup_{z \in \mathcal{C}} a^T z \quad (499)$$

When point x is finite and set \mathcal{C} contains finite points, under this projection interpretation, if the supporting hyperplane is a separating hyperplane then the support function is finite. From Example E.5.0.0.8, projection $P_{\underline{\partial\mathcal{H}}_-} x$ of x on any given supporting hyperplane $\underline{\partial\mathcal{H}}_-$ is

$$P_{\underline{\partial\mathcal{H}}_-} x = x - a(a^T a)^{-1}(a^T x - \sigma_{\mathcal{C}}(a)) \quad (1899)$$

^{E.15} $-(\mathcal{C} - Px)^*$ is the normal cone to set \mathcal{C} at point Px . (§E.10.3.2)




Figure 143: Dual interpretation of projection of point x on convex set \mathcal{C} in \mathbb{R}^2 . (a) $\kappa = (a^T a)^{-1} (a^T x - \sigma_{\mathcal{C}}(a))$ (b) Minimum distance from x to \mathcal{C} is found by maximizing distance to all hyperplanes supporting \mathcal{C} and separating it from x . A convex problem for any convex set, distance of maximization is unique.

With reference to Figure 143, identifying

$$\mathcal{H}_+ = \{y \in \mathbb{R}^n \mid a^\top y \geq \sigma_{\mathcal{C}}(a)\} \quad (99)$$

then

$$\begin{aligned} \|x - P_{\mathcal{C}}x\| &= \sup_{\partial\mathcal{H}_- \mid x \in \mathcal{H}_+} \|x - P_{\partial\mathcal{H}_-}x\| = \sup_{a \mid x \in \mathcal{H}_+} \|a(a^\top a)^{-1}(a^\top x - \sigma_{\mathcal{C}}(a))\| \\ &= \sup_{a \mid x \in \mathcal{H}_+} \frac{1}{\|a\|} |a^\top x - \sigma_{\mathcal{C}}(a)| \end{aligned} \quad (1900)$$

which can be expressed as a convex optimization, for arbitrary positive constant τ

$$\begin{aligned} \|x - P_{\mathcal{C}}x\| &= \frac{1}{\tau} \underset{a}{\text{maximize}} \quad a^\top x - \sigma_{\mathcal{C}}(a) \\ &\text{subject to} \quad \|a\| \leq \tau \end{aligned} \quad (1901)$$

The unique minimum-distance projection on convex set \mathcal{C} is therefore

$$P_{\mathcal{C}}x = x - a^*(a^{*\top}x - \sigma_{\mathcal{C}}(a^*)) \frac{1}{\tau^2} \quad (1902)$$

where optimally $\|a^*\| = \tau$.

E.9.1.1.1 Exercise. *Dual projection technique on polyhedron.*

Test that projection paradigm from Figure 143 on any convex polyhedral set. ▼

E.9.1.2 Dual interpretation of projection on cone

In the circumstance set \mathcal{C} is a closed convex cone \mathcal{K} and there exists a hyperplane separating given point x from \mathcal{K} , then optimal $\sigma_{\mathcal{K}}(a^*)$ takes value 0 [177, §C.2.3.1]. So problem (1901) for projection of x on \mathcal{K} becomes

$$\begin{aligned} \|x - P_{\mathcal{K}}x\| &= \frac{1}{\tau} \underset{a}{\text{maximize}} \quad a^\top x \\ &\text{subject to} \quad \|a\| \leq \tau \\ &\quad a \in \mathcal{K}^\circ \end{aligned} \quad (1903)$$

The norm inequality in (1903) can be handled by Schur complement (§3.1.7.2). Normals a to all hyperplanes supporting \mathcal{K} belong to the polar cone $\mathcal{K}^\circ = -\mathcal{K}^*$ by definition: (289)

$$a \in \mathcal{K}^\circ \Leftrightarrow \langle a, x \rangle \leq 0 \text{ for all } x \in \mathcal{K} \quad (1904)$$

Projection on cone \mathcal{K} is

$$P_{\mathcal{K}}x = \left(I - \frac{1}{\tau^2}a^*a^{*\top}\right)x \quad (1905)$$

whereas projection on the polar cone $-\mathcal{K}^*$ is (§E.9.2.2.1)

$$P_{\mathcal{K}^\circ}x = x - P_{\mathcal{K}}x = \frac{1}{\tau^2}a^*a^{*\top}x \quad (1906)$$

Negating vector a , this maximization problem (1903) becomes a minimization (the same problem) and the polar cone becomes the dual cone:

$$\begin{aligned} \|x - P_{\mathcal{K}}x\| &= -\frac{1}{\tau} \underset{a}{\text{minimize}} \quad a^\top x \\ &\text{subject to} \quad \|a\| \leq \tau \\ &\quad a \in \mathcal{K}^* \end{aligned} \quad (1907)$$

E.9.2 Projection on cone

When convex set \mathcal{C} is a cone, there is a finer statement of optimality conditions:

E.9.2.0.1 Theorem. *Unique projection on cone.* [177, §A.3.2]

Let $\mathcal{K} \subseteq \mathbb{R}^n$ be a closed convex cone, and \mathcal{K}^* its dual (§2.13.1). Then Px is the unique minimum-distance projection of $x \in \mathbb{R}^n$ on \mathcal{K} if and only if

$$Px \in \mathcal{K}, \quad \langle Px - x, Px \rangle = 0, \quad Px - x \in \mathcal{K}^* \quad (1908)$$

◇

In words, Px is the unique minimum-distance projection of x on \mathcal{K} if and only if

- 1) projection Px lies in \mathcal{K}
- 2) direction $Px - x$ is orthogonal to the projection Px
- 3) direction $Px - x$ lies in the dual cone \mathcal{K}^* .

As the theorem is stated, it admits projection on \mathcal{K} having empty interior; *id est*, on convex cones in a proper subspace of \mathbb{R}^n .

Projection on \mathcal{K} of any point $x \in -\mathcal{K}^*$, belonging to the negative dual cone, is the origin. By (1908): the set of all points reaching the origin, when projecting on \mathcal{K} , constitutes the negative dual cone; a.k.a, the polar cone

$$\mathcal{K}^\circ = -\mathcal{K}^* = \{x \in \mathbb{R}^n \mid Px = \mathbf{0}\} \quad (1909)$$

E.9.2.1 Relationship to subspace projection

Conditions 1 and 2 of the theorem are common with orthogonal projection on a subspace $\mathcal{R}(P)$: Condition 1 is the most basic requirement; namely, the projection $Px \in \mathcal{R}(P)$ belongs to the subspace. Invoking perpendicularity condition (1794), we recall the second requirement for projection on a subspace:

$$Px - x \perp \mathcal{R}(P) \quad \text{or} \quad Px - x \in \mathcal{R}(P)^\perp \quad (1910)$$

which corresponds to condition 2. Yet condition 3 is a generalization of subspace projection; *id est*, for unique minimum-distance projection on a closed convex cone \mathcal{K} , polar cone $-\mathcal{K}^*$ plays the role $\mathcal{R}(P)^\perp$ plays for subspace projection ($P_{\mathcal{R}}x = x - P_{\mathcal{R}^\perp}x$). Indeed, $-\mathcal{K}^*$ is the algebraic complement in the orthogonal vector sum (p.714) [238] [177, §A.3.2.5]

$$\mathcal{K} \boxplus -\mathcal{K}^* = \mathbb{R}^n \Leftrightarrow \text{cone } \mathcal{K} \text{ is closed and convex} \quad (1911)$$

Also, given unique minimum-distance projection Px on \mathcal{K} satisfying Theorem E.9.2.0.1, then by projection on the algebraic complement via $I - P$ in §E.2 we have

$$-\mathcal{K}^* = \{x - Px \mid x \in \mathbb{R}^n\} = \{x \in \mathbb{R}^n \mid Px = \mathbf{0}\} \quad (1912)$$

consequent to Moreau (1915). Recalling any subspace is a closed convex cone^{E.16}

$$\mathcal{K} = \mathcal{R}(P) \Leftrightarrow -\mathcal{K}^* = \mathcal{R}(P)^\perp \quad (1913)$$

meaning, when a cone is a subspace $\mathcal{R}(P)$, then the dual cone becomes its orthogonal complement $\mathcal{R}(P)^\perp$. [53, §2.6.1] In this circumstance, condition 3 becomes coincident with condition 2.

The properties of projection on cones following in §E.9.2.2 further generalize to subspaces by: (4)

$$\mathcal{K} = \mathcal{R}(P) \Leftrightarrow -\mathcal{K} = \mathcal{R}(P) \quad (1914)$$

^{E.16} but a proper subspace is not a proper cone (§2.7.2.2.1).

E.9.2.2 Salient properties: Projection Px on closed convex cone \mathcal{K}

[177, §A.3.2] [92, §5.6] For $x, x_1, x_2 \in \mathbb{R}^n$

1. $P_{\mathcal{K}}(\alpha x) = \alpha P_{\mathcal{K}}x \quad \forall \alpha \geq 0$ (nonnegative homogeneity)
2. $\|P_{\mathcal{K}}x\| \leq \|x\|$
3. $P_{\mathcal{K}}x = \mathbf{0} \Leftrightarrow x \in -\mathcal{K}^*$
4. $P_{\mathcal{K}}(-x) = -P_{-\mathcal{K}}x$
5. (Jean-Jacques Moreau (1962)) [238]

$$\begin{aligned} x = x_1 + x_2, \quad x_1 \in \mathcal{K}, \quad x_2 \in -\mathcal{K}^*, \quad x_1 \perp x_2 \\ \Leftrightarrow \\ x_1 = P_{\mathcal{K}}x, \quad x_2 = P_{-\mathcal{K}^*}x \end{aligned} \quad (1915)$$

6. $\mathcal{K} = \{x - P_{-\mathcal{K}^*}x \mid x \in \mathbb{R}^n\} = \{x \in \mathbb{R}^n \mid P_{-\mathcal{K}^*}x = \mathbf{0}\}$
7. $-\mathcal{K}^* = \{x - P_{\mathcal{K}}x \mid x \in \mathbb{R}^n\} = \{x \in \mathbb{R}^n \mid P_{\mathcal{K}}x = \mathbf{0}\}$ (1912)

E.9.2.2.1 Corollary. *$I - P$ for cones.* (confer §E.2)

Denote by $\mathcal{K} \subseteq \mathbb{R}^n$ a closed convex cone, and call \mathcal{K}^* its dual. Then $x - P_{-\mathcal{K}^*}x$ is the unique minimum-distance projection of $x \in \mathbb{R}^n$ on \mathcal{K} if and only if $P_{-\mathcal{K}^*}x$ is the unique minimum-distance projection of x on $-\mathcal{K}^*$ the polar cone. \diamond

Proof. Assume $x_1 = P_{\mathcal{K}}x$. Then by Theorem E.9.2.0.1 we have

$$x_1 \in \mathcal{K}, \quad x_1 - x \perp x_1, \quad x_1 - x \in \mathcal{K}^* \quad (1916)$$

Now assume $x - x_1 = P_{-\mathcal{K}^*}x$. Then we have

$$x - x_1 \in -\mathcal{K}^*, \quad -x_1 \perp x - x_1, \quad -x_1 \in -\mathcal{K} \quad (1917)$$

But these two assumptions are apparently identical. We must therefore have

$$x - P_{-\mathcal{K}^*}x = x_1 = P_{\mathcal{K}}x \quad (1918)$$

\blacklozenge

E.9.2.2.2 Corollary. *Unique projection via dual or normal cone.*
 [92, §4.7] (§E.10.3.2, confer Theorem E.9.1.0.3) Given point $x \in \mathbb{R}^n$ and closed convex cone $\mathcal{K} \subseteq \mathbb{R}^n$, the following are equivalent statements:

1. point Px is the unique minimum-distance projection of x on \mathcal{K}
2. $Px \in \mathcal{K}$, $x - Px \in -(\mathcal{K} - Px)^* = -\mathcal{K}^* \cap (Px)^\perp$
3. $Px \in \mathcal{K}$, $\langle x - Px, Px \rangle = 0$, $\langle x - Px, y \rangle \leq 0 \quad \forall y \in \mathcal{K}$

◇

E.9.2.2.3 Example. *Unique projection on nonnegative orthant.*
 (confer (1235)) From Theorem E.9.2.0.1, to project matrix $H \in \mathbb{R}^{m \times n}$ on the self-dual orthant (§2.13.5.1) of nonnegative matrices $\mathbb{R}_+^{m \times n}$ in isomorphic \mathbb{R}^{mn} , the necessary and sufficient conditions are:

$$\begin{aligned} H^* &\geq \mathbf{0} \\ \text{tr}((H^* - H)^T H^*) &= 0 \\ H^* - H &\geq \mathbf{0} \end{aligned} \tag{1919}$$

where the inequalities denote entrywise comparison. The optimal solution H^* is simply H having all its negative entries zeroed;

$$H_{ij}^* = \max\{H_{ij}, 0\}, \quad i, j \in \{1 \dots m\} \times \{1 \dots n\} \tag{1920}$$

Now suppose the nonnegative orthant is translated by $T \in \mathbb{R}^{m \times n}$; *id est*, consider $\mathbb{R}_+^{m \times n} + T$. Then projection on the translated orthant is [92, §4.8]

$$H_{ij}^* = \max\{H_{ij}, T_{ij}\} \tag{1921}$$

□

E.9.2.2.4 Example. *Unique projection on truncated convex cone.*
 Consider the problem of projecting a point x on a closed convex cone that is artificially bounded; really, a bounded convex polyhedron having a vertex at the origin:

$$\begin{aligned} &\underset{y \in \mathbb{R}^N}{\text{minimize}} && \|x - Ay\|_2 \\ &\text{subject to} && y \succeq 0 \\ &&& \|y\|_\infty \leq 1 \end{aligned} \tag{1922}$$

where the convex cone has vertex-description (§2.12.2.0.1), for $A \in \mathbb{R}^{n \times N}$

$$\mathcal{K} = \{Ay \mid y \succeq 0\} \quad (1923)$$

and where $\|y\|_\infty \leq 1$ is the artificial bound. This is a convex optimization problem having no known closed-form solution, in general. It arises, for example, in the fitting of hearing aids designed around a programmable graphic equalizer (a filter bank whose only adjustable parameters are gain per band each bounded above by unity). [84] The problem is equivalent to a Schur-form semidefinite program (§3.1.7.2)

$$\begin{aligned} & \underset{y \in \mathbb{R}^N, t \in \mathbb{R}}{\text{minimize}} && t \\ & \text{subject to} && \begin{bmatrix} tI & x - Ay \\ (x - Ay)^T & t \end{bmatrix} \succeq 0 \\ & && 0 \preceq y \preceq \mathbf{1} \end{aligned} \quad (1924)$$

□

E.9.3 nonexpansivity

E.9.3.0.1 Theorem. *Nonexpansivity.* [154, §2] [92, §5.3]
When $\mathcal{C} \subset \mathbb{R}^n$ is an arbitrary closed convex set, projector P projecting on \mathcal{C} is nonexpansive in the sense: for any vectors $x, y \in \mathbb{R}^n$

$$\|Px - Py\| \leq \|x - y\| \quad (1925)$$

with equality when $x - Px = y - Py$. ^{E.17}

◇

Proof. [47]

$$\begin{aligned} \|x - y\|^2 &= \|Px - Py\|^2 + \|(I - P)x - (I - P)y\|^2 \\ &\quad + 2\langle x - Px, Px - Py \rangle + 2\langle y - Py, Py - Px \rangle \end{aligned} \quad (1926)$$

Nonnegativity of the last two terms follows directly from the *unique minimum-distance projection theorem* (§E.9.1.0.2). ♦

^{E.17}This condition for equality corrects an error in [69] (where the norm is applied to each side of the condition given here) easily revealed by counter-example.

The foregoing proof reveals another flavor of nonexpansivity; for each and every $x, y \in \mathbb{R}^n$

$$\|Px - Py\|^2 + \|(I - P)x - (I - P)y\|^2 \leq \|x - y\|^2 \quad (1927)$$

Deutsch shows yet another: [92, §5.5]

$$\|Px - Py\|^2 \leq \langle x - y, Px - Py \rangle \quad (1928)$$

E.9.4 Easy projections

- Projecting any matrix $H \in \mathbb{R}^{n \times n}$ in the Euclidean/Frobenius sense orthogonally on the subspace of symmetric matrices \mathbb{S}^n in isomorphic \mathbb{R}^{n^2} amounts to taking the symmetric part of H ; (§2.2.2.0.1) *id est*, $(H + H^T)/2$ is the projection.
- To project any $H \in \mathbb{R}^{n \times n}$ orthogonally on the symmetric hollow subspace \mathbb{S}_h^n in isomorphic \mathbb{R}^{n^2} (§2.2.3.0.1, §7.0.1), we may take the symmetric part and then zero all entries along the main diagonal, or *vice versa* (because this is projection on the intersection of two subspaces); *id est*, $(H + H^T)/2 - \delta^2(H)$.
- To project a matrix on the nonnegative orthant $\mathbb{R}_+^{m \times n}$, simply clip all negative entries to 0. Likewise, projection on the nonpositive orthant $\mathbb{R}_-^{m \times n}$ sees all positive entries clipped to 0. Projection on other orthants is equally simple with appropriate clipping.
- Projecting on hyperplane, halfspace, slab: §E.5.0.0.8.
- Projection of $y \in \mathbb{R}^n$ on Euclidean ball $\mathcal{B} = \{x \in \mathbb{R}^n \mid \|x - a\| \leq c\}$: for $y \neq a$, $P_{\mathcal{B}}y = \frac{c}{\|y - a\|}(y - a) + a$.
- Clipping in excess of $|1|$ each entry of a point $x \in \mathbb{R}^n$ is equivalent to unique minimum-distance projection of x on a hypercube centered at the origin. (*confer* §E.10.3.2)
- Projection of $x \in \mathbb{R}^n$ on a (rectangular) *hyperbox*: [53, §8.1.1]

$$\mathcal{C} = \{y \in \mathbb{R}^n \mid l \preceq y \preceq u, l \prec u\} \quad (1929)$$

$$P(x)_{k=0\dots n} = \begin{cases} l_k, & x_k \leq l_k \\ x_k, & l_k \leq x_k \leq u_k \\ u_k, & x_k \geq u_k \end{cases} \quad (1930)$$

- Projection of x on the set of all cardinality- k vectors $\{y \mid \text{card } y \leq k\}$ keeps k entries of greatest magnitude and clips to 0 those remaining.
- Unique minimum-distance projection of $H \in \mathbb{S}^n$ on the positive semidefinite cone \mathbb{S}_+^n in the Euclidean/Frobenius sense is accomplished by eigen decomposition (diagonalization) followed by clipping all negative eigenvalues to 0.
- Unique minimum-distance projection on the generally nonconvex subset of all matrices belonging to \mathbb{S}_+^n having rank not exceeding ρ (§2.9.2.1) is accomplished by clipping all negative eigenvalues to 0 and zeroing the smallest nonnegative eigenvalues keeping only ρ largest. (§7.1.2)
- Unique minimum-distance projection, in the Euclidean/Frobenius sense, of $H \in \mathbb{R}^{m \times n}$ on the generally nonconvex subset of all $m \times n$ matrices of rank no greater than k is the singular value decomposition (§A.6) of H having all singular values beyond the k^{th} zeroed. This is also a solution to projection in the sense of spectral norm. [291, p.79, p.208]
- Projection on \mathcal{K} of any point $x \in -\mathcal{K}^*$, belonging to the polar cone, is equivalent to projection on the origin. (§E.9.2)
- $P_{\mathbb{S}_+^N \cap \mathbb{S}_c^N} = P_{\mathbb{S}_+^N} P_{\mathbb{S}_c^N}$ (1179)
- $P_{\mathbb{R}_+^{N \times N} \cap \mathbb{S}_h^N} = P_{\mathbb{R}_+^{N \times N}} P_{\mathbb{S}_h^N}$ (§7.0.1.1)
- $P_{\mathbb{R}_+^{N \times N} \cap \mathbb{S}^N} = P_{\mathbb{R}_+^{N \times N}} P_{\mathbb{S}^N}$ (§E.9.5)

E.9.4.0.1 Exercise. *Projection on spectral norm ball.*

Find the unique minimum-distance projection on the convex set of all $m \times n$ matrices whose largest singular value does not exceed 1; *id est*, on $\{X \in \mathbb{R}^{m \times n} \mid \|X\|_2 \leq 1\}$ the spectral norm ball (§2.3.2.0.5). ▼




Figure 144: Closed convex set \mathcal{C} belongs to subspace \mathbb{R}^n (shown bounded in sketch and drawn without proper perspective). Point y is unique minimum-distance projection of x on \mathcal{C} ; equivalent to product of orthogonal projection of x on \mathbb{R}^n and minimum-distance projection of result z on \mathcal{C} .

E.9.4.1 notes

Projection on Lorentz (second-order) cone: [53, exer.8.3(c)].

Deutsch [95] provides an algorithm for projection on polyhedral cones.

Youla [351, §2.5] lists eleven “useful projections”, of square-integrable uni- and bivariate real functions on various convex sets, in closed form.

Unique minimum-distance projection on an ellipsoid: Example 4.6.0.0.1.

Numerical algorithms for projection on the nonnegative simplex and 1-norm ball are in [109].

E.9.5 Projection on convex set in subspace

Suppose a convex set \mathcal{C} is contained in some subspace \mathbb{R}^n . Then unique minimum-distance projection of any point in $\mathbb{R}^n \oplus \mathbb{R}^{n\perp}$ on \mathcal{C} can be accomplished by first projecting orthogonally on that subspace, and then projecting the result on \mathcal{C} ; [92, §5.14] *id est*, the ordered product of two individual projections that is not commutable.

Proof. (\Leftarrow) To show that, suppose unique minimum-distance projection $P_{\mathcal{C}}x$ on $\mathcal{C} \subset \mathbb{R}^n$ is y as illustrated in Figure 144;

$$\|x - y\| \leq \|x - q\| \quad \forall q \in \mathcal{C} \quad (1931)$$

Further suppose $P_{\mathbb{R}^n}x$ equals z . By the *Pythagorean theorem*

$$\|x - y\|^2 = \|x - z\|^2 + \|z - y\|^2 \quad (1932)$$

because $x - z \perp z - y$. (1794) [220, §3.3] Then point $y = P_{\mathcal{C}}x$ is the same as $P_{\mathcal{C}}z$ because

$$\|z - y\|^2 = \|x - y\|^2 - \|x - z\|^2 \leq \|z - q\|^2 = \|x - q\|^2 - \|x - z\|^2 \quad \forall q \in \mathcal{C} \quad (1933)$$

which holds by assumption (1931).

(\Rightarrow) Now suppose $z = P_{\mathbb{R}^n}x$ and

$$\|z - y\| \leq \|z - q\| \quad \forall q \in \mathcal{C} \quad (1934)$$

meaning $y = P_{\mathcal{C}}z$. Then point y is identical to $P_{\mathcal{C}}x$ because

$$\|x - y\|^2 = \|x - z\|^2 + \|z - y\|^2 \leq \|x - q\|^2 = \|x - z\|^2 + \|z - q\|^2 \quad \forall q \in \mathcal{C} \quad (1935)$$

by assumption (1934). \blacklozenge

This proof is extensible via translation argument. (§E.4) Unique minimum-distance projection on a convex set contained in an affine subset is, therefore, similarly accomplished.

Projecting matrix $H \in \mathbb{R}^{n \times n}$ on convex cone $\mathcal{K} = \mathbb{S}^n \cap \mathbb{R}_+^{n \times n}$ in isomorphic \mathbb{R}^{n^2} can be accomplished, for example, by first projecting on \mathbb{S}^n and only then projecting the result on $\mathbb{R}_+^{n \times n}$ (confer §7.0.1). This is because that projection product is equivalent to projection on the subset of the nonnegative orthant in the symmetric matrix subspace.

E.10 Alternating projection

Alternating projection is an iterative technique for finding a point in the intersection of a number of arbitrary closed convex sets \mathcal{C}_k , or for finding the distance between two nonintersecting closed convex sets. Because it can sometimes be difficult or inefficient to compute the intersection or express it analytically, one naturally asks whether it is possible to instead project (unique minimum-distance) alternately on the individual \mathcal{C}_k ; often easier and what motivates adoption of this technique. Once a cycle of alternating projections (an *iteration*) is complete, we then *iterate* (repeat the cycle) until convergence. If the intersection of two closed convex sets is empty, then by *convergence* we mean the *iterate* (the result after a cycle of alternating projections) settles to a point of minimum distance separating the sets.

While alternating projection can find the point in the nonempty intersection closest to a given point b , it does not necessarily find the closest point. Finding that closest point is made dependable by an elegantly simple enhancement via correction to the alternating projection technique: this *Dykstra algorithm* (1973) for projection on the intersection is one of the most beautiful projection algorithms ever discovered. It is accurately interpreted as the discovery of what alternating projection originally sought to accomplish: unique minimum-distance projection on the nonempty intersection of a number of arbitrary closed convex sets \mathcal{C}_k . Alternating projection is, in fact, a special case of the Dykstra algorithm whose discussion we defer until §E.10.3.

E.10.0.1 commutative projectors

Given two arbitrary convex sets \mathcal{C}_1 and \mathcal{C}_2 and their respective minimum-distance projection operators P_1 and P_2 , if projectors commute for each and every $x \in \mathbb{R}^n$ then it is easy to show $P_1P_2x \in \mathcal{C}_1 \cap \mathcal{C}_2$ and $P_2P_1x \in \mathcal{C}_1 \cap \mathcal{C}_2$. When projectors commute ($P_1P_2 = P_2P_1$), a point in the intersection can be found in a finite number of steps; while commutativity is a sufficient condition, it is not necessary (§6.8.1.1.1 for example).

When \mathcal{C}_1 and \mathcal{C}_2 are subspaces, in particular, projectors P_1 and P_2 commute if and only if $P_1P_2 = P_{\mathcal{C}_1 \cap \mathcal{C}_2}$ or iff $P_2P_1 = P_{\mathcal{C}_1 \cap \mathcal{C}_2}$ or iff P_1P_2 is the orthogonal projection on a Euclidean subspace. [92, lem.9.2] Subspace projectors will commute, for example, when $P_1(\mathcal{C}_2) \subset \mathcal{C}_2$ or $P_2(\mathcal{C}_1) \subset \mathcal{C}_1$ or $\mathcal{C}_1 \subset \mathcal{C}_2$ or $\mathcal{C}_2 \subset \mathcal{C}_1$ or $\mathcal{C}_1 \perp \mathcal{C}_2$. When subspace projectors commute, this




Figure 145: First several alternating projections in von Neumann-style projection (1946), of point b , converging on closest point Pb in intersection of two closed convex sets in \mathbb{R}^2 : C_1 and C_2 are partially drawn in vicinity of their intersection. Pointed normal cone \mathcal{K}^\perp (1975) is translated to Pb , the unique minimum-distance projection of b on intersection. For this particular example, it is possible to start anywhere in a large neighborhood of b and still converge to Pb . Alternating projections are themselves robust with respect to significant noise because they belong to translated normal cone.

means we can find a point in the intersection of those subspaces in a finite number of steps; we find, in fact, the closest point.

E.10.0.1.1 Theorem. *Kronecker projector.* [290, §2.7]
Given any projection matrices P_1 and P_2 (subspace projectors), then

$$P_1 \otimes P_2 \quad \text{and} \quad P_1 \otimes I \quad (1936)$$

are projection matrices. The product preserves symmetry if present. \diamond

E.10.0.2 noncommutative projectors

Typically, one considers the method of alternating projection when projectors do not commute; *id est*, when $P_1 P_2 \neq P_2 P_1$.

The iconic example for noncommutative projectors illustrated in Figure 145 shows the iterates converging to the closest point in the intersection of two arbitrary convex sets. Yet simple examples like Figure 146 reveal that noncommutative alternating projection does not




Figure 146: The sets $\{\mathcal{C}_k\}$ in this example comprise two halfspaces \mathcal{H}_1 and \mathcal{H}_2 . The von Neumann-style alternating projection in \mathbb{R}^2 quickly converges to P_1P_2b (feasibility). The unique minimum-distance projection on the intersection is, of course, Pb .

always yield the closest point, although we shall show it always yields some point in the intersection or a point that attains the distance between two convex sets.

Alternating projection is also known as *successive projection* [157] [154] [55], *cyclic projection* [128] [230, §3.2], *successive approximation* [69], or *projection on convex sets* [288] [289, §6.4]. It is traced back to von Neumann (1933) [329] and later Wiener [335] who showed that higher iterates of a product of two orthogonal projections on subspaces converge at each point in the ambient space to the unique minimum-distance projection on the intersection of the two subspaces. More precisely, if \mathcal{R}_1 and \mathcal{R}_2 are closed subspaces of a Euclidean space and P_1 and P_2 respectively denote orthogonal projection on \mathcal{R}_1 and \mathcal{R}_2 , then for each vector b in that space,

$$\lim_{i \rightarrow \infty} (P_1P_2)^i b = P_{\mathcal{R}_1 \cap \mathcal{R}_2} b \quad (1937)$$

Deutsch [92, thm.9.8, thm.9.35] shows rate of convergence for subspaces to be *geometric* [350, §1.4.4]; bounded above by $\kappa^{2^{i+1}} \|b\|$, $i=0, 1, 2, \dots$, where

$0 \leq \kappa < 1$:

$$\|(P_1 P_2)^i b - P_{\mathcal{R}_1 \cap \mathcal{R}_2} b\| \leq \kappa^{2i+1} \|b\| \quad (1938)$$

This means convergence can be slow when κ is close to 1. Rate of convergence on intersecting halfspaces is also geometric. [93] [262]

This von Neumann sense of alternating projection may be applied to convex sets that are not subspaces, although convergence is not necessarily to the unique minimum-distance projection on the intersection. Figure 145 illustrates one application where convergence is reasonably geometric and the result is the unique minimum-distance projection. Figure 146, in contrast, demonstrates convergence in one iteration to a fixed point (of the projection product)^{E.18} in the intersection of two halfspaces; a.k.a, feasibility problem. It was Dykstra who in 1983 [110] (§E.10.3) first solved this projection problem.

E.10.0.3 the bullets

Alternating projection has, therefore, various meaning dependent on the application or field of study; it may be interpreted to be: a distance problem, a feasibility problem (von Neumann), or a projection problem (Dykstra):

- **Distance.** Figure 147(a)(b). Find a unique point of projection $P_1 b \in \mathcal{C}_1$ that attains the distance between any two closed convex sets \mathcal{C}_1 and \mathcal{C}_2 ;

$$\|P_1 b - b\| = \text{dist}(\mathcal{C}_1, \mathcal{C}_2) \triangleq \inf_{z \in \mathcal{C}_2} \|P_1 z - z\| \quad (1939)$$

- **Feasibility.** Figure 147(c), $\bigcap \mathcal{C}_k \neq \emptyset$. Given a number L of indexed closed convex sets $\mathcal{C}_k \subset \mathbb{R}^n$, find any fixed point in their intersection by iterating (i) a projection product starting from b ;

$$\left(\prod_{i=1}^{\infty} \prod_{k=1}^L P_k \right) b \in \bigcap_{k=1}^L \mathcal{C}_k \quad (1940)$$

- **Optimization.** Figure 147(c), $\bigcap \mathcal{C}_k \neq \emptyset$. Given a number of indexed closed convex sets $\mathcal{C}_k \subset \mathbb{R}^n$, uniquely project a given point b on $\bigcap \mathcal{C}_k$;

$$\|Pb - b\| = \inf_{x \in \bigcap \mathcal{C}_k} \|x - b\| \quad (1941)$$

^{E.18} *Fixed point* of a mapping $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a point x whose image is identical under the map; *id est*, $Tx = x$.




Figure 147:

(a) **(distance)** Intersection of two convex sets in \mathbb{R}^2 is empty. Method of alternating projection would be applied to find that point in \mathcal{C}_1 nearest \mathcal{C}_2 .

(b) **(distance)** Given $b \in \mathcal{C}_2$, then $P_1b \in \mathcal{C}_1$ is nearest b iff $(y - P_1b)^T (b - P_1b) \leq 0 \forall y \in \mathcal{C}_1$ by the *unique minimum-distance projection theorem* (§E.9.1.0.2). When P_1b attains the distance between the two sets, hyperplane $\{y \mid (b - P_1b)^T (y - P_1b) = 0\}$ separates \mathcal{C}_1 from \mathcal{C}_2 . [53, §2.5.1]

(c) **(0 distance)** Intersection is nonempty.

(optimization) We may want the point Pb in $\bigcap \mathcal{C}_k$ nearest point b .

(feasibility) We may instead be satisfied with a fixed point of the projection product P_1P_2b in $\bigcap \mathcal{C}_k$.

E.10.1 Distance and existence

Existence of a fixed point is established:

E.10.1.0.1 Theorem. *Distance.* [69]

Given any two closed convex sets \mathcal{C}_1 and \mathcal{C}_2 in \mathbb{R}^n , then $P_1 b \in \mathcal{C}_1$ is a fixed point of the projection product $P_1 P_2$ if and only if $P_1 b$ is a point of \mathcal{C}_1 nearest \mathcal{C}_2 . \diamond

Proof. (\Rightarrow) Given fixed point $a = P_1 P_2 a \in \mathcal{C}_1$ with $b \triangleq P_2 a \in \mathcal{C}_2$ in tandem so that $a = P_1 b$, then by the *unique minimum-distance projection theorem* (§E.9.1.0.2)

$$\begin{aligned} (b - a)^T(u - a) &\leq 0 \quad \forall u \in \mathcal{C}_1 \\ (a - b)^T(v - b) &\leq 0 \quad \forall v \in \mathcal{C}_2 \\ &\Leftrightarrow \\ \|a - b\| &\leq \|u - v\| \quad \forall u \in \mathcal{C}_1 \text{ and } \forall v \in \mathcal{C}_2 \end{aligned} \quad (1942)$$

by Schwarz inequality $\|\langle x, y \rangle\| \leq \|x\| \|y\|$ [201] [271].

(\Leftarrow) Suppose $a \in \mathcal{C}_1$ and $\|a - P_2 a\| \leq \|u - P_2 u\| \forall u \in \mathcal{C}_1$. Now suppose we choose $u = P_1 P_2 a$. Then

$$\|u - P_2 u\| = \|P_1 P_2 a - P_2 P_1 P_2 a\| \leq \|a - P_2 a\| \Leftrightarrow a = P_1 P_2 a \quad (1943)$$

Thus $a = P_1 b$ (with $b = P_2 a \in \mathcal{C}_2$) is a fixed point in \mathcal{C}_1 of the projection product $P_1 P_2$. ^{E.19} \blacklozenge

E.10.2 Feasibility and convergence

The set of all fixed points of any nonexpansive mapping is a closed convex set. [136, lem.3.4] [27, §1] The projection product $P_1 P_2$ is nonexpansive by Theorem E.9.3.0.1 because, for any vectors $x, a \in \mathbb{R}^n$

$$\|P_1 P_2 x - P_1 P_2 a\| \leq \|P_2 x - P_2 a\| \leq \|x - a\| \quad (1944)$$

If the intersection of two closed convex sets $\mathcal{C}_1 \cap \mathcal{C}_2$ is empty, then the iterates converge to a point of minimum distance, a fixed point of the projection product. Otherwise, convergence is to some fixed point in their intersection

^{E.19}Point $b = P_2 a$ can be shown, similarly, to be a fixed point of the product $P_2 P_1$.

(a feasible point) whose existence is guaranteed by virtue of the fact that each and every point in the convex intersection is in one-to-one correspondence with fixed points of the nonexpansive projection product.

Bauschke & Borwein [27, §2] argue that any sequence monotonic in the sense of Fejér is convergent. ^{E.20}

E.10.2.0.1 Definition. *Fejér monotonicity.* [239]

Given closed convex set $\mathcal{C} \neq \emptyset$, then a sequence $x_i \in \mathbb{R}^n$, $i=0, 1, 2, \dots$, is monotonic in the sense of Fejér with respect to \mathcal{C} iff

$$\|x_{i+1} - c\| \leq \|x_i - c\| \quad \text{for all } i \geq 0 \quad \mathbf{and} \quad \text{each and every } c \in \mathcal{C} \quad (1945)$$

△

Given $x_0 \stackrel{\Delta}{=} b$, if we express each iteration of alternating projection by

$$x_{i+1} = P_1 P_2 x_i, \quad i=0, 1, 2, \dots \quad (1946)$$

and define any fixed point $a = P_1 P_2 a$, then sequence x_i is Fejér monotone with respect to fixed point a because

$$\|P_1 P_2 x_i - a\| \leq \|x_i - a\| \quad \forall i \geq 0 \quad (1947)$$

by nonexpansivity. The nonincreasing sequence $\|P_1 P_2 x_i - a\|$ is bounded below hence convergent because any bounded monotonic sequence in \mathbb{R} is convergent; [228, §1.2] [37, §1.1] $P_1 P_2 x_{i+1} = P_1 P_2 x_i = x_{i+1}$. Sequence x_i therefore converges to some fixed point. If the intersection $\mathcal{C}_1 \cap \mathcal{C}_2$ is nonempty, convergence is to some point there by the *distance theorem*. Otherwise, x_i converges to a point in \mathcal{C}_1 of minimum distance to \mathcal{C}_2 .

E.10.2.0.2 Example. *Hyperplane/orthant intersection.*

Find a feasible point (1940) belonging to the nonempty intersection of two convex sets: given $A \in \mathbb{R}^{m \times n}$, $\beta \in \mathcal{R}(A)$

$$\mathcal{C}_1 \cap \mathcal{C}_2 = \mathbb{R}_+^n \cap \mathcal{A} = \{y \mid y \succeq 0\} \cap \{y \mid Ay = \beta\} \subset \mathbb{R}^n \quad (1948)$$

the nonnegative orthant with affine subset \mathcal{A} an intersection of hyperplanes. Projection of an iterate $x_i \in \mathbb{R}^n$ on \mathcal{A} is calculated

$$P_2 x_i = x_i - A^T (A A^T)^{-1} (A x_i - \beta) \quad (1839)$$

^{E.20}Other authors prove convergence by different means; e.g., [154] [55].




Figure 148: From Example E.10.2.0.2 in \mathbb{R}^2 , showing von Neumann-style alternating projection to find feasible point belonging to intersection of nonnegative orthant with hyperplane \mathcal{A} . Point Pb lies at intersection of hyperplane with ordinate axis. In this particular example, the feasible point found is coincidentally optimal. Rate of convergence depends upon angle θ ; as it becomes more acute, convergence slows. [154, §3]




Figure 149: Geometric convergence of iterates in norm, for Example E.10.2.0.2 in \mathbb{R}^{1000} .

while, thereafter, projection of the result on the orthant is simply

$$x_{i+1} = P_1 P_2 x_i = \max\{\mathbf{0}, P_2 x_i\} \quad (1949)$$

where the maximum is entrywise (§E.9.2.2.3).

One realization of this problem in \mathbb{R}^2 is illustrated in Figure 148: For $A = \begin{bmatrix} 1 & 1 \end{bmatrix}$, $\beta = 1$, and $x_0 = b = \begin{bmatrix} -3 & 1/2 \end{bmatrix}^T$, the iterates converge to the feasible point $Pb = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$.

To give a more palpable sense of convergence in higher dimension, we do this example again but now we compute an alternating projection for the case $A \in \mathbb{R}^{400 \times 1000}$, $\beta \in \mathbb{R}^{400}$, and $b \in \mathbb{R}^{1000}$, all of whose entries are independently and randomly set to a uniformly distributed real number in the interval $[-1, 1]$. Convergence is illustrated in Figure 149. \square

This application of alternating projection to feasibility is extensible to any finite number of closed convex sets.

E.10.2.0.3 Example. Under- and over-projection. [50, §3]

Consider the following variation of alternating projection: We begin with some point $x_0 \in \mathbb{R}^n$ then project that point on convex set \mathcal{C} and then project that same point x_0 on convex set \mathcal{D} . To the first iterate we assign $x_1 = (P_{\mathcal{C}}(x_0) + P_{\mathcal{D}}(x_0))\frac{1}{2}$. More generally,

$$x_{i+1} = (P_{\mathcal{C}}(x_i) + P_{\mathcal{D}}(x_i))\frac{1}{2}, \quad i = 0, 1, 2, \dots \quad (1950)$$

Because the Cartesian product of convex sets remains convex, (§2.1.8) we can reformulate this problem.

Consider the convex set

$$\mathcal{S} \triangleq \begin{bmatrix} \mathcal{C} \\ \mathcal{D} \end{bmatrix} \quad (1951)$$

representing Cartesian product $\mathcal{C} \times \mathcal{D}$. Now, those two projections $P_{\mathcal{C}}$ and $P_{\mathcal{D}}$ are equivalent to one projection on the Cartesian product; *id est*,

$$P_{\mathcal{S}} \left(\begin{bmatrix} x_i \\ x_i \end{bmatrix} \right) = \begin{bmatrix} P_{\mathcal{C}}(x_i) \\ P_{\mathcal{D}}(x_i) \end{bmatrix} \quad (1952)$$

Define the subspace

$$\mathcal{R} \triangleq \left\{ v \in \begin{bmatrix} \mathbb{R}^n \\ \mathbb{R}^n \end{bmatrix} \mid [I \quad -I]v = \mathbf{0} \right\} \quad (1953)$$

By the results in Example [E.5.0.0.6](#)

$$P_{\mathcal{RS}}\left(\begin{bmatrix} x_i \\ x_i \end{bmatrix}\right) = P_{\mathcal{R}}\left(\begin{bmatrix} P_{\mathcal{C}}(x_i) \\ P_{\mathcal{D}}(x_i) \end{bmatrix}\right) = \begin{bmatrix} P_{\mathcal{C}}(x_i) + P_{\mathcal{D}}(x_i) \\ P_{\mathcal{C}}(x_i) + P_{\mathcal{D}}(x_i) \end{bmatrix} \frac{1}{2} \quad (1954)$$

This means the proposed variation of alternating projection is equivalent to an alternation of projection on convex sets \mathcal{S} and \mathcal{R} . If \mathcal{S} and \mathcal{R} intersect, these iterations will converge to a point in their intersection; hence, to a point in the intersection of \mathcal{C} and \mathcal{D} .

We need not apply equal weighting to the projections, as supposed in (1950). In that case, definition of \mathcal{R} would change accordingly. \square

E.10.2.1 Relative measure of convergence

Inspired by Fejér monotonicity, the alternating projection algorithm from the example of convergence illustrated by Figure [149](#) employs a redundant sequence: The first sequence (indexed by j) estimates point $(\prod_{j=1}^{\infty} \prod_{k=1}^L P_k)b$ in the presumably nonempty intersection of L convex sets, then the quantity

$$\left\| x_i - \left(\prod_{j=1}^{\infty} \prod_{k=1}^L P_k \right) b \right\| \quad (1955)$$

in second sequence x_i is observed per iteration i for convergence. *A priori* knowledge of a feasible point (1940) is both impractical and antithetical. We need another measure:

Nonexpansivity implies

$$\left\| \left(\prod_{\ell=1}^L P_{\ell} \right) x_{k,i-1} - \left(\prod_{\ell=1}^L P_{\ell} \right) x_{ki} \right\| = \|x_{ki} - x_{k,i+1}\| \leq \|x_{k,i-1} - x_{ki}\| \quad (1956)$$

where

$$x_{ki} \triangleq P_k x_{k+1,i} \in \mathbb{R}^n, \quad x_{L+1,i} \triangleq x_{1,i-1} \quad (1957)$$

x_{ki} represents unique minimum-distance projection of $x_{k+1,i}$ on convex set k at iteration i . So a good convergence measure is total monotonic sequence

$$\varepsilon_i \triangleq \sum_{k=1}^L \|x_{ki} - x_{k,i+1}\|, \quad i=0, 1, 2, \dots \quad (1958)$$

where $\lim_{i \rightarrow \infty} \varepsilon_i = 0$ whether or not the intersection is nonempty.

E.10.2.1.1 Example. *Affine subset \cap positive semidefinite cone.*

Consider the problem of finding $X \in \mathbb{S}^n$ that satisfies

$$X \succeq 0, \quad \langle A_j, X \rangle = b_j, \quad j=1 \dots m \quad (1959)$$

given nonzero $A_j \in \mathbb{S}^n$ and real b_j . Here we take \mathcal{C}_1 to be the positive semidefinite cone \mathbb{S}_+^n while \mathcal{C}_2 is the affine subset of \mathbb{S}^n

$$\begin{aligned} \mathcal{C}_2 &= \mathcal{A} \triangleq \{X \mid \text{tr}(A_j X) = b_j, \quad j=1 \dots m\} \subseteq \mathbb{S}^n \\ &= \{X \mid \begin{bmatrix} \text{svec}(A_1)^T \\ \vdots \\ \text{svec}(A_m)^T \end{bmatrix} \text{svec} X = b\} \\ &\triangleq \{X \mid A \text{svec} X = b\} \end{aligned} \quad (1960)$$

where $b = [b_j] \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n(n+1)/2}$, and symmetric vectorization svec is defined by (50). Projection of iterate $X_i \in \mathbb{S}^n$ on \mathcal{A} is: (§E.5.0.0.6)

$$P_2 \text{svec} X_i = \text{svec} X_i - A^\dagger (A \text{svec} X_i - b) \quad (1961)$$

Euclidean distance from X_i to \mathcal{A} is therefore

$$\text{dist}(X_i, \mathcal{A}) = \|X_i - P_2 X_i\|_F = \|A^\dagger (A \text{svec} X_i - b)\|_2 \quad (1962)$$

Projection of $P_2 X_i \triangleq \sum_j \lambda_j q_j q_j^T$ on the positive semidefinite cone (§7.1.2) is found from its eigen decomposition (§A.5.2);

$$P_1 P_2 X_i = \sum_{j=1}^n \max\{0, \lambda_j\} q_j q_j^T \quad (1963)$$

Distance from $P_2 X_i$ to the positive semidefinite cone is therefore

$$\text{dist}(P_2 X_i, \mathbb{S}_+^n) = \|P_2 X_i - P_1 P_2 X_i\|_F = \sqrt{\sum_{j=1}^n \min\{0, \lambda_j\}^2} \quad (1964)$$

When the intersection is empty $\mathcal{A} \cap \mathbb{S}_+^n = \emptyset$, the iterates converge to that positive semidefinite matrix closest to \mathcal{A} in the Euclidean sense. Otherwise, convergence is to some point in the nonempty intersection.

Barvinok (§2.9.3.0.1) shows that if a point feasible with (1959) exists, then there exists an $X \in \mathcal{A} \cap \mathbb{S}_+^n$ such that

$$\text{rank } X \leq \left\lfloor \frac{\sqrt{8m+1}-1}{2} \right\rfloor \quad (246)$$

□

E.10.2.1.2 Example. *Semidefinite matrix completion.*

Continuing Example E.10.2.1.1: When $m \leq n(n+1)/2$ and the A_j matrices are distinct members of the standard orthonormal basis $\{E_{\ell q} \in \mathbb{S}^n\}$ (53)

$$\{A_j \in \mathbb{S}^n, j=1 \dots m\} \subseteq \{E_{\ell q}\} = \left\{ \begin{array}{ll} e_\ell e_\ell^T, & \ell = q = 1 \dots n \\ \frac{1}{\sqrt{2}}(e_\ell e_q^T + e_q e_\ell^T), & 1 \leq \ell < q \leq n \end{array} \right\} \quad (1965)$$

and when the constants b_j are set to constrained entries of variable $X \triangleq [X_{\ell q}] \in \mathbb{S}^n$

$$\{b_j, j=1 \dots m\} \subseteq \left\{ \begin{array}{ll} X_{\ell q}, & \ell = q = 1 \dots n \\ X_{\ell q} \sqrt{2}, & 1 \leq \ell < q \leq n \end{array} \right\} = \{\langle X, E_{\ell q} \rangle\} \quad (1966)$$

then the equality constraints in (1959) fix individual entries of $X \in \mathbb{S}^n$. Thus the feasibility problem becomes a *positive semidefinite matrix completion problem*. Projection of iterate $X_i \in \mathbb{S}^n$ on \mathcal{A} simplifies to (*confer*(1961))

$$P_2 \text{svec } X_i = \text{svec } X_i - A^T(A \text{svec } X_i - b) \quad (1967)$$

From this we can see that orthogonal projection is achieved simply by setting corresponding entries of $P_2 X_i$ to the known entries of X , while the entries of $P_2 X_i$ remaining are set to corresponding entries of the current iterate X_i .

Using this technique, we find a positive semidefinite completion for

$$\begin{bmatrix} 4 & 3 & ? & 2 \\ 3 & 4 & 3 & ? \\ ? & 3 & 4 & 3 \\ 2 & ? & 3 & 4 \end{bmatrix} \quad (1968)$$

Initializing the unknown entries to 0, they all converge geometrically to 1.5858 (rounded) after about 42 iterations.




Figure 150: Distance (*confer*(1964)) between positive semidefinite cone and iterate (1967) in affine subset \mathcal{A} (1960) for Laurent's completion problem; initially, decreasing geometrically.

Laurent gives a problem for which no positive semidefinite completion exists: [208]

$$\begin{bmatrix} 1 & 1 & ? & 0 \\ 1 & 1 & 1 & ? \\ ? & 1 & 1 & 1 \\ 0 & ? & 1 & 1 \end{bmatrix} \quad (1969)$$

Initializing unknowns to 0, by alternating projection we find the constrained matrix closest to the positive semidefinite cone,

$$\begin{bmatrix} 1 & 1 & 0.5454 & 0 \\ 1 & 1 & 1 & 0.5454 \\ 0.5454 & 1 & 1 & 1 \\ 0 & 0.5454 & 1 & 1 \end{bmatrix} \quad (1970)$$

and we find the positive semidefinite matrix closest to the affine subset \mathcal{A} (1960):

$$\begin{bmatrix} 1.0521 & 0.9409 & 0.5454 & 0.0292 \\ 0.9409 & 1.0980 & 0.9451 & 0.5454 \\ 0.5454 & 0.9451 & 1.0980 & 0.9409 \\ 0.0292 & 0.5454 & 0.9409 & 1.0521 \end{bmatrix} \quad (1971)$$

These matrices (1970) and (1971) attain the Euclidean distance $\text{dist}(\mathcal{A}, \mathbb{S}_+^n)$. Convergence is illustrated in Figure 150. \square




Figure 151: \mathcal{H}_1 and \mathcal{H}_2 are the same halfspaces as in Figure 146. Dykstra's alternating projection algorithm generates the alternations $b, x_{21}, x_{11}, x_{22}, x_{12}, x_{12} \dots x_{12}$. The path illustrated from b to x_{12} in \mathbb{R}^2 terminates at the desired result, Pb in Figure 146. The $\{y_{ki}\}$ correspond to the first two difference vectors drawn (in the first iteration $i=1$), then oscillate between zero and a negative vector thereafter. These alternations are not so robust in presence of noise as for the example in Figure 145.

E.10.3 Optimization and projection

Unique projection on the nonempty intersection of arbitrary convex sets to find the closest point therein is a convex optimization problem. The first successful application of alternating projection to this problem is attributed to Dykstra [110] [54] who in 1983 provided an elegant algorithm that prevails today. In 1988, Han [157] rediscovered the algorithm and provided a primal–dual convergence proof. A synopsis of the history of alternating projection^{E.21} can be found in [56] where it becomes apparent that Dykstra's work is seminal.

^{E.21}For a synopsis of alternating projection applied to distance geometry, see [313, §3.1].

E.10.3.1 Dykstra's algorithm

Assume we are given some point $b \in \mathbb{R}^n$ and closed convex sets $\{\mathcal{C}_k \subset \mathbb{R}^n \mid k=1 \dots L\}$. Let $x_{ki} \in \mathbb{R}^n$ and $y_{ki} \in \mathbb{R}^n$ respectively denote a *primal* and *dual vector* (whose meaning can be deduced from Figure 151 and Figure 152) associated with set k at iteration i . Initialize

$$y_{k0} = 0 \quad \forall k=1 \dots L \quad \text{and} \quad x_{1,0} = b \quad (1972)$$

Denoting by $P_k t$ the unique minimum-distance projection of t on \mathcal{C}_k , and for convenience $x_{L+1,i} = x_{1,i-1}$ (1957), iterate x_{1i} calculation proceeds:^{E.22}

$$\begin{aligned} & \text{for } i=1, 2, \dots \text{until convergence } \{ \\ & \quad \text{for } k=L \dots 1 \{ \\ & \quad \quad t = x_{k+1,i} - y_{k,i-1} \\ & \quad \quad x_{ki} = P_k t \\ & \quad \quad y_{ki} = P_k t - t \\ & \quad \quad \} \\ & \quad \} \end{aligned} \quad (1973)$$

Assuming a nonempty intersection, then the iterates converge to the unique minimum-distance projection of point b on that intersection; [92, §9.24]

$$Pb = \lim_{i \rightarrow \infty} x_{1i} \quad (1974)$$

In the case all the \mathcal{C}_k are affine, then calculation of y_{ki} is superfluous and the algorithm becomes identical to alternating projection. [92, §9.26] [128, §1] Dykstra's algorithm is so simple, elegant, and represents such a tiny increment in computational intensity over alternating projection, it is nearly always arguably cost effective.

E.10.3.2 Normal cone

Glunt [135, §4] observes that the overall effect of Dykstra's iterative procedure is to drive t toward the translated normal cone to $\bigcap \mathcal{C}_k$ at the solution Pb (translated to Pb). The normal cone gets its name from its graphical construction; which is, loosely speaking, to draw the outward-normals at Pb (Definition E.9.1.0.1) to all the convex sets \mathcal{C}_k touching Pb . The relative interior of the normal cone subtends these normal vectors.

^{E.22}We reverse order of projection ($k=L \dots 1$) in the algorithm for continuity of exposition.




Figure 152: Two examples (truncated): Normal cone to $\mathcal{H}_1 \cap \mathcal{H}_2$ at the origin, and at point Pb on the boundary. \mathcal{H}_1 and \mathcal{H}_2 are the same halfspaces from Figure 151. The normal cone at the origin $\mathcal{K}_{\mathcal{H}_1 \cap \mathcal{H}_2}^\perp(\mathbf{0})$ is simply $-\mathcal{K}^*$.

E.10.3.2.1 Definition. *Normal cone.* [237] [37, p.261] [177, §A.5.2] [48, §2.1] [270, §3] [271, p.15] The normal cone to any set $\mathcal{S} \subseteq \mathbb{R}^n$ at any particular point $a \in \mathbb{R}^n$ is defined as the closed cone

$$\mathcal{K}_{\mathcal{S}}^\perp(a) \triangleq \{z \in \mathbb{R}^n \mid z^\top(y - a) \leq 0 \quad \forall y \in \mathcal{S}\} = -(\mathcal{S} - a)^* \quad (1975)$$

an intersection of halfspaces about the origin in \mathbb{R}^n , hence convex regardless of convexity of \mathcal{S} ; it is the negative dual cone to the translate $\mathcal{S} - a$; the set of all vectors normal to \mathcal{S} at a . \triangle

Examples of normal cone construction are illustrated in Figure 62 and Figure 152: The normal cone at the origin in Figure 152 is the vector sum (§2.1.8) of two normal cones; [48, §3.3, exer.10] for $\mathcal{H}_1 \cap \text{int } \mathcal{H}_2 \neq \emptyset$

$$\mathcal{K}_{\mathcal{H}_1 \cap \mathcal{H}_2}^\perp(\mathbf{0}) = \mathcal{K}_{\mathcal{H}_1}^\perp(\mathbf{0}) + \mathcal{K}_{\mathcal{H}_2}^\perp(\mathbf{0}) \quad (1976)$$

This formula applies more generally to other points in the intersection.

The normal cone to any affine set \mathcal{A} at $\alpha \in \mathcal{A}$, for example, is the orthogonal complement of $\mathcal{A} - \alpha$. Projection of any point in the translated





Figure 153: A few renderings (including next page) of normal cone $\mathcal{K}_{\mathcal{E}^3}^\perp$ to ellipsope \mathcal{E}^3 (Figure 109) at point $\mathbf{11}^T$, projected on \mathbb{R}^3 . In [209, fig.2], normal cone is claimed circular in this dimension (severe numerical artifacts corrupt boundary and make interior corporeal, drawn truncated).



normal cone $\mathcal{K}_{\mathcal{C}}^{\perp}(a \in \mathcal{C}) + a$ on convex set \mathcal{C} is identical to a ; in other words, point a is that point in \mathcal{C} closest to any point belonging to the translated normal cone $\mathcal{K}_{\mathcal{C}}^{\perp}(a) + a$; *e.g.*, Theorem [E.4.0.0.1](#).

When set \mathcal{S} is a convex cone \mathcal{K} , then the normal cone to \mathcal{K} at the origin

$$\mathcal{K}_{\mathcal{K}}^{\perp}(\mathbf{0}) = -\mathcal{K}^* \quad (1977)$$

is the negative dual cone. Any point belonging to $-\mathcal{K}^*$, projected on \mathcal{K} , projects on the origin. More generally, [[92](#), §4.5]

$$\mathcal{K}_{\mathcal{K}}^{\perp}(a) = -(\mathcal{K} - a)^* \quad (1978)$$

$$\mathcal{K}_{\mathcal{K}}^{\perp}(a \in \mathcal{K}) = -\mathcal{K}^* \cap a^{\perp} \quad (1979)$$

The normal cone to $\bigcap \mathcal{C}_k$ at Pb in Figure [146](#) is ray $\{\xi(b - Pb) \mid \xi \geq 0\}$ illustrated in Figure [152](#). Applying Dykstra's algorithm to that example, convergence to the desired result is achieved in two iterations as illustrated in Figure [151](#). Yet applying Dykstra's algorithm to the example in Figure [145](#) does not improve rate of convergence, unfortunately, because the given point b and all the alternating projections already belong to the translated normal cone at the vertex of intersection.

E.10.3.3 speculation

From these few examples we surmise, unique minimum-distance projection on *blunt* (not sharp or acute, informally) polyhedral cones having nonempty interior may be found by Dykstra's algorithm in few iterations.

Appendix F

Notation and a few definitions

- b (italic $abcdefghijklmnopqrstuvwxy$) column vector, scalar, logical condition
- b_i i^{th} entry of vector $b = [b_i, i = 1 \dots n]$ or i^{th} b vector from a set or list $\{b_j, j = 1 \dots n\}$ or i^{th} iterate of vector b
- $b_{i:j}$ or $b(i:j)$, truncated vector comprising i^{th} through j^{th} entry of vector b
- $b_k(i:j)$ or $b_{i:j,k}$, truncated vector comprising i^{th} through j^{th} entry of vector b_k
- b^T vector transpose
- b^H Hermitian (complex conjugate) transpose b^{*T}
- A^{-2T} matrix transpose of squared inverse
- A^{T1} first of various transpositions of a cubix or quartix A (p.622, p.627)
- A matrix, scalar, or logical condition
(italic $ABCDEFGHIJKLMNPOQRSTUVWXYZ$)
- skinny* a skinny matrix; meaning, more rows than columns: $\begin{bmatrix} \\ \\ \\ \end{bmatrix}$. When there are more equations than unknowns, we say that the system $Ax = b$ is overdetermined. [139, §5.3]
- fat* a fat matrix; meaning, more columns than rows: $\begin{bmatrix} & & \end{bmatrix}$
underdetermined

\mathcal{A}	some set (calligraphic $ABCDEFGHIJKLMNOPQRSTUVWXYZ$)
$\mathcal{F}(\mathcal{C} \ni A)$	smallest face (152) that contains element A of set \mathcal{C}
$\mathcal{G}(\mathcal{K})$	generators (§2.8.1.2) of set \mathcal{K} ; any collection of points and directions whose hull constructs \mathcal{K}
A^{-1}	inverse of matrix A
A^\dagger	Moore-Penrose pseudoinverse of matrix A
$\sqrt{\quad}$	positive square root
$\sqrt[\ell]{\quad}$	positive ℓ^{th} root
$A^{1/2}$ and \sqrt{A}	$A^{1/2}$ is any matrix such that $A^{1/2}A^{1/2} = A$. For $A \in \mathbb{S}_+^n$, $\sqrt{A} \in \mathbb{S}_+^n$ is unique and $\sqrt{A}\sqrt{A} = A$. [48, §1.2] (§A.5.2.1)
$\sqrt[D]{\quad}$	$\triangleq [\sqrt{d_{ij}}]$. (1262) <i>Hadamard positive square root</i> : $D = \sqrt[D]{\quad} \circ \sqrt[D]{\quad}$
\mathfrak{A}	Euler Fraktur $\mathfrak{A} \mathfrak{B} \mathfrak{C} \mathfrak{D} \mathfrak{E} \mathfrak{F} \mathfrak{G} \mathfrak{H} \mathfrak{I} \mathfrak{J} \mathfrak{K} \mathfrak{L} \mathfrak{M} \mathfrak{N} \mathfrak{O} \mathfrak{P} \mathfrak{Q} \mathfrak{R} \mathfrak{S} \mathfrak{T} \mathfrak{U} \mathfrak{V} \mathfrak{W} \mathfrak{X} \mathfrak{Y} \mathfrak{Z}$
\mathcal{E}	member of elliptope \mathcal{E}_t (1012) parametrized by scalar t
\mathcal{E}	elliptope (991)
E	elementary matrix
E_{ij}	member of standard orthonormal basis for symmetric (53) or symmetric hollow (68) matrices
A_{ij}	or $A(i, j)$, ij^{th} entry of matrix $A = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} \cdot & \cdot & \cdot \end{matrix} & \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} \end{matrix}$ or rank-one matrix $a_i a_j^T$ (§4.7)
A_i	i^{th} matrix from a set or i^{th} principal submatrix or i^{th} iterate of A
$A(i, :)$	i^{th} row of matrix A
$A(:, j)$	j^{th} column of matrix A [139, §1.1.8]

$A_{i:j,k:\ell}$	or $A(i:j, k:\ell)$, submatrix taken from i^{th} through j^{th} row and k^{th} through ℓ^{th} column
$A(ij)$	A is a function of i and j
<i>e.g.</i>	<i>exempli gratia</i> , from the Latin meaning <i>for sake of example</i>
no.	<i>number</i> , from the Latin <i>numero</i>
a.i.	affinely independent (§2.4.2.3)
c.i.	conically independent (§2.10)
l.i.	linearly independent
w.r.t	<i>with respect to</i>
a.k.a	<i>also known as</i>
Re	real part
Im	imaginary part
i or j	$\sqrt{-1}$
\subset \supset \cap \cup	standard set theory, <i>subset</i> , <i>superset</i> , <i>intersection</i> , <i>union</i>
\in	membership, <i>element belongs to</i> , or <i>element is a member of</i>
\ni	membership, <i>contains</i> as in $\mathcal{C} \ni y$ (\mathcal{C} contains element y)
\ni	<i>such that</i>
\exists	<i>there exists</i>
\therefore	<i>therefore</i>
\forall	<i>for all</i> , or <i>over all</i>
\propto	<i>proportional to</i>
∞	infinity
\equiv	<i>equivalent to</i>

- \triangleq defined equal to, or equal by definition
- \approx approximately equal to
- \simeq isomorphic to or with
- \cong congruent to or with
- Hadamard quotient as in, for $x, y \in \mathbb{R}^n$, $\frac{x}{y} \triangleq [x_i/y_i, i=1 \dots n] \in \mathbb{R}^n$
- Hadamard product of matrices: $x \circ y \triangleq [x_i y_i, i=1 \dots n] \in \mathbb{R}^n$
- ⊗ Kronecker product of matrices (§D.1.2.1)
- ⊕ vector sum of sets $\mathcal{X} = \mathcal{Y} \oplus \mathcal{Z}$ where every element $x \in \mathcal{X}$ has unique expression $x = y + z$ where $y \in \mathcal{Y}$ and $z \in \mathcal{Z}$; [271, p.19] then the summands are algebraic complements. $\mathcal{X} = \mathcal{Y} \oplus \mathcal{Z} \Rightarrow \mathcal{X} = \mathcal{Y} + \mathcal{Z}$. Now assume \mathcal{Y} and \mathcal{Z} are nontrivial subspaces. $\mathcal{X} = \mathcal{Y} + \mathcal{Z} \Rightarrow \mathcal{X} = \mathcal{Y} \oplus \mathcal{Z} \Leftrightarrow \mathcal{Y} \cap \mathcal{Z} = \mathbf{0}$ [272, §1.2] [92, §5.8]. Each element from a vector sum (+) of subspaces has a unique representation (⊕) when a basis from each subspace is linearly independent of bases from all the other subspaces.
- ⊖ likewise, the vector difference of sets
- ⊞ orthogonal vector sum of sets $\mathcal{X} = \mathcal{Y} \boxplus \mathcal{Z}$ where every element $x \in \mathcal{X}$ has unique orthogonal expression $x = y + z$ where $y \in \mathcal{Y}$, $z \in \mathcal{Z}$, and $y \perp z$. [293, p.51] $\mathcal{X} = \mathcal{Y} \boxplus \mathcal{Z} \Rightarrow \mathcal{X} = \mathcal{Y} + \mathcal{Z}$. If $\mathcal{Z} \subseteq \mathcal{Y}^\perp$ then $\mathcal{X} = \mathcal{Y} \boxplus \mathcal{Z} \Leftrightarrow \mathcal{X} = \mathcal{Y} \oplus \mathcal{Z}$. [92, §5.8] If $\mathcal{Z} = \mathcal{Y}^\perp$ then the summands are orthogonal complements.
- ± plus or minus or plus and minus
- ⊥ as in $A \perp B$ meaning A is orthogonal to B (and vice versa), where A and B are sets, vectors, or matrices. When A and B are vectors (or matrices under Frobenius norm), $A \perp B \Leftrightarrow \langle A, B \rangle = 0 \Leftrightarrow \|A + B\|^2 = \|A\|^2 + \|B\|^2$
- \ as in $\setminus \mathcal{A}$ means logical not \mathcal{A} , or relative complement of set \mathcal{A} ; id est, $\setminus \mathcal{A} = \{x \notin \mathcal{A}\}$; e.g., $\mathcal{B} \setminus \mathcal{A} \triangleq \{x \in \mathcal{B} \mid x \notin \mathcal{A}\} \equiv \mathcal{B} \cap \setminus \mathcal{A}$

- \Rightarrow or \Leftarrow sufficiency or necessity, *implies*; e.g., $A \Rightarrow B \Leftrightarrow \neg A \Leftarrow \neg B$
 \Leftrightarrow *if and only if* (iff) or *corresponds with* or *necessary and sufficient* or *logical equivalence*
is as in A *is* B means $A \Rightarrow B$; conventional usage of English language imposed by logicians
 \nRightarrow or \nLeftarrow *does not imply*
 \leftarrow *is replaced with*; substitution, assignment
 \rightarrow *goes to*, or *approaches*, or *maps to*
 $t \rightarrow 0^+$ t *goes to 0 from above*; meaning, *from the positive* [177, p.2]
 \vdots \ddots \dots as in $1 \dots 1$ meaning *continuation*, e.g., a sequence of ones in a row
 \dots as in $i=1 \dots N$ meaning, i is a sequence of successive integers beginning with 1 and ending with N
 $:$ as in $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ meaning f *is a mapping*, or sequence of successive integers specified by bounds as in $i:j$ (if $j < i$ then sequence is descending)
 $f : \mathcal{M} \rightarrow \mathcal{R}$ meaning f *is a mapping from ambient space* \mathcal{M} *to ambient* \mathcal{R} , not necessarily denoting either domain or range
 $|$ as in $f(x) | x \in \mathcal{C}$ means *with the condition(s)* or *such that* or *evaluated for*, or as in $\{f(x) | x \in \mathcal{C}\}$ means *evaluated for each and every* x *belonging to set* \mathcal{C}
 $g|_{x_p}$ *expression* g *evaluated at* x_p
 A, B as in, for example, $A, B \in \mathbb{S}^N$ means $A \in \mathbb{S}^N$ and $B \in \mathbb{S}^N$
 (A, B) *open interval* between A and B in \mathbb{R} , or *variable pair* perhaps of disparate dimension
 $[A, B]$ *closed interval* or *line segment* between A and B in \mathbb{R}
 $()$ *hierarchal, parenthetical, optional*

$\{ \}$	curly braces denote a <i>set</i> or <i>list</i> , e.g., $\{Xa \mid a \succeq 0\}$ the set comprising Xa evaluated for each and every $a \succeq 0$ where membership of a to some space is implicit, a <i>union</i>
$\langle \rangle$	angle brackets denote <i>vector inner-product</i> (27) (32)
$[\]$	matrix or vector, or quote insertion, or citation
$[d_{ij}]$	matrix whose ij^{th} entry is d_{ij}
$[x_i]$	vector whose i^{th} entry is x_i
x_p	particular value of x
x_0	particular instance of x , or initial value of a sequence x_i
x_1	first entry of vector x , or first element of a set or list $\{x_i\}$
x_ε	<i>extreme point</i>
x_+	vector x whose negative entries are replaced with 0; $x_+ = \frac{1}{2}(x + x)$ (481) or <i>clipped vector</i> x or <i>nonnegative part of</i> x
x_-	$x_- \triangleq \frac{1}{2}(x - x)$ or <i>nonpositive part of</i> $x = x_+ + x_-$
\tilde{x}	known data
x^*	optimal value of variable x
x^*	<i>complex conjugate</i> or <i>dual variable</i> or <i>extreme direction of dual cone</i>
f^*	<i>convex conjugate function</i>
$P_C x$ or Px	projection of point x on set \mathcal{C} , P is operator or idempotent matrix
$P_k x$	projection of point x on set \mathcal{C}_k or on range of implicit vector
$\delta(A)$	(§A.1) <i>vector made from the main diagonal of</i> A if A is a matrix; otherwise, <i>diagonal matrix made from vector</i> A
$\delta^2(A)$	$\equiv \delta(\delta(A))$. For vector or diagonal matrix Λ , $\delta^2(\Lambda) = \Lambda$
$\delta(A)^2$	$= \delta(A)\delta(A)$ where A is a vector

$\lambda_i(X)$	i^{th} entry of vector λ is function of X
$\lambda(X)_i$	i^{th} entry of vector-valued function of X
$\lambda(A)$	vector of eigenvalues of matrix A , (1374) typically arranged in nonincreasing order
$\sigma(A)$	vector of singular values of matrix A (always arranged in nonincreasing order), or <i>support function</i>
Σ	diagonal matrix of singular values, not necessarily square
\sum	sum
$\pi(\gamma)$	nonlinear <i>permutation operator</i> (or <i>presorting function</i>) arranges vector γ into nonincreasing order (§7.1.3)
Ξ	permutation matrix
Π	doublet or permutation matrix or operator
\prod	product
$\psi(Z)$	signum-like <i>step function</i> that returns a scalar for matrix argument (650), it returns a vector for vector argument (1477)
D	symmetric hollow matrix of distance-square, or <i>Euclidean distance matrix</i>
\mathbf{D}	Euclidean distance matrix operator
$\mathbf{D}^T(X)$	adjoint operator
$\mathbf{D}(X)^T$	transpose of $\mathbf{D}(X)$
$\mathbf{D}^{-1}(X)$	inverse operator
$\mathbf{D}(X)^{-1}$	inverse of $\mathbf{D}(X)$
D^*	optimal value of variable D
D^*	dual to variable D
D°	polar variable D

∂	<i>partial derivative</i> or <i>matrix of distance-square squared</i> (1300) or <i>boundary</i> of set \mathcal{K} as in $\partial\mathcal{K}$
∂y	partial differential of y
$\sqrt{d_{ij}}$	(absolute) distance scalar
d_{ij}	distance-square scalar, EDM entry
\mathbf{V}	geometric centering operator, $\mathbf{V}(D) = -VDV^{\frac{1}{2}}$ (911)
$\mathbf{V}_{\mathcal{N}}$	$\mathbf{V}_{\mathcal{N}}(D) = -V_{\mathcal{N}}^{\text{T}}DV_{\mathcal{N}}$ (925)
V	$N \times N$ symmetric elementary, auxiliary, projector, geometric centering matrix, $\mathcal{R}(V) = \mathcal{N}(\mathbf{1}^{\text{T}})$, $\mathcal{N}(V) = \mathcal{R}(\mathbf{1})$, $V^2 = V$ (§B.4.1)
$V_{\mathcal{N}}$	$N \times N - 1$ Schoenberg auxiliary matrix, $\mathcal{R}(V_{\mathcal{N}}) = \mathcal{N}(\mathbf{1}^{\text{T}})$, $\mathcal{N}(V_{\mathcal{N}}^{\text{T}}) = \mathcal{R}(\mathbf{1})$ (§B.4.2)
$V_{\mathcal{X}}$	$V_{\mathcal{X}}V_{\mathcal{X}}^{\text{T}} \equiv V^{\text{T}}X^{\text{T}}XV$ (1105)
X	point list ((69) having cardinality N) arranged columnar in $\mathbb{R}^{n \times N}$, or set of generators, or extreme directions, or matrix variable
G	Gram matrix $X^{\text{T}}X$
r	affine dimension
n	Euclidean dimension of list X , or integer
N	cardinality of list X , or integer
dom	function domain
<i>on</i>	<i>function $f(x)$ on \mathcal{A}</i> means \mathcal{A} is dom f , or <i>projection of x on \mathcal{A}</i> means \mathcal{A} is Euclidean body on which projection of x is made
<i>onto</i>	<i>function $f(x)$ maps onto \mathcal{M}</i> means f over its domain is a surjection with respect to \mathcal{M}
<i>one-to-one</i>	injective map or unique correspondence between sets
epi	function epigraph

span	as in $\text{span } A = \mathcal{R}(A) = \{Ax \mid x \in \mathbb{R}^n\}$ when A is a matrix
$\mathcal{R}(A)$	the subspace: <i>range of A</i> , or span basis $\mathcal{R}(A)$; $\mathcal{R}(A) \perp \mathcal{N}(A^T)$
basis $\mathcal{R}(A)$	<i>columnar basis for range of A</i> , or a <i>minimal set constituting generators for the vertex-description of $\mathcal{R}(A)$</i> , or a <i>linearly independent set of vectors spanning $\mathcal{R}(A)$</i>
$\mathcal{N}(A)$	the subspace: <i>nullspace of A</i> ; $\mathcal{N}(A) \perp \mathcal{R}(A^T)$
\mathbb{R}^n	Euclidean n -dimensional real vector space, $\mathbb{R}^0 = \mathbf{0}$, $\mathbb{R}^1 = \mathbb{R}$ (nonnegative integer n)
$\mathbb{R}^{m \times n}$	Euclidean vector space of m by n dimensional real matrices
\times	Cartesian product. $\mathbb{R}^{m \times n - m} \triangleq \mathbb{R}^{m \times (n-m)}$
$\begin{bmatrix} \mathbb{R}^m \\ \mathbb{R}^n \end{bmatrix}$	$\mathbb{R}^m \times \mathbb{R}^n = \mathbb{R}^{m+n}$
\mathbb{C}^n or $\mathbb{C}^{n \times n}$	Euclidean complex vector space of respective dimension n and $n \times n$
\mathbb{R}_+^n or $\mathbb{R}_+^{n \times n}$	nonnegative orthant in Euclidean vector space of respective dimension n and $n \times n$
\mathbb{R}_-^n or $\mathbb{R}_-^{n \times n}$	nonpositive orthant in Euclidean vector space of respective dimension n and $n \times n$
\mathbb{S}^n	subspace comprising all (real) symmetric $n \times n$ matrices, the <i>symmetric matrix subspace</i>
$\mathbb{S}^{n\perp}$	orthogonal complement of \mathbb{S}^n in $\mathbb{R}^{n \times n}$, the antisymmetric matrices
\mathbb{S}_+^n	convex cone comprising all (real) symmetric positive semidefinite $n \times n$ matrices, the <i>positive semidefinite cone</i>
$\text{int } \mathbb{S}_+^n$	interior of convex cone comprising all (real) symmetric positive semidefinite $n \times n$ matrices; <i>id est</i> , positive definite matrices
$\mathbb{S}_+(\rho)$	convex set of all positive semidefinite $n \times n$ matrices whose rank equals or exceeds ρ

\mathbb{EDM}^N	cone of $N \times N$ Euclidean distance matrices in the symmetric hollow subspace
$\sqrt{\mathbb{EDM}^N}$	nonconvex cone of $N \times N$ Euclidean absolute distance matrices in the symmetric hollow subspace (§6.3)
PSD	positive semidefinite
SDP	semidefinite program
SVD	singular value decomposition
SNR	signal to noise ratio
dB	decibel
EDM	Euclidean distance matrix
\mathbb{S}_1^n	subspace comprising all symmetric $n \times n$ matrices having all zeros in first row and column (1889)
\mathbb{S}_h^n	subspace comprising all symmetric hollow $n \times n$ matrices ($\mathbf{0}$ main diagonal), the <i>symmetric hollow subspace</i> (60)
$\mathbb{S}_h^{n\perp}$	orthogonal complement of \mathbb{S}_h^n in \mathbb{S}^n (61), the set of all diagonal matrices
\mathbb{S}_c^n	subspace comprising all geometrically centered symmetric $n \times n$ matrices; <i>geometric center subspace</i> $\mathbb{S}_c^N \triangleq \{Y \in \mathbb{S}^N \mid Y\mathbf{1} = \mathbf{0}\}$ (1885)
$\mathbb{S}_c^{n\perp}$	orthogonal complement of \mathbb{S}_c^n in \mathbb{S}^n (1887)
$\mathbb{R}_c^{m \times n}$	subspace comprising all geometrically centered $m \times n$ matrices
X^\perp	basis $\mathcal{N}(X^T)$
x^\perp	$\mathcal{N}(x^T)$, $\{y \mid x^T y = 0\}$
$\mathcal{R}(P)^\perp$	$\mathcal{N}(P^T)$
\mathcal{R}^\perp	$= \{y \in \mathbb{R}^n \mid \langle x, y \rangle = 0 \ \forall x \in \mathcal{R}\}$ (338); orthogonal complement of $\mathcal{R} \subseteq \mathbb{R}^n$
\mathcal{K}^\perp	normal cone
\mathcal{K}	<i>cone</i>

\mathcal{K}^*	<i>dual cone</i>
\mathcal{K}°	<i>polar cone; $\mathcal{K}^* = -\mathcal{K}^\circ$</i>
$\mathcal{K}_{\mathcal{M}+}$	<i>monotone nonnegative cone</i>
$\mathcal{K}_{\mathcal{M}}$	<i>monotone cone</i>
\mathcal{K}_λ	<i>spectral cone</i>
$\mathcal{K}_{\lambda\delta}^*$	<i>cone of majorization</i>
\mathcal{H}	<i>halfspace</i>
\mathcal{H}_-	<i>halfspace described using an outward-normal (98) to the hyperplane partially bounding it</i>
\mathcal{H}_+	<i>halfspace described using an inward-normal (99) to the hyperplane partially bounding it</i>
$\partial\mathcal{H}$	<i>hyperplane; <i>id est</i>, partial boundary of halfspace</i>
$\underline{\partial\mathcal{H}}$	<i>supporting hyperplane</i>
$\underline{\partial\mathcal{H}}_-$	<i>a supporting hyperplane having outward-normal with respect to set it supports</i>
$\underline{\partial\mathcal{H}}_+$	<i>a supporting hyperplane having inward-normal with respect to set it supports</i>
\underline{d}	<i>vector of distance-square</i>
\underline{d}_{ij}	<i>lower bound on distance-square d_{ij}</i>
\overline{d}_{ij}	<i>upper bound on distance-square d_{ij}</i>
\overline{AB}	<i>closed line segment between points A and B</i>
AB	<i>matrix multiplication of A and B</i>
\overline{C}	<i>closure of set C</i>
<i>decomposition</i>	<i>orthonormal (1799) page 654, biorthogonal (1776) page 648</i>

<i>expansion</i>	<i>orthogonal</i> (1809) page 657, <i>biorthogonal</i> (363) page 176
<i>vector</i>	<i>column vector</i> in \mathbb{R}^n
<i>entry</i>	<i>scalar element or real variable constituting a vector or matrix</i>
<i>cubic</i>	member of $\mathbb{R}^{M \times N \times L}$
<i>quartic</i>	member of $\mathbb{R}^{M \times N \times L \times K}$
<i>feasible set</i>	most simply, <i>the set of all variable values satisfying all constraints of an optimization problem</i>
<i>solution set</i>	most simply, <i>the set of all optimal solutions to an optimization problem; a subset of the feasible set and not necessarily a single point</i>
<i>optimal solution</i>	solution to an optimization problem, distinguished throughout from <i>feasible solution</i> which may or may not be optimal
<i>natural order</i>	with reference to stacking columns in a vectorization means <i>a vector made from superposing column 1 on top of column 2 then superposing the result on column 3 and so on; as in a vector made from entries of the main diagonal $\delta(A)$ means taken from left to right and top to bottom</i>
<i>operator</i>	mapping to a vector space (a multidimensional function)
<i>tight</i>	with reference to a bound means <i>a bound that can be met</i> , with reference to an inequality means <i>equality is achievable</i>
g'	first derivative of possibly multidimensional function with respect to real argument
g''	second derivative with respect to real argument
$\xrightarrow{Y} dg$	first directional derivative of possibly multidimensional function g in direction $Y \in \mathbb{R}^{K \times L}$ (maintains dimensions of g)
$\xrightarrow{Y} dg^2$	second directional derivative of g in direction Y
∇	<i>gradient</i> from calculus, ∇f is shorthand for $\nabla_x f(x)$. $\nabla f(y)$ means $\nabla_y f(y)$ or <i>gradient $\nabla_x f(y)$ of $f(x)$ with respect to x evaluated at y</i>

∇^2	<i>second-order gradient</i>
Δ	distance scalar (Figure 23), or first-order difference matrix (770), or infinitesimal difference operator (§D.1.4)
Δ_{ijk}	triangle made by vertices i , j , and k
I	Roman numeral
I	identity matrix
\mathcal{I}	<i>index set</i> , a discrete set of indices
\emptyset	<i>empty set</i> , an implicit member of every set
0	real zero
$\mathbf{0}$	<i>origin</i> or vector or matrix of zeros
O	<i>sort-index matrix</i>
\mathcal{O}	<i>order of magnitude</i> information required, or <i>computational intensity</i> : $\mathcal{O}(N)$ is first order, $\mathcal{O}(N^2)$ is second, and so on
1	real one
$\mathbf{1}$	vector of ones
e_i	vector whose i^{th} entry is 1 (otherwise 0), i^{th} member of the standard basis for \mathbb{R}^n (54)
max	<i>maximum</i> [177, §0.1.1] or <i>largest element of a totally ordered set</i>
maximize x	<i>find the maximum of a function with respect to variable x</i>
arg	<i>argument</i> of operator or function, or <i>variable of optimization</i>
sup \mathcal{X}	<i>supremum</i> of totally ordered set \mathcal{X} , <i>least upper bound</i> , may or may not belong to set [177, §0.1.1]
arg sup $f(x)$	<i>argument x at supremum of function f</i> ; not necessarily unique or a member of function domain
subject to	specifies constraints to an optimization problem

min	<i>minimum</i> [177, §0.1.1] or <i>smallest element of a totally ordered set</i>
minimize x	<i>find the function minimum with respect to variable x</i>
inf \mathcal{X}	<i>infimum of totally ordered set \mathcal{X}, greatest lower bound, may or may not belong to set</i> [177, §0.1.1]
arg inf $f(x)$	<i>argument x at infimum of function f; not necessarily unique or a member of function domain</i>
iff	<i>if and only if, necessary and sufficient; also the meaning indiscriminately attached to appearance of the word “if” in the statement of a mathematical definition, [111, p.106] [228, p.4] an esoteric practice worthy of abolition because of ambiguity thus conferred</i>
rel	<i>relative</i>
int	<i>interior</i>
lim	<i>limit</i>
sgn	<i>signum function or sign</i>
round	<i>round to nearest integer</i>
mod	<i>modulus function</i>
tr	<i>matrix trace</i>
rank	<i>as in rank A, rank of matrix A; $\dim \mathcal{R}(A)$</i>
dim	<i>dimension, $\dim \mathbb{R}^n = n$, $\dim(x \in \mathbb{R}^n) = n$, $\dim \mathcal{R}(x \in \mathbb{R}^n) = 1$, $\dim \mathcal{R}(A \in \mathbb{R}^{m \times n}) = \text{rank}(A)$</i>
aff	<i>affine hull</i>
dim aff	<i>affine dimension</i>
card	<i>cardinality, number of nonzero entries $\text{card } x \triangleq \ x\ _0$ or N is cardinality of list $X \in \mathbb{R}^{n \times N}$ (p.291)</i>
conv	<i>convex hull</i>

cenv	convex envelope
cone	conic hull
content	of high-dimensional bounded polyhedron, volume in 3 dimensions, area in 2, and so on
cof	matrix of cofactors corresponding to matrix argument
dist	distance between point or set arguments; <i>e.g.</i> , $\text{dist}(x, \mathcal{B})$
vec	vectorization of $m \times n$ matrix, Euclidean dimension mn (31)
svec	vectorization of symmetric $n \times n$ matrix, Euclidean dimension $n(n+1)/2$ (50)
dvec	vectorization of symmetric hollow $n \times n$ matrix, Euclidean dimension $n(n-1)/2$ (67)
$\sphericalangle(x, y)$	<i>angle</i> between vectors x and y , or <i>dihedral angle</i> between affine subsets
\succeq	generalized inequality; <i>e.g.</i> , $A \succeq 0$ means: <i>vector or matrix A must be expressible in a biorthogonal expansion having nonnegative coordinates with respect to extreme directions of some implicit pointed closed convex cone \mathcal{K} (§2.13.2.0.1, §2.13.7.1.1), or comparison to the origin with respect to some implicit pointed closed convex cone (2.7.2.2), or (when $\mathcal{K} = \mathbb{S}_+^n$) matrix A belongs to the positive semidefinite cone of symmetric matrices (§2.9.0.1), or (when $\mathcal{K} = \mathbb{R}_+^n$) vector A belongs to the nonnegative orthant (each vector entry is nonnegative, §2.3.1.1)</i>
\succ	strict generalized inequality, membership to cone interior
$\not\succeq$	not positive definite
\geq	scalar inequality, <i>greater than or equal to</i> ; comparison of scalars, or entrywise comparison of vectors or matrices with respect to \mathbb{R}_+
<i>nonnegative</i>	for $\alpha \in \mathbb{R}^n$, $\alpha \succeq 0$
$>$	<i>greater than</i>

<i>positive</i>	for $\alpha \in \mathbb{R}^n$, $\alpha \succ 0$; <i>id est</i> , no zero entries when with respect to nonnegative orthant, no vector on boundary with respect to pointed closed convex cone \mathcal{K}
$\lfloor \cdot \rfloor$	floor function, $\lfloor x \rfloor$ is greatest integer not exceeding x
$ \cdot $	entrywise absolute value of scalars, vectors, and matrices
\log	natural (or Napierian) logarithm
\det	matrix determinant
$\ x\ $	vector 2-norm or <i>Euclidean norm</i> $\ x\ _2$
$\ x\ _\ell$	$= \sqrt[\ell]{\sum_{j=1}^n x_j ^\ell}$ vector ℓ -norm for $\ell \geq 1$
	$\triangleq \sum_{j=1}^n x_j ^\ell$ vector ℓ -norm for $0 \leq \ell < 1$ (violates §3.1.3 no.3)
$\ x\ _\infty$	$= \max\{ x_j \mid \forall j\}$ <i>infinity-norm</i>
$\ x\ _2^2$	$= x^T x = \langle x, x \rangle$
$\ x\ _1$	$= \mathbf{1}^T x $ 1-norm, <i>dual infinity-norm</i>
$\ x\ _0$	0-norm (§4.5.1), <i>cardinality of vector x</i> , $\text{card } x \equiv \ x\ _0$, $0^0 \triangleq 0$
$\ X\ _2$	$= \sup_{\ a\ =1} \ Xa\ _2 = \sigma_1 = \sqrt{\lambda(X^T X)_1}$ matrix 2-norm (<i>spectral norm</i>), <i>largest singular value</i> , $\ Xa\ _2 \leq \ X\ _2 \ a\ _2$ [139, p.56], $\ \delta(x)\ _2 = \ x\ _\infty$
$\ X\ $	$= \ X\ _F$ <i>Frobenius' matrix norm</i>

Bibliography

- [1] Suliman Al-Homidan and Henry Wolkowicz. Approximate and exact completion problems for Euclidean distance matrices using semidefinite programming, April 2004.
orion.math.uwaterloo.ca/~hwolkowi/henry/reports/edmapr04.ps
- [2] Faiz A. Al-Khayyal and James E. Falk. Jointly constrained biconvex programming. *Mathematics of Operations Research*, 8(2):273–286, May 1983.
<http://www.convexoptimization.com/TTOOLS/Falk.pdf>
- [3] Abdo Y. Alfakih. On the uniqueness of Euclidean distance matrix completions. *Linear Algebra and its Applications*, 370:1–14, 2003.
- [4] Abdo Y. Alfakih. On the uniqueness of Euclidean distance matrix completions: the case of points in general position. *Linear Algebra and its Applications*, 397:265–277, 2005.
- [5] Abdo Y. Alfakih, Amir Khandani, and Henry Wolkowicz. Solving Euclidean distance matrix completion problems via semidefinite programming. *Computational Optimization and Applications*, 12(1):13–30, January 1999.
<http://citeseer.ist.psu.edu/alfakih97solving.html>
- [6] Abdo Y. Alfakih and Henry Wolkowicz. On the embeddability of weighted graphs in Euclidean spaces. Research Report CORR 98-12, Department of Combinatorics and Optimization, University of Waterloo, May 1998.
<http://citeseer.ist.psu.edu/alfakih98embeddability.html>
Erratum: p.394 herein.
- [7] Abdo Y. Alfakih and Henry Wolkowicz. Matrix completion problems. In Henry Wolkowicz, Romesh Saigal, and Lieven Vandenbergh, editors, *Handbook of Semidefinite Programming: Theory, Algorithms, and Applications*, chapter 18. Kluwer, 2000.
- [8] Abdo Y. Alfakih and Henry Wolkowicz. Two theorems on Euclidean distance matrices and Gale transform. *Linear Algebra and its Applications*, 340:149–154, 2002.
- [9] Farid Alizadeh. *Combinatorial Optimization with Interior Point Methods and Semi-Definite Matrices*. PhD thesis, University of Minnesota, Computer Science Department, Minneapolis Minnesota USA, October 1991.

- [10] Farid Alizadeh. Interior point methods in semidefinite programming with applications to combinatorial optimization. *SIAM Journal on Optimization*, 5(1):13–51, February 1995.
- [11] Kurt Anstreicher and Henry Wolkowicz. On Lagrangian relaxation of quadratic matrix constraints. *SIAM Journal on Matrix Analysis and Applications*, 22(1):41–55, 2000.
- [12] Howard Anton. *Elementary Linear Algebra*. Wiley, second edition, 1977.
- [13] James Aspnes, David Goldenberg, and Yang Richard Yang. On the computational complexity of sensor network localization. In *Proceedings of the First International Workshop on Algorithmic Aspects of Wireless Sensor Networks (ALGOSENSORS)*, volume 3121 of *Lecture Notes in Computer Science*, pages 32–44, Turku Finland, July 2004. Springer-Verlag.
cs-www.cs.yale.edu/homes/aspnes/localization-abstract.html
- [14] D. Avis and K. Fukuda. A pivoting algorithm for convex hulls and vertex enumeration of arrangements and polyhedra. *Discrete and Computational Geometry*, 8:295–313, 1992.
- [15] Christine Bachoc and Frank Vallentin. New upper bounds for kissing numbers from semidefinite programming. *ArXiv.org*, October 2006.
<http://arxiv.org/abs/math/0608426>
- [16] Mihály Bakonyi and Charles R. Johnson. The Euclidean distance matrix completion problem. *SIAM Journal on Matrix Analysis and Applications*, 16(2):646–654, April 1995.
- [17] Keith Ball. An elementary introduction to modern convex geometry. In Silvio Levy, editor, *Flavors of Geometry*, volume 31, chapter 1, pages 1–58. MSRI Publications, 1997.
www.msri.org/publications/books/Book31/files/ball.pdf
- [18] Richard G. Baraniuk. Compressive sensing [lecture notes]. *IEEE Signal Processing Magazine*, 24(4):118–121, July 2007.
<http://www.convoptimization.com/TOOLS/GeometryCardinality.pdf>
- [19] George Phillip Barker. Theory of cones. *Linear Algebra and its Applications*, 39:263–291, 1981.
- [20] George Phillip Barker and David Carlson. Cones of diagonally dominant matrices. *Pacific Journal of Mathematics*, 57(1):15–32, 1975.
- [21] George Phillip Barker and James Foran. Self-dual cones in Euclidean spaces. *Linear Algebra and its Applications*, 13:147–155, 1976.
- [22] Dror Baron, Michael B. Wakin, Marco F. Duarte, Shriram Sarvotham, and Richard G. Baraniuk. Distributed compressed sensing. Technical Report ECE-0612, Rice University, Electrical and Computer Engineering Department, December 2006.
<http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.85.4789>

- [23] Alexander I. Barvinok. Problems of distance geometry and convex properties of quadratic maps. *Discrete & Computational Geometry*, 13(2):189–202, 1995.
- [24] Alexander I. Barvinok. A remark on the rank of positive semidefinite matrices subject to affine constraints. *Discrete & Computational Geometry*, 25(1):23–31, 2001.
<http://citeseer.ist.psu.edu/304448.html>
- [25] Alexander I. Barvinok. *A Course in Convexity*. American Mathematical Society, 2002.
- [26] Alexander I. Barvinok. Approximating orthogonal matrices by permutation matrices. *Pure and Applied Mathematics Quarterly*, 2:943–961, 2006.
- [27] Heinz H. Bauschke and Jonathan M. Borwein. On projection algorithms for solving convex feasibility problems. *SIAM Review*, 38(3):367–426, September 1996.
- [28] Steven R. Bell. *The Cauchy Transform, Potential Theory, and Conformal Mapping*. CRC Press, 1992.
- [29] Jean Bellissard and Bruno Iochum. Homogeneous and facially homogeneous self-dual cones. *Linear Algebra and its Applications*, 19:1–16, 1978.
- [30] Adi Ben-Israel. Linear equations and inequalities on finite dimensional, real or complex, vector spaces: A unified theory. *Journal of Mathematical Analysis and Applications*, 27:367–389, 1969.
- [31] Aharon Ben-Tal and Arkadi Nemirovski. *Lectures on Modern Convex Optimization: Analysis, Algorithms, and Engineering Applications*. SIAM, 2001.
- [32] John J. Benedetto and Paulo J.S.G. Ferreira editors. *Modern Sampling Theory: Mathematics and Applications*. Birkhäuser, 2001.
- [33] Christian R. Berger, Javier Areta, Krishna Pattipati, and Peter Willett. Compressed sensing – A look beyond linear programming. In *Proceedings of the IEEE International Conference on Acoustics, Speech, and Signal Processing*, pages 3857–3860, 2008.
- [34] Radu Berinde, Anna C. Gilbert, Piotr Indyk, Howard J. Karloff, and Martin J. Strauss. Combining geometry and combinatorics: A unified approach to sparse signal recovery, April 2008.
<http://arxiv.org/abs/0804.4666>
- [35] Abraham Berman. *Cones, Matrices, and Mathematical Programming*, volume 79 of *Lecture Notes in Economics and Mathematical Systems*. Springer-Verlag, 1973.
- [36] Dimitri P. Bertsekas. *Nonlinear Programming*. Athena Scientific, second edition, 1999.
- [37] Dimitri P. Bertsekas, Angelia Nedić, and Asuman E. Ozdaglar. *Convex Analysis and Optimization*. Athena Scientific, 2003.
- [38] Rajendra Bhatia. *Matrix Analysis*. Springer-Verlag, 1997.

- [39] Pratik Biswas, Tzu-Chen Liang, Kim-Chuan Toh, Yinyu Ye, and Ta-Chung Wang. Semidefinite programming approaches for sensor network localization with noisy distance measurements. *IEEE Transactions on Automation Science and Engineering*, 3(4):360–371, October 2006.
- [40] Pratik Biswas, Tzu-Chen Liang, Ta-Chung Wang, and Yinyu Ye. Semidefinite programming based algorithms for sensor network localization. *ACM Transactions on Sensor Networks*, 2(2):188–220, May 2006.
http://www.stanford.edu/~yyye/combined_rev3.pdf
- [41] Pratik Biswas, Kim-Chuan Toh, and Yinyu Ye. A distributed SDP approach for large-scale noisy anchor-free graph realization with applications to molecular conformation. *SIAM Journal on Scientific Computing*, 30(3):1251–1277, March 2008.
<http://www.stanford.edu/~yyye/SISC-molecule-revised-2.pdf>
- [42] Pratik Biswas and Yinyu Ye. Semidefinite programming for ad hoc wireless sensor network localization. In *Proceedings of the Third International Symposium on Information Processing in Sensor Networks (IPSN)*, pages 46–54. IEEE, April 2004.
<http://www.stanford.edu/~yyye/adhocn2.pdf>
- [43] Leonard M. Blumenthal. On the four-point property. *Bulletin of the American Mathematical Society*, 39:423–426, 1933.
- [44] Leonard M. Blumenthal. *Theory and Applications of Distance Geometry*. Oxford University Press, 1953.
- [45] A. W. Bojanczyk and A. Lutoborski. The Procrustes problem for orthogonal Stiefel matrices. *SIAM Journal on Scientific Computing*, 21(4):1291–1304, February 1998. Date is electronic publication:
<http://citeseer.ist.psu.edu/500185.html>
- [46] Ingwer Borg and Patrick Groenen. *Modern Multidimensional Scaling*. Springer-Verlag, 1997.
- [47] Jonathan M. Borwein and Heinz Bauschke. Projection algorithms and monotone operators, 1998.
<http://oldweb.cecm.sfu.ca/personal/jborwein/projections4.pdf>
- [48] Jonathan M. Borwein and Adrian S. Lewis. *Convex Analysis and Nonlinear Optimization: Theory and Examples*. Springer-Verlag, 2000.
<http://www.convexoptimization.com/TOOLS/Borwein.pdf>
- [49] Richard Bouldin. The pseudo-inverse of a product. *SIAM Journal on Applied Mathematics*, 24(4):489–495, June 1973.
<http://www.convexoptimization.com/TOOLS/Pseudoinverse.pdf>
- [50] Stephen Boyd and Jon Dattorro. Alternating projections, 2003.
http://www.stanford.edu/class/ee392o/alt_proj.pdf
- [51] Stephen Boyd, Laurent El Ghaoui, Eric Feron, and Venkataramanan Balakrishnan. *Linear Matrix Inequalities in System and Control Theory*. SIAM, 1994.

- [52] Stephen Boyd, Seung-Jean Kim, Lieven Vandenbergh, and Arash Hassibi. A tutorial on geometric programming. *Optimization and Engineering*, 8(1):1389–4420, March 2007.
http://www.stanford.edu/~boyd/papers/gp_tutorial.html
- [53] Stephen Boyd and Lieven Vandenbergh. *Convex Optimization*. Cambridge University Press, 2004.
<http://www.stanford.edu/~boyd/cvxbook>
- [54] James P. Boyle and Richard L. Dykstra. A method for finding projections onto the intersection of convex sets in Hilbert spaces. In R. Dykstra, T. Robertson, and F. T. Wright, editors, *Advances in Order Restricted Statistical Inference*, pages 28–47. Springer-Verlag, 1986.
- [55] Lev M. Brègman. The method of successive projection for finding a common point of convex sets. *Soviet Mathematics*, 162(3):487–490, 1965. AMS translation of Doklady Akademii Nauk SSSR, 6:688-692.
- [56] Lev M. Brègman, Yair Censor, Simeon Reich, and Yael Zepkowitz-Malachi. Finding the projection of a point onto the intersection of convex sets via projections onto halfspaces, 2003.
http://www.optimization-online.org/DB_FILE/2003/06/669.pdf
- [57] Mike Brookes. Matrix reference manual: Matrix calculus, 2002.
<http://www.ee.ic.ac.uk/hp/staff/dmb/matrix/intro.html>
- [58] Richard A. Brualdi. *Combinatorial Matrix Classes*. Cambridge University Press, 2006.
- [59] Emmanuel J. Candès. Compressive sampling. In *Proceedings of the International Congress of Mathematicians, Madrid*, volume III, pages 1433–1452. European Mathematical Society, August 2006.
<http://www.acm.caltech.edu/~emmanuel/papers/CompressiveSampling.pdf>
- [60] Emmanuel J. Candès and Justin K. Romberg. ℓ_1 -MAGIC : Recovery of sparse signals via convex programming, October 2005.
<http://www.acm.caltech.edu/l1magic/downloads/l1magic.pdf>
- [61] Emmanuel J. Candès, Justin K. Romberg, and Terence Tao. Robust uncertainty principles: Exact signal reconstruction from highly incomplete frequency information. *IEEE Transactions on Information Theory*, 52(2):489–509, February 2006.
<http://arxiv.org/abs/math.NA/0409186> (2004 DRAFT)
<http://www.acm.caltech.edu/~emmanuel/papers/ExactRecovery.pdf>
- [62] Emmanuel J. Candès, Michael B. Wakin, and Stephen P. Boyd. Enhancing sparsity by reweighted ℓ_1 minimization. *Journal of Fourier Analysis and Applications*, 14(5-6):877–905, October 2008.
<http://arxiv.org/abs/0711.1612v1>

- [63] Michael Carter, Holly Jin, Michael Saunders, and Yinyu Ye. SPASELOC: An adaptive subproblem algorithm for scalable wireless sensor network localization. *SIAM Journal on Optimization*, 17(4):1102–1128, December 2006.
<http://www.convexoptimization.com/TOOLS/JinSIAM.pdf>
Erratum: p.284 herein.
- [64] Lawrence Cayton and Sanjoy Dasgupta. Robust Euclidean embedding. In *Proceedings of the 23rd International Conference on Machine Learning (ICML)*, Pittsburgh Pennsylvania USA, 2006.
icml2006.org/icml.documents/camera-ready/022_Robust_Euclidean_Emb.pdf
- [65] Yves Chabrillac and Jean-Pierre Crouzeix. Definiteness and semidefiniteness of quadratic forms revisited. *Linear Algebra and its Applications*, 63:283–292, 1984.
- [66] Manmohan K. Chandraker, Sameer Agarwal, Fredrik Kahl, David Nistér, and David J. Kriegman. Autocalibration via rank-constrained estimation of the absolute quadric. In *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition*, June 2007.
<http://vision.ucsd.edu/~manu/cvpr07-quadric.pdf>
- [67] Rick Chartrand. Exact reconstruction of sparse signals via nonconvex minimization. *IEEE Signal Processing Letters*, 14(10):707–710, October 2007.
math.lanl.gov/Research/Publications/Docs/chartrand-2007-exact.pdf
- [68] Chi-Tsong Chen. *Linear System Theory and Design*. Oxford University Press, 1999.
- [69] Ward Cheney and Allen A. Goldstein. Proximity maps for convex sets. *Proceedings of the American Mathematical Society*, 10:448–450, 1959.
- [70] Mung Chiang. *Geometric Programming for Communication Systems*. Now, 2005.
- [71] Stéphane Chrétien. An alternating l1 approach to the compressed sensing problem, 2008.
http://arxiv.org/PS_cache/arxiv/pdf/0809/0809.0660v1.pdf
- [72] Steven Chu. Autobiography from *Les Prix Nobel*, 1997.
nobelprize.org/physics/laureates/1997/chu-autobio.html
- [73] Ruel V. Churchill and James Ward Brown. *Complex Variables and Applications*. McGraw-Hill, fifth edition, 1990.
- [74] Jon F. Claerbout and Francis Muir. Robust modeling of erratic data. *Geophysics*, 38:826–844, 1973.
- [75] John B. Conway. *A Course in Functional Analysis*. Springer-Verlag, second edition, 1990.
- [76] Richard W. Cottle, Jong-Shi Pang, and Richard E. Stone. *The Linear Complementarity Problem*. Academic Press, 1992.
- [77] G. M. Crippen and T. F. Havel. *Distance Geometry and Molecular Conformation*. Wiley, 1988.

- [78] Frank Critchley. *Multidimensional scaling: a critical examination and some new proposals*. PhD thesis, University of Oxford, Nuffield College, 1980.
- [79] Frank Critchley. On certain linear mappings between inner-product and squared-distance matrices. *Linear Algebra and its Applications*, 105:91–107, 1988.
- [80] Ronald E. Crochiere and Lawrence R. Rabiner. *Multirate Digital Signal Processing*. Prentice-Hall, 1983.
- [81] Joachim Dahl, Bernard H. Fleury, and Lieven Vandenberghe. Approximate maximum-likelihood estimation using semidefinite programming. In *Proceedings of the IEEE International Conference on Acoustics, Speech, and Signal Processing*, volume VI, pages 721–724, April 2003.
- [82] George B. Dantzig. *Linear Programming and Extensions*. Princeton University Press, 1998.
- [83] Alexandre d’Aspremont, Laurent El Ghaoui, Michael I. Jordan, and Gert R. G. Lanckriet. A direct formulation for sparse PCA using semidefinite programming. In Lawrence K. Saul, Yair Weiss, and Léon Bottou, editors, *Advances in Neural Information Processing Systems 17*, pages 41–48. MIT Press, Cambridge Massachusetts USA, 2005.
http://books.nips.cc/papers/files/nips17/NIPS2004_0645.pdf
- [84] Jon Dattorro. Constrained least squares fit of a filter bank to an arbitrary magnitude frequency response, 1991.
<http://www.stanford.edu/~dattorro/Hearing.htm>
- [85] Joel Dawson, Stephen Boyd, Mar Hershenson, and Thomas Lee. Optimal allocation of local feedback in multistage amplifiers via geometric programming. *IEEE Transactions on Circuits and Systems I: Fundamental Theory and Applications*, 48(1):1–11, January 2001.
http://www.stanford.edu/~boyd/papers/fdbk_alloc.html
- [86] Etienne de Klerk. *Aspects of Semidefinite Programming: Interior Point Algorithms and Selected Applications*. Kluwer Academic Publishers, 2002.
- [87] Jan de Leeuw. Fitting distances by least squares. UCLA Statistics Series Technical Report No. 130, Interdivisional Program in Statistics, UCLA, Los Angeles California USA, 1993.
<http://citeseer.ist.psu.edu/deleeuw93fitting.html>
- [88] Jan de Leeuw. Multidimensional scaling. In *International Encyclopedia of the Social & Behavioral Sciences*. Elsevier, 2001.
<http://preprints.stat.ucla.edu/274/274.pdf>
- [89] Jan de Leeuw. Unidimensional scaling. In Brian S. Everitt and David C. Howell, editors, *Encyclopedia of Statistics in Behavioral Science*, volume 4, pages 2095–2097. Wiley, 2005.
<http://www.convexoptimization.com/T00LS/deLeeuw2005.pdf>

- [90] Jan de Leeuw and Willem Heiser. Theory of multidimensional scaling. In P. R. Krishnaiah and L. N. Kanal, editors, *Handbook of Statistics*, volume 2, chapter 13, pages 285–316. North-Holland Publishing, Amsterdam, 1982.
- [91] Erik D. Demaine, Francisco Gomez-Martin, Henk Meijer, David Rappaport, Perouz Taslakian, Godfried T. Toussaint, Terry Winograd, and David R. Wood. The distance geometry of music, 2007.
<http://arxiv.org/abs/0705.4085>
- [92] Frank Deutsch. *Best Approximation in Inner Product Spaces*. Springer-Verlag, 2001.
- [93] Frank Deutsch and Hein Hundal. The rate of convergence of Dykstra’s cyclic projections algorithm: The polyhedral case. *Numerical Functional Analysis and Optimization*, 15:537–565, 1994.
- [94] Frank Deutsch and Peter H. Maserick. Applications of the Hahn-Banach theorem in approximation theory. *SIAM Review*, 9(3):516–530, July 1967.
- [95] Frank Deutsch, John H. McCabe, and George M. Phillips. Some algorithms for computing best approximations from convex cones. *SIAM Journal on Numerical Analysis*, 12(3):390–403, June 1975.
- [96] Michel Marie Deza and Monique Laurent. *Geometry of Cuts and Metrics*. Springer-Verlag, 1997.
- [97] Carolyn Pillers Dobler. A matrix approach to finding a set of generators and finding the polar (dual) of a class of polyhedral cones. *SIAM Journal on Matrix Analysis and Applications*, 15(3):796–803, July 1994. Erratum: p.195 herein.
- [98] Elizabeth D. Dolan, Robert Fourer, Jorge J. Moré, and Todd S. Munson. Optimization on the NEOS server. *SIAM News*, 35(6):4,8,9, August 2002.
- [99] Bruce Randall Donald. 3-D structure in chemistry and molecular biology, 1998.
<http://www.cs.duke.edu/brd/Teaching/Previous/Bio>
- [100] David L. Donoho. Neighborly polytopes and sparse solution of underdetermined linear equations. Technical Report 2005-04, Stanford University, Department of Statistics, January 2005.
www-stat.stanford.edu/~donoho/Reports/2005/NPaSSULE-01-28-05.pdf
- [101] David L. Donoho. Compressed sensing. *IEEE Transactions on Information Theory*, 52(4):1289–1306, April 2006.
www-stat.stanford.edu/~donoho/Reports/2004/CompressedSensing091604.pdf
- [102] David L. Donoho. High-dimensional centrally symmetric polytopes with neighborliness proportional to dimension. *Discrete & Computational Geometry*, 35(4):617–652, May 2006.
- [103] David L. Donoho, Michael Elad, and Vladimir Temlyakov. Stable recovery of sparse overcomplete representations in the presence of noise. *IEEE Transactions on Information Theory*, 52(1):6–18, January 2006.
www-stat.stanford.edu/~donoho/Reports/2004/StableSparse-Donoho-et-al.pdf

- [104] David L. Donoho and Philip B. Stark. Uncertainty principles and signal recovery. *SIAM Journal on Applied Mathematics*, 49(3):906–931, June 1989.
- [105] David L. Donoho and Jared Tanner. Neighborliness of randomly projected simplices in high dimensions. *Proceedings of the National Academy of Sciences*, 102(27):9452–9457, July 2005.
- [106] David L. Donoho and Jared Tanner. Sparse nonnegative solution of underdetermined linear equations by linear programming. *Proceedings of the National Academy of Sciences*, 102(27):9446–9451, July 2005.
- [107] David L. Donoho and Jared Tanner. Counting faces of randomly projected polytopes when the projection radically lowers dimension. *Journal of the American Mathematical Society*, 22(1):1–53, January 2009.
- [108] Miguel Nuno Ferreira Fialho dos Anjos. *New Convex Relaxations for the Maximum Cut and VLSI Layout Problems*. PhD thesis, University of Waterloo, Ontario Canada, Department of Combinatorics and Optimization, 2001.
cheetah.vlsi.uwaterloo.ca/~anjos/MFAnjosPhDThesis.pdf
- [109] John Duchi, Shai Shalev-Shwartz, Yoram Singer, and Tushar Chandra. Efficient projections onto the ℓ_1 -ball for learning in high dimensions. In *Proceedings of the 25th International Conference on Machine Learning (ICML)*, pages 272–279, Helsinki Finland, July 2008. Association for Computing Machinery (ACM).
<http://icml2008.cs.helsinki.fi/papers/361.pdf>
- [110] Richard L. Dykstra. An algorithm for restricted least squares regression. *Journal of the American Statistical Association*, 78(384):837–842, 1983.
- [111] Peter J. Eccles. *An Introduction to Mathematical Reasoning: numbers, sets and functions*. Cambridge University Press, 1997.
- [112] Carl Eckart and Gale Young. The approximation of one matrix by another of lower rank. *Psychometrika*, 1(3):211–218, September 1936.
www.convexoptimization.com/TOOLS/eckart&young.1936.pdf
- [113] Alan Edelman, Tomás A. Arias, and Steven T. Smith. The geometry of algorithms with orthogonality constraints. *SIAM Journal on Matrix Analysis and Applications*, 20(2):303–353, 1998.
- [114] John J. Edgell. Graphics calculator applications on 4-D constructs, 1996.
<http://archives.math.utk.edu/ICTCM/EP-9/C47/pdf/paper.pdf>
- [115] Ivar Ekeland and Roger Témam. *Convex Analysis and Variational Problems*. SIAM, 1999.
- [116] Julius Farkas. Theorie der einfachen Ungleichungen. *Journal für die reine und angewandte Mathematik*, 124:1–27, 1902.
dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?ht=VIEW&did=D261364
- [117] Maryam Fazel. *Matrix Rank Minimization with Applications*. PhD thesis, Stanford University, Department of Electrical Engineering, March 2002.
<http://www.cds.caltech.edu/~maryam/thesis-final.pdf>

- [118] Maryam Fazel, Haitham Hindi, and Stephen P. Boyd. A rank minimization heuristic with application to minimum order system approximation. In *Proceedings of the American Control Conference*, volume 6, pages 4734–4739. American Automatic Control Council (AACC), June 2001.
<http://www.cds.caltech.edu/~maryam/nucnorm.html>
- [119] Maryam Fazel, Haitham Hindi, and Stephen P. Boyd. Log-det heuristic for matrix rank minimization with applications to Hankel and Euclidean distance matrices. In *Proceedings of the American Control Conference*. American Automatic Control Council (AACC), June 2003.
http://www.cds.caltech.edu/~maryam/acc03_final.pdf
- [120] Maryam Fazel, Haitham Hindi, and Stephen P. Boyd. Rank minimization and applications in system theory. In *Proceedings of the American Control Conference*. American Automatic Control Council (AACC), June 2004.
<http://www.cds.caltech.edu/~maryam/acc04-tutorial.pdf>
- [121] J. T. Feddema, R. H. Byrne, J. J. Harrington, D. M. Kilman, C. L. Lewis, R. D. Robinett, B. P. Van Leeuwen, and J. G. Young. Advanced mobile networking, sensing, and controls. Sandia Report SAND2005-1661, Sandia National Laboratories, Albuquerque New Mexico USA, March 2005.
www.prod.sandia.gov/cgi-bin/techlib/access-control.pl/2005/051661.pdf
- [122] César Fernández, Thomas Szyperski, Thierry Bruyère, Paul Ramage, Egon Mösinger, and Kurt Wüthrich. NMR solution structure of the pathogenesis-related protein P14a. *Journal of Molecular Biology*, 266:576–593, 1997.
- [123] Richard Phillips Feynman, Robert B. Leighton, and Matthew L. Sands. *The Feynman Lectures on Physics: Commemorative Issue*, volume I. Addison-Wesley, 1989.
- [124] P. A. Fillmore and J. P. Williams. Some convexity theorems for matrices. *Glasgow Mathematical Journal*, 12:110–117, 1971.
- [125] Paul Finsler. Über das Vorkommen definiten und semidefiniten Formen in Scharen quadratischer Formen. *Commentarii Mathematici Helvetici*, 9:188–192, 1937.
- [126] Anders Forsgren, Philip E. Gill, and Margaret H. Wright. Interior methods for nonlinear optimization. *SIAM Review*, 44(4):525–597, 2002.
- [127] Shmuel Friedland and Anatoli Torokhti. Generalized rank-constrained matrix approximations. *SIAM Journal on Matrix Analysis and Applications*, 29(2):656–659, March 2007.
<http://www2.math.uic.edu/~friedlan/frtor8.11.06.pdf>
- [128] Norbert Gaffke and Rudolf Mathar. A cyclic projection algorithm via duality. *Metrika*, 36:29–54, 1989.
- [129] Jérôme Galtier. Semi-definite programming as a simple extension to linear programming: convex optimization with queueing, equity and other telecom functionals. In *3ème Rencontres Francophones sur les Aspects Algorithmiques des*

- Télécommunications (AlgoTel)*. INRIA, France, 2001.
www-sop.inria.fr/mascotte/Jerome.Galtier/misc/Galtier01b.pdf
- [130] Laurent El Ghaoui. EE 227A: Convex Optimization and Applications, Lecture 11 – October 3. University of California, Berkeley, Fall 2006. Scribe: Nikhil Shetty.
<http://www.convexoptimization.com/TOOLS/Ghaoui.pdf>
- [131] Laurent El Ghaoui and Silviu-Iulian Niculescu, editors. *Advances in Linear Matrix Inequality Methods in Control*. SIAM, 2000.
- [132] Philip E. Gill, Walter Murray, and Margaret H. Wright. *Numerical Linear Algebra and Optimization*, volume 1. Addison-Wesley, 1991.
- [133] Philip E. Gill, Walter Murray, and Margaret H. Wright. *Practical Optimization*. Academic Press, 1999.
- [134] James Gleik. *Isaac Newton*. Pantheon Books, 2003.
- [135] W. Glunt, Tom L. Hayden, S. Hong, and J. Wells. An alternating projection algorithm for computing the nearest Euclidean distance matrix. *SIAM Journal on Matrix Analysis and Applications*, 11(4):589–600, 1990.
- [136] K. Goebel and W. A. Kirk. *Topics in Metric Fixed Point Theory*. Cambridge University Press, 1990.
- [137] Michel X. Goemans and David P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of the Association for Computing Machinery*, 42(6):1115–1145, November 1995.
<http://www-math.mit.edu/~goemans/maxcut-jacm.pdf>
- [138] D. Goldfarb and K. Scheinberg. Interior point trajectories in semidefinite programming. *SIAM Journal on Optimization*, 8(4):871–886, 1998.
- [139] Gene H. Golub and Charles F. Van Loan. *Matrix Computations*. Johns Hopkins, third edition, 1996.
- [140] P. Gordan. Ueber die auflösung linearer gleichungen mit reellen coefficienten. *Mathematische Annalen*, 6:23–28, 1873.
- [141] John Clifford Gower. Euclidean distance geometry. *The Mathematical Scientist*, 7:1–14, 1982.
<http://www.convexoptimization.com/TOOLS/Gower2.pdf>
- [142] John Clifford Gower. Properties of Euclidean and non-Euclidean distance matrices. *Linear Algebra and its Applications*, 67:81–97, 1985.
<http://www.convexoptimization.com/TOOLS/Gower1.pdf>
- [143] John Clifford Gower and Garnt B. Dijksterhuis. *Procrustes Problems*. Oxford University Press, 2004.
- [144] John Clifford Gower and David J. Hand. *Biplots*. Chapman & Hall, 1996.

- [145] Alexander Graham. *Kronecker Products and Matrix Calculus with Applications*. Ellis Horwood Limited, 1981.
- [146] Michael Grant, Stephen Boyd, and Yinyu Ye. *cvx*: MATLAB software for disciplined convex programming, 2007.
<http://www.stanford.edu/~boyd/cvx>
- [147] Robert M. Gray. Toeplitz and circulant matrices: A review. *Foundations and Trends in Communications and Information Theory*, 2(3):155–239, 2006.
<http://www-ee.stanford.edu/~gray/toeplitz.pdf>
- [148] T. N. E. Greville. Note on the generalized inverse of a matrix product. *SIAM Review*, 8:518–521, 1966.
- [149] Rémi Gribonval and Morten Nielsen. Highly sparse representations from dictionaries are unique and independent of the sparseness measure, October 2003.
<http://www.math.aau.dk/research/reports/R-2003-16.pdf>
- [150] Rémi Gribonval and Morten Nielsen. Sparse representations in unions of bases. *IEEE Transactions on Information Theory*, 49(12):1320–1325, December 2003.
<http://www.math.aau.dk/~mnielsen/papers.htm>
- [151] Karolos M. Grigoriadis and Eric B. Beran. Alternating projection algorithms for linear matrix inequalities problems with rank constraints. In Laurent El Ghaoui and Silviu-Iulian Niculescu, editors, *Advances in Linear Matrix Inequality Methods in Control*, chapter 13, pages 251–267. SIAM, 2000.
- [152] Peter Gritzmann and Victor Klee. On the complexity of some basic problems in computational convexity: II. volume and mixed volumes. Technical Report TR:94-31, DIMACS, Rutgers University, 1994.
<ftp://dimacs.rutgers.edu/pub/dimacs/TechnicalReports/TechReports/1994/94-31.ps>
- [153] Peter Gritzmann and Victor Klee. On the complexity of some basic problems in computational convexity: II. volume and mixed volumes. In T. Bisztriczky, P. McMullen, R. Schneider, and A. Ivić Weiss, editors, *Polytopes: Abstract, Convex and Computational*, pages 373–466. Kluwer Academic Publishers, 1994.
- [154] L. G. Gubin, B. T. Polyak, and E. V. Raik. The method of projections for finding the common point of convex sets. *U.S.S.R. Computational Mathematics and Mathematical Physics*, 7(6):1–24, 1967.
- [155] Osman Güler and Yinyu Ye. Convergence behavior of interior-point algorithms. *Mathematical Programming*, 60(2):215–228, 1993.
- [156] P. R. Halmos. Positive approximants of operators. *Indiana University Mathematics Journal*, 21:951–960, 1972.
- [157] Shih-Ping Han. A successive projection method. *Mathematical Programming*, 40:1–14, 1988.
- [158] Godfrey H. Hardy, John E. Littlewood, and George Pólya. *Inequalities*. Cambridge University Press, second edition, 1952.

- [159] Arash Hassibi and Mar Hershenson. Automated optimal design of switched-capacitor filters. In *Proceedings of the Conference on Design, Automation, and Test in Europe*, page 1111, March 2002.
- [160] Johan Håstad. Some optimal inapproximability results. In *Proceedings of the Twenty-Ninth Annual ACM Symposium on Theory of Computing (STOC)*, pages 1–10, El Paso Texas USA, 1997. Association for Computing Machinery (ACM). <http://citeseer.ist.psu.edu/280448.html>, 1999.
- [161] Johan Håstad. Some optimal inapproximability results. *Journal of the Association for Computing Machinery*, 48(4):798–859, July 2001.
- [162] Timothy F. Havel and Kurt Wüthrich. An evaluation of the combined use of nuclear magnetic resonance and distance geometry for the determination of protein conformations in solution. *Journal of Molecular Biology*, 182:281–294, 1985.
- [163] Tom L. Hayden and Jim Wells. Approximation by matrices positive semidefinite on a subspace. *Linear Algebra and its Applications*, 109:115–130, 1988.
- [164] Tom L. Hayden, Jim Wells, Wei-Min Liu, and Pablo Tarazaga. The cone of distance matrices. *Linear Algebra and its Applications*, 144:153–169, 1991.
- [165] Uwe Helmke and John B. Moore. *Optimization and Dynamical Systems*. Springer-Verlag, 1994.
- [166] Bruce Hendrickson. Conditions for unique graph realizations. *SIAM Journal on Computing*, 21(1):65–84, February 1992.
- [167] T. Herrmann, Peter Güntert, and Kurt Wüthrich. Protein NMR structure determination with automated NOE assignment using the new software CANDID and the torsion angle dynamics algorithm DYANA. *Journal of Molecular Biology*, 319(1):209–227, May 2002.
- [168] Mar Hershenson. Design of pipeline analog-to-digital converters via geometric programming. In *Proceedings of the IEEE/ACM International Conference on Computer Aided Design (ICCAD)*, pages 317–324, November 2002.
- [169] Mar Hershenson. Efficient description of the design space of analog circuits. In *Proceedings of the 40th ACM/IEEE Design Automation Conference*, pages 970–973, June 2003.
- [170] Mar Hershenson, Stephen Boyd, and Thomas Lee. Optimal design of a CMOS OpAmp via geometric programming. *IEEE Transactions on Computer-Aided Design of Integrated Circuits and Systems*, 20(1):1–21, January 2001. <http://www.stanford.edu/~boyd/papers/opamp.html>
- [171] Mar Hershenson, Dave Colleran, Arash Hassibi, and Navraj Nandra. Synthesizable full custom mixed-signal IP. *Electronics Design Automation Consortium (EDA)*, 2002. http://www.eda.org/edps/edp02/PAPERS/edp02-s6_2.pdf

- [172] Mar Hershenson, Sunderarajan S. Mohan, Stephen Boyd, and Thomas Lee. Optimization of inductor circuits via geometric programming. In *Proceedings of the 36th ACM/IEEE Design Automation Conference*, pages 994–998, 1999.
http://www.stanford.edu/~boyd/papers/inductor_opt.html
- [173] Nick Higham. Matrix Procrustes problems, 1995.
<http://www.ma.man.ac.uk/~higham/talks>
Lecture notes.
- [174] Richard D. Hill and Steven R. Waters. On the cone of positive semidefinite matrices. *Linear Algebra and its Applications*, 90:81–88, 1987.
- [175] Jean-Baptiste Hiriart-Urruty. Ensembles de Tchebychev *vs.* ensembles convexes: l'état de la situation vu via l'analyse convexe non lisse. *Annales des Sciences Mathématiques du Québec*, 22(1):47–62, 1998.
- [176] Jean-Baptiste Hiriart-Urruty and Claude Lemaréchal. *Convex Analysis and Minimization Algorithms II: Advanced Theory and Bundle Methods*. Springer-Verlag, second edition, 1996.
- [177] Jean-Baptiste Hiriart-Urruty and Claude Lemaréchal. *Fundamentals of Convex Analysis*. Springer-Verlag, 2001.
- [178] Alan J. Hoffman and Helmut W. Wielandt. The variation of the spectrum of a normal matrix. *Duke Mathematical Journal*, 20:37–40, 1953.
- [179] Alfred Horn. Doubly stochastic matrices and the diagonal of a rotation matrix. *American Journal of Mathematics*, 76(3):620–630, July 1954.
<http://www.convexoptimization.com/TOOLS/AHorn.pdf>
- [180] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 1987.
- [181] Roger A. Horn and Charles R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 1994.
- [182] Alston S. Householder. *The Theory of Matrices in Numerical Analysis*. Dover, 1975.
- [183] Hong-Xuan Huang, Zhi-An Liang, and Panos M. Pardalos. Some properties for the Euclidean distance matrix and positive semidefinite matrix completion problems. *Journal of Global Optimization*, 25(1):3–21, January 2003.
- [184] Robert J. Marks II, editor. *Advanced Topics in Shannon Sampling and Interpolation Theory*. Springer-Verlag, 1993.
- [185] 5W Infographic. Wireless 911. *Technology Review*, 107(5):78–79, June 2004.
<http://www.technologyreview.com>
- [186] Nathan Jacobson. *Lectures in Abstract Algebra, vol. II - Linear Algebra*. Van Nostrand, 1953.

- [187] Viren Jain and Lawrence K. Saul. Exploratory analysis and visualization of speech and music by locally linear embedding. In *Proceedings of the IEEE International Conference on Acoustics, Speech, and Signal Processing*, volume 3, pages 984–987, 2004.
http://www.cs.ucsd.edu/~saul/papers/lle_icassp04.pdf
- [188] Joakim Jaldén. Bi-criterion ℓ_1/ℓ_2 -norm optimization. Master’s thesis, Royal Institute of Technology (KTH), Department of Signals Sensors and Systems, Stockholm Sweden, September 2002.
www.ee.kth.se/php/modules/publications/reports/2002/IR-SB-EX-0221.pdf
- [189] Joakim Jaldén, Cristoff Martin, and Björn Ottersten. Semidefinite programming for detection in linear systems – Optimality conditions and space-time decoding. In *Proceedings of the IEEE International Conference on Acoustics, Speech, and Signal Processing*, volume VI, April 2003.
<http://www.s3.kth.se/signal/reports/03/IR-S3-SB-0309.pdf>
- [190] Florian Jarre. Convex analysis on symmetric matrices. In Henry Wolkowicz, Romesh Saigal, and Lieven Vandenbergh, editors, *Handbook of Semidefinite Programming: Theory, Algorithms, and Applications*, chapter 2. Kluwer, 2000.
- [191] Holly Hui Jin. *Scalable Sensor Localization Algorithms for Wireless Sensor Networks*. PhD thesis, University of Toronto, Graduate Department of Mechanical and Industrial Engineering, 2005.
www.stanford.edu/group/SOL/dissertations/holly-thesis.pdf
- [192] Charles R. Johnson and Pablo Tarazaga. Connections between the real positive semidefinite and distance matrix completion problems. *Linear Algebra and its Applications*, 223/224:375–391, 1995.
- [193] Charles R. Johnson and Pablo Tarazaga. Binary representation of normalized symmetric and correlation matrices. *Linear and Multilinear Algebra*, 52(5):359–366, 2004.
- [194] George B. Thomas, Jr. *Calculus and Analytic Geometry*. Addison-Wesley, fourth edition, 1972.
- [195] Mark Kahrs and Karlheinz Brandenburg, editors. *Applications of Digital Signal Processing to Audio and Acoustics*. Kluwer Academic Publishers, 1998.
- [196] Thomas Kailath. *Linear Systems*. Prentice-Hall, 1980.
- [197] Tosio Kato. *Perturbation Theory for Linear Operators*. Springer-Verlag, 1966.
- [198] Paul J. Kelly and Norman E. Ladd. *Geometry*. Scott, Foresman and Company, 1965.
- [199] Yoonsoo Kim and Mehran Mesbahi. On the rank minimization problem. In *Proceedings of the American Control Conference*, volume 3, pages 2015–2020. American Automatic Control Council (AACC), June 2004.

- [200] Ron Kimmel. *Numerical Geometry of Images: Theory, Algorithms, and Applications*. Springer-Verlag, 2003.
- [201] Erwin Kreyszig. *Introductory Functional Analysis with Applications*. Wiley, 1989.
- [202] Takahito Kuno, Yasutoshi Yajima, and Hiroshi Konno. An outer approximation method for minimizing the product of several convex functions on a convex set. *Journal of Global Optimization*, 3(3):325–335, September 1993.
- [203] Amy N. Langville and Carl D. Meyer. *Google's PageRank and Beyond: The Science of Search Engine Rankings*. Princeton University Press, 2006.
- [204] Jean B. Lasserre. A new Farkas lemma for positive semidefinite matrices. *IEEE Transactions on Automatic Control*, 40(6):1131–1133, June 1995.
- [205] Jean B. Lasserre and Eduardo S. Zeron. A Laplace transform algorithm for the volume of a convex polytope. *Journal of the Association for Computing Machinery*, 48(6):1126–1140, November 2001.
<http://arxiv.org/abs/math/0106168> (title revision).
- [206] Monique Laurent. A connection between positive semidefinite and Euclidean distance matrix completion problems. *Linear Algebra and its Applications*, 273:9–22, 1998.
- [207] Monique Laurent. A tour d'horizon on positive semidefinite and Euclidean distance matrix completion problems. In Panos M. Pardalos and Henry Wolkowicz, editors, *Topics in Semidefinite and Interior-Point Methods*, pages 51–76. American Mathematical Society, 1998.
- [208] Monique Laurent. Matrix completion problems. In Christodoulos A. Floudas and Panos M. Pardalos, editors, *Encyclopedia of Optimization*, volume III (Interior-M), pages 221–229. Kluwer, 2001.
<http://homepages.cwi.nl/~monique/files/opt.ps>
- [209] Monique Laurent and Svatopluk Poljak. On a positive semidefinite relaxation of the cut polytope. *Linear Algebra and its Applications*, 223/224:439–461, 1995.
- [210] Monique Laurent and Svatopluk Poljak. On the facial structure of the set of correlation matrices. *SIAM Journal on Matrix Analysis and Applications*, 17(3):530–547, July 1996.
- [211] Monique Laurent and Franz Rendl. Semidefinite programming and integer programming. *Optimization Online*, 2002.
http://www.optimization-online.org/DB_HTML/2002/12/585.html
- [212] Monique Laurent and Franz Rendl. Semidefinite programming and integer programming. In K. Aardal, George L. Nemhauser, and R. Weismantel, editors, *Discrete Optimization*, volume 12 of *Handbooks in Operations Research and Management Science*, chapter 8, pages 393–514. Elsevier, 2005.
- [213] Charles L. Lawson and Richard J. Hanson. *Solving Least Squares Problems*. SIAM, 1995.

- [214] Jung Rye Lee. The law of cosines in a tetrahedron. *Journal of the Korea Society of Mathematical Education Series B: The Pure and Applied Mathematics*, 4(1):1–6, 1997.
- [215] Vladimir L. Levin. Quasi-convex functions and quasi-monotone operators. *Journal of Convex Analysis*, 2(1/2):167–172, 1995.
- [216] Scott Nathan Levine. *Audio Representations for Data Compression and Compressed Domain Processing*. PhD thesis, Stanford University, Department of Electrical Engineering, 1999.
<http://www-ccrma.stanford.edu/~scott1/thesis/thesis.pdf>
- [217] Adrian S. Lewis. Eigenvalue-constrained faces. *Linear Algebra and its Applications*, 269:159–181, 1998.
- [218] Anhua Lin. *Projection algorithms in nonlinear programming*. PhD thesis, Johns Hopkins University, 2003.
- [219] Miguel Sousa Lobo, Lieven Vandenberghe, Stephen Boyd, and Hervé Lebret. Applications of second-order cone programming. *Linear Algebra and its Applications*, 284:193–228, November 1998. Special Issue on Linear Algebra in Control, Signals and Image Processing.
<http://www.stanford.edu/~boyd/socp.html>
- [220] David G. Luenberger. *Optimization by Vector Space Methods*. Wiley, 1969.
- [221] David G. Luenberger. *Introduction to Dynamic Systems: Theory, Models, & Applications*. Wiley, 1979.
- [222] David G. Luenberger. *Linear and Nonlinear Programming*. Addison-Wesley, second edition, 1989.
- [223] Zhi-Quan Luo, Jos F. Sturm, and Shuzhong Zhang. Superlinear convergence of a symmetric primal-dual path following algorithm for semidefinite programming. *SIAM Journal on Optimization*, 8(1):59–81, 1998.
- [224] Zhi-Quan Luo and Wei Yu. An introduction to convex optimization for communications and signal processing. *IEEE Journal On Selected Areas In Communications*, 24(8):1426–1438, August 2006.
- [225] Morris Marden. *Geometry of Polynomials*. American Mathematical Society, second edition, 1985.
- [226] K. V. Mardia. Some properties of classical multi-dimensional scaling. *Communications in Statistics: Theory and Methods*, A7(13):1233–1241, 1978.
- [227] K. V. Mardia, J. T. Kent, and J. M. Bibby. *Multivariate Analysis*. Academic Press, 1979.
- [228] Jerrold E. Marsden and Michael J. Hoffman. *Elementary Classical Analysis*. Freeman, second edition, 1995.
- [229] Rudolf Mathar. The best Euclidean fit to a given distance matrix in prescribed dimensions. *Linear Algebra and its Applications*, 67:1–6, 1985.

- [230] Rudolf Mathar. *Multidimensionale Skalierung*. B. G. Teubner Stuttgart, 1997.
- [231] Nathan S. Mendelsohn and A. Lloyd Dulmage. The convex hull of sub-permutation matrices. *Proceedings of the American Mathematical Society*, 9(2):253–254, April 1958.
<http://www.convexoptimization.com/TOOLS/permu.pdf>
- [232] Mehran Mesbahi and G. P. Papavassilopoulos. On the rank minimization problem over a positive semi-definite linear matrix inequality. *IEEE Transactions on Automatic Control*, 42(2):239–243, February 1997.
- [233] Mehran Mesbahi and G. P. Papavassilopoulos. Solving a class of rank minimization problems via semi-definite programs, with applications to the fixed order output feedback synthesis. In *Proceedings of the American Control Conference*, volume 1, pages 77–80. American Automatic Control Council (AACC), June 1997.
- [234] Sunderarajan S. Mohan, Mar Hershenson, Stephen Boyd, and Thomas Lee. Simple accurate expressions for planar spiral inductances. *IEEE Journal of Solid-State Circuits*, 34(10):1419–1424, October 1999.
http://www.stanford.edu/~boyd/papers/inductance_expressions.html
- [235] Sunderarajan S. Mohan, Mar Hershenson, Stephen Boyd, and Thomas Lee. Bandwidth extension in CMOS with optimized on-chip inductors. *IEEE Journal of Solid-State Circuits*, 35(3):346–355, March 2000.
http://www.stanford.edu/~boyd/papers/bandwidth_ext.html
- [236] E. H. Moore. On the reciprocal of the general algebraic matrix. *Bulletin of the American Mathematical Society*, 26:394–395, 1920. Abstract.
- [237] B. S. Mordukhovich. Maximum principle in the problem of time optimal response with nonsmooth constraints. *Journal of Applied Mathematics and Mechanics*, 40:960–969, 1976.
- [238] Jean-Jacques Moreau. Décomposition orthogonale d’un espace Hilbertien selon deux cônes mutuellement polaires. *Comptes Rendus de l’Académie des Sciences, Paris*, 255:238–240, 1962.
- [239] T. S. Motzkin and I. J. Schoenberg. The relaxation method for linear inequalities. *Canadian Journal of Mathematics*, 6:393–404, 1954.
- [240] Neil Muller, Lourenço Magaia, and B. M. Herbst. Singular value decomposition, eigenfaces, and 3D reconstructions. *SIAM Review*, 46(3):518–545, September 2004.
- [241] Katta G. Murty and Feng-Tien Yu. *Linear Complementarity, Linear and Nonlinear Programming*. Heldermann Verlag, Internet edition, 1988.
ioe.engin.umich.edu/people/fac/books/murty/linear_complementarity_webbook
- [242] Oleg R. Musin. An extension of Delsarte’s method. The kissing problem in three and four dimensions. In *Proceedings of the 2004 COE Workshop on Sphere Packings*, Kyushu University Japan, 2005.
<http://arxiv.org/abs/math.MG/0512649>

- [243] Stephen G. Nash and Ariela Sofer. *Linear and Nonlinear Programming*. McGraw-Hill, 1996.
- [244] Yurii Nesterov and Arkadii Nemirovskii. *Interior-Point Polynomial Algorithms in Convex Programming*. SIAM, 1994.
- [245] L. Nirenberg. *Functional Analysis*. New York University, New York, 1961. Lectures given in 1960-1961, notes by Lesley Sibner.
- [246] Jorge Nocedal and Stephen J. Wright. *Numerical Optimization*. Springer-Verlag, 1999.
- [247] Royal Swedish Academy of Sciences. Nobel prize in chemistry, 2002.
<http://nobelprize.org/chemistry/laureates/2002/public.html>
- [248] C. S. Ogilvy. *Excursions in Geometry*. Dover, 1990. Citation: *Proceedings of the CUPM Geometry Conference*, Mathematical Association of America, No.16 (1967), p.21.
- [249] Ricardo Oliveira, João Costeira, and João Xavier. Optimal point correspondence through the use of rank constraints. In *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition*, volume 2, pages 1016–1021, June 2005.
- [250] Önder Filiz and Aylin Yener. Rank constrained temporal-spatial filters for CDMA systems with base station antenna arrays. In *Proceedings of the Johns Hopkins University Conference on Information Sciences and Systems*, March 2003.
labs.ee.psu.edu/labs/wcan/Publications/filiz-yener_ciss03.pdf
- [251] Alan V. Oppenheim and Ronald W. Schaffer. *Discrete-Time Signal Processing*. Prentice-Hall, 1989.
- [252] Robert Orsi, Uwe Helmke, and John B. Moore. A Newton-like method for solving rank constrained linear matrix inequalities, April 2006.
users.rsise.anu.edu.au/~robert/publications/OHM06_extended_version.pdf
- [253] Brad Osgood. Notes on the Ahlfors mapping of a multiply connected domain, 2000.
www-ee.stanford.edu/~osgood/Ahlfors-Bergman-Szego.pdf
- [254] M. L. Overton and R. S. Womersley. On the sum of the largest eigenvalues of a symmetric matrix. *SIAM Journal on Matrix Analysis and Applications*, 13:41–45, 1992.
- [255] Pythagoras Papadimitriou. *Parallel Solution of SVD-Related Problems, With Applications*. PhD thesis, Department of Mathematics, University of Manchester, October 1993.
- [256] Panos M. Pardalos and Henry Wolkowicz, editors. *Topics in Semidefinite and Interior-Point Methods*. American Mathematical Society, 1998.

- [257] Gábor Pataki. Cone-LP's and semidefinite programs: Geometry and a simplex-type method. In William H. Cunningham, S. Thomas McCormick, and Maurice Queyranne, editors, *Integer Programming and Combinatorial Optimization, Proceedings of the 5th International IPCO Conference, Vancouver, British Columbia, Canada, June 3-5, 1996*, volume 1084 of *Lecture Notes in Computer Science*, pages 162–174. Springer-Verlag, 1996.
- [258] Gábor Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Mathematics of Operations Research*, 23(2):339–358, 1998.
<http://citeseer.ist.psu.edu/pataki97rank.html>
Erratum: p.605 herein.
- [259] Gábor Pataki. The geometry of semidefinite programming. In Henry Wolkowicz, Romesh Saigal, and Lieven Vandenbergh, editors, *Handbook of Semidefinite Programming: Theory, Algorithms, and Applications*, chapter 3. Kluwer, 2000.
- [260] Teemu Pennanen and Jonathan Eckstein. Generalized Jacobians of vector-valued convex functions. Technical Report RRR 6-97, RUTCOR, Rutgers University, May 1997.
<http://rutcor.rutgers.edu/pub/rrr/reports97/06.ps>
- [261] Roger Penrose. A generalized inverse for matrices. In *Proceedings of the Cambridge Philosophical Society*, volume 51, pages 406–413, 1955.
- [262] Chris Perkins. A convergence analysis of Dykstra's algorithm for polyhedral sets. *SIAM Journal on Numerical Analysis*, 40(2):792–804, 2002.
- [263] Florian Pfender and Günter M. Ziegler. Kissing numbers, sphere packings, and some unexpected proofs. *Notices of the American Mathematical Society*, 51(8):873–883, September 2004.
<http://www.convexoptimization.com/TOOLS/Pfender.pdf>
- [264] Benjamin Recht. *Convex Modeling with Priors*. PhD thesis, Massachusetts Institute of Technology, Media Arts and Sciences Department, 2006.
www.media.mit.edu/physics/publications/theses/06.06.Recht.pdf
- [265] Benjamin Recht, Maryam Fazel, and Pablo A. Parrilo. Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization. *Optimization Online*, June 2007.
<http://arxiv.org/abs/0706.4138>
- [266] Alex Reznik. Problem 3.41, from Sergio Verdú. *Multuser Detection*. Cambridge University Press, 1998.
www.ee.princeton.edu/~verdu/mud/solutions/3/3.41.areznik.pdf, 2001.
- [267] Sara Robinson. Hooked on meshing, researcher creates award-winning triangulation program. *SIAM News*, 36(9), November 2003.
<http://www.siam.org/news/news.php?id=370>
- [268] R. Tyrrell Rockafellar. *Conjugate Duality and Optimization*. SIAM, 1974.

- [269] R. Tyrrell Rockafellar. Lagrange multipliers in optimization. *SIAM-American Mathematical Society Proceedings*, 9:145–168, 1976.
- [270] R. Tyrrell Rockafellar. Lagrange multipliers and optimality. *SIAM Review*, 35(2):183–238, June 1993.
- [271] R. Tyrrell Rockafellar. *Convex Analysis*. Princeton University Press, 1997. First published in 1970.
- [272] C. K. Rushforth. Signal restoration, functional analysis, and Fredholm integral equations of the first kind. In Henry Stark, editor, *Image Recovery: Theory and Application*, chapter 1, pages 1–27. Academic Press, 1987.
- [273] Peter Santos. Application platforms and synthesizable analog IP. In *Proceedings of the Embedded Systems Conference*, San Francisco California USA, 2003.
<http://www.convexoptimization.com/TOOLS/Santos.pdf>
- [274] Shankar Sastry. *Nonlinear Systems: Analysis, Stability, and Control*. Springer-Verlag, 1999.
- [275] Uwe Schäfer. A linear complementarity problem with a P-matrix. *SIAM Review*, 46(2):189–201, June 2004.
- [276] Isaac J. Schoenberg. Remarks to Maurice Fréchet’s article “Sur la définition axiomatique d’une classe d’espace distanciés vectoriellement applicable sur l’espace de Hilbert”. *Annals of Mathematics*, 36(3):724–732, July 1935.
<http://www.convexoptimization.com/TOOLS/Schoenberg2.pdf>
- [277] Isaac J. Schoenberg. Metric spaces and positive definite functions. *Transactions of the American Mathematical Society*, 44:522–536, 1938.
<http://www.convexoptimization.com/TOOLS/Schoenberg3.pdf>
- [278] Peter H. Schönemann. A generalized solution of the orthogonal Procrustes problem. *Psychometrika*, 31(1):1–10, 1966.
- [279] Peter H. Schönemann, Tim Dorcey, and K. Kienapple. Subadditive concatenation in dissimilarity judgements. *Perception and Psychophysics*, 38:1–17, 1985.
- [280] Alexander Schrijver. On the history of combinatorial optimization (till 1960). In K. Aardal, G. L. Nemhauser, and R. Weismantel, editors, *Handbook of Discrete Optimization*, pages 1–68. Elsevier, 2005.
<http://citeseer.ist.psu.edu/466877.html>
- [281] Seymour Sherman. Doubly stochastic matrices and complex vector spaces. *American Journal of Mathematics*, 77(2):245–246, April 1955.
<http://www.convexoptimization.com/TOOLS/Sherman.pdf>
- [282] Joshua A. Singer. *Log-Penalized Linear Regression*. PhD thesis, Stanford University, Department of Electrical Engineering, June 2004.
<http://www.convexoptimization.com/TOOLS/Josh.pdf>

- [283] Anthony Man-Cho So and Yinyu Ye. A semidefinite programming approach to tensegrity theory and realizability of graphs. In *Proceedings of the Seventeenth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 766–775, Miami Florida USA, January 2006. Association for Computing Machinery (ACM). <http://www.stanford.edu/~yyye/d-real-soda2.pdf>
- [284] Anthony Man-Cho So and Yinyu Ye. Theory of semidefinite programming for sensor network localization. *Mathematical Programming: Series A and B*, 109(2):367–384, January 2007. <http://www.stanford.edu/~yyye/local-theory.pdf>
- [285] Steve Spain. Polaris Wireless corp., 2005. <http://www.polariswireless.com>
Personal communication.
- [286] Nathan Srebro, Jason D. M. Rennie, and Tommi S. Jaakkola. Maximum-margin matrix factorization. In Lawrence K. Saul, Yair Weiss, and Léon Bottou, editors, *Advances in Neural Information Processing Systems 17 (NIPS 2004)*, pages 1329–1336. MIT Press, 2005.
- [287] Wolfram Stadler, editor. *Multicriteria Optimization in Engineering and in the Sciences*. Springer-Verlag, 1988.
- [288] Henry Stark, editor. *Image Recovery: Theory and Application*. Academic Press, 1987.
- [289] Henry Stark. Polar, spiral, and generalized sampling and interpolation. In Robert J. Marks II, editor, *Advanced Topics in Shannon Sampling and Interpolation Theory*, chapter 6, pages 185–218. Springer-Verlag, 1993.
- [290] Willi-Hans Steeb. *Matrix Calculus and Kronecker Product with Applications and C++ Programs*. World Scientific Publishing Co., 1997.
- [291] Gilbert W. Stewart and Ji-guang Sun. *Matrix Perturbation Theory*. Academic Press, 1990.
- [292] Josef Stoer. On the characterization of least upper bound norms in matrix space. *Numerische Mathematik*, 6(1):302–314, December 1964. <http://www.convexoptimization.com/TOOLS/Stoer.pdf>
- [293] Josef Stoer and Christoph Witzgall. *Convexity and Optimization in Finite Dimensions I*. Springer-Verlag, 1970.
- [294] Gilbert Strang. *Linear Algebra and its Applications*. Harcourt Brace, third edition, 1988.
- [295] Gilbert Strang. *Calculus*. Wellesley-Cambridge Press, 1992.
- [296] Gilbert Strang. *Introduction to Linear Algebra*. Wellesley-Cambridge Press, second edition, 1998.
- [297] Gilbert Strang. Course 18.06: Linear algebra, 2004. <http://web.mit.edu/18.06/www>

- [298] Stefan Straszewicz. Über exponierte Punkte abgeschlossener Punktfolgen. *Fundamenta Mathematicae*, 24:139–143, 1935.
<http://www.convexoptimization.com/TOOLS/Straszewicz.pdf>
- [299] Jos F. Sturm and Shuzhong Zhang. On cones of nonnegative quadratic functions. *Optimization Online*, April 2001.
www.optimization-online.org/DB_HTML/2001/05/324.html
- [300] George P. H. Styan. A review and some extensions of Takemura’s generalizations of Cochran’s theorem. Technical Report 56, Stanford University, Department of Statistics, September 1982.
- [301] George P. H. Styan and Akimichi Takemura. Rank additivity and matrix polynomials. Technical Report 57, Stanford University, Department of Statistics, September 1982.
- [302] Jun Sun, Stephen Boyd, Lin Xiao, and Persi Diaconis. The fastest mixing Markov process on a graph and a connection to a maximum variance unfolding problem. *SIAM Review*, 48(4):681–699, December 2006.
- [303] Chen Han Sung and Bit-Shun Tam. A study of projectionally exposed cones. *Linear Algebra and its Applications*, 139:225–252, 1990.
- [304] Yoshio Takane. On the relations among four methods of multidimensional scaling. *Behaviormetrika*, 4:29–43, 1977.
<http://takane.brinkster.net/Yoshio/p008.pdf>
- [305] Akimichi Takemura. On generalizations of Cochran’s theorem and projection matrices. Technical Report 44, Stanford University, Department of Statistics, August 1980.
- [306] Dharmpal Takhar, Jason N. Laska, Michael B. Wakin, Marco F. Duarte, Dror Baron, Shriram Sarvotham, Kevin F. Kelly, and Richard G. Baraniuk. A new compressive imaging camera architecture using optical-domain compression. In *Proceedings of the SPIE Conference on Computational Imaging IV*, volume 6065, February 2006.
<http://www.dsp.rice.edu/cs/cscam-SPIEJan06.pdf>
- [307] Peng Hui Tan and Lars K. Rasmussen. The application of semidefinite programming for detection in CDMA. *IEEE Journal on Selected Areas in Communications*, 19(8), August 2001.
- [308] Pablo Tarazaga. Faces of the cone of Euclidean distance matrices: Characterizations, structure and induced geometry. *Linear Algebra and its Applications*, 408:1–13, 2005.
- [309] Warren S. Torgerson. *Theory and Methods of Scaling*. Wiley, 1958.
- [310] Lloyd N. Trefethen and David Bau, III. *Numerical Linear Algebra*. SIAM, 1997.
- [311] Michael W. Trosset. Applications of multidimensional scaling to molecular conformation. *Computing Science and Statistics*, 29:148–152, 1998.

- [312] Michael W. Trosset. Distance matrix completion by numerical optimization. *Computational Optimization and Applications*, 17(1):11–22, October 2000.
- [313] Michael W. Trosset. Extensions of classical multidimensional scaling: Computational theory. www.math.wm.edu/~trosset/r.mds.html, 2001. Revision of technical report entitled “Computing distances between convex sets and subsets of the positive semidefinite matrices” first published in 1997.
- [314] Michael W. Trosset and Rudolf Mathar. On the existence of nonglobal minimizers of the STRESS criterion for metric multidimensional scaling. In *Proceedings of the Statistical Computing Section*, pages 158–162. American Statistical Association, 1997.
- [315] Joshua Trzasko and Armando Manduca. Highly undersampled magnetic resonance image reconstruction via homotopic ℓ_0 -minimization. *IEEE Transactions on Medical Imaging*, 28(1):106–121, January 2009.
www.convexoptimization.com/TOOLS/SparseRecon.pdf
- [316] Joshua Trzasko, Armando Manduca, and Eric Borisch. Sparse MRI reconstruction via multiscale L_0 -continuation. In *Proceedings of the IEEE 14th Workshop on Statistical Signal Processing*, pages 176–180, August 2007.
<http://www.convexoptimization.com/TOOLS/LOMRI.pdf>
- [317] P. P. Vaidyanathan. *Multirate Systems and Filter Banks*. Prentice-Hall, 1993.
- [318] Jan van Tiel. *Convex Analysis, an Introductory Text*. Wiley, 1984.
- [319] Lieven Vandenberghe and Stephen Boyd. Semidefinite programming. *SIAM Review*, 38(1):49–95, March 1996.
- [320] Lieven Vandenberghe and Stephen Boyd. Connections between semi-infinite and semidefinite programming. In R. Reemtsen and J.-J. Rückmann, editors, *Semi-Infinite Programming*, chapter 8, pages 277–294. Kluwer Academic Publishers, 1998.
- [321] Lieven Vandenberghe and Stephen Boyd. Applications of semidefinite programming. *Applied Numerical Mathematics*, 29(3):283–299, March 1999.
- [322] Lieven Vandenberghe, Stephen Boyd, and Shao-Po Wu. Determinant maximization with linear matrix inequality constraints. *SIAM Journal on Matrix Analysis and Applications*, 19(2):499–533, April 1998.
- [323] Robert J. Vanderbei. *Convex optimization: Interior-point methods and applications*, 1999.
<http://orfe.princeton.edu/~rvdb/pdf/talks/pumath/talk.pdf>
- [324] Richard S. Varga. *Geršgorin and His Circles*. Springer-Verlag, 2004.
- [325] Martin Vetterli and Jelena Kovačević. *Wavelets and Subband Coding*. Prentice-Hall, 1995.

- [326] Martin Vetterli, Pina Marziliano, and Thierry Blu. Sampling signals with finite rate of innovation. *IEEE Transactions on Signal Processing*, 50(6):1417–1428, June 2002.
<http://bigwww.epfl.ch/publications/vetterli0201.pdf>
<http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.19.5630>
- [327] È. B. Vinberg. The theory of convex homogeneous cones. *Transactions of the Moscow Mathematical Society*, 12:340–403, 1963. American Mathematical Society and London Mathematical Society joint translation, 1965.
- [328] Marie A. Vitulli. A brief history of linear algebra and matrix theory, 2004.
darkwing.uoregon.edu/~vitulli/441.sp04/LinAlgHistory.html
- [329] John von Neumann. *Functional Operators, Volume II: The Geometry of Orthogonal Spaces*. Princeton University Press, 1950. Reprinted from mimeographed lecture notes first distributed in 1933.
- [330] Michael B. Wakin, Jason N. Laska, Marco F. Duarte, Dror Baron, Shriram Sarvotham, Dharmpal Takhar, Kevin F. Kelly, and Richard G. Baraniuk. An architecture for compressive imaging. In *Proceedings of the IEEE International Conference on Image Processing (ICIP)*, pages 1273–1276, October 2006.
<http://www.dsp.rice.edu/cs/CSCam-ICIP06.pdf>
- [331] Roger Webster. *Convexity*. Oxford University Press, 1994.
- [332] Kilian Q. Weinberger and Lawrence K. Saul. Unsupervised learning of image manifolds by semidefinite programming. In *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition*, volume 2, pages 988–995, 2004.
http://www.cs.ucsd.edu/~saul/papers/sde_cvpr04.pdf
- [333] Eric W. Weisstein. Mathworld – A Wolfram Web Resource, 2007.
<http://mathworld.wolfram.com/search>
- [334] Bernard Widrow and Samuel D. Stearns. *Adaptive Signal Processing*. Prentice-Hall, 1985.
- [335] Norbert Wiener. On factorization of matrices. *Commentarii Mathematici Helvetici*, 29:97–111, 1955.
- [336] Ami Wiesel, Yonina C. Eldar, and Shlomo Shamai (Shitz). Semidefinite relaxation for detection of 16-QAM signaling in MIMO channels. *IEEE Signal Processing Letters*, 12(9):653–656, September 2005.
- [337] Michael P. Williamson, Timothy F. Havel, and Kurt Wüthrich. Solution conformation of proteinase inhibitor IIA from bull seminal plasma by ^1H nuclear magnetic resonance and distance geometry. *Journal of Molecular Biology*, 182:295–315, 1985.
- [338] Willie W. Wong. Cayley-Menger determinant and generalized N-dimensional Pythagorean theorem, November 2003. Application of Linear Algebra: Notes on Talk given to Princeton University Math Club.
<http://www.princeton.edu/~wwong/papers/gp-r.pdf>

- [339] William Wootton, Edwin F. Beckenbach, and Frank J. Fleming. *Modern Analytic Geometry*. Houghton Mifflin, 1975.
- [340] Margaret H. Wright. The interior-point revolution in optimization: History, recent developments, and lasting consequences. *Bulletin of the American Mathematical Society*, 42(1):39–56, January 2005.
- [341] Stephen J. Wright. *Primal-Dual Interior-Point Methods*. SIAM, 1997.
- [342] Shao-Po Wu. *max-det Programming with Applications in Magnitude Filter Design*. A dissertation submitted to the department of Electrical Engineering, Stanford University, December 1997.
- [343] Shao-Po Wu and Stephen Boyd. `sdpsol`: A parser/solver for semidefinite programming and determinant maximization problems with matrix structure, 1995.
http://www.stanford.edu/~boyd/old_software/SDPSOL.html
- [344] Shao-Po Wu and Stephen Boyd. `sdpsol`: A parser/solver for semidefinite programs with matrix structure. In Laurent El Ghaoui and Silviu-Iulian Niculescu, editors, *Advances in Linear Matrix Inequality Methods in Control*, chapter 4, pages 79–91. SIAM, 2000.
<http://www.stanford.edu/~boyd/sdpsol.html>
- [345] Shao-Po Wu, Stephen Boyd, and Lieven Vandenberghe. FIR filter design via spectral factorization and convex optimization. In Biswa Nath Datta, editor, *Applied and Computational Control, Signals, and Circuits*, volume 1, chapter 5, pages 215–246. Birkhäuser, 1998.
<http://www.stanford.edu/~boyd/papers/magdes.html>
- [346] Naoki Yamamoto and Maryam Fazel. Computational approach to quantum encoder design for purity optimization. *Physical Review A*, 76(1), July 2007.
<http://arxiv.org/abs/quant-ph/0606106>
- [347] David D. Yao, Shuzhong Zhang, and Xun Yu Zhou. Stochastic linear-quadratic control via primal-dual semidefinite programming. *SIAM Review*, 46(1):87–111, March 2004. Erratum: p.215 herein.
- [348] Yinyu Ye. Semidefinite programming for Euclidean distance geometric optimization. Lecture notes, 2003.
<http://www.stanford.edu/class/ee392o/EE392o-yinyu-ye.pdf>
- [349] Yinyu Ye. Convergence behavior of central paths for convex homogeneous self-dual cones, 1996.
<http://www.stanford.edu/~yyye/yyye/ye.ps>
- [350] Yinyu Ye. *Interior Point Algorithms: Theory and Analysis*. Wiley, 1997.
- [351] D. C. Youla. Mathematical theory of image restoration by the method of convex projection. In Henry Stark, editor, *Image Recovery: Theory and Application*, chapter 2, pages 29–77. Academic Press, 1987.

- [352] Fuzhen Zhang. *Matrix Theory: Basic Results and Techniques*. Springer-Verlag, 1999.
- [353] Günter M. Ziegler. Kissing numbers: Surprises in dimension four. *Emissary*, pages 4–5, Spring 2004.
<http://www.msri.org/communications/emissary>

Index

- 0-norm, 209, 265, 298, 300, 301, 307, 308, 327, 331, 338, 726
 - Boolean, 266
- 1-norm, 205–211, 240, 311, 420, 726
 - ball, *see* ball
 - Boolean, 266
 - nonnegative, 209, 301, 307, 308
- 2-norm, 46, 204–206, 312, 726
 - ball, *see* ball, Euclidean
 - Boolean, 266
 - matrix, 66, 504, 517, 573, 582, 589, 590, 688, 726
 - inequality, 726
- $Ax = b$, 45, 82, 207, 208, 270, 278, 298, 300, 310, 312, 343, 644, 645
 - Boolean, 265
 - nonnegative, 83, 84, 209, 301, 307, 308
- V , 590
- δ , 541
- ∞ -norm, 205, 206, 211
 - ball, *see* ball
- λ , 542, 717
- π , 210, 299, 511, 515, 536, 717
- ψ , 274, 574
- σ , 542, 717
- $\sqrt{\text{EDM}}$, 455
- \succeq , 725
- k -largest norm, 211
- active, 281
- adaptive, 333, 338, 340
- additivity, 98, 109, 556
- adjacency, 48
- adjoint
 - operator, 46, 152, 157, 488
 - self, 363, 541
- affine, 215
- algebra, 57, 75
- boundary, 90
- combination, 63, 68, 74, 138
- dimension, 21, 36, 56, 397, 399, 400, 409, 489
 - complementary, 489
 - low, 373
 - minimization, 522, 534
 - Précis, 400
 - reduction, 446
 - spectral projection, 535
- hull, *see* hull
- independence, 67, 73, 74, 130, 134
 - preservation, 74
- map, 75
- set, 36, 56, 58
 - parallel, 36
- subset, 58, 68
 - hyperplane intersection, 82
 - nullspace basis span, 82
 - projector, 647, 658
- transformation, 45, 75, 232, 248, 492
 - positive semidefinite cone, 110
- affinity, 36
- algebra
 - affine, 57, 75
 - cone, 152
 - face, 88, 89
 - fundamental theorem, 566
 - linear, 81
 - interior, 38
 - linear, 81, 541
 - projection, 658
- algebraic
 - complement, 90, 144
 - multiplicity, 567

- Alizadeh, 272
- alternation, 438
- alternative, *see* theorem
- ambient, 481, 494
 - vector space, 34
- anchor, 284, 369, 376
- angle, 80
 - acute, 68
 - alternating projection, 698
 - brackets, 46, 47
 - constraint, 362–364, 380
 - dihedral, 27, 80, 381, 443, 597
 - EDM cone, 116
 - halfspace, 69
 - inequality, 354, 441
 - matrix, 385
 - interpoint, 359
 - obtuse, 68
 - positive semidefinite cone, 116
 - right, 118, 147, 167, 168
 - torsion, 27, 380
- antidiagonal, 389
- antisymmetric, 52
 - antihollow, 55, 501
- antisymmetry, 96, 556
- artificial intelligence, 23
- asymmetry, 33, 68
- axioms of the metric, 351
- axis of revolution, 117, 118, 596
- ball
 - 1-norm, 207, 208
 - nonnegative, 309
 - ∞ -norm, 207
 - Euclidean, 37, 60, 207, 208, 365, 647, 687
 - nuclear norm, 61, 63
 - packing, 365
 - spectral norm, 66, 688
- Barvinok, 33, 106, 127, 130, 133, 257, 348
- base, 90
 - station, 377
- basis, 53, 175, 719
 - discrete cosine transform, 303
 - nonorthogonal, 175, 659
 - projection on, 659
- orthogonal
 - projection, 659
- orthonormal, 53, 85, 656
- pursuit, 330
- standard
 - matrix, 54, 56, 712
 - vector, 54, 139, 360, 723
- bees, 27, 365
- bijection, *see* bijective
- bijective, 46, 111, 610
 - Gram form, 391
 - inner-product form, 395
 - invertible, 50, 644
 - isometry, 50, 437
 - linear, 48, 50, 53, 56, 132, 169, 515, 595
- biorthogonal
 - decomposition, 648, 656
 - expansion, 144, 171, 175, 649
 - EDM, 469
 - projection, 660
 - unique, 173, 175, 178, 470, 586, 661
- biorthogonality, 173, 176
 - condition, 171, 568, 586
- bound
 - lower, 504
 - polyhedral, 453
- boundary, 38, 89, 473
 - point, 87
 - classical, 38
 - cone, 99
 - EDM, 451, 466
 - membership, 157, 164, 262
 - PSD, 38, 107, 112, 124–127, 262, 478
 - ray, 99
 - conventional, 89
 - extreme, 104
 - of halfspace, 68
 - of hypercube slice, 340
 - of open set, 39
 - of point, 89
 - of ray, 90
 - relative, 37, 60, 89
- bounded, 60
- bowl, 624, 625
- box, 78, 687

- bridge, 381, 410, 460, 505, 521
- calculus
 - matrix, 613
 - of inequalities, 7
 - of variations, 161
- Candès, 329
- Carathéodory's theorem, 156, 670
- cardinality, 356
 - one, 207, 309
 - convex envelope, *see* convex
 - geometry, 207, 309
 - minimization, 207, 208, 300, 327, 338
 - Boolean, 265
 - nonnegative, 209, 298, 301, 307, 308
 - signed, 310
 - problem, *see* cardinality minimization
 - quasiconcavity, 298
 - regularization, 304, 327
- Cartesian
 - axes, 36, 59, 76, 93, 100, 142, 153
 - cone, 144, 156
 - coordinates, 432
 - product, 44, 151, 258, 699, 719
 - cone, 95
- Cayley-Menger
 - determinant, 443, 445
 - form, 401, 422, 423, 511, 535
- cellular, 378
 - telephone network, 377
- center
 - geometric, 361, 387, 430
 - gravity, 387
 - mass, 387
- central path, 258
- certificate, 165, 200, 262
 - null, 157
- chain rule, 620
 - two argument, 620
- Chu, 7
- clipping, 212, 500, 687, 688, 716
- closed, *see* set
- coefficient
 - binomial, 211
 - projection, 171, 476, 665, 672
- cofactor, 444
- compaction, 286, 295, 362, 456, 525
- comparable, 96, 98, 172
- complement
 - algebraic, 90, 144
 - projection on, 652
 - orthogonal, 52
 - relative, 85, 714
- complementarity, 205, 264
 - linear, 191
 - maximal, 252, 257, 280
 - problem, 191
 - strict, 265
- complementary
 - affine dimension, 489
 - dimension, 81
 - eigenvalues, 652
 - halfspaces, 70
 - inertia, 424, 562
 - slackness, 263
 - subspace, 52
- composition
 - EDM, 418
- compressed sensing, 207, 208, 329, 330, 332, 339, 341
 - nonnegative, 209, 301, 307–309
 - problem, 209, 300, 301, 307, 308
 - Procrustes, 318
- compressive sampling, *see* compressed
- concave, 201, 202
- condition
 - biorthogonality, 171, 568, 586
 - convexity
 - first order, 233, 235, 238, 240
 - second order, 237, 242
 - optimality
 - first order, 161, 188–190
 - semidefinite program, 264
 - unconstrained, 161, 228, 229, 674
- cone, 90, 725
 - algebra, 152
 - blade, 155
 - boundary, *see* boundary
 - circular, 94, 117, 118
 - right, 117
 - convex, 44, 93, 95, 102

- dual, 144, 192, 194, 195, 427
 - algorithm, 192
 - construction, 146, 147
 - examples, 154
 - facet, 171, 196
 - formula, 162, 163, 179, 186, 187
 - halfspace-description, 162
 - in subspace, 176, 181
 - linear inequality, 157
 - Lorentz, 154, 168
 - membership relation, 154, 156
 - pointed, 139, 151
 - properties, 151
 - unique, 144, 170
 - vertex-description, 179, 192
- EDM, 357, 449, 451, 452, 454
 - $\sqrt{\text{EDM}}$, 455
 - boundary, 451, 466
 - circular, 119
 - construction, 467
 - convexity, 455
 - decomposition, 490
 - dual, 481, 482, 492, 496
 - extreme direction, 469
 - face, 468, 470
 - interior, 451, 466
 - one-dimensional, 104
 - positive semidefinite, 470, 478
 - projection, 518, 529, 533
- empty set, 93
- face, 99, 171, 196
 - smallest, 95, 114, 128, 164, 165, 255, 468, 477, 483, 493
- halfline, 95
- halfspace, 148
- ice cream, *see* cone, Lorentz
- interior, *see* interior
- intersection, 95
- invariance, 95, 132
- Lorentz, 94, 118, 137, 154
 - dual, 154, 168
- majorization, 545
- membership, 60, 96, 107, 188, 725
 - discretized, 163
 - relation, 154, 156, 180, 259, 260, 262
- monotone, 184, 185
 - nonnegative, 182, 183
- negative, 172, 661
- nonconvex, 91–94
- normal, 188, 189, 475, 679, 685, 705–708
 - elliptope, 707
 - membership relation, 188
 - orthant, 188
- one-dimensional, 90, 94, 130, 139
 - EDM, 104
 - PSD, 102
- origin, 66, 93, 95, 96, 99, 102, 130
- orthogonal, 153
- pointed, 95, 96, 102, 156, 173, 186
 - closed convex, 96, 132
 - dual, 139, 151
 - polyhedral, 100, 139, 170
- polar, 90, 144, 475, 683, 721
- polyhedral, 66, 130, 136–138
 - dual, 131, 167, 170, 171
 - equilateral, 167
 - facet, 171, 196
 - halfspace-description, 136
 - majorization, 537
 - pointed, 100, 139, 170
 - proper, 131
 - vertex-description, 66, 138, 170
- positive semidefinite, 41, 106, 108, 109, 485
 - boundary, 107, 112, 124, 262, 478
 - circular, 117, 212
 - dimension, 114
 - dual, 166
 - extreme direction, 102, 108, 116
 - face, 114, 135, 477, 483, 493
 - hyperplane, supporting, 114, 115
 - inscription, 121
 - interior, 107, 260
 - inverse image, 110
 - one-dimensional, 102
 - optimization, 256
 - polyhedral, 122
 - rank, 114, 253
 - visualization, 254

- proper, 98, 192
- quadratic, 94
- ray, 90, 94, 139
- recession, 145
- second order, *see* cone, Lorentz
- self-dual, 157, 166, 167
- shrouded, 167, 496
- simplicial, 67, 141–143, 172
 - dual, 172, 180
- spectral, 422, 424, 511
 - dual, 427
 - orthant, 514
- subspace, 94
- supporting hyperplane, 95, 96, 171
- tangent, 475
- two-dimensional, 130
- unique, 106, 144, 451
- vertex, 96
- congruence, 390, 421
- conic
 - combination, 68, 138
 - constraint, 190, 212
 - coordinates, 144, 175, 178, 196
 - hull, *see* hull
 - independence, 25, 67, 129–134
 - versus* dimension, 130
 - dual cone, 181
 - extreme direction, 132
 - preservation, 132
 - problem, 150, 190, 250
 - section, 119
 - circular cone, 120
- conservation
 - dimension, 400, 575
- constellation, 20
- constraint
 - angle, 362–364, 380
 - binary, 265, 319
 - cardinality, 300, 327, 338
 - nonnegative, 209, 298, 301, 307, 308
 - signed, 310
 - conic, 190, 212
 - equality, 161
 - affine, 250
 - inequality, 280
- matrix
 - Gram, 362, 363
 - orthogonal, *see* Procrustes, 346
 - orthonormal, 314
 - norm, 312, 321, 574
 - spectral, 66
 - polynomial, 212, 319
 - rank, *see* rank
 - sort, 436
 - tractable, 250
- content, 445, 725
- contour plot, 189, 236
- contraction, 66, 339
- convergence, 283, 437, 691
 - geometric, 693
 - local, 283, 300, 311
- convex, 19
 - combination, 34, 68, 138
 - envelope
 - bilinear, 247
 - cardinality, 270, 523
 - rank, 292, 376, 522, 523
 - equivalent, 293
 - form, 7, 249
 - geometry, 33
 - fundamental, 70, 75, 80, 349, 361
 - hull, *see* hull
 - iteration
 - cardinality, 209, 298–301, 310, 311
 - rank, 281–283, 527
 - rank one, 314, 345, 538
 - stall, 302, 313, 320
 - optimization, 7, 204, 250, 369
 - art, 8, 271, 286, 540
 - fundamental, 249
 - polyhedron, *see* polyhedron
 - problem, 150, 161, 190, 201, 206, 217, 224, 252, 259, 516, 677
 - geometry, 19, 25
 - nonlinear, 249
 - statement as solution, 7, 249
 - tractable, 250
 - program, *see* problem
 - quadratic, *see* function
- convexity, 201

- condition
 - first order, 233, 235, 238, 240
 - second order, 237, 242
- invariance, 43
- log, 247
- coordinates
 - conic, 144, 175, 178, 196
- Crippen & Havel, 381
- criterion
 - EDM, 476, 480
 - dual, 488
 - Finsler, 480
 - metric *versus* matrix, 402
 - Schoenberg, 28, 361, 423, 493
- cubix, 614, 617, 722
- curvature, 237, 245
- decomposition, 721
 - biorthogonal, 648, 656
 - dyad, 580
 - eigen, 566
 - orthonormal, 654, 656
 - singular value, 569
 - compact, 569
 - full, 571
 - subcompact, 570
- deconvolution, 330
- Delsarte method, 366
- dense, 88
- derivative, 228
 - directional, 235, 613, 621, 625
 - dimension, 624
 - gradient, 625
 - second, 626
 - gradient, 631
 - table, 634
- description
 - conversion, 141
 - halfspace, 67, 70
 - affine, 82
 - dual cone, 162
 - vertex, 67, 68, 103
 - affine, 82
 - of halfspace, 133
 - polyhedral cone, 66, 138, 170
 - polyhedron, 60
- determinant, 440, 552, 565, 634, 640
 - Cayley-Menger, 443, 445
 - inequality, 557
 - Kronecker, 544
 - product, 554
- diagonal, 541
 - δ , 541
 - binary, 648
 - dominance, 123
 - inequality, 557
 - matrix, 566
 - commutative, 555
 - diagonalization, 569
 - values
 - eigen, 566
 - singular, 569
- diagonalizable, 173, 566
 - simultaneously, 115, 264, 283, 553, 555, 578
- diagonalization, 40, 566
 - diagonal matrix, 569
 - expansion by, 173
 - symmetric, 568
- diamond, 389
- differential
 - Fréchet, 622
- diffusion, 456
- dilation, 434
- dimension, 36
 - affine, *see* affine
 - complementary, 81
 - conservation, 400, 575
 - embedding, 57
 - positive semidefinite cone
 - face, 114
 - Précis, 400
 - rank, 129, 399, 400, 724
- direction, 90, 101, 621
 - matrix, 281, 282, 293, 294, 327, 367, 528
 - vector, 299, 300
- disc, 61
- discretization, 162, 203, 237, 493
- distance
 - Euclidean, 356

- geometry, 20, 377
 - matrix, *see* EDM
 - origin to affine, 72, 207, 208, 647, 663
 - nonnegative, 209, 308
 - property, 351
 - taxicab, 206–208, 240
- doublet, *see* matrix
- dual
 - affine dimension, 489
 - feasible set, 258
 - norm, 154, 600
 - of dual, 151, 268, 325
 - problem, 148, 199, 210, 251, 263–265, 325
 - projection, 678, 680
 - on cone, 681
 - on convex set, 678, 680
 - on EDM cone, 533
 - optimization, 679
 - strong, 150, 263, 610
- duality gap, 149, 150, 260, 269, 325
- dvec, 56
- dyad, *see* matrix
- Dykstra algorithm, 691, 704, 705
- edge, 86
- EDM, 26, 349, 350, 357, 495
 - $\sqrt{\text{EDM}}$, 455
 - closest, 509
 - composition, 417–420, 452, 453, 455
 - construction, 464
 - criterion, 476, 480
 - dual, 488
 - definition, 356, 462
 - Gram form, 359, 473
 - inner-product form, 382
 - interpoint angle, 359
 - relative-angle form, 384
 - dual, 489
 - exponential entry, 417
 - graph, 353, 366, 370, 373, 374, 458
 - membership relation, 493
 - projection, 479
 - range, 465
 - test, numerical, 354, 361
 - unique, 416, 429, 430, 510
- eigen, 566
 - decomposition, 566
 - matrix, 107, 568
 - spectrum, 422, 511
 - ordered, 424
 - unordered, 426
 - value, 401, 551, 600
 - λ , 542, 717
 - coefficients, 667, 670
 - EDM, 421, 434, 463
 - interlaced, 407, 421, 559
 - intertwined, 407
 - inverse, 568
 - Kronecker, 544
 - largest, 409, 601, 603, 604
 - left, 566
 - of sum, 558
 - principal, 327
 - product, 553, 555, 556, 606
 - real, 546, 566, 568
 - Schur, 563
 - smallest, 409, 418, 601
 - sum of, 48, 552, 603–605
 - symmetric, 568
 - transpose, 567
 - unique, 117, 566
 - zero, 575, 583
- vector, 602, 603
 - distinct, 567
 - EDM, 463
 - Kronecker, 544
 - left, 566
 - normal, 568
 - principal, 327, 328
 - real, 566, 569, 576
 - symmetric, 568
 - unique, 567
- elbow, 503, 504
- element, 202
 - minimal, 98
 - minimum, 98
- ellipsoid, 34, 39, 41, 45, 312
- elliptope, 268, 270, 369, 411, 412, 420, 473, 474
- vertex, 269, 412, 413

- embedding, 397
 - dimension, 57
- empty, 37
 - interior, 37
 - set, 36, 37, 89, 599
 - closed, 38
 - cone, 93
 - hull, 58, 61, 66
 - open, 38
- entry, 35, 202, 711, 716–718, 722
 - smallest/largest, 210
- epigraph, 219, 241, 247
 - convex function, 219
 - form, 225
- errata, 195, 215, 284, 394, 605
- Euclidean
 - ball, 37, 60, 207, 208, 365, 647, 687
 - distance, 350, 356
 - geometry, 20, 370, 377, 673
 - matrix, *see* EDM
 - metric, 351, 404
 - fifth property, 351, 354, 355, 439, 441, 447
 - norm, *see* 2-norm
 - projection, 125
 - space, 34, 719
- exclusive
 - mutually, 160
- expansion, 176, 722
 - biorthogonal, 144, 171, 175, 649
 - EDM, 469
 - projection, 660
 - unique, 173, 175, 178, 470, 586, 661
 - implied by diagonalization, 173
 - orthogonal, 54, 144, 174, 655, 657
 - w.r.t orthant, 175
- exponential, 641
- exposed, 85
 - direction, 103
 - extreme, 87, 96, 135
 - face, 86, 135
 - point, 87, 88, 103
 - density, 88
- extreme, 85
 - boundary, 104
- direction, 101, 105, 132
 - distinct, 101
- dual cone, 131, 171, 176, 178, 181, 193–195
 - EDM cone, 104, 469, 487
 - one-dimensional, 102, 104
 - origin, 102
 - PSD cone, 102, 108, 116
 - shared, 195
 - supporting hyperplane, 171
 - unique, 101
- exposed, 87, 96, 135
- point, 65, 85, 87, 88, 257
 - cone, 96, 128
 - Fantope, 282
 - hypercube slice, 300
- ray, 101, 473
- face, 40, 86
 - algebra, 88, 89
 - exposed, *see* exposed
 - isomorphic, 115, 364, 468
 - polyhedron, 135, 136
 - recognition, 23
 - smallest, 89, 164, 165, *see* cone
 - transitivity, 88
- facet, 88, 171, 196
- Fan, 603
- Fantope, 61, 62, 117, 223, 282
 - extreme point, 282
- Farkas' lemma, 154, 159
 - positive definite, 260
 - not, 262
 - positive semidefinite, 259
- fat, 82, 711
 - full-rank, 82
- finitely generated, 136
- floor, 726
- Forsgren & Gill, 249
- Fourier
 - transform, discrete, 50, 333
- Frobenius, *see* norm
- full, 37
 - dimensional, 37, 98
 - rank, 82
- function, *see* operator

- affine, 44, 215, 230
 - supremum, 218
- bilinear, 247
- composition, 232, 248
- concave, 201, 247, 525, 601
- continuity, 202, 239, 247
- convex, 201, 202, 358, 385
 - nonlinear, 202, 217
 - simultaneously, 221–223, 242
 - strictly, 203, 204, 232, 237, 243, 244, 519, 524, 530, 624, 625
- differentiable, 202, 217, 243, 247
- fractional, 222, 233, 238, 242
 - inverted, 212, 563
 - power, 213
 - projector, 222
- invertible, 45, 50
- linear, *see* operator
- matrix, 239
 - inverted, *see* function, fractional
- monotonic, 201, 210, 211, 231, 247
- multidimensional, 201, 227, 242, 722
 - affine, 215
- objective, *see* objective
- presorting, 210, 299, 511, 515, 536, 717
- product, 221, 233, 246, 247
- pseudofractional, 221
- quadratic, 204, 225, 234, 356, 383, 671
 - nonnegative, 562
 - strictly, 204, 237, 243, 530, 624, 625
- quasiconcave, 246, 247, 274
 - cardinality, 298
 - rank, 124
- quasiconvex, 124, 219, 245, 247
 - continuity, 246, 247
- quasilinear, 238, 247, 574
- quotient, *see* function, fractional
- ratio, *see* function, fractional
- smooth, 202
- sorting, 210, 299, 511, 515, 536, 717
- step, 574
 - matrix, 274
- stress, 238, 506, 527
- support, 78, 217
- vector, 202
- fundamental
 - convex
 - geometry, 70, 75, 80, 349, 361
 - optimization, 249
 - metric property, 351, 404
 - semidefiniteness test, 166, 546
 - subspaces, 45, 80, 644, 654
 - theorem algebra, 566
 - linear, 81
- Gâteaux differential, 622
- Gaffke, 482
- generators, 102
 - c.i., 132
 - hull, 138
 - affine, 73, 664
 - conic, 133, 134
 - convex, 60
 - l.i., 133
 - minimal set, 60, 73, 133, 134, 664
- geometric
 - center, 361, 387, 430
 - subspace, 391, 392, 450, 460, 484, 522, 578, 675
 - centering, 387
 - operator, 393
 - Hahn-Banach theorem, 70, 75, 80
 - mean, 641
 - multiplicity, 567
- Geršgorin, 122
- gimbal, 597
- global positioning system, 22, 284
- Gower, 349
- gradient, 161, 189, 217, 227, 613, 632
 - conjugate, 339
 - derivative, 631
 - directional, 625
 - dimension, 227
 - first order, 631
 - image, 336
 - monotonic, 232
 - norm
 - Frobenius, 637
 - product, 617
 - second order, 632, 633
 - sparsity, 331

- table, 634
 - zero, 161, 228, 229, 674
- Gram matrix, 291, 359, 369
 - constraint, 362, 363
- Hadamard
 - product, 47, 518, 542, 543, 620, 714
 - quotient, 634, 714
 - square root, 712
- Hahn-Banach theorem, 70, 75, 80
- halfline, 90
- halfspace, 68, 69, 148
 - boundary, 68
 - description, 67, 70
 - affine, 82
 - polyhedral cone, 136
 - vertex-, 133, 134
 - interior, 90
 - polarity, 68
- Hayden & Wells, 449, 463, 499
- Hessian, 228, 613
- hexagon, 364
- homogeneity, 98, 205, 247, 357
- honeycomb, 27
- Horn & Johnson, 547, 548
- horn, flared, 94
- hull, 56
 - affine, 36, 56–59, 138
 - empty set, 58
 - point, 37, 58
 - positive semidefinite cone, 58
 - rank-1 correlation matrices, 58
 - unique, 57
- conic, 59, 66, 138
 - empty set, 66
- convex, 57, 59, 60, 138, 350
 - dyads, 61
 - empty set, 61
 - extreme directions, 102
 - orthogonal matrices, 65
 - orthonormal matrices, 65
 - outer product, 61
 - permutation matrices, 65
 - positive semidefinite cone, 116
 - projection matrices, 61
 - rank-one matrices, 61, 64
 - rank-one symmetric matrices, 116
 - unique, 60
- hyperboloid, 579
- hyperbox, 78, 687
- hypercube, 78, 207, 687
 - nonnegative, 270
 - slice
 - nonnegative, 300, 340
- hyperdimensional, 618
- hyperdisc, 625
- hyperplane, 68–70, 230
 - hypersphere radius, 71
 - independent, 82
 - movement, 71, 84, 209
 - normal, 70
 - separating, 80, 159
 - strictly, 80
 - supporting, 75, 77, 114, 115, 189, 234–236
 - nontrivially, 75
 - polarity, 75
 - strictly, 78, 95, 96, 234
 - unique, 233, 234
 - vertex-description, 73
- hypersphere, 61, 71, 365, 413
 - circumhypersphere, 401
- hypograph, 220, 247
- idempotent, *see* matrix
- iff, 724
- image, 44
 - inverse, 44, 110, 169
 - cone, 152
 - magnetic resonance, 330
- indefinite, 420
- independence
 - affine, 67, 73, 74, 130, 134
 - preservation, 74
 - conic, 25, 67, 129–134
 - versus* dimension, 130
 - dual cone, 181
 - extreme direction, 132
 - preservation, 132
 - linear, 35, 67, 130, 133, 585
 - preservation, 35
- inequality

- angle, 354, 441
 - matrix, 385
- Bessel, 655
- determinant, 557
- diagonal, 557
- generalized, 25, 98, 144, 154
 - dual, 156, 166
 - partial order, 96
- Hardy-Littlewood-Pólya, 512
- Jensen, 233
- linear, 33, 68, 157
 - matrix, 25, 168, 169, 258, 259, 262
- norm, 267, 726
 - triangle, 205
- rank, 557
- singular value, 66
- spectral, 66, 422
- trace, 557
- triangle, 351, 383, 404, 405, 440
 - norm, 205
 - strict, 406
 - unique, 439
 - variation, 161
- inertia, 421, 562
 - complementary, 424, 562
 - Sylvester's law, 554
- infimum, 248, 599, 653, 674, 724
- inflection, 237
- injective, 49, 50, 98, 198, 391, 718
 - invertible, 50, 643, 644
 - non, 51
 - nullspace, 50
- innovation, 30
 - rate, 329
- interior, 37, 38
 - point, 37
 - method, 252, 369, 491
 - algebra, 38
 - cone
 - EDM, 451, 466
 - membership, 156
 - positive semidefinite, 107, 260
 - empty, 37
 - halfspace, 90
 - of point, 37, 38
 - ray, 90
 - relative, 37, 89
- intersection, 43
 - cone, 95
 - halfspaces, 70
 - hyperplane, 82
 - convex set, 76
 - line with boundary, 39
 - nullspaces, 85
 - planes, 83
 - positive semidefinite cone
 - affine, 127, 262
 - geometric center subspace, 578
 - line, 371
 - subspaces, 85
 - tangential, 41
- invariance, 386
 - cone, 95, 132
 - Gram form, 387
 - inner-product form, 390
 - orthogonal, 50, 437
 - reflection, 388
 - rotation, 388
 - set, 419
 - translation, 386
- inversion
 - conic coordinates, 198
 - Gram form, 393
 - injective, 50
 - matrix, 584, 630
 - nullspace, 45, 50
 - operator, 45, 50
- is, 715
- isedm(), 354, 361
- isometry, 50, 437, 506
- isomorphic, 48, 52, 54, 714
 - face, 115, 364, 468
 - isometrically, 40, 49, 111, 166
 - non, 578
 - range, 49
- isomorphism, 48, 168, 394, 396, 450
 - isometric, 50, 437
 - symmetric hollow subspace, 56
 - symmetric matrix subspace, 53
- iterate, 691, 701, 702, 705

- iteration
 - alternating projection, 531, 691, 697, 705
 - convex, *see* convex
- Jacobian, 228
- K-convexity, 202
- Karhunen–Loève transform, 428
- kissing, 149, 207, 208, 307–309, 365, 647
 - number, 366
- KKT conditions, 509
- Kronecker product, 111, 542–544, 610, 618, 636
 - eigen, 544
 - inverse, 333
 - positive semidefinite, 356, 556
 - projector, 692
 - pseudoinverse, 645
 - trace, 544
 - vector, 620
 - vectorization, 543
- Lagrange multiplier, 161
 - sign, 199
- Lagrangian, 199, 325, 337
- largest entries, 210
- Lasso, 330
- lattice, 287–290, 296, 297
 - regular, 27, 286
- Laurent, 417
- law of cosines, 382
- least
 - 1-norm, 207, 208, 266, 307
 - energy, 362, 524
 - norm, 207, 267, 646
 - squares, 284, 285, 646
- Legendre-Fenchel transform, 522
- Lemaréchal, 19
- line, 230, 242
 - tangential, 42
- linear
 - algebra, 81, 541
 - bijection, *see* bijective
 - complementarity, 191
 - function, *see* operator
 - independence, 35, 67, 130, 133, 585
 - preservation, 35
 - inequality, 33, 68, 157
 - matrix, 25, 168, 169, 258, 259, 262
 - operator, 44, 54, 202, 215, 217, 359, 393, 395, 437, 471, 541, 676
 - injective, 50
 - noninjective, 51
 - projector, 651, 656, 658, 679
 - program, 79, 83, 84, 150, 205, 217, 249, 251, 252, 366
 - dual, 150, 251
 - prototypical, 150, 251
 - regression, 330
 - transformation, *see* transformation
- localization, 22, 371
 - sensor network, 284
 - standardized test, 286
 - unique, 21, 284, 286, 370, 373
 - wireless, 377
- log, 640
 - det, 245, 377, 525, 631, 639
- machine learning, 23
- majorization, 545
 - cone, 545
 - symmetric hollow, 545
- manifold, 23, 24, 65, 66, 114, 456, 457, 504, 538, 595, 606
- map
 - affine, 75
 - invertible, 45, 50
 - isotonic, 434
 - linear
 - of cone, 132
 - postive semidefinite cone
 - Schur-form, 109
 - USA, 26, 431, 432
- Markov process, 462
- mater, 614
- Mathar, 482
- Matlab, 31, 129, 266, 295, 302, 307, 313, 326, 367, 432, 435, 573
 - MRI, 329, 331, 334, 339, 340
- matrix, 614
 - angle

- interpoint, 359
- relative, 384
- auxiliary, 361, 590, 594
 - orthonormal, 593
 - projector, 590
 - Schoenberg, 358, 592
 - table, 594
- binary, *see* matrix, Boolean
- Boolean, 269, 413
- bordered, 407, 421
- calculus, 613
- circulant, 244, 568, 591
 - permutation, 334
 - symmetric, 334
- completion, *see* problem
- correlation, 58, 412, 413
- covariance, 361
- determinant, *see* determinant
- diagonal, *see* diagonal
- direction, 281, 282, 293, 294, 327, 367, 528
- distance
 - Euclidean, *see* EDM
 - taxicab, 240, 420
- doublet, 387, 587, 675
 - nullspace, 587, 588
 - range, 587, 588
- dyad, 580, 582
 - decomposition, 580
 - independence, 584, 585
 - negative, 582
 - nullspace, 582, 583
 - projection on, 666, 668
 - projector, 583, 648, 665, 666
 - range, 582, 583
 - sum, 566, 569, 586
 - symmetric, 116, 584
- EDM, *see* EDM
- elementary, 419, 588, 590
 - nullspace, 588, 589
 - range, 588, 589
- exponential, 244, 641
- fat, 51, 82, 711
- Fourier, 50, 333
- Gale, 373
- geometric centering, 361, 387, 451, 460, 590
- Gram, 291, 359, 369
 - constraint, 362, 363
- Hermitian, 546
- Householder, 589
 - auxiliary, 591
- idempotent, 651
 - nonsymmetric, 647
 - range, 647, 648
 - symmetric, 653, 656
- inverse, 243, 557, 568, 584, 630
 - injective, 644
- Jordan form, 551, 567
- measurement, 500
- normal, 49, 552, 568, 570
- normalization, 270
- nullspace, *see* nullspace
- orthogonal, 315, 316, 388, 569, 595
 - manifold, 595
 - product, 595
 - symmetric, 596
- orthonormal, 50, 61, 65, 282, 314, 388, 571, 593, 603, 645, 655
 - manifold, 66
- partitioned, 561, 562, 564
- permutation, 65, 315, 316, 403, 544, 569, 595
 - circulant, 334
 - extreme point, 65
 - product, 595
 - symmetric, 334, 590, 609
- positive definite
 - inverse, 557
 - positive real numbers, 239
- positive semidefinite, 546, 549, 551
 - difference, 127
 - from extreme directions, 117
 - Kronecker, 356, 556
 - nonsymmetric, 551
 - square root, 557, 569, 712
- product, *see* product
- projection, 111, 591, 654
 - Kronecker, 692
 - nonorthogonal, 647

- orthogonal, 653
- product, 691
- pseudoinverse, 111, 176, 229, 267, 561, 643
 - Kronecker, 645
 - nullspace, 643, 644
 - product, 644
 - range, 643, 644
 - SVD, 573
 - transpose, 173
 - unique, 643
- quotient
 - Hadamard, 634, 714
- range, *see* range
- rank, *see* rank
- reflection, 388, 389, 596, 609
- rotation, 388, 595
 - of range, 596
- rotation of, 598
- Schur-form, 561
- simple, 581
- skinny, 49, 176, 711
- sort index, 433
- square root, 712, *see* matrix, positive semidefinite
- squared, 243
- standard basis, 54, 56, 712
- Stiefel, *see* matrix, orthonormal
- stochastic, 65, 315
- sum
 - eigenvalue, 558
 - rank, 124, 553
- symmetric, 52, 568
 - real numbers, 239
 - subspace of, 52
- symmetrized, 547
- trace, *see* trace
- unitary, 595
 - symmetric, 333, 335
- max cut problem, 322
- maximum variance unfolding, 456, 462
- membership relation, 154, 180, 259
 - boundary, 157, 164, 262
 - discretized, 162, 493
 - EDM, 493
 - in subspace, 180
 - interior, 156, 260
 - normal cone, 188
- metric, 50, 240, 350
 - postulate, 351
 - property, 351, 404
 - fifth, 351, 354, 355, 439
 - space, 351
- minimal
 - element, 98
 - set, 60, 73, 133, 134, 664
- minimax problem, 148
- minimization, 161, 188, 286, 625
 - cardinality, 207, 208, 300, 327, 338
 - Boolean, 265
 - nonnegative, 209, 298, 301, 307, 308
 - signed, 310
 - fractional, inverted, *see* function
 - norm, *see* problem
 - Frobenius, 225, 229, 674
 - on hypercube, 78
 - rank, 281, 314, 327, 343, 345, 527, 535
 - trace, 371, 522, 563
- minimum
 - cardinality, *see* minimization
 - element, 98
 - global
 - unique, 201, 625
 - value
 - unique, 204
- MIT, 341
- molecular conformation, 22, 380
- monotonic, 224, 231, 283, 697, 700
 - Fejér, 697
 - function, 201, 210, 211, 231, 247
 - gradient, 232
- Moore-Penrose conditions, 643
- MRI, 329, 330
- Muller, 573
- multidimensional
 - function, 201, 227, 242
 - affine, 215
 - scaling, 22, 428
 - ordinal, 434
- multilateration, 284, 377

- multipath, 377
- multiplicity, 553, 567
- nearest neighbors, 458
- neighborhood graph, 457, 458
- nesting, 406
- Newton, 365, 613
- nondegeneracy, 351
- nonexpansive, 655, 686
- nonnegative, 402, 725
 - part, 212, 500, 687, 688, 716
- nonnegativity, 205, 351
- nonvertical, 230
- norm, 204–206, 298, 726
 - ∞ -, *see* ∞ -norm
 - k largest, 211
 - ball, *see* ball
 - constraint, 312, 321, 574
 - spectral, 66
 - dual, 154, 600
 - Euclidean, *see* 2-norm
 - Frobenius, 48, 204
 - gradient, 637
 - minimization, 225, 229, 674
 - inequality, 267, 726
 - triangle, 205
 - Ky Fan, 600
 - least, 207, 267, 646
 - nuclear, 600
 - ball, 61, 63
 - one-, *see* 1-norm
 - property, 205
 - regularization, 322, 646
 - spectral, 504, 517, 573, 582, 589, 590, 600, 688, 726
 - ball, 66, 688
 - inequality, 66, 726
 - two-, *see* 2-norm
 - zero, 574
 - zero-, *see* 0-norm
- normal, 70, 189, 678
 - cone, *see* cone
 - equation, 646
 - facet, 171
 - inward, 68, 481
 - outward, 68
 - vector, 678
- Notation, 711
- novelty, 30
- NP-hard, 527
- nuclear magnetic resonance, 380
- nullspace, 81, 591, 676
 - basis, 82
 - doublet, 587, 588
 - dyad, 582, 583
 - elementary matrix, 588, 589
 - form, 80
 - hyperplane, 69
 - injective, 50
 - intersection, 85
 - orthogonal complement, 719
 - product, 554, 582, 586, 654
 - pseudoinverse, 643, 644
- objective, 79
 - convex, 249
 - strictly, 519, 524
 - function, 79, 150, 161, 188, 190
 - linear, 217, 292, 371
 - multidimensional, 97, 98, 204, 298, 337
 - nonlinear, 217
 - polynomial, 319
 - quadratic, 225
 - real, 204
 - value, 263, 277
 - unique, 251
- offset, 386
- on, 718
- one-to-one, 718, *see* injective
- onto, 718, *see* surjective
- open, *see* set
- operator, 722
 - adjoint, 46, 152, 157, 488, 541
 - invertible, 45, 50
 - linear, 44, 54, 202, 215, 217, 359, 393, 395, 437, 471, 541, 676
 - injective, 50
 - noninjective, 51
 - projector, 651, 656, 658, 679
 - nonlinear, 202, 217
 - nullspace, 387, 391, 393, 395, 676

- permutation, 210, 299, 511, 515, 536, 717
- unitary, 50
- optimality
 - condition
 - first order, 161, 188–190
 - semidefinite program, 264
 - unconstrained, 161, 228, 229, 674
 - conic constraint, 190
 - equality constraint, 161
 - global, 249
 - perturbation, 277
- optimization, 19, 249
 - combinatorial, 65, 79, 210, 265, 315, 319, 322, 413, 540
 - convex, 7, 19, 204, 250, 369
 - art, 8, 271, 286, 540
 - fundamental, 249
 - multiobjective, 97, 98, 204, 298, 337
 - tractable, 250
 - vector, 97, 98, 204, 298, 337
- order
 - natural, 46, 541, 722
 - nonincreasing, 513, 569, 570, 717
 - of projection, 500
 - partial, 60, 94, 96–98, 107, 109, 164, 171, 172, 556, 725
 - generalized inequality, 96
 - property, 96, 98
 - total, 96, 97, 723, 724
- origin, 34
 - cone, 66, 93, 95, 96, 99, 102, 130
 - projection of, 207, 208, 647, 663
 - projection on, 683, 688
 - translation to, 398
- Orion nebula, 20
- orthant, 35, 141, 175, 188
 - nonnegative, 157, 487
- orthogonal, 46
 - complement, 52
 - equivalence, 598
 - expansion, 54, 144, 174, 655, 657
 - invariance, 50, 437
- orthonormal, 46
 - decomposition, 654
 - matrix, *see* matrix
- orthonormality condition, 174
- overdetermined, 711
- PageRank, 462
- parallel, 36, 657
- pattern recognition, 22
- Penrose conditions, 643
- pentahedron, 444
- pentatope, 444
- perpendicular, 46, 653, 674
- perturbation, 272, 274
 - optimality, 277
- phantom, 330
- plane, 68
 - segment, 101
- point
 - boundary, 87
 - of hypercube slice, 340
 - exposed, *see* exposed
 - fixed, 283, 320, 339, 676, 694–697
 - inflection, 237
- polychoron, 90, 444
- polyhedron, 40, 57, 79, 135, 410
 - bounded, 60, 65, 209
 - face, 135, 136
 - halfspace-description, 135
 - norm ball, 208
 - stochastic matrices, 65
 - unbounded, 36, 40, 137
 - vertex-description, 60, 138
 - vertices, 65, 137, 208
- polynomial, 562
 - constraint, 212, 319
 - objective, 319
 - quadratic, 562
- polytope, 135
- positive, 726
 - strictly, 403
- precision
 - numerical, 136, 252, 269, 295, 337, 366, 529
- primal
 - feasible set, 258
 - problem, 150, 251
 - via dual, 265

- principal
 - component analysis, 327, 428
 - eigenvalue, 327
 - eigenvector, 327, 328
 - submatrix, 369, 439, 444, 468, 559
 - leading, 404, 406
 - theorem, 559
- principle
 - halfspace-description, 70
 - separating hyperplane, 80
 - supporting hyperplane, 75
- problem
 - 1-norm, 206–210, 266, 311
 - nonnegative, 209, 301, 307, 308
 - Boolean, 65, 79, 265, 319, 413
 - cardinality, 300, 327, 338
 - Boolean, 265
 - nonnegative, 209, 298, 301, 307, 308
 - signed, 310
 - combinatorial, 65, 79, 210, 265, 315, 319, 322, 413, 540
 - completion, 352, 409, 416, 447, 456
 - geometry, 456
 - semidefinite, 702, 703
 - compressed sensing, *see* compressed
 - concave, 201, 250
 - conic, 150, 190, 250
 - convex, 150, 161, 190, 201, 206, 217, 224, 252, 259, 516, 677
 - geometry, 19, 25
 - nonlinear, 249
 - statement as solution, 7, 249
 - tractable, 250
 - dual, *see* dual
 - equivalent, 282
 - feasibility, 139, 164, 165, 191
 - semidefinite, 255, 262, 345
 - fractional, inverted, *see* function
 - max cut, 322
 - minimax, 148
 - nonlinear, 369
 - convex, 249
 - prevalent, 505
 - primal, 150, 251
 - via dual, 265
 - Procrustes, *see* Procrustes
 - proximity, 499, 501, 504, 507, 518, 529
 - EDM, nonconvex, 514
 - in spectral norm, 517
 - rank heuristic, 524, 526
 - SDP, Gram form, 521
 - semidefinite program, 510, 520, 530
 - same, 282
 - stress, 506, 527
 - tractable, 250
 - procedure
 - rank reduction, 273
 - Procrustes
 - combinatorial, 315
 - compressed sensing, 318
 - diagonal, 612
 - linear program, 609
 - maximization, 609
 - orthogonal, 606
 - two sided, 608, 610
 - orthonormal, 314
 - solution
 - unique, 607
 - symmetric, 611
 - translation, 607
 - product, 246, 619
 - Cartesian, 44, 151, 258, 699, 719
 - cone, 95
 - commutative, 244, 264, 553–555, 578, 691
 - determinant, 554
 - direct, 618
 - eigenvalue, 553, 555, 556, 606
 - matrix, 555, 556
 - fractional, *see* function
 - function, *see* function
 - gradient, 617
 - Hadamard, 47, 518, 542, 543, 620, 714
 - inner
 - matrix, 359
 - vector, 46, 247, 252, 329, 383, 666, 716
 - vectorized matrix, 46, 551
 - Kronecker, 111, 542–544, 610, 618, 636
 - eigen, 544

- inverse, 333
- positive semidefinite, 356, 556
- projector, 692
- pseudoinverse, 645
- vector, 620
- vectorization, 543
- nullspace, 554, 582, 586, 654
- orthogonal, 595
- outer, 61
- permutation, 595
- positive definite, 548
 - nonsymmetric, 548
- positive semidefinite, 555, 556, 559
 - nonsymmetric, 555
- projector, 486, 666, 691
- pseudofractional, *see* function
- pseudoinverse, 644
 - Kronecker, 645
- range, 554, 582, 586, 654
- rank, 553, 554
- tensor, 618
- trace, *see* trace
- zero, 577
- program, 79, 249
 - convex, *see* problem
 - geometric, 8
 - integer, 65, 79, 413
 - linear, 79, 83, 84, 150, 205, 217, 249, 251, 252, 366
 - dual, 150, 251
 - prototypical, 150, 251
 - nonlinear
 - convex, 249
 - prototypical, 151
 - quadratic, 252, 437
 - semidefinite, 151, 205, 217, 249, 251, 317
 - dual, 151, 251
 - equality constraint, 109
 - Gram form, 532
 - prototypical, 151, 251
 - Schur-form, 212, 225, 520, 531, 686
- projection, 48, 643
 - algebra, 658
 - alternating, 365, 691, 692
 - convergence, 696, 698, 700, 703
 - distance, 694, 695
 - Dijkstra algorithm, 691, 704, 705
 - feasibility, 694–696
 - iteration, 691, 705
 - on affine/psd cone, 701
 - on EDM cone, 531
 - on halfspaces, 693
 - on orthant/hyperplane, 697, 698
 - optimization, 694, 695, 704
 - over/under, 699
- biorthogonal expansion, 660
- coefficient, 171, 476, 665, 672
- cyclic, 693
- direction, 653, 656, 668
 - nonorthogonal, 650, 651, 665
- dual, *see* dual, projection
- easy, 687
- Euclidean, 125, 501, 504, 508, 510, 518, 529, 678
- matrix, *see* matrix
- minimum-distance, 651, 654, 674, 677
- nonorthogonal, 517, 650, 651, 665
 - eigenvalue coefficients, 667
- oblique, 207, 208, 506, 647, *see* projection, nonorthogonal
- of convex set, 46
- of matrix, 673, 675
- of origin, 207, 208, 647, 663
- of PSD cone, 113
- of range/rowspace, 677
- on 1-norm ball, 689
- on affine, 46, 658, 659, 663
 - hyperplane, 663, 665
 - of origin, 207, 208, 647, 663
 - vertex-description, 664
- on algebraic complement, 652
- on basis
 - nonorthogonal, 659
 - orthogonal, 659
- on box, 687
- on cardinality k , 688
- on cone, 682, 684, 688
 - EDM, 518, 519, 529, 534, 535
 - polyhedral, 689

- second-order (Lorentz), 689
- truncated, 685
- on cone, PSD, 125, 507, 509, 510, 688
 - geometric center subspace, 486
 - rank constrained, 228, 510, 516, 602
- on convex set, 677
 - in affine subset, 689
 - in subspace, 689
- on convex sets, 693
- on dyad, 666, 668
- on ellipsoid, 312
- on Euclidean ball, 687
- on function domain, 220
- on geometric center subspace, 486, 675
- on halfspace, 665
- on hollow symmetric matrices, 508, 687
- on hyperbox, 687
- on hypercube, 687
- on hyperplane, 663, 665, *see* projection,
 - on affine
- on hypersphere, 687
- on intersection, 689
- on matrix
 - elementary, 662
 - rank constrained, 688
 - subspace, 673
 - vectorized, 665, 666, 668
- on origin, 683, 688
- on orthant, 513, 685, 687
 - in subspace, 513
- on row space, 51
- on simplex, 689
- on slab, 665
- on spectral norm ball, 688
- on subspace, 46, 654, 683
- on symmetric matrices, 687
- on vector
 - cardinality k , 688
 - nonorthogonal, 665
 - orthogonal, 668
- one-dimensional, 668
- order of, 500, 502, 689
- orthogonal, 668
 - eigenvalue coefficients, 670
 - semidefiniteness test as, 671
- spectral, 460, 511, 514
 - norm, 517
 - unique, 514
- successive, 693
- two sided, 673, 675, 677
- umbral, 46
- unique, 651, 654, 674, 677, 678
- vectorization, 476, 675
- projector
 - biorthogonal, 657
 - commutative, 691
 - non, 486, 689, 692
 - complement, 653
 - direction, 653, 656, 668
 - nonorthogonal, 650, 651, 665
 - dyad, 583, 648, 665, 666
 - Kronecker, 692
 - linear operator, 51, 651, 656, 658, 679
 - nonorthogonal, 651, 658, 665
 - orthogonal, 654
 - orthonormal, 657
 - product, 486, 666, 691
 - range, 654
- pyramid, 43, 445, 446
- quadrant, 35
- quadrature, 388
- quartix, 616, 722
- quasi-, *see* function
- quotient
 - Hadamard, 634, 714
 - Rayleigh, 672
- range, 59, 81, 674
 - doublet, 587, 588
 - dyad, 582, 583
 - elementary matrix, 588, 589
 - form, 80
 - idempotent, 647, 648
 - orthogonal complement, 719
 - product, 554, 582, 586, 654
 - pseudoinverse, 643, 644
 - rotation, 596
- rank, 554, 570, 571, 724
 - $-\rho$ subset, 112

- constraint, 281, 314, 319, 327, 343, 345, 527, 535
 - projection on matrices, 688
 - projection on PSD cone, 228, 510, 516, 602
- convex envelope, *see* convex
- convex iteration, *see* convex
- diagonalizable, 553
- dimension, 129, 399, 400, 724
 - affine, 401, 527
- full, 82
- heuristic, 524, 526, 527
- inequality, 557
- log det, 525, 526
- lowest, 255
- minimization, 281, 314, 327, 343, 345, 527, 535
- one, 413, 560, 580, 582, 584
 - convex iteration, 314, 345, 538
 - hull, *see* hull, convex
 - modification, 584, 588
 - subset, 228
 - symmetric, 116
- partitioned matrix, 564
- positive semidefinite cone, 253
 - face, 114
- Précis, 400
- product, 553, 554
- projection matrix, 652, 655
- quasiconcavity, 124
- reduction, 253, 272, 273, 278
 - procedure, 273
- regularization, 298, 314, 322, 344, 345, 527, 538
- Schur-form block, 564
- sum, 124
 - positive semidefinite, 124, 553
- trace, 523, 524
 - heuristic, 522
- ray, 90
 - boundary of, 90
 - cone, 90, 94, 139
 - boundary, 99
 - EDM, 104
 - positive semidefinite, 102
 - extreme, 101, 473
 - interior, 90
- Rayleigh
 - quotient, 672
- realizable, 20, 364, 506, 507
- reconstruction, 428
 - isometric, 373, 432
 - isotonic, 432, 434
 - unique, 353, 373, 374, 390, 391
- recursion, 339, 491, 541, 559
- reflection, 388, 389
 - invariant, 388
 - vector, 596
- reflexivity, 96, 556
- regular
 - lattice, 27, 286
 - simplex, 444
 - tetrahedron, 441, 455
- regularization
 - cardinality, 304, 327
 - norm, 322, 646
 - rank, 298, 314, 322, 344, 345, 527, 538
- relative, 37
 - angle, 382, 384
 - inequality, 354, 385, 441
 - complement, 85, 714
- relaxation, 65, 79, 210, 268, 270, 371, 413, 435
- Riemann, 252
- robotics, 23, 28
- rotation, 388, 597
 - invariant, 388
 - of matrix, 598
 - of range, 596
 - vector, 388, 595
- round, 724
- saddle value, 150
- scaling, 428
 - multidimensional, 22, 428
 - ordinal, 434
 - unidimensional, 527
- Schoenberg
 - auxiliary matrix, 358, 592
 - criterion, 28, 361, 423, 493, 673
- Schur

- form, 109, 561
 - anomaly, 226
 - convex set, 110
 - rank of block, 564
 - semidefinite program, 212, 225, 520, 531, 686
 - sparse, 563
 - complement, 292, 371, 561, 562, 564
- second order
 - cone, 94
 - program, 252
- self-distance, 351
- semidefinite
 - domain, 546
 - program, *see* program
 - symmetry *versus*, 546
- sensor, 284
 - network localization, 284, 369, 376
- separation, 80
- sequence, 697
- set
 - Cartesian product, 43
 - closed, 34, 37, 38, 112, 152
 - connected, 34
 - convex, 34
 - invariance, 43
 - projection on, 677
 - difference, 44
 - empty, 36, 37, 89, 599
 - closed, 38
 - cone, 93
 - hull, 58, 61, 66
 - open, 38
 - feasible, 43, 161, 188, 204, 249, 722
 - dual, 258
 - primal, 258
 - intersection, 43
 - invariant, 419
 - level, 189, 215, 219, 227, 228, 236, 247, 250
 - affine, 217
 - minimal, 60, 73, 133, 134, 664
 - nonconvex, 38
 - open, 34, 36, 38, 39
 - projection of, 46
 - Schur-form, 109
 - smooth, 411
 - solution, 204, 722
 - sublevel, 220, 235, 241, 246, 562
 - quasiconvex function, 246
 - sum, 43
 - superlevel, 220, 246
 - union, 43
- shape, 21
- Sherman-Morrison, 584, 630
- shift, 386
- shroud, 389, 390, 474
 - cone, 167, 496
- SIAM, 250
- signal
 - dropout, 302
 - processing
 - digital, 302, 330, 341, 428
- simplex, 141
 - area, 443
 - content, 445
 - nonnegative, 308, 309
 - regular, 444
 - unit, 139, 140
 - volume, 445
- singular value, 570, 600
 - σ , 542, 717
 - decomposition
 - compact, 569
 - full, 571
 - geometrical, 572
 - pseudoinverse, 573
 - subcompact, 570
 - symmetric, 574
 - inequality, 66
 - largest, 688, 726
 - normal matrix, 49, 570
 - problem, 517
 - sum of, 48, 63, 600, 601
 - symmetric matrix, 574
 - triangle inequality, 63
- skinny, 176, 711
- slab, 36, 293, 665
- slack variable, 251
- Slater's condition, 150, 199, 260, 263

- slice, 118, 168, 377, 624, 625
- smallest entries, 210
- solid, 43, 135, 140, 441
- solution
 - analytical, 225, 249, 599
 - feasible, 204, 722
 - numerical, 294
 - optimal, 79, 83, 84, 204, 722
 - problem statement as, 7, 249
 - set, 204, 722
 - trivial, 35
 - unique, 43, 203, 204, 255, 371, 517, 519, 524, 534
 - vertex, 79, 209, 210, 318
- sort
 - function, 210, 299, 511, 515, 536, 717
 - index matrix, 433
 - largest entries, 211
 - monotone nonnegative, 513
 - smallest entries, 210
- span, 59, 719
- sparsity, 207, 208, 265, 298, 300, 310, 327, 329
 - gradient, 331
 - nonnegative, 209, 301, 308, 309
- spectral
 - cone, 422, 424, 511
 - dual, 427
 - orthant, 514
 - inequality, 66, 422
 - norm, 504, 517, 573, 582, 589, 590, 600, 688, 726
 - ball, 66, 688
 - inequality, 66, 726
 - projection, *see* projection
- sphere packing, 365
- steepest descent, 624
- strict
 - complementarity, 265
 - feasibility, 260
 - positivity, 403
 - triangle inequality, 406
- strong
 - dual, 150, 263, 610
 - duality, 264
- subject to, 723
- submatrix
 - principal, 369, 439, 444, 468, 559
 - leading, 404, 406
 - theorem, 559
- subspace, 34
 - complementary, 52
 - fundamental, 45, 644, 654
 - geometric center, 391, 392, 450, 460, 484, 522, 578, 675
 - orthogonal complement, 114, 675
 - hollow, 54, 392, 485
 - intersection, 85
 - hyperplanes, 82
 - nullspace basis
 - span, 82
 - proper, 34
 - representation, 80
 - symmetric hollow, 54
 - tangent, 114
 - translation invariant, 387, 675
- successive
 - approximation, 693
 - projection, 693
- sum, 43
 - eigenvalues of, 558
 - matrix
 - rank, 124, 553
 - of eigenvalues, 48, 552
 - of singular values, 48, 63, 600, 601
 - vector, 43
 - unique, 52, 144, 585
- supremum, 218, 723
- surjective, 50, 391, 393, 395, 512, 718
 - linear, 395, 396, 413, 415
- svec, 53
- Swiss roll, 24
- symmetry, 351
- tangent
 - line, 41, 371
 - subspace, 114
- tangential, 41, 78, 372
- Taylor series, 613, 630
- tetrahedron, 140, 308
 - angle inequality, 441

- regular, 441, 455
- theorem
 - 0 eigenvalues, 575
 - alternating projection
 - distance, 696
 - alternative, 157, 158, 160, 165
 - EDM, 495
 - semidefinite, 261
 - weak, 160
 - Barvinok, 127
 - Bunt-Motzkin, 678
 - Carathéodory, 156, 670
 - cone
 - faces, 99
 - intersection, 95
 - conic coordinates, 196
 - convexity condition, 233
 - decomposition
 - dyad, 580
 - directional derivative, 625
 - discretized membership, 163
 - dual cone intersection, 192
 - duality, weak, 263
 - EDM, 413
 - eigenvalue
 - of difference, 558
 - order, 557
 - sum, 558
 - zero, 575
 - elliptope vertices, 413
 - exposed, 104
 - extreme existence, 86
 - extremes, 102, 103
 - Farkas' lemma, 154, 159
 - not positive definite, 262
 - positive definite, 260
 - positive semidefinite, 259
 - fundamental
 - algebra, 566
 - algebra, linear, 81
 - convex optimization, 249
 - generalized inequality and membership, 154
 - Geršgorin discs, 122
 - gradient monotonicity, 232
 - Hahn-Banach, 70, 75, 80
 - halfspaces, 70
 - Hardy-Littlewood-Pólya, 512
 - hypersphere, 413
 - inequalities, 512
 - intersection, 43
 - inverse image, 44, 47
 - line, 242
 - linearly independent dyads, 585
 - majorization, 545
 - mean value, 630
 - monotone nonnegative sort, 513
 - nonexpansivity, 686
 - pointed cones, 96
 - positive semidefinite, 550
 - cone subsets, 125
 - matrix sum, 124
 - principal submatrix, 559
 - symmetric, 560
 - projection
 - algebraic complement, 684
 - minimum-distance, 678
 - on affine, 659
 - on cone, 682
 - on PSD geometric intersection, 486
 - on subspace, 46
 - via dual cone, 685
 - via normal cone, 679
 - projector
 - rank/trace, 655
 - semidefinite, 560
 - proper-cone boundary, 99
 - Pythagorean, 383, 690
 - range of dyad sum, 586
 - rank
 - affine dimension, 401
 - partitioned matrix, 564
 - Schur-form block, 564
 - trace, 652
 - real eigenvector, 566
 - sparse sampling, 330
 - Tour, 417
- tight, 353, 722
- trace, 541, 542, 600, 601, 637
 - /rank gap, 523

- eigenvalues, 552
- heuristic, 283, 522, 524
- inequality, 556, 557
- Kronecker, 544
- maximization, 460
- minimization, 371, 522, 563
- of product, 47, 553, 556
- product, 544, 556
- projection matrix, 652
- zero, 577
- trajectory, 419, 480
 - sensor, 375
- transformation
 - affine, 45, 75, 232, 248, 492
 - positive semidefinite cone, 110
 - Fourier, discrete, 50, 333
 - invertible, 50
 - linear, 35, 74, 132, 451
 - cone, 95, 132, 152
 - dual, 157
 - injective, 50, 98
 - polyhedron, 136
 - rigid, 21, 373
 - similarity, 670
- transitivity, 88, 96, 556
- translation invariant, 386
 - subspace, 387, 675
- trefoil, 458, 459, 461
- triangle, 352
 - inequality, 351, 383, 404, 405, 440
 - norm, 205
 - strict, 406
 - unique, 439
- triangulation, 21
- trilateration, 21, 43, 370
 - tandem, 376
- unbounded
 - below, 79, 159
- underdetermined, 343, 711
- unfolding, 456, 462
- unfurling, 456
- unimodal, 245
- unique, 8
 - eigenvalues, 566
 - extreme direction, 101
 - localization, 284, 286
 - minimum element, 98
 - solution, 203, 372
- unraveling, 456, 459, 461
- untieing, 459, 461
- value
 - absolute, 205, 238
 - minimum
 - unique, 204
 - objective, 263, 277
 - unique, 251
- variable
 - matrix, 73
 - slack, 251
- variation
 - calculus, 161
 - total, 336
- vec, 47, 541, 543
- vector, 34, 722
 - binary, 58, 270, 319, 411
 - direction, 299, 300
 - dual, 705
 - normal, 678
 - parallel, 657
 - Perron, 421
 - primal, 705
 - product
 - inner, 46, 247, 252, 329, 383, 551, 666, 716
 - Kronecker, 620
 - quadrature, 388
 - reflection, 596
 - rotation, 388, 595
 - space, 34, 719
 - standard basis, 54, 139, 360, 723
- vectorization, 47
 - projection, 476
 - symmetric, 53
 - hollow, 56
- Venn, 137, 502
- vertex, 39, 43, 87, 88, 96
 - description, 67, 68, 103
 - affine, 82
 - halfspace, 134
 - polyhedral cone, 66, 138, 170

- polyhedron, 60
- solution, 79, 209, 210, 318
- Vitulli, 581
- vortex, 467

- Wikimization*, 31, 129, 192, 273, 307, 321,
326, 340, 348, 432, 434, 435, 580
- Wüthrich, 22
- wavelet, 330, 332
- wedge, 100, 154
- wireless location, 22, 284, 377
- Wolkowicz, 609
- womb, 614

- zero, 574
 - definite, 578
 - entry, 574
 - gradient, 161, 228, 229, 674
 - matrix product, 577
 - norm-, 574
 - trace, 577



Form

$$\|D^x - s(D^x) - H\|$$

min

$$\|D^x - H\|_p$$

$$s(D^x) - x^T r_0$$




$$s(A) \quad 1 - s(A)$$

$$\frac{s + s^T + p(c)}{s - s^T + p(c)}$$

$$s^T(D) \neq s^T(D^x)$$

$$s - (D^x + D^x)$$

$$\Delta \neq s^T(D^x) \quad (D^x)$$



Convex Optimization & Euclidean Distance Geometry, Dattorro



Μεβοο Publishing USA